

LONDON  
SCHOOL of  
HYGIENE  
& TROPICAL  
MEDICINE



Hiep, Nguyen Xuan; Ngondi, Jeremiah M; Anh, Vu Tuan; Dat, Tran Minh; An, Tran Van; Dung, Nguyen Chi; Thang, Nguyen Duy; Chu, Brian K; Willis, Rebecca; Bakhtiari, Ana; Pavluck, Alexandre L; Johnson, James; Sidwell, Joshua; Brady, Molly; Henry, Rob; Mosher, Aryc; Porco, Travis C; Lietman, Thomas M; Rotondo, Lisa A; Lewallen, Susan; Courtright, Paul; Solomon, Anthony W; Aboe, Agatha; Adamu, Liknaw; Alemayehu, Wondu; Alemu, Menbere; Alexander, Neal DE; Bakhtiari, Ana; Bero, Berhanu; Brooker, Simon J; Bush, Simon; Chu, Brian K; Courtright, Paul; Dejene, Michael; Emerson, Paul M; Flueckiger, Rebecca M; Foster, Allen; Gadisa, Solomon; Gass, Katherine; Gebre, Teshome; Habtamu, Zelalem; Haddad, Danny; Harvey, Erik; Haslam, Dominic; Kalua, Khumbo; Kello, Amir B; King, Jonathan D; Le Mesurier, Richard; Lewallen, Susan; Lietman, Thomas M; MacArthur, Chad; Macleod, Colin; Mariotti, Silvio P; Massey, Anna; Mathieu, Els; McCullagh, Siobhain; Mekasha, Addis; Millar, Tom; Mpyet, Caleb; Munoz, Beatriz; Ngondi, Jeremiah; Ogden, Stephanie; Pavluck, Alex; Pearce, Joseph; Resnikoff, Serge; Sarah, Virginia; Sarr, Boubacar; Sisay, Alemayehu; Smith, Jennifer L; Solomon, Anthony W; Thomson, Jo; West, Sheila K; Willis, Rebecca (2018) Trachoma in Viet Nam: results of 11 surveillance surveys conducted with the Global Trachoma Mapping Project. *OPHTHALMIC EPIDEMIOLOGY*, 25 (sup1). pp. 93-102. ISSN 0928-6586 DOI: <https://doi.org/10.1080/09286586.2018.1477964>


Downloaded from: <http://researchonline.lshtm.ac.uk/4651139/>

DOI: [10.1080/09286586.2018.1477964](https://doi.org/10.1080/09286586.2018.1477964)

#### Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact [researchonline@lshtm.ac.uk](mailto:researchonline@lshtm.ac.uk).

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>



## Trachoma in Viet Nam: results of 11 surveillance surveys conducted with the Global Trachoma Mapping Project

Nguyen Xuan Hiep, Jeremiah M. Ngondi, Vu Tuan Anh, Tran Minh Dat, Tran Van An, Nguyen Chi Dung, Nguyen Duy Thang, Brian K. Chu, Rebecca Willis, Ana Bakhtiari, Alexandre L. Pavluck, James Johnson, Joshua Sidwell, Molly Brady, Rob Henry, Aryc Mosher, Travis C. Porco, Thomas M. Lietman, Lisa A. Rotondo, Susan Lewallen, Paul Courtright, Anthony W. Solomon & for the Global Trachoma Mapping Project

To cite this article: Nguyen Xuan Hiep, Jeremiah M. Ngondi, Vu Tuan Anh, Tran Minh Dat, Tran Van An, Nguyen Chi Dung, Nguyen Duy Thang, Brian K. Chu, Rebecca Willis, Ana Bakhtiari, Alexandre L. Pavluck, James Johnson, Joshua Sidwell, Molly Brady, Rob Henry, Aryc Mosher, Travis C. Porco, Thomas M. Lietman, Lisa A. Rotondo, Susan Lewallen, Paul Courtright, Anthony W. Solomon & for the Global Trachoma Mapping Project (2018) Trachoma in Viet Nam: results of 11 surveillance surveys conducted with the Global Trachoma Mapping Project, *Ophthalmic Epidemiology*, 25:sup1, 93-102, DOI: [10.1080/09286586.2018.1477964](https://doi.org/10.1080/09286586.2018.1477964)

To link to this article: <https://doi.org/10.1080/09286586.2018.1477964>



© 2018 The Author(s). Published with license by Taylor & Francis



Published online: 31 Dec 2018.



Submit your article to this journal [↗](#)



Article views: 59



View Crossmark data [↗](#)



Citing articles: 3 View citing articles [↗](#)

## Trachoma in Viet Nam: results of 11 surveillance surveys conducted with the Global Trachoma Mapping Project

Nguyen Xuan Hiep<sup>a</sup>, Jeremiah M. Ngondi<sup>b</sup>, Vu Tuan Anh<sup>a</sup>, Tran Minh Dat<sup>a</sup>, Tran Van An<sup>a</sup>, Nguyen Chi Dung<sup>a</sup>, Nguyen Duy Thang<sup>a</sup>, Brian K. Chu<sup>c</sup>, Rebecca Willis<sup>c</sup>, Ana Bakhtiari<sup>c</sup>, Alexandre L. Pavluck<sup>c</sup>, James Johnson<sup>d</sup>, Joshua Sidwell<sup>e</sup>, Molly Brady<sup>e</sup>, Rob Henry<sup>f</sup>, Ayc Moshfer<sup>f</sup>, Travis C. Porco<sup>g,h,i</sup>, Thomas M. Lietman<sup>g,h,i</sup>, Lisa A. Rotondo<sup>e</sup>, Susan Lewallen<sup>j</sup>, Paul Courtright<sup>j</sup>, and Anthony W. Solomon<sup>k,l</sup>, for the Global Trachoma Mapping Project\*

<sup>a</sup>Viet Nam Institute of Ophthalmology, Ministry of Health, Hanoi, Viet Nam; <sup>b</sup>RTI International, Dar es Salaam, Tanzania; <sup>c</sup>Task Force for Global Health, Decatur, GA, USA; <sup>d</sup>FHI 360, Bangkok, Thailand; <sup>e</sup>RTI International, Washington, DC, USA; <sup>f</sup>United States Agency for International Development, Washington, DC, USA; <sup>g</sup>F.I. Proctor Foundation, San Francisco, CA, USA; <sup>h</sup>Department of Ophthalmology, San Francisco, CA, USA; <sup>i</sup>Department of Epidemiology & Biostatistics, University of California, San Francisco, CA, USA; <sup>j</sup>Kilimanjaro Center for Community Ophthalmology, Division of Ophthalmology, University of Cape Town, Cape Town, South Africa; <sup>k</sup>Clinical Research Department, London School of Hygiene & Tropical Medicine, London, UK; <sup>l</sup>London Centre for Neglected Tropical Disease Research, London, UK

### ABSTRACT

**Purpose:** Following interventions against trachoma in Viet Nam, impact surveys conducted in 2003–2011 suggested that trachoma was no longer a public health problem. In 2014, we undertook surveillance surveys to estimate prevalence of trachomatous inflammation—follicular (TF) and trichiasis.

**Methods:** A population-based prevalence survey was undertaken in 11 evaluation units (EUs) encompassing 24 districts, using Global Trachoma Mapping Project methods. A two-stage cluster sampling design was used in each EU, whereby 20 clusters and 60 children per cluster were sampled. Consenting eligible participants (children aged 1–9 years and adults aged  $\geq 50$  years) were examined for trachoma.

**Results:** A total of 9391 households were surveyed, and 20,185 participants (98.8% of those enumerated) were examined for trachoma. EU-level TF prevalence in 1–9-year-olds ranged from 0% to 1.6%. In one cluster (in Hà Giang Province), the percentage of children with TF was 10.3%. The overall pattern of cluster-level percentages of children with TF, however, was consistent with an exponential distribution, which would be consistent with trachoma disappearing. Among people aged  $\geq 50$  years, prevalence of trichiasis by EU ranged from 0% to 0.75%; these estimates are equivalent to 0–0.13% in all ages. The prevalence of trichiasis unknown to the health system among people aged  $\geq 50$  years, by EU, ranged from 0% to 0.17%, which is equivalent to 0–0.03% in all ages.

**Conclusion:** Findings suggest that trachoma is no longer a public health problem in any of the 11 EUs surveyed. However, given the high proportion of children with TF in one cluster in Hà Giang Province, further investigations will be undertaken.

### ARTICLE HISTORY

Received 10 September 2017  
Revised 18 April 2018  
Accepted 11 May 2018

### KEYWORDS

Global Trachoma Mapping Project; SAFE strategy; surveillance survey; Trachoma; trichiasis; Viet Nam

## Introduction

Trachoma is a neglected tropical disease that causes blindness.<sup>1</sup> In 2016, it was estimated that worldwide, 200 million people were at risk, and 3.2 million people needed surgery to avoid trachomatous blindness, in 42 countries.<sup>2</sup> According to criteria set by the World Health Organization (WHO), trachoma is a public health problem where the prevalence of trachomatous inflammation—follicular (TF) is  $\geq 5\%$  in 1–9-year-olds, and/or the prevalence of trachomatous trichiasis unknown to the health system is  $\geq 0.2\%$  in  $\geq 15$ -year-

olds (approximately equivalent to  $\geq 0.1\%$  in the all-ages population).<sup>3</sup> Global political commitment for elimination of trachoma as a public health problem was made manifest by a 1998 World Health Assembly resolution; the target date for achievement of this goal is December 2020.<sup>4,5</sup>

Viet Nam has been fighting trachoma for more than 60 years. In 1950, the Ophthalmologic Institute of Hanoi (which later became the Viet Nam National Institute of Ophthalmology (VNIO)) estimated that more than 70% of the population on

**CONTACT** Jeremiah M. Ngondi  [jngondi@rti.org](mailto:jngondi@rti.org)

\*See Appendix.

Color versions of one or more of the figures in the article can be found online at [www.tandfonline.com/iope](http://www.tandfonline.com/iope).

© 2018 The Author(s). Published with license by Taylor & Francis

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

the outskirts of Hanoi suffered from active trachoma at some time during their lives.<sup>6</sup> In 1951, more than 200,000 people in North Viet Nam were treated for trachoma by mobile teams.<sup>6</sup> Following sustained trachoma control efforts by the government, the national prevalence had declined to 7% in 1995.<sup>7</sup> With the 1996 launch of the WHO Alliance for the Global Elimination of Trachoma by 2020 (GET2020), the government of Viet Nam and its partners started implementing the “SAFE strategy” against trachoma, i.e. surgery for trichiasis, antibiotics to clear infection with the causative organism (*Chlamydia trachomatis*), and facial cleanliness and environmental improvement to reduce transmission.<sup>8</sup> In 2000–2001, 67 district-based surveys were undertaken in 27 suspected-endemic provinces to identify local needs for SAFE. After deployment of SAFE interventions, impact surveys were undertaken at four time points: 2003, 2004, 2005, and 2009–2011. These surveys suggested that trachoma was no longer a public health problem, according to WHO criteria. However, WHO’s standard operating procedures for validation of elimination as a public health problem<sup>9</sup> had not yet been released. In 2014, given the need to complete a second round of surveys to establish that recrudescence of active trachoma had not occurred<sup>10</sup>, we undertook a series of pre-validation trachoma surveillance surveys in Viet Nam, aiming to estimate the prevalence of trachomatous inflammation—follicular (TF) in children aged 1–9 years and the prevalence of

trichiasis unknown to the health system in adults, in previously trachoma-endemic districts.

## Methods

We implemented population-based prevalence surveys (PBPSs) using the systems and methods of the WHO-recommended Global Trachoma Mapping Project (GTMP), including version 2 of the training system,<sup>11,12</sup> with five exceptions: (1) graders were certified to participate in fieldwork if they passed a slide-based test, rather than an inter-grader agreement exercise using live subjects; (2) we did not collect data on access to water and sanitation; (3) examination was limited to 1–9-year-olds and ≥50-year-olds, unlike other GTMP projects where all consenting individuals aged ≥1 year living in sampled households were examined; (4) a fixed number of children (rather than a fixed number of households) was recruited in each sampled cluster; and (5) for individuals diagnosed with trichiasis, we asked questions about previous management of trichiasis through surgery or epilation.

Individual-level demographic and examination data were entered into the GTMP-LINKS application on Android smartphones, transmitted to the Cloud, then processed, and reviewed by VNIO (the agency designated by the Ministry of Health for oversight of ophthalmic issues in Viet Nam), as described elsewhere.<sup>12</sup> Our principal outcome measures were the evaluation unit (EU)-level TF prevalence in 1–9-year-

**Table 1.** Characteristics of survey population by evaluation unit, Global Trachoma Mapping Project, Viet Nam, 2014.

Province	Districts in evaluation unit	Population (2009) <sup>13</sup>	Evaluation Unit	Justification for survey
Vinh Phuc	Vinh Tuong	189,000	1	Both districts had baseline prevalence of TF >10% in 2001 and both had interventions
	Me Linh	192,000	2	
	Yen Lac	145,000		
	Binh Xuyen	108,000		
Ha Tinh	Lap Thach	118,000	3	Both districts >10% in 2001 baseline and both had intervention
	Can Loc	128,000		
Hoa Binh	Nghi Xuan	98,000	4	Both districts >10% in 2001 baseline and both had intervention
	Kim Boi	114,000		
Hung Yen	Yen Thuy	60,000	5	District >10% in 2001 baseline and had intervention
	Van Lam	113,000		
Yen Bai	Van Chan	144,000	6	Both districts >10% in 2001 baseline and both had intervention
	Yen Bai town	91,000		
	Luc Yen	102,000	7	Baseline >10% but no intervention
	Van Yen	27,000		
Vinh Long	Tam Binh	154,000	8	Baseline 9.5% and 10.1% but no intervention
	Vung Liem	159,000		
Hà Giang	Bac Quang	45,000	9	2005 MoH survey of communes between 7.0–17.2%; intervention in 2 districts
	Yen Minh	78,000		
	Bac Me	47,000		
Ninh Thuan	Ninh Hai	89,000	10	Baseline survey 5.3–5.8% in southern Viet Nam (intervention in one)
	Ninh Phuoc	135,000		
Binh Thuan	Ham Tan	69,000	11	Baseline surveys 7.8–7.9%, no intervention
	Tuy Phong	140,000		
	Bac Binh	117,000		

olds, and the EU-level prevalence of trichiasis unknown to the health system in  $\geq 50$ -year-olds.

We surveyed 11 EUs (covering 24 districts) selected in consultation with programme partners. The full list of EUs and the rationale for conducting a survey in each one form Table 1.

### Sample size calculations

We powered each survey to estimate the EU-level TF prevalence in 1–9-year-olds, using an expected prevalence of 10%, and aiming to have 95% confidence of estimating that prevalence with absolute precision of  $\pm 3\%$ . Our design effect, based on previous surveys, was 2.65, which resulted in a sample size estimate of 1019 children.<sup>12</sup> We additionally wished to estimate trichiasis prevalence in adults, but our sample size was calculated only using parameters relating to TF in children.

### Sample selection

#### Selection of clusters

We used two-stage cluster sampling, categorizing villages (*thon*) as our first-stage clusters. A list of *thon* in each EU was obtained from the district authorities. We systematically selected 20 *thon* with probability proportional to population size, using a computer-generated random starting point.

#### Selection of households and participants

In selected *thon*, we used compact segment sampling to choose one segment (*xom*) at random. Within that *xom*, a household was visited if at least one child aged 1–9 years, or at least one adult aged  $\geq 50$  years, lived there (i.e., had slept there for the past month). If individuals in relevant age ranges were away from home but were expected to be available later the same day, an attempt was made to revisit the household. Once 60 children had been enrolled, the cluster was considered complete. If 60 children could not be enrolled in a single *xom*, the next-nearest *xom* was also visited.

### Trachoma grading

Graders participating in the surveys had obtained a kappa for diagnosing TF of  $\geq 0.9$  in an inter-grader agreement test based on 50 conjunctival photographs of children with and without active trachoma.<sup>11</sup> Examination was undertaken according to the criteria set out in the WHO simplified trachoma grading scheme.<sup>13</sup> Graders used binocular loupes with 2.5 $\times$  magnification, and the sun (or if necessary, a torch) for illumination. When trichiasis was diagnosed in

an eye, the subject was asked if health workers had previously recommended surgery or epilation for that eye.

### Data analysis

We conducted analyses in R (R Foundation for Statistical Computing, Vienna, Austria) and Structured Query Language. Cluster (*xom*)-level data on TF in 1–9-year-olds were adjusted for the age of those examined, using 1-year age bands and the data from the most recent Viet Nam census as the reference dataset.<sup>14</sup> Similarly, cluster-level data on trichiasis were adjusted for the age and gender of those examined, using 5-year age bands and the most recent census data.<sup>12</sup> The EU-level prevalence of each sign was calculated as the mean of the adjusted cluster-level proportions. We calculated confidence intervals by bootstrapping, with replacement, over 10,000 replications. Prevalence estimates for trichiasis in those aged  $\geq 50$  years were used to calculate estimates for the all-ages population by multiplying the prevalence in  $\geq 50$ -year-olds by 0.173: the proportion of the national population aged  $\geq 50$  years, based on the 2009 national census.<sup>14</sup>

Although we powered our surveys to estimate TF prevalence at EU level, our compact segment sampling approach (involving examination of 60 consenting 1–9-year-olds in a block of adjacent households in one or two *xom*) means that the proportion of children with TF in a cluster estimates the true prevalence for that group of households. To visualize the distribution of TF by cluster across the entire surveyed area, the means of the northings and eastings for all households within a cluster (obtained by GPS) were calculated to provide a single set of coordinates; results were plotted using QGIS (<http://www.qgis.org/en/site/>). Using Mathematica 11.1 (Wolfram Research, Champaign, IL, USA), we fit the cluster-level TF prevalence data to a zero-inflated negative binomial, parametrized so the mean of the distribution scaled with the sample size in that community.

### Ethical considerations

Ethical approval was granted by the ethics committees of the London School of Hygiene & Tropical Medicine, London, UK (reference 6319) and the Biomedical Ethics Board of National Institute of Ophthalmology, Hanoi, Viet Nam. We obtained informed verbal consent for examination from each participant or (for children) from their parent or guardian. Individuals with conjunctivitis, whether meeting the definition of active trachoma or not, were provided with two tubes of 1% tetracycline eye ointment; individuals with trichiasis were referred to a local surgeon.

**Table 2.** Characteristics of survey population by evaluation unit, Global Trachoma Mapping Project, Viet Nam, 2014.

Province	Districts in evaluation unit	Number of households surveyed	Children aged 1–9 years			Adults aged ≥50 years		
			Number enumerated	Proportion examined (%)	Proportion male (%)	Number enumerated	Proportion examined (%)	Proportion male (%)
Vinh Phuc	Yen Lac, Me Linh	595	1,206	97.4	52.8	758	99.5	41.8
Vinh Phuc	Vinh Tuong, Binh Xuyen, Lap Thach	573	1,211	99.8	51.5	748	100.0	44.1
Ha Tinh	Can Loc, Nghi Xuan	846	1,212	98.2	52.6	482	98.8	37.1
Hoa Binh	Kim Boi, Yen Thuy	910	1,193	99.3	53.6	317	100.0	35.3
Hung Yen	Van Lam	905	1,218	98.3	54.2	1,192	99.3	41.9
Yen Bai	Van Chan, Yen Bai town	890	1,199	98.3	50.0	472	99.8	39.0
Yen Bai	Luc Yen, Van Yen	867	1,208	99.7	53.6	417	98.1	42.0
Vinh Long	Tam Binh, Vung Liem	1,342	1,214	98.9	48.6	1,609	99.9	42.3
Ha Giang	Bac Quang, Yen Minh, Bac Me	725	1,142	96.6	49.9	290	99.3	49.3
Ninh Thuan	Ninh Hai, Ninh Phuoc	1,011	1,208	97.9	49.5	882	95.8	42.0
Binh Thuan	Ham Tan, Tuy Phong, Bac Binh	727	1,203	99.9	51.5	58	100.0	29.8
Total		9,391	13,214	98.6	51.6	7,225	99.1	41.6

## Results

Fieldwork was undertaken from May to August, 2014. In 9391 households, survey teams examined a total of 20,185 residents (98.8% of those enumerated), composed of 13,028 1–9-year-olds and 7,157 ≥ 50-year-olds. Table 2 summarizes the characteristics of the population sampled by EU. Overall, 52% of 1–9-year-olds examined and 42% of ≥50-year-olds examined were male. The mean ages of those examined were 4.8 years (standard deviation 2.6) for the 1–9-year-old group, and 63.4 years (10.4) for the ≥50-year-old group.

## Prevalence of trachoma

The prevalence of TF in 1–9-year-olds was <5% in all 11 EUs and ranged from 0% to 1.6% (Table 3, Figure 1). Among people aged ≥50 years, prevalence of any trichiasis by EU ranged from 0% to 0.75%; these estimates are equivalent to prevalences of any trichiasis in all ages of 0–0.13%. Of 136 people identified as having trichiasis, 125 (92%) were known to the health system, meaning that management for eyes with trichiasis had previously been (1) offered but refused; (2) accepted and a surgical date set; or (3) received, but disease had subsequently recurred. Accounting for

**Table 3.** Prevalence of trachoma by evaluation unit, Global Trachoma Mapping Project, Viet Nam, 2014.

Province	Districts in Evaluation Unit	Prevalence of TF in children aged 1–9 years: % (95% CI) <sup>1</sup>	Trichiasis cases unknown to health system			
			All trichiasis cases		Trichiasis cases unknown to health system	
			Prevalence in adults ≥50 years: % (95% CI) <sup>2</sup>	Prevalence in all ages: % <sup>3</sup>	Prevalence in adults ≥50 years: % (95% CI) <sup>2</sup>	Prevalence in all ages: % <sup>3</sup>
Vinh Phuc	Yen Lac, Me Linh	0	0.75 (0.41–1.15)	0.13	0.17 (0–0.39)	0.03
Vinh Phuc	Vinh Tuong, Binh Xuyen, Lap Thach	0.2 (0–0.5)	0.72 (0.49–1.02)	0.13	0.01 (0–0.04)	0.00
Ha Tinh	Can Loc, Nghi Xuan	0.0 (0–0.1)	0.08 (0–0.21)	0.01	0	0.00
Hoa Binh	Kim Boi, Yen Thuy	0	0.17 (0–0.41)	0.03	0	0.00
Hung Yen	Van Lam	0	0.35 (0.14–0.58)	0.06	0.05 (0–0.12)	0.01
Yen Bai	Van Chan, Yen Bai town	0.1 (0–0.4)	0.22 (0.06–0.37)	0.04	0	0.00
Yen Bai	Luc Yen, Van Yen	0.2 (0–0.7)	0.04 (0–0.10)	0.01	0	0.00
Vinh Long	Tam Binh, Vung Liem	0	0	0.00	0	0.00
Ha Giang	Bac Quang, Yen Minh, Bac Me	1.5 (0.5–2.9)	0.30 (0.05–0.57)	0.05	0	0.00
Ninh Thuan	Ninh Hai, Ninh Phuoc	0.1 (0–0.4)	0.45 (0.22 – 0.79)	0.08	0	0.00
Binh Thuan	Ham Tan, Tuy Phong, Bac Binh	0.2 (0–0.5)	0.23 (0.06–0.50)	0.04	0	0.00


CI, confidence interval.

<sup>1</sup>Adjusted for age, in one-year bands.

<sup>2</sup>Adjusted for gender and age, in five-year bands.

<sup>3</sup>The all-ages (population level) estimate was derived by multiplying the prevalence in ≥50-year-olds by 0.173: the proportion of the 2009 national population aged ≥50 years.





**Figure 1.** Prevalence of (a) trichomatous inflammation—follicular (TF) in children aged 1–9 years and (b) trichiasis unknown to the health system in all ages, Global Trachoma Mapping Project, Viet Nam, 2014.

previous management of trichiasis, the prevalence of trichiasis unknown to the health system among people aged  $\geq 50$  years, by EU, ranged from 0% to 0.17%, which was equivalent to 0–0.03% in all ages (Table 3, Figure 1).

### **Distribution of trichiasis by age, gender**

The age- and gender-specific distribution of trichiasis (in all EUs pooled together) is shown in Figure 2. The age/gender-specific prevalence of trichiasis known to the health system increased with age; however, there were no statistically significant differences among males compared to females (Figure 2(a)). There were very few cases of trichiasis unknown to the health system (Figure 2(b)).

### **Distribution of TF at cluster level**


Of the 220 clusters surveyed, the proportion of 1–9-year-olds with TF was 0% in 200 clusters; 1.6 to 3.5% in 19 clusters, and 10.3% in one cluster in Yên Minh district of Hà Giang Province (Figure 3). Using maximum likelihood estimation, we calculated a shape parameter for the negative binomial distribution of 0.17 (95% CI 0.07–388, bootstrap percentile,  $n = 999$ ),

the very wide confidence intervals commensurate with the fact that only 20 clusters contained cases of TF.

## **Discussion**

This series of pre-validation trachoma surveillance surveys showed that all 11 of the formerly endemic EUs studied now have TF prevalence estimates below the 5% elimination threshold. While in two of 11 EUs, our data suggest that the prevalence of trichiasis in all ages was  $> 0.1\%$  in 2014, most people with trichiasis appear to be “known to the health system”. The prevalence of trichiasis unknown to the health system in all ages was below the elimination threshold in each EU.

These surveys were epidemiologically robust. Whereas previous trachoma surveys in Viet Nam used the assurance sampling trachoma rapid assessment (ASTRA) methodology,<sup>15</sup> we used PBPS methods recommended by WHO.<sup>12</sup> Unlike ASTRA, PBPS provide point prevalence estimates for TF and trichiasis,<sup>16</sup> which are needed to determine if elimination thresholds have been reached.<sup>9</sup> A second point of difference was the fact that individuals with trichiasis were asked about previous management recommendations from health workers: the first GTMP-supported survey in which this was




**Figure 2.** Age- and gender-specific prevalence of trichiasis (A) known, and (B) unknown to the health system in those aged  $\geq 50$  years; data from all evaluation units combined, Global Trachoma Mapping Project, Viet Nam, 2014.

done. These questions were added in order to help the country determine whether EUs had reached elimination thresholds for trichiasis, which are couched in terms of the prevalence of trichiasis “unknown to the health system” – that is, the prevalence of trichiasis, counting only those individuals with the condition who have not previously been offered an operation or epilation for it.<sup>17</sup> Following successful implementation of the questions in Viet Nam, their use was adopted throughout the GTMP.

There were some weaknesses in our work in Viet Nam that should be acknowledged. Although we took our methodological template from previous WHO guidance<sup>18</sup> and worked within the framework of the GTMP,<sup>12</sup> we did not examine individuals aged 10–49 years, in order to try to save time for our fieldworkers, who went house-to-house to recruit survey subjects in steeply mountainous terrain. The relatively small numbers of  $\geq 50$ -year-olds examined in each EU resulted in relatively wide confidence intervals around the trichiasis prevalence estimates

(Table 3). The lack of examination of adults aged  $< 50$  years potentially limits the comparison of our findings to elimination targets, since in many trachoma-endemic settings, trichiasis can be seen in people less than 50 years old. We have assumed that the prevalence of trichiasis in people aged  $< 50$  years was negligible. To a certain extent, this assumption is supported by the age distribution of trichiasis cases that the teams identified in the field (Figure 2), which underpinned our transformation of the trichiasis prevalence estimates in  $\geq 50$ -year-olds to all-ages estimates. Our assessment of the trichiasis situation is further limited by the fact that we did not record the presence or absence of scars in the tarsal conjunctivae of eyes noted to have trichiasis, because the recommendation to do so was released after our surveys were complete.<sup>19</sup> However, this point may be moot for Viet Nam, because the data at hand, allowing for the limitations noted above, suggest that elimination threshold prevalences for trichiasis have been reached.





**Figure 3.** Geographic distribution of cluster-level prevalence of TF in children aged 1–9 years, Global Trachoma Mapping Project, Viet Nam, 2014.

We also sought to trim fieldwork time by not collecting household-level data on access to water and sanitation, which was routinely integrated within trachoma surveys in other GTMP sub-projects.<sup>20–26</sup> Due to a paucity of cases of active trachoma, our graders did not undertake live subject inter-grader agreement exercises to qualify for survey deployment, but were instead assessed using a standard set of 50 conjunctival photographs. During the GTMP, Lao PDR<sup>27</sup> and Cambodia<sup>28</sup> circumvented a similar lack of local cases by sending trainee graders to Ethiopia for field training and certification, but this was not possible for Viet Nam. It is possible that this lack of live-subject testing makes our data less accurate. Finally, in most EUs, males were relatively under-represented in the sample of  $\geq 50$ -year-olds examined (Table 2). This is a source of potential bias that would tend to lead to overestimation of trichiasis prevalence (because trichiasis occurs more frequently in women than in men)<sup>29</sup>; age- and gender-based standardization will have partially, but not completely, compensated. In all other respects, the usual quality assurance and quality control measures of the GTMP<sup>30</sup> were deployed, and within the limitations we have identified, we are confident of the strength of the data we present here.<sup>31</sup>

Notwithstanding the wide confidence intervals for the distribution shape parameter, the cluster-level distribution of TF cases was consistent with a geometric distribution. We expect to see this pattern when an

infectious disease is disappearing<sup>32,33</sup>; it was not consistent with a Poisson distribution. Occasional higher prevalence communities are expected with a geometric distribution. These do not necessarily represent persistent outliers. Selecting a community in the tail of the geometric distribution with a high prevalence is a predictable event. These outliers do not necessarily have more transmission potential in the future, and may regress towards the mean on future visits. While a single cluster in Hà Giang Province had more than one in 10 children with TF, it may not necessarily be a focus of disease of public health significance. Nonetheless, further investigations will be undertaken in Hà Giang.

In the last few decades, Viet Nam has made huge progress against trachoma. Momentum was accelerated following national adoption of the SAFE strategy in the late 1990s. From 2000 to 2008, 83,830 surgeries for trichiasis were performed, more than 2.1 million azithromycin (Zithromax®, Pfizer, New York, NY) treatments were distributed to people in 850 endemic communes of 21 northern and central coastal provinces.<sup>34</sup> The F and E components of SAFE were also implemented. Evaluations completed in 2004 and 2005 concluded that (1) health promotion activities were excellent and implementation of water and sanitation improvement activities were underway<sup>35</sup> and (2) that F & E had significantly contributed to reductions in the prevalence of active trachoma.<sup>36</sup> National-level estimates for 2015 suggested

that 91% of households had access to drinking water within a 30 minute round-trip, while 81% of households had access to improved sanitation.<sup>37</sup>

The data presented here suggest that Viet Nam is on track to meet GET2020 targets, and we recommend that the country now starts populating a dossier on trachoma's elimination as a public health problem.<sup>9</sup> This dossier could serve as a valuable information repository whilst further work is undertaken to investigate (and, if necessary, manage) trachoma in Hà Giang and its surrounds. Members of the WHO Alliance for GET2020 are ready and willing to support.

### Conflict of interest

None of the following authors have any proprietary interests or conflicts of interest related to this submission: Nguyen Xuan Hiep, Jeremiah M. Ngondi, Vu Tuan Anh, Tran Minh Dat, Cung Hong Son, Tran Van An, Nguyen Chi Dung, Nguyen Duy Thang, Brian K. Chu, Rebecca Willis, Ana Bakhtiari, Alexandre L. Pavluck, James Johnson, Joshua Sidwell, Molly Brady, Rob Henry, Aryc Mosher, Travis C. Porco, Thomas M. Lietman, Lisa Rotondo, Susan Lewallen, Paul Courtright, Anthony W. Solomon.

### Funding

The fieldwork described in this paper was generously supported by the American People through the United States Agency for International Development (USAID) via its END in Asia project, implemented by FHI360 under cooperative agreement number OAA-A-10-00051, and its ENVISION project, implemented by RTI International under cooperative agreement number AID-OAA-A-11-00048. The Global Trachoma Mapping Project, which provided logistical, epidemiological, and data management support, was funded by a grant from the United Kingdom's Department for International Development [ARIES: 203145] to Sightsavers, which led a consortium of non-governmental organizations and academic institutions to complete baseline trachoma mapping worldwide. A committee established in March 2012 to examine issues surrounding completion of global trachoma mapping was initially supported by a grant from Pfizer to the International Trachoma Initiative. AWS was a Wellcome Trust Intermediate Clinical Fellow (098521) at the London School of Hygiene & Tropical Medicine. Other than the personal contributions of RH and AM, who contributed as study investigators whilst employed by USAID (and are therefore co-authors of this article), the funders of this study had no role in the study design, data collection, data analysis, data interpretation, or the writing of the report. The authors alone are responsible for the views expressed in this article and they do not necessarily represent the views, decisions or policies of the institutions with which they are affiliated.

### References

1. World Health Organization. *Neglected Tropical Diseases: Hidden Successes, Emerging Opportunities*. Geneva: World Health Organization; 2006. [http://apps.who.int/iris/bitstream/10665/69367/1/WHO\\_CDS\\_NTD\\_2006.2\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/69367/1/WHO_CDS_NTD_2006.2_eng.pdf).
2. World Health Organization Alliance for the Global Elimination of Trachoma by 2020. Eliminating trachoma: accelerating towards 2020. 2016. [http://www.who.int/trachoma/news/News\\_Trachoma\\_Towards\\_2020/en/](http://www.who.int/trachoma/news/News_Trachoma_Towards_2020/en/). Accessed September 12, 2016.
3. World Health Organization. Report of the 2nd Global Scientific Meeting on Trachoma, Geneva, 25–27 August, 2003 (WHO/PBD/GET 03.1). Geneva: World Health Organization; 2003.
4. World Health Organization. *Future Approaches to Trachoma Control: Report of a Global Scientific Meeting*, Geneva, 17–20 June 1996; 1997. <http://www.who.int/iris/handle/10665/63413>. Accessed September 12, 2016.
5. World Health Assembly. Global elimination of blinding trachoma. 51st World Health Assembly. Geneva, 16 May 1998, Resolution WHA51.11: 1998. <http://www.who.int/blindness/causes/WHA51.11/en/>. Accessed December 14, 2015.
6. Braff E, Winklestein W, Winkelstein W. Field treatment of trachoma in North Vietnam. *Public Health Rep* 1896-1970. 1952;67(12):1233. doi:10.2307/4588340.
7. Hoffman PR. ITI Expands Trachoma Control Program in Vietnam, Launches New Program in Nepal. <http://www.hoffmanpr.com/press-release/iti-expands-trachoma-control-program-in-vietnam-launches-new-program-in-nepal/>. Accessed May 8, 2017.
8. Mabey DC, Solomon AW, Foster A. Trachoma. *Lancet*. 2003; 362:223–229.
9. World Health Organization. *Validation of Elimination of Trachoma as a Public Health Problem*. Geneva, Switzerland: WHO; 2016. [http://www.who.int/neglected\\_diseases/resources/who\\_htm\\_ntd\\_2016.8/en/](http://www.who.int/neglected_diseases/resources/who_htm_ntd_2016.8/en/). Accessed May 9, 2017.
10. World Health Organization Strategic and Technical Advisory Group on Neglected Tropical Diseases. *Technical Consultation on Trachoma Surveillance. September 11–12, 2014, Task Force for Global Health, Decatur, USA (WHO/HTM/NTD/2015.02)*. Geneva, Switzerland: World Health Organization; 2015. [http://apps.who.int/iris/bitstream/10665/174085/1/WHO-HTM\\_NTD\\_2015.02\\_en.pdf](http://apps.who.int/iris/bitstream/10665/174085/1/WHO-HTM_NTD_2015.02_en.pdf).
11. Courtright P, Gass K, Lewallen S, et al. Global trachoma mapping project: training for mapping of trachoma (version 2). London: International Coalition for Trachoma Control; 2013. <http://www.trachomacoalition.org/node/357>.
12. Solomon AW, Pavluck AL, Courtright P, et al. The global trachoma mapping project: methodology of a 34-country population-based study. *Ophthalmic Epidemiol*. 2015;22(3):214–225. doi:10.3109/09286586.2015.1037401.
13. Thylefors B, Dawson CR, Jones BR, West SK, Taylor HR. A simple system for the assessment of trachoma

- and its complications. *Bull World Health Organ.* 1987;65:477–483.
14. General Statistics Office of Viet Nam. *The 2009 Vietnam Population and Housing Census: Completed Results*. Ha Noi, Viet Nam; 2010. [http://www.gso.gov.vn/default\\_en.aspx?tabid=515&idmid=5&ItemID=10799](http://www.gso.gov.vn/default_en.aspx?tabid=515&idmid=5&ItemID=10799).
  15. Myatt M, Mai NP, Quynh NQ, et al. Using lot quality-assurance sampling and area sampling to identify priority areas for trachoma control: Viet Nam. *Bull World Health Organ.* 2005;83(10):756–763.
  16. Ngondi J, Reacher M, Matthews F, Brayne C, Emerson P. Trachoma survey methods: a literature review. *Bull World Health Organ.* 2009;87(2):143–151. doi:10.2471/BLT.07.046326.
  17. World Health Organization. *Report of the 3rd Global Scientific Meeting on Trachoma, Johns Hopkins University, Baltimore, MA, 19-20 July 2010*. Geneva: World Health Organization; 2010. <http://www.who.int/blindness/publications/3RDGLOBALSCIENTIFICMEETINGONTRACHOMA.pdf>.
  18. Solomon A, Zondervan M, Kuper H, Buchan J, Mabey D, Foster A. *Trachoma Control: A Guide for Programme Managers*. Geneva, Switzerland: WHO; 2006.
  19. World Health Organization Alliance for the Global Elimination of Trachoma by 2020. *Second Global Scientific Meeting on Trachomatous Trichiasis, Cape Town, 4-6 November 2015 (WHO/HTM/NTD/2016.5)*. Geneva: World Health Organization; 2016. [http://www.who.int/trachoma/resources/who\\_htm\\_ntd\\_2016.5/en/](http://www.who.int/trachoma/resources/who_htm_ntd_2016.5/en/).
  20. Freeman MC, Ogden S, Jacobson J, et al. Integration of water, sanitation, and hygiene for the prevention and control of neglected tropical diseases: a rationale for inter-sectoral collaboration. *PLoS Negl Trop Dis.* 2013;7(9):e2439. doi:10.1371/journal.pntd.0002439.
  21. Kalua K, Chisambi A, Chinyanya D, et al. Completion of baseline trachoma mapping in malawi: results of eight population-based prevalence surveys conducted with the global trachoma mapping project. *Ophthalmic Epidemiol.* 2016;23(sup1):32–38. doi:10.1080/09286586.2016.1230224.
  22. Mwingira UJ, Kabona G, Kamugisha M, et al. Progress of trachoma mapping in mainland Tanzania: results of baseline surveys from 2012 to 2014. *Ophthalmic Epidemiol.* 2016;23(6):373–380. doi:10.1080/09286586.2016.1236974.
  23. Elshafie BE, Osman KH, Macleod C, et al. The epidemiology of trachoma in Darfur States and Khartoum State, Sudan: results of 32 population-based prevalence surveys. *Ophthalmic Epidemiol.* 2016;23(6):381–391. doi:10.1080/09286586.2016.1243718.
  24. Ko R, Macleod C, Pahau D, et al. Population-based trachoma mapping in six evaluation units of Papua New Guinea. *Ophthalmic Epidemiol.* 2016;23(sup1):22–31. doi:10.1080/09286586.2016.1235715.
  25. Adera TH, Macleod C, Endriyas M, et al. Prevalence of and risk factors for trachoma in Southern Nations, Nationalities, and Peoples' Region, Ethiopia: results of 40 population-based prevalence surveys carried out with the global trachoma mapping project. *Ophthalmic Epidemiol.* 2016;23(sup1):84–93. doi:10.1080/09286586.2016.1247876.
  26. Mpyet C, Muhammad N, Adamu MD, et al. Prevalence of trachoma in Kano State, Nigeria: results of 44 local government area-level surveys. *Ophthalmic Epidemiol.* 2017;24(3):195–203. doi:10.1080/09286586.2016.1265657.
  27. Southisombath K, Sisalermsak S, Chansan P, et al. National trachoma assessment in the Lao People's Democratic Republic in 2013–2014. *Ophthalmic Epidemiol.* 2016;23(sup1):8–14. doi:10.1080/09286586.2016.1236973.
  28. Meng N, Seiha D, Thorn P, et al. Assessment of trachoma in Cambodia: trachoma is not a public health problem. *Ophthalmic Epidemiol.* 2016;23(sup1):3–7. doi:10.1080/09286586.2016.1230223.
  29. Cromwell EA, Courtright P, King JD, Rotondo LA, Ngondi J, Emerson PM. The excess burden of trachomatous trichiasis in women: a systematic review and meta-analysis. *Trans R Soc Trop Med Hyg.* 2009;103(10):985–992. doi:10.1016/j.trstmh.2009.03.012.
  30. Solomon A, Willis R, Pavluck AL, Alemayehu W. Quality Assurance and Quality Control in the Global Trachoma Mapping Project. *Am J Trop Med Hyg.* 2018;99(4):858–863. doi:10.4269/ajtmh.18-0082.
  31. Engels D. The global trachoma mapping project: a catalyst for progress against neglected tropical diseases. *Ophthalmic Epidemiol.* 2016;23(sup1):1–2. doi:10.1080/09286586.2016.1257139.
  32. Lietman TM, Gebre T, Abdou A, et al. The distribution of the prevalence of ocular chlamydial infection in communities where trachoma is disappearing. *Epidemics.* 2015;11:85–91. doi:10.1016/j.epidem.2015.03.003.
  33. Rahman SA, West SK, Mkocho H, et al. The distribution of ocular chlamydia prevalence across tanzanian communities where trachoma is declining. *PLoS Negl Trop Dis.* 2015;9(3):e0003682. doi:10.1371/journal.pntd.0003682.
  34. International Trachoma Initiative. *Bring on the light: the coming defeat of blinding trachoma. International Trachoma Initiative 10th Anniversary Report*. 2010. [http://www.pfizer.com/sites/default/files/responsibility/ITI\\_10th\\_anniversary\\_report.pdf](http://www.pfizer.com/sites/default/files/responsibility/ITI_10th_anniversary_report.pdf). Accessed May 10, 2017.
  35. Kuper H, Solomon AW, Buchan JC, Zondervan M, Mabey D, Foster A. Participatory evaluations of trachoma control programmes in eight countries. *Trop Med Int Health.* 2005;10(8):764–772. doi:10.1111/j.1365-3156.2005.01451.x.
  36. Khandekar R, Ton TKT, Do Thi P. Impact of face washing and environmental improvement on reduction of active trachoma in Vietnam—a public health intervention study. *Ophthalmic Epidemiol.* 2006;13(1):43–52. doi:10.1080/09286580500477507.
  37. WHO and UNICEF JMP for WASH. *Estimates on the use of water, sanitation and hygiene in Viet Nam*. 2017. <https://washdata.org/data>.

## Appendix

The Global Trachoma Mapping Project Investigators are: Agatha Aboe (1,11), Liknaw Adamu (4), Wondu Alemayehu (4,5), Menbere Alemu (4), Neal D. E. Alexander (9), Ana Bakhtiari (2,9), Berhanu Bero (4), Simon J. Brooker (1,6), Simon Bush (7,8), Brian K. Chu (2,9), Paul Courtright (1,3,4,7,11), Michael Dejene (3), Paul M. Emerson (1,6,7), Rebecca M. Flueckiger (2), Allen Foster (1,7), Solomon Gadisa (4), Katherine Gass (6,9), Teshome

Gebre (4), Zelalem Habtamu (4), Danny Haddad (1,6,7,8), Erik Harvey (1,6,10), Dominic Haslam (8), Khumbo Kalua (5), Amir B. Kello (4,5), Jonathan D. King (6,10,11), Richard Le Mesurier (4,7), Susan Lewallen (4,11), Thomas M. Lietman (10), Chad MacArthur (6,11), Colin Macleod (3,9), Silvio P. Mariotti (7,11), Anna Massey (8), Els Mathieu (6,11), Siobhain McCullagh (8), Addis Mekasha (4), Tom Millar (4,8), Caleb Mpyet (3,5), Beatriz Muñoz (6,9), Jeremiah Ngondi (1,3,6,11), Stephanie Ogden (6), Alex Pavluck (2,4,10), Joseph Pearce (10), Serge Resnikoff (1), Virginia Sarah (4), Boubacar Sarr (5),

Alemayehu Sisay (4), Jennifer L. Smith (11), Anthony W. Solomon (1,2,3,4,5,6,7,8,9,10,11), Jo Thomson (4); Sheila K. West (1,10,11), Rebecca Willis (2,9). Key: (1) Advisory Committee, (2) Information Technology, Geographical Information Systems, and Data Processing, (3) Epidemiological Support, (4) Ethiopia Pilot Team, (5) Master Grader Trainers, (6) Methodologies Working Group, (7) Prioritisation Working Group, (8) Proposal Development, Finances and Logistics, (9) Statistics and Data Analysis, (10) Tools Working Group, (11) Training Working Group.