

**PREVALENCE OF OVERWEIGHT AND OBESITY AMONG
ADOLESCENTS IN BANGLADESH: DOES FOOD HABITS AND
PHYSICAL ACTIVITIES HAVE GENDER DIFFERENTIAL
EFFECT?**

Journal:	<i>Journal of Biosocial Science</i>
Manuscript ID	JBS-4581.R2
Manuscript Type:	Research Article
Date Submitted by the Author:	29-Jan-2019
Complete List of Authors:	Khan, Md. Mostaured; University of Rajshahi, Population Science and Human Resource Development Karim, Masud ; University of Rajshahi, Population Science and Human Resource Development Islam, Ahmed Zohirul; University of Rajshahi, Population Science & Human Resource Development Islam, Md. Rafiqul; University of Rajshahi, Population Science and Human Resource Development Khan, Hafiz ; University of West London, The Graduate School Khalilullah, Md. Ibrahim ; University of Rajshahi, Population Science and Human Resource Development
Keywords:	Obesity, Child Health and Nutrition

SCHOLARONE™
Manuscripts

**PREVALENCE OF OVERWEIGHT AND OBESITY AMONG ADOLESCENTS
IN BANGLADESH: DOES FOOD HABITS AND PHYSICAL ACTIVITIES HAVE
GENDER DIFFERENTIAL EFFECT?**

Authors: Md. Mostaufed Ali Khan¹, Masud Karim¹, Ahmed Zohirul Islam¹, Md. Rafiqul Islam^{1*}, Hafiz T. A. Khan² and Md. Ibrahim Khalilullah¹

¹ Department of Population Science and Human Resource Development,
University of Rajshahi, Rajshahi-6205, Bangladesh.

² The Graduate School,
University of West London,
St Mary's Road, Ealing, London W5 5RF, United Kingdom.

****Corresponding Author:***

Md. Rafiqul Islam
Professor, Department of Population Science and Human Resource Development,
University of Rajshahi, Rajshahi-6205, Bangladesh.

Email address of the corresponding author: rafique_pops@yahoo.com

Keywords: Obesity; Adolescent; Food habit; Physical activities; Bangladesh.

1 **Abstract**

2 The aim of the study is to examine the gender differential outcomes of food habits and
3 physical activities on obesity among school-aged adolescents in Bangladesh. Nationally
4 representative data extracted from the 2014 Global School-based Student Health Survey
5 (GSHS) were utilized. The information related to physical and mental health was collected
6 from 2989 school-aged adolescents in Bangladesh. To fulfill the study aims, an exploratory
7 data analysis and multivariate logistic regression model were employed. Female adolescents
8 were at a lower risk of being overweight or obese (AOR = 0.573) with a prevalence of 7.4%
9 compared to males (9.9%). The results showed that high consumption of vegetables (both:
10 AOR = 0.454; males: AOR = 0.504; and females: AOR = 0.432), high soft drink
11 consumption (both: AOR=2.357; males: AOR = 2.929; and females: AOR = 1.677), high fast
12 food consumption (both: AOR = 2.777; males: AOR = 6.064; and females: AOR = 1.695),
13 sleep disturbance (both: AOR = 0.675; males: AOR = 0.590; and females: AOR = 0.555), and
14 regular walking or cycling to school (both: AOR = 0.472; males: AOR = 0.430; and females:
15 AOR = 0.557) were vital influencing factors for being overweight or obese among
16 adolescents across both sexes. Sedentary activities during leisure time were also identified as
17 significant predictors of being overweight and obesity for males. Regular fruit and vegetable
18 consumption, the avoidance of soft drinks and fast food, an increase in vigorous physical
19 activity, regular attendance at physical education classes and less sedentary activities in
20 leisure time could all help reduce the risk of being overweight or obese for both sexes.

21

22

23

24

Introduction

25 Obese and overweight children and adolescents represents one of the biggest
26 challenges to face public health in the 21st century and is greatly affecting many low and
27 middle-income countries (LMICs) (De Onis *et al.*, 2010; Peng *et al.*, 2017). The prevalence
28 of obesity across the world has nearly trebled since 1975 (WHO, 2018). In 2016, over 1.9
29 billion adults and 340 million adolescents worldwide, including children, were found to be
30 overweight or obese (WHO, 2018). A high risk of obesity was observed particularly for
31 Asians and Pacific Islanders (Young *et al.*, 2017) although in South Asian countries,
32 malnutrition (stunting, wasting and underweight) among children is a more hazardous
33 situation. Problems with obesity are also a matter of vital concern in many developing
34 countries including Bangladesh due to its flourishing economy (Shafique *et al.*, 2007). Since
35 the year 2000, the increase in Body Mass Index (BMI) in East and South Asian countries in
36 particular has accelerated swiftly for both sexes (Collaboration, 2017). Rapid urbanization
37 and industrialization, plus economic development and globalization of food production are
38 some of the important causal factors for this situation emerging in the developing world.

39 Previous research has identified the many negative aspects of being overweight or
40 obese on the health and growth of children and adolescents that can extend into adulthood
41 and increase the risk of developing chronic diseases such as cardiovascular disease (Singh *et*
42 *al.*, 2013), chronic kidney disease (Singh *et al.*, 2013), diabetes, many cancers (Lauby-
43 Secretan *et al.*, 2016), and disabilities (Dereń *et al.*, 2018). Furthermore, being overweight
44 and obesity are significantly related to mortality (Di Angelantonio *et al.*, 2016; Flegal *et al.*,
45 2013).

46 Although there is a growing body of studies that have examined the various risk
47 factors of being overweight and obese, there is no specific study that is focused on the gender

48 differential of obesity as a whole. Some studies have mentioned that the diverse food habits
49 and physical activities of children have a significant impact on their weight (Virtanen *et al.*,
50 2015), as do other metabolic and socio-demographic factors (Hossain *et al.*, 2018). These
51 factors include insufficient physical activity (Li *et al.*, 2017), shortened duration of sleep at
52 night (Brug *et al.*, 2012), physical education (PE) at school (Naiman *et al.*, 2015) and
53 physical activity (PA) facilities (Hood *et al.*, 2014). Diverse food habits involving the
54 consumption of fast food (Davis & Carpenter, 2009; Rosenheck, 2008), low level of fruit and
55 vegetable intake and high fat and sugar intake (Epstein *et al.*, 2012), food insecurity (Lyons *et*
56 *al.*, 2008; Robaina & Martin, 2013) and poor diet quality (Robaina & Martin, 2013), were
57 also found to be important determinants for overweight and obese children and adolescents.

58 In most developing countries, epidemiological studies on school-level risk factors for
59 obesity are still inadequate and any differences in terms of gender are unknown. Males and
60 females display differences in fat stores, dissimilation in anatomical fat distribution, and also
61 in high food intake and low physical activity (Reue, 2017) that gives strength to this study. In
62 Bangladesh, gender discrimination exists in all sectors including health and nutrition
63 (Hossain *et al.*, 2018; Shafique *et al.*, 2007). A number of attempts have been made to
64 uncover the risk factors of being overweight or obese but there has not been any research on
65 the gender differential risk factors among children and adolescents. This study focuses on the
66 prevalence of obesity and on ascertaining the gender differential outcomes of food habits and
67 physical activity on overweight and obese school-aged adolescents in Bangladesh.

68 **Methods**

69 ***Study design and Sampling procedure***

70 This study has used data extracted from the Global School-based Student Health
71 Survey (GSHS) 2014. The survey collected data from school-age adolescents (usually aged

72 11-17 years) in 43 developing countries including Bangladesh and was administrated by the
73 World Health Organization (WHO) in collaboration with the Center for Disease Control
74 (CDC). Data were collected using a clustered sampling technique and a standardized
75 scientific sample selection process, conventional school-based methodology, and a
76 combination of core questionnaire modules with expanded questions plus country-specific
77 questionnaires utilized by the survey. The school response rate was 90–100% with the student
78 response rate ranged between 76–96% and the overall response rate at 69–96% for each of
79 the countries. In Bangladesh, information related to dietary behaviors, hygiene, drug, tobacco
80 and alcohol use, sexual behaviors, mental health, physical activity etc. and was collected by
81 GSHS in 2014 from 2989 adolescents. Full clarifications of the study, including the core
82 questionnaire used with items selected from pertinent modules, are available at the websites
83 of CDC and WHO (C. WHO, 2017).

84 ***Calculation of BMI***

85 The respondent's BMI was calculated applying the following formula:

$$86 \quad BMI = \frac{Weight (kg)}{Height^2 (m)}$$

87 As all the respondents were up to 18 years of age, they were classed as being overweight if
88 their calculated BMI exceeded the standardized value for age and sex at +1SD of Z scores of
89 BMI (equivalent to BMI 25 kg/m² at 19 years of age). They were classed as being obese if
90 their calculated BMI exceeded the standardized value for age and sex at +2SD of Z scores of
91 BMI (equivalent to BMI 30 kg/m² at 19 years of age) on the basis of BMI interpretation
92 provided by WHO (Onis *et al.*, 2007; WHO, 2015).

93

94

95 *Outcome and Explanatory variables*

96 To achieve the objective of the study, being overweight or obese were considered as
97 dependent or outcome variables. The outcome variable was addressed as follows:

$$98 \quad Y = \textit{Overweight or obese} = \begin{cases} 1, & \text{Yes} \\ 0, & \text{No} \end{cases}$$

99 Several explanatory variables related to food insecurity, food habits, depression and physical
100 activities were treated as risk factors for being overweight and obese with variables selected
101 in accordance with their importance based on previous research. The information was
102 categorized according to the recommendation provided by WHO (WHO, 2012). A complete
103 list of explanatory variables is shown in table 1. (**Table 1**)

104 *Statistical analysis*

105 Any association between the state of being overweight and obese and different
106 explanatory variables were assessed by Chi-square tests (usually, set at $p < 0.05$ level of
107 significance). As the outcome variable of this study had two categories, the binary logistic
108 regression model was fitted to measure the impact of selected explanatory variables on the
109 outcome variable. In this study, the odds ratios (ORs) were estimated to assess the strength of
110 association between the outcome variable and the explanatory variables, and 95% confidence
111 intervals (CIs) were exerted to examine the level of significance. The data were analyzed
112 using the computer program SPSS in Windows version 23.0 (SPSS Inc., Chicago, IL).

113 **Results**

114 Table 2 represents the characteristics of the respondents. The mean age was 14.2 (± 0.98)
115 years, mean height was 1.563 meters (± 0.087), mean weight was 45.88 kilograms (± 7.868)
116 and mean calculated BMI was 18.78 kg/m² (± 2.87). The prevalence of being overweight and
117 obese was 9.9% for males and 7.4% for females. (**Table 2**)

118 Table 3 illustrates the association between being overweight and obesity and
119 selected explanatory variables were pursued by applying a Chi-square test to observe the
120 significance. In this study, the frequency of respondents to experience hunger, fruit and
121 vegetable eating, consumption of soft drinks, fast food eating, sleep disturbance, physical
122 activity (PA), physical education (PE) attendance were significantly related to being
123 overweight or obesity for adolescents of both sexes. A high consumption of fast food had the
124 highest prevalence of being overweight and obesity for males (25.3%), while the highest
125 prevalence observed for females (13.1%) was among those that never attended PE classes.
126 Male respondents with higher fruit (2.4%) and vegetable (5.4%) eating habits displayed a
127 lower prevalence of being overweight or obese. Similarly, there were only 2.5% and 4.6% of
128 females with high fruit and vegetable eating habits that were either overweight or obese.
129 Alternatively, 18.7% of males and 12.4% of females who consumed soft drinks at a high
130 frequency were overweight and obese. Only 5.9% of male and 4.9% of female respondents
131 who were vigorously physically active were found to be overweight or obese. The frequency
132 of being overweight and obese was lower among male and female respondents who walked
133 or cycled to school (male: 5.5% and female: 4.8%) and attended PE classes regularly (male:
134 8.2% and female: 7.4%). There was a significant association among male respondents
135 between a high amount of sitting or sedentary activities leading to a high prevalence of being
136 overweight and obesity (19.2%). **(Table 3)**

137 *Effect of food habits and physical activities on being overweight and obesity*

138 Table 4 illustrates the effects of adolescent food habits and physical activities for
139 being overweight and obese in Bangladesh. The occurrence of either state was decreased for
140 female adolescents (AOR = 0.573, CI: 0.403-0.816) compared to male adolescents. Regular
141 feelings of hunger at 2.789 (AOR = 2.789, CI: 1.733-4.489) times, highly accelerates the risk
142 of being overweight and obese than among those that never feel hunger. A high consumption

143 of fruit (AOR = 0.454, CI: 0.205-0.997) and vegetables (AOR = 0.475, CI: 0.294-0.768)
144 significantly diminished the risk of adolescents being overweight or obese. However, a high
145 consumption of soft drinks (AOR = 2.357, CI: 1.544-3.597) and fast food (AOR = 2.777, CI:
146 1.755-4.392) significantly increased the risks. Adolescents with frequent sleep disturbances
147 (AOR = 0.675, CI: 0.481-0.947) were found less likely to be overweight or obese. This was
148 also the case for those adolescents that walked or cycled to school (AOR = 0.472, CI: 0.327-
149 0.682) and attended regular PE classes (AOR = 0.592, CI: 0.327-0.682) when compared to
150 those that never walked or cycled or attended PE.

151 The fitted model of Cox and Snell R^2 , and Nagelkerke R^2 was shown to be 61.0%
152 and 81.3%, respectively and was estimated from the linear relationship between the
153 independent variables. The overall model was significant when all independent variables
154 were controlled for age. (Table 4)

155 *Gender differential effect of food habits on being overweight and obesity*

156 Table 5 shows the results of the logistic regression model of the gender differential
157 influence of food patterns on being overweight and obesity among school-aged adolescents in
158 Bangladesh. The likelihood of either of these states was decreased for males who sometimes
159 went hungry (AOR = 1.399, CI: 1.036-1.891) or went hungry most of the time (AOR =
160 2.759, CI: 1.846-4.125) than it was for respondents that never went hungry. The risk of being
161 overweight or obese was also decreased for males that ate a lot of fruit (AOR = 0.372, CI:
162 0.203-0.683). The occurrence was decreased for males with a high frequency of vegetable
163 (AOR = 0.504, CI: 0.333-0.764) eating and reduced for females with an average (AOR =
164 0.582, CI: 0.372-0.910) or high vegetable eating habit (AOR = 0.432, CI: 0.248-0.753)
165 compared to males and females with a low vegetable eating habit. However, males with an
166 average soft drink consumption habit were at a higher risk of being overweight or obese
167 (AOR = 2.583, CI: 1.855-3.597) as were adolescents with a high soft drinks' consumption

168 habit (male: AOR = 2.929, CI: 2.086-4.112; female: AOR = 1.677, CI: 1.022-2.753)
169 compared to adolescents whose weekly consumption of soft drinks was lower. High
170 consumption of fast food significantly increased the chances of ending up overweight or
171 obese for both sexes (male: AOR = 6.064, CI: 4.327-8.499; and female: AOR = 1.695, CI:
172 1.011-3.174) as it was for males with an average fast food eating habit (AOR = 1.503, CI:
173 1.084-2.083).

174 The fitted model of Cox and Snell R^2 , and Nagelkerke R^2 was shown to be 56.1%
175 and 74.7% respectively of the variance for males, and 58.9% and 78.6% respectively of the
176 variance for females and was estimated from the linear relationship between the independent
177 variables. The overall model was significant when all independent variables were controlled
178 for age. (**Table 5**)

179 ***Gender differential effect of physical activities on being overweight and obesity***

180 The results of the logistic regression model shown in Table 6 illustrate the effect of
181 PA on being overweight and obesity among school-aged adolescents. Sleep disturbance was
182 found to have a significantly decreased association with obesity (male: AOR = 0.590, CI:
183 0.455-0.766; and female: AOR = 0.555, CI: 0.369-0.837). As would be expected, being
184 overweight or obese decreased among vigorously physically active males (AOR = 0.751, CI:
185 0.592-0.991) as it was for respondents who took part in moderate PA. The risk of being
186 overweight or obese was reduced for both males and females who occasionally walked or
187 cycled to school (male: AOR = 0.265, CI: 0.171-0.410; and female: AOR = 0.453, CI: 0.205-
188 0.924) or who regularly walked or cycled to school (male: AOR = 0.430, CI: 0.322-0.576;
189 and female: AOR = 0.557, CI: 0.359-0.866) compared to respondents that never walked or
190 cycled to school. The likelihood of being overweight or obese decreased among males and
191 females that occasionally attended PE classes (male: AOR = 0.420, CI: 0.281-0.627; and
192 female: AOR = 0.445, CI: 0.266-0.745) or for males that regularly attended such classes

193 (male: AOR = 0.488, CI: 0.330-0.722) compared to males and females that never attended
194 PE. The risk of being overweight or obese was increased by 3.404 (AOR = 3.404, CI: 2.343-
195 4.945) times higher for males with high sitting or sedentary activities compared to males with
196 moderate sitting activities.

197 In the fitted model, the Cox and Snell R^2 , and Nagelkerke R^2 was 54.1% and 77.2%
198 respectively. The variances of males and for females was 60.0% and 80.0% respectively, that
199 can be estimated from the linear relationship between the independent variables. The overall
200 model was significant when all explanatory variables were included. (Table 6)

201 Discussion

202 The prevalence of being overweight and obesity is showing an increasing trend in
203 Bangladesh (Biswas *et al.*, 2017). It has not yet become an alarming situation for adolescents
204 but it is increasing day-by-day. The findings of this study indicate that the risk of males being
205 overweight or obese is notably higher than it is for females. This study findings also show
206 that male adolescents with high food insecurity are at an increased risk of being overweight
207 or obese that is consistent with earlier studies (Robaina & Martin, 2013; Sanjeevi *et al.*,
208 2018). As well as Sanjeevi *et al.* (2018) showing a uniformity with the results of our study,
209 these authors concluded that food insecurity is associated with a less conducive
210 multidimensional home environmental subscale score and poor diet quality that, in turn, is
211 related to greater BMI. Lohman *et al.* (2016) also identified a gender differential outcome of
212 household food insecurity for being overweight or obese. In Bangladesh, less importance is
213 generally given to female children than to males in all sectors.

214 Dietary behavior and different food patterns have a diverse impact on being
215 overweight or obesity (Rautiainen *et al.*, 2015; Virtanen *et al.*, 2015). This study has
216 identified a significantly lower risk of both these states in adolescents with a high fruit and
217 vegetable eating habit. A gender differential effect was seen again when such eating habits

218 significantly decreased the risk for male adolescents, but in the case of females, no significant
219 effect due to fruit-eating was found but a highly significant effect was identified with average
220 to higher vegetable consumption. Previous research supports this hypothesis by identifying
221 that there is a significant positive impact of regular fruit and vegetable intake among children
222 and adolescents (Epstein *et al.*, 2012; Field *et al.*, 2003). According to Rohde *et al.* (2017),
223 responsible intake of macronutrients, energy, fruit and vegetables can help restrain excessive
224 weight gain among children. More precisely, fruit and vegetables provide fiber, are low in
225 calories and rich in minerals and vitamins that help to keep a person healthy and energized.

226 This study shows that a high consumption of soft drinks and fast food increases the
227 risk for both male and female adolescents of becoming overweight or obese. As a
228 consequence, adolescents are at a high risk of experiencing problems with their weight
229 regardless of their intake of junk food. A number of previous studies have shown that high
230 consumption of soft drinks and fast food has a highly negative effect on obesity in
231 adolescents and young children (Davis & Carpenter, 2009; Moore *et al.*, 2009; Rosenheck,
232 2008). These types of food and drink contain more fat and sugar, and fewer vitamins and
233 minerals than healthier alternatives and therefore can lead to poor weight management and
234 body metabolism leading further to a risk of obesity (Lucan & DiNicolantonio, 2015). In
235 recent experiments, researchers have shown that reducing soft drink and fast food
236 consumption in adolescents has been successful in lessening the prevalence of obesity
237 (Cantoral *et al.*, 2016; Hu, 2013; Laxy *et al.*, 2015). In addition, a high intake of artificially
238 sweetened soft beverages enhances the risk of obesity-related cancers (Hodge *et al.*, 2018). In
239 Bangladesh, the quality of soft drinks and fast food is much poorer than in developed
240 countries that is perhaps a similar situation in other developing countries.

241 The study results show there is a lower risk of obesity among adolescents of both
242 sexes who often face sleep problems due to depression, that is inconsistent with findings from

243 several previous studies (Brug *et al.*, 2012; Mannan *et al.*, 2016; Nielsen *et al.*, 2011). In
244 addition, (Nielsen *et al.*, 2011) observed a significant link between short duration of sleep and
245 being overweight or obesity among young adults including children. A meta-analysis and
246 systematic review of longitudinal studies conducted by Mannan *et al.* (2016) revealed a 70%
247 greater risk of depressed male and female adolescents being overweight or obese.

248 Physical activity (PA) is an emerging determinant of weight for both children and
249 adults. The study found there was a much lower risk of male adolescents being overweight or
250 obese if they were vigorously active compared to those who were moderately active. This
251 hypothesis is supported by a few earlier studies such as Ogden *et al.* (2016). Also showing a
252 consistency with the study findings, Chaput *et al.* (2018) noticed there was a lower risk of
253 obesity by level of sedentary behavior in children who were vigorously physically active but
254 found no significant effect of PA in case of overweight or obese males or females. Males and
255 females that regularly walked or cycled to school were at a very low risk of being overweight
256 or obese. Walking and cycling carry twofold advantages: they help protect the environment
257 and prevent excessive weight gain by increasing body metabolism. Responsible parents
258 should therefore encourage their children to regularly walk or cycle to school.

259 The attendance of adolescents at PE has also been identified as a feasible predictor of
260 being overweight and obesity in Bangladesh. The respondents of both sexes that regularly
261 attended PE had a very low risk of developing weight problems compared to those that never
262 attended PE (Naiman *et al.*, 2015). Spending time at PE can help reduce the gap between
263 actual and recommended physical activity for children and adolescents (Fernandes & Sturm,
264 2010) and also help increase the number of days per week spent in vigorous PA (Jinsook,
265 2012). So, it can be seen that PE indirectly affects excessive weight gain among adolescents
266 and children. Unfortunately, however, PE facilities in Bangladesh are very poor and there is
267 poor awareness of the benefits of PE among parents plus a lack of strict regulations.

268 Sedentary activities of adolescents show a negative impact on being overweight or obesity for
269 male adolescents but show no significant effects on female adolescents. Those male
270 adolescents with high levels of sitting activity per day are almost at three times higher risk of
271 becoming overweight or obese. This finding is supported by previous research related to
272 adults (Chau *et al.*, 2012; Ng *et al.*, 2017). In Australia, the risk of being overweight or obese
273 increases significantly among workers with mostly sitting jobs than workers with mostly
274 standing jobs (Chau *et al.*, 2012). However, there are no studies that describe the effect of
275 sitting behaviors as a cause of weight problems or obesity among children and adolescents.
276 Adolescent leisure time activities per day such as watching TV, gossiping, playing computer
277 games etc. increase their risk of becoming overweight or obese especially for male
278 adolescents.

279 This study had several limitations. For example, a secondary source of data was used
280 for analysis and thus some important variables are missing in relation to being overweight
281 and obesity. Nevertheless, an attempt has been made to depict a compact demonstration of
282 the effect of adolescent eating behaviors and PA on being overweight and obese. Future
283 studies could be undertaken to collect data covering variables involved in differences
284 between rural-urban areas.

285 In conclusion, findings from the study suggest that the levels of being overweight or
286 obesity among school-aged adolescents in Bangladesh have yet to be improved. This research
287 discovered gender differences in food practice and physical activity among adolescents that
288 affect their overweight or obesity levels and therefore indicates improvements are needed in
289 their behavior around eating and physical activity. A regular consumption of healthy food,
290 particularly a diet rich in fruit and vegetables and avoidance of soft drinks and fast food
291 especially for males, are necessary components for helping to lessen the risk of adolescents in
292 Bangladesh being overweight or obese. Increasing levels of physical activity, cutting back on

293 high levels of leisure time sitting activities, especially among males, and encouraging
294 adolescents of both sexes to regularly walk or cycle to school, can all help to cut the risk of
295 developing weight problems. Policy in this area should focus on the need for regular
296 attendance at PE classes to help improve the health of school-age adolescents. The
297 implementation of such policies would help decrease the risk of adolescent obesity in
298 Bangladesh and, in turn, be helpful for prolonging their good health.

299 **Abbreviations**

300 GSHS, Global School-based Student Health Survey; BMI, Body Mass Index; SD, Standard
301 Deviation; PE, Physical Education; PA, Physical Activity; WHO, World Health
302 Organization; CDC, Center for Disease Control; CI, Confidence Interval; OR, Odds Ratio.

303 **Acknowledgments**

304 The authors are thankful to the Office of the Global School-based Student Health Survey
305 (GSHS), Department of Chronic Diseases and Health Promotion, World Health Organization,
306 jointly worked with Center for Disease Control (CDC), for permitting us to use the datasets
307 for this analysis.

308 **Ethical Approval**

309 The study was ethically approved and the statement was obtained from the Ministry of Health
310 and Family Welfare, Dhaka, Bangladesh. The World Health Organization (WHO) financially
311 and technically supports this survey with the collaboration of the Center for Disease Control
312 (CDC). Further, written permission was obtained from each participating school and from all
313 classroom teachers.

314 **Conflicts of interest**

315 The authors have no competing interests to declare.

316

Funding

317 The authors received no funding grant from any funding agency, commercial entity or not-
318 for-profit organization for this study.

319 References

- 320 **Biswas T, Uddin MJ, Mamun AA, Pervin S and Garnett SP** (2017). Increasing prevalence of overweight and
321 obesity in Bangladeshi women of reproductive age: Findings from 2004 to 2014. *PLOS ONE*, **12**(7),
322 e0181080.
- 323 **Brug J, van Stralen MM, te Velde SJ, Chinapaw MJ, De Bourdeaudhuij I et al.** (2012). Differences in
324 weight status and energy-balance related behaviors among schoolchildren across Europe: the
325 ENERGY-project. *PLOS ONE*, **7**(4), e34742.
- 326 **Cantoral A, Téllez-Rojo MM, Ettinger A, Hu H, Hernández-Ávila M and Peterson K** (2016). Early
327 introduction and cumulative consumption of sugar-sweetened beverages during the pre-school period
328 and risk of obesity at 8–14 years of age. *Pediatric Obesity*, **11**(1), 68-74.
- 329 **Chaput JP, Barnes J, Tremblay M, Fogelholm M, Hu G et al.** (2018). Thresholds of physical activity
330 associated with obesity by level of sedentary behaviour in children. *Pediatric Obesity*, **13**, 450–457.
- 331 **Chau JY, van der Ploeg HP, Merom D, Chey T and Bauman AE** (2012). Cross-sectional associations
332 between occupational and leisure-time sitting, physical activity and obesity in working adults.
333 *Preventive Medicine*, **54**(3), 195-200.
- 334 **Collaboration NRF** (2017). Worldwide trends in body-mass index, underweight, overweight, and obesity from
335 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128·9 million
336 children, adolescents, and adults. *The Lancet*, **390**(10113), 2627-2642.
- 337 **Davis B and Carpenter C** (2009). Proximity of fast-food restaurants to schools and adolescent obesity.
338 *American Journal of Public Health*, **99**(3), 505-510.
- 339 **De Onis M, Blössner M and Borghi E** (2010). Global prevalence and trends of overweight and obesity among
340 preschool children-. *The American journal of clinical nutrition*, **92**(5), 1257-1264.
- 341 **Dereń K, Nyankovskyy S, Nyankovska O, Łuszczki E, Wyszynska J et al.** (2018). The prevalence of
342 underweight, overweight and obesity in children and adolescents from Ukraine. *Scientific Reports*,
343 **8**(1), 3625.
- 344 **Di Angelantonio E, Bhupathiraju SN, Wormser D, Gao P, Kaptoge S et al.** (2016). Body-mass index and
345 all-cause mortality: individual-participant-data meta-analysis of 239 prospective studies in four
346 continents. *The Lancet*, **388**(10046), 776-786.
- 347 **Epstein LH, Gordy CC, Raynor HA, Beddome M, Kilanowski CK and Paluch R** (2012). Increasing fruit
348 and vegetable intake and decreasing fat and sugar intake in families at risk for childhood obesity.
349 *Obesity*, **9**(3), 171-178.
- 350 **Fernandes M and Sturm R** (2010). Facility provision in elementary schools: Correlates with physical
351 education, recess, and obesity. *Preventive Medicine*, **50** (Suppl 1), S30-S35.
- 352 **Field AE, Gillman MW, Rosner B, Rockett HR and Colditz GA** (2003). Association between fruit and
353 vegetable intake and change in body mass index among a large sample of children and adolescents in
354 the United States. *International journal of obesity*, **27**(7), 821-826.
- 355 **Flegal KM, Kit BK, Orpana H and Graubard BI** (2013). Association of all-cause mortality with overweight
356 and obesity using standard body mass index categories: A systematic review and meta-analysis. *JAMA*,
357 **309**(1), 71-82.
- 358 **Hodge AM, Bassett JK, Milne RL, English DR and Giles GG** (2018). Consumption of sugar-sweetened and
359 artificially sweetened soft drinks and risk of obesity-related cancers. *Public health nutrition*, **21**(9),
360 1618-1626.

- 361 **Hood NE, Colabianchi N, Terry-McElrath YM, O'Malley PM and Johnston LD** (2014). Physical activity
362 breaks and facilities in US secondary schools. *Journal of School Health*, **84**(11), 697-705.
- 363 **Hossain MM, Islam MR, Sarker ASR, Khan MMA and Taneepanichskul S** (2018). Prevalence and
364 determinants risk factors of underweight and overweight among women in Bangladesh. *Obesity*
365 *Medicine*, **11**, 1-5
- 366 **Hu FB** (2013). Resolved: there is sufficient scientific evidence that decreasing sugar-sweetened beverage
367 consumption will reduce the prevalence of obesity and obesity-related diseases. *Obesity reviews*, **14**(8),
368 606-619.
- 369 **Jinsook K** (2012). Are Physical Education-Related State Policies and Schools' Physical Education Requirement
370 Related to Children's Physical Activity and Obesity? *Journal of School Health*, **82**(6), 268-276.
- 371 **Lauby-Secretan B, Scoccianti C, Loomis D, Grosse Y, Bianchini F and Straif K** (2016). Body fatness and
372 cancer—viewpoint of the IARC Working Group. *New England Journal of Medicine*, **375**(8), 794-798.
- 373 **Laxy M, Malecki KC, Givens ML, Walsh MC and Nieto FJ** (2015). The association between neighborhood
374 economic hardship, the retail food environment, fast food intake, and obesity: findings from the Survey
375 of the Health of Wisconsin. *BMC public health*, **15**(1), 237.
- 376 **Li M, Xue H, Wen M, Wang W and Wang Y** (2017). Nutrition and physical activity related school
377 environment/policy factors and child obesity in China: a nationally representative study of 8573
378 students in 110 middle schools. *Pediatric Obesity*, **12**(6), 485-493.
- 379 **Lohman B J, Gillette MT and Neppel TK** (2016). Harsh Parenting and Food Insecurity in Adolescence: The
380 Association With Emerging Adult Obesity. *Journal of Adolescent Health*, **59**(1), 123-127.
- 381 **Lucan SC and DiNicolantonio JJ** (2015). How calorie-focused thinking about obesity and related diseases
382 may mislead and harm public health. An alternative. *Public health nutrition*, **18**(4), 571-581.
- 383 **Lyons AA, Park J and Nelson CH** (2008). Food Insecurity and Obesity: A Comparison of Self-Reported and
384 Measured Height and Weight. *American Journal of Public Health*, **98**(4), 751-757.
- 385 **Mannan M, Mamun A, Doi S and Clavarino A** (2016). Prospective Associations between Depression and
386 Obesity for Adolescent Males and Females- A Systematic Review and Meta-Analysis of Longitudinal
387 Studies. *PLOS ONE*, **11**(6), e0157240.
- 388 **Moore LV, Diez Roux AV, Nettleton JA, Jacobs DR and Franco M** (2009). Fast-food consumption, diet
389 quality, and neighborhood exposure to fast food: the multi-ethnic study of atherosclerosis. *American*
390 *journal of epidemiology*, **170**(1), 29-36.
- 391 **Naiman DI, Leatherdale ST, Gotay C and Mâsse LC** (2015). School factors associated with the provision of
392 physical education and levels of physical activity among elementary school students in Ontario. *Can J*
393 *Public Health*, **106**(5), 290-296.
- 394 **Ng N, Santosa A and Kowal P** (2017). Sitting time and obesity among older adults in low- and middle-income
395 countriesNawi Ng. *European Journal of Public Health*, **27**(suppl_3), Doi: 10.1093/eurpub/ckx186.042
- 396 **Nielsen L, Danielsen K and Sørensen T** (2011). Short sleep duration as a possible cause of obesity: critical
397 analysis of the epidemiological evidence. *Obesity reviews*, **12**(2), 78-92.
- 398 **Ogden CL, Carroll MD and Lawman HG** (2016). Trends in obesity prevalence among children and
399 adolescents in the united states, 1988-1994 through 2013-2014. *JAMA*, **315**(21), 2292-2299.
- 400 **Onis Md, Onyango AW, Borghi E, Siyam A, Nishida C and Siekmann J** (2007). Development of a WHO
401 growth reference for school-aged children and adolescents. *Bulletin of the World health Organization*,
402 **85**(9), 660-667.
- 403 **Peng W, Goldsmith R and Berry EM** (2017). Demographic and lifestyle factors associated with adherence to
404 the Mediterranean diet in relation to overweight/obesity among Israeli adolescents: findings from the
405 Mabat Israeli national youth health and nutrition survey. *Public health nutrition*, **20**(5), 883-892.
- 406 **Rautiainen S, Wang L, Lee IM, Manson JE, Buring JE and Sesso HD** (2015). Higher Intake of Fruit, but
407 Not Vegetables or Fiber, at Baseline Is Associated with Lower Risk of Becoming Overweight or Obese
408 in Middle-Aged and Older Women of Normal BMI at Baseline. *The Journal of Nutrition*, **145**(5), 960-
409 968.
- 410 **Reue K** (2017). Sex differences in obesity: X chromosome dosage as a risk factor for increased food intake,
411 adiposity and co-morbidities. *Physiology & Behavior*, **176**, 174-182.

- 412 **Robaina KA and Martin KS** (2013). Food insecurity, poor diet quality, and obesity among food pantry
413 participants in Hartford, CT. *Journal of nutrition education and behavior*, **45**(2), 159-164.
- 414 **Rohde JF, Larsen SC, Ängquist L, Olsen NJ, Stougaard M, Mortensen EL et al.** (2017). Effects of the
415 Healthy Start randomized intervention on dietary intake among obesity-prone normal-weight children.
416 *Public health nutrition*, **20**(16), 2988-2997.
- 417 **Rosenheck R** (2008). Fast food consumption and increased caloric intake: a systematic review of a trajectory
418 towards weight gain and obesity risk. *Obesity reviews*, **9**(6), 535-547.
- 419 **Sanjeevi N, Freeland-Graves J and Hersh M** (2018). Food insecurity, diet quality and body mass index of
420 women participating in the Supplemental Nutrition Assistance Program: The role of intrapersonal,
421 home environment, community and social factors. *Appetite*, **125**, 109-117.
- 422 **Shafique S, Akhter N, Stallkamp G, de Pee S, Panagides D and Bloem MW** (2007). Trends of under-and
423 overweight among rural and urban poor women indicate the double burden of malnutrition in
424 Bangladesh. *International journal of epidemiology*, **36**(2), 449-457.
- 425 **Singh GM, Danaei G, Farzadfar F, Stevens GA, Woodward M, Wormser D et al.** (2013). The Age-Specific
426 Quantitative Effects of Metabolic Risk Factors on Cardiovascular Diseases and Diabetes: A Pooled
427 Analysis. *PLOS ONE*, **8**(7), e65174.
- 428 **Virtanen M, Kivimäki H, Ervasti J, Oksanen T, Pentti J, Kouvonen A et al.** (2015). Fast-food outlets and
429 grocery stores near school and adolescents' eating habits and overweight in Finland. *European Journal*
430 *of Public Health*, **25**(4), 650-655.
- 431 **WHO** (2012). Global recommendations on physical activity for health. *World Health Organization*. WHO,
432 Geneva, Switzerland.
- 433 **WHO** (2015). BMI-for-age (5-19 years). *World Health Organization*. WHO, Geneva, Switzerland. URL:
434 http://www.who.int/growthref/who2007_bmi_for_age/en/ (accessed 12th May, 2018)
- 435 **WHO** (2018). Obesity and overweight. *World Health Organization*, WHO, Geneva, Switzerland. URL:
436 <http://www.who.int/mediacentre/factsheets/fs311/en/> (accessed 15th May, 2018)
- 437 **WHO and CDC** (2017). URL: <http://www.cdc.gov/GSHS/>; <http://www.who.int/chp/gshs/en/>
- 438 **Young DR, Koebnick C and Hsu JWY** (2017). Sociodemographic associations of 4-year overweight and
439 obese incidence among a racially diverse cohort of healthy weight 18-year-olds. *Pediatric Obesity*,
440 **12**(6), 502-510.

441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456

457 **Table 1.** The complete list of explanatory variables.

Variables	Category	Measurements	Duration	Variable Type
How often went hungry	1 = Never		During the past 30 days	Food insecurity
	2 = Rarely			
	3 = Most of the time			
Fruit eating Vegetable eating Soft drinks Fast food eating	1 = Low	less than one time per day	During the past 30 days	Food habits
	2 = Average	1-2 times per day		
	3 = High	More than 2 times per day		
Sleep disturbance	1 = Never		During the past 30 days	Depression
	2 = Often			
Physical activity	1 = Moderate	≤2 days	60 Minutes per day During the past 7 days	Physical activity
	2 = Vigorous	> 2 days		
Walk or bike to school	1 = Never	0 days	During the past 7 days	
	2 = occasionally	1-3 days		
Physical education attendance	3 = Regularly	> 3 days		
Sitting activities (includes TV watching, playing computer games etc.)	1 = Moderate	< 5 hours per day		
	2 = High	≥ 5 hours per day		

458

459

460 **Table 2.** The characteristics of study population.

Characteristics	n	Minimum	Maximum	Mean (SD)
Age (in years)	2980	11	17	14.2 (±0.98)
Height (m)	2703	1.27	1.9	1.563 (±0.087)
Weight (kg)	2703	28	102	45.88 (±7.868)
BMI	2703	13.05	39.67	18.78 (±2.87)
Gender	n	Overweight and obese (%)		
Male	1192	9.9%		
Female	1788	7.4%		

461 **Note:** All percentages are weighted.

462

463 **Table 3.** Percentage distribution of overweight/obesity among school-aged adolescents in Bangladesh according to their food habits and physical activities.
464

Risk Factors	Overweight or obese					χ^2_{cal} (<i>p-value</i>)
	Male		χ^2_{cal} (<i>p-value</i>)	Female		
	No (%)	Yes (%)		No (%)	Yes (%)	
How often went hungry:						
Never	333(93.8%)	22(6.2%)	15.419	573(93.2%)	42(6.8%)	8.840
Sometimes	507(92.5%)	41(7.5%)	(0.000)	728(94.4%)	43(5.6%)	(0.012)
Most of the time	75(81.5%)	17(18.5%)		226(89.0%)	28(11.0%)	
Fruit eating:						
Low	440(91.3%)	42(8.7%)	8.720	754(93.2%)	55(6.8%)	11.295
Average	322(90.2%)	35(9.8%)	(0.013)	561(91.1%)	55(8.9%)	(0.004)
High	160(97.6%)	5(2.4%)		1552(97.5%)	6(2.5%)	
Vegetable eating:						
Low	218(90.1%)	24(9.9%)	5.656	319(88.1%)	43(11.9%)	18.840
Average	318(90.4%)	41(9.6%)	(0.05)	666(93.5%)	46(6.5%)	(<0.0001)
High	318(94.6%)	18(5.4%)		561(95.4%)	27(4.6%)	

Soft drinks:						
Low	488(94.6%)	28(5.4%)	28.429	842(93.0%)	63(7.0%)	13.590
Average	299(92.6%)	24(7.4%)	(<0.0001)	515(95.2%)	26(4.8%)	(0.001)
High	126(81.3%)	29(18.7%)		183(87.6%)	26(12.4%)	
Fast food eating:						
Low	533(94.2%)	33(5.8%)	42.451	1009(92.7%)	79(7.3%)	9.912
Average	314(92.6%)	25(7.4%)	(<0.0001)	409(95.8%)	18(4.2%)	(0.007)
High	74(74.7%)	25(25.3%)		131(88.5%)	17(11.5%)	
Sleep disturbance:						
Never	356(88.6%)	46(11.4%)	9.502	645(90.8%)	65(9.2%)	9.072
Often	577(94.0%)	37(6.0%)	(0.002)	917(94.6%)	52(5.4%)	(0.003)
Physical activity:						
Moderate	281(87.0%)	42(13.0%)	14.509	567(90.4%)	60(9.6%)	13.772
Vigorous	635(94.1%)	40(5.9%)	(<0.0001)	959(95.1%)	49(4.9%)	(<0.0001)
Walk or bike to school:						
Never	175(81.8%)	39(18.2%)	35.484	486(89.3%)	58(10.7%)	19.068
Occasionally	181(94.3%)	11(5.7%)	(<0.0001)	161(94.7%)	9(5.3%)	(<0.0001)
Regularly	564(94.5%)	33(5.5%)		886(95.2%)	45(4.8%)	
PE attendance:						
Never	61(80.3%)	15(19.7%)	16.001	106(86.9%)	16(13.1%)	8.673
Occasionally	401(93.9%)	26(6.1%)	(<0.0001)	783(94.1%)	49(5.9%)	(0.013)
Regularly	436(91.8%)	39(8.2%)		615(92.6%)	49(7.4%)	
Sitting activities:						
Moderate	831(92.4%)	68(7.6%)	12.567	1451(93.1%)	107(6.9%)	0.000
High	63(80.8%)	15(19.2%)	(<0.0001)	68(93.2%)	5(6.8%)	(0.995)

465 **Note:** Significant at $p < 0.05$ level; PE, Physical Education

466

467 **Table 4.** Logistic regression analysis to estimate the effect of adolescents' food habits and physical activities on
468 the state of overweight/obesity in Bangladesh, 2014.

Risk Factors	Overweight or obese	
	AOR	95% CI
Gender		
Male ^(RC)	1.00
Female	0.573**	(0.403-0.816)
How often went hungry:		
Never ^(RC)	1.00
Sometimes	1.475*	(1.001-2.175)
Most of the time	2.789**	(1.733-4.489)
Fruit eating:		
Low ^(RC)	1.00
Average	1.070	(0.749-1.530)
High	0.454*	(0.205-0.997)
Vegetable eating:		
Low ^(RC)	1.00
Average	0.625*	(0.420-0.930)
High	0.475**	(0.294-0.768)
Soft drinks:		
Low ^(RC)	1.00
Average	1.244	(0.814-1.900)
High	2.357**	(1.544-3.597)
Fast food eating:		
Low ^(RC)	1.00
Average	1.244	(0.804-1.923)
High	2.777**	(1.755-4.392)

Sleep disturbance:		
Never ^(RC)	1.00
Often	0.675*	(0.481-0.947)
Physical activity:		
Moderate ^(RC)	1.00
Vigorous	0.850	(0.597-1.209)
Walk or bike to school:		
Never ^(RC)	1.00
Occasionally	0.334**	(0.185-0.601)
Regularly	0.472**	(0.327-0.682)
PE attendance:		
Never ^(RC)	1.00
Occasionally	0.483**	(0.284-0.822)
Regularly	0.592*	(0.327-0.682)
Sitting activities:		
Moderate ^(RC)	1.00
High	1.551	(0.860-2.797)
Model summary:		
Model Chi-Square (p-value)	2220.687 (<0.001)	
-2Log Likelihood	1048.196	
Cox and Snell R-square	0.610	
Nagelkerke R-square	0.813	

469 **Note:** Sample are weighted and controlled by age. “(RC)” denotes reference category; AOR, adjusted odds ratio;
 470 Significant at ** $p < 0.01$, and * $p < 0.05$ level.

471

472 **Table 5.** Logistic regression analysis to estimate the gender differential effect of food habits on adolescents’
 473 overweight/obesity in Bangladesh, 2014.

Risk Factors	Male		Female	
	AOR	95% CI	AOR	95% CI
How often went hungry:				
Never ^(RC)	1.00	1.00
Sometimes	1.399*	(1.036-1.891)	1.235	(0.791-1.928)
Most of the time	2.759**	(1.846-4.125)	1.540	(0.904-2.623)
Fruit eating:				
Low ^(RC)	1.00	1.00
Average	1.201	(0.901-1.601)	1.163	(0.763-1.773)
High	0.372**	(0.203-0.683)	0.645	(0.258-1.613)
Vegetable eating:				
Low ^(RC)	1.00	1.00
Average	1.043	(0.753-1.446)	0.582*	(0.372-0.910)
High	0.504**	(0.333-0.764)	0.432**	(0.248-0.753)
Soft drinks:				
Low ^(RC)	1.00	1.00
Average	2.583**	(1.855-3.597)	0.785	(0.470-1.312)
High	2.929**	(2.086-4.112)	1.677*	(1.022-2.753)
Fast food eating:				
Low ^(RC)	1.00	1.00
Average	1.503*	(1.084-2.083)	0.737	(0.411-1.321)
High	6.064**	(4.327-8.499)	1.695*	(1.011-3.174)
Model summary:				
Model Chi-Square (p-value)	2327.11 (<0.001)		1401.33 (<0.001)	
-2Log Likelihood	1596.663		781.801	
Cox and Snell R-square	0.561		0.589	
Nagelkerke R-square	0.747		0.786	

474 **Table 6.** Logistic regression analysis to estimate the gender differential effect of physical activities on
 475 adolescents' overweight/obesity in Bangladesh, 2014.

Risk Factors	Male		Female	
	AOR	95% CI	AOR	95% CI
Sleep disturbance:				
Never ^(RC)	1.00	1.00
Often	0.590**	(0.455-0.766)	0.555**	(0.369-0.837)
Physical activity:				
Moderate ^(RC)	1.00	1.00
Vigorous	0.751*	(0.562-0.991)	0.809	(0.532-1.229)
Walk or bike to school:				
Never ^(RC)	1.00	1.00
Occasionally	0.265**	(0.171-0.410)	0.453*	(0.205-0.924)
Regularly	0.430**	(0.322-0.576)	0.557**	(0.359-0.866)
PE attendance:				
Never ^(RC)	1.00	1.00
Occasionally	0.420**	(0.281-0.627)	0.445**	(0.266-0.745)
Regularly	0.488**	(0.330-0.722)	0.754	(0.461-1.234)
Sitting activities:				
Moderate ^(RC)	1.00	1.00
High	3.404**	(2.343-4.945)	0.469	(0.154-1.427)
Model summary:				
Model Chi-Square (p-value)	2102.016 (<0.001)		1403.618 (<0.001)	
-2Log Likelihood	1638.223		718.459	
Cox and Snell R-square	0.541		0.600	
Nagelkerke R-square	0.772		0.800	

476 **Note:** Sample are weighted and controlled by age. "(RC)" denotes reference category; AOR, adjusted odds ratio;
 477 Significant at ** $p < 0.01$, and * $p < 0.05$ level.