



PRIFYSGOL
BANGOR
UNIVERSITY

Use of a discrete choice experiment approach to elicit patients' preferences for hip fracture rehabilitation services as part of a feasibility study

Charles, Joanna; Roberts, Jessica; Din, Nafees; Williams, Nefyn; Yeo, Seow Tien; Edwards, Rhiannon

The Lancet

DOI:
[10.1016/S0140-6736\(16\)32271-1](https://doi.org/10.1016/S0140-6736(16)32271-1)

Published: 25/11/2016

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):
Charles, J., Roberts, J., Din, N., Williams, N., Yeo, S. T., & Edwards, R. (2016). Use of a discrete choice experiment approach to elicit patients' preferences for hip fracture rehabilitation services as part of a feasibility study. *The Lancet*, 388(Supplement 2), S35.
[https://doi.org/10.1016/S0140-6736\(16\)32271-1](https://doi.org/10.1016/S0140-6736(16)32271-1)

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Title of manuscript

Using a Discrete Choice Experiment (DCE) approach to elicit patients' preferences for hip fracture rehabilitation services as part of a feasibility study.

Names and affiliations of authors

Dr Joanna M Charles (PhD), Centre for Health Economics & Medicines Evaluation, Bangor University, Bangor, Gwynedd, UK

Dr Jessica L Roberts (PhD), School of Healthcare Sciences, Bangor University, UK

Dr Nafees Ud Din (MSc), School of Healthcare Sciences, Bangor University, UK

Dr Nefyn H Williams (PhD), School of Healthcare Sciences, Bangor University, UK

Ms Seow Tien Yeo (MSc), Centre for Health Economics & Medicines Evaluation, Bangor University, Bangor, Gwynedd, UK

Prof Rhiannon T Edwards (DPhil), Centre for Health Economics & Medicines Evaluation, Bangor University, Bangor, Gwynedd, UK

Corresponding Author

Dr Joanna Charles, Centre for Health Economics & Medicines Evaluation, Ardudwy, Normal Site, Bangor University, Bangor, Gwynedd, LL57 2PZ, UK. **E-mail:** j.charles@bangor.ac.uk

Background

Hip fracture is a common health issue in older age. Hip fracture is associated with consequences for the individual such as mortality and frailty. Multidisciplinary rehabilitation has been proposed as a method to help older people recover after a hip fracture. Previous research exploring patients' preferences for hip fracture rehabilitation has been conducted in Australia. However, there is little research in a UK context to inform rehabilitation services.

Methods

As part of a larger feasibility study conducted in North Wales, a Discrete Choice Experiment (DCE) questionnaire was administered to patients who had recently experienced hip fracture and were randomised to receive either usual care or a newly developed multidisciplinary rehabilitation intervention. DCEs are a stated preference technique for eliciting individuals' preferences about goods and services. Individuals make trade-offs between choices, and these trade-offs provide information about individuals' relative preferences for different attributes of a service. Findings from a systematic review, patient focus groups and healthcare professional survey conducted to develop the intervention were used to design the attributes and levels for the DCE questionnaire. Participants who consented to take part in the larger feasibility study were asked to complete the DCE questionnaire at their three-month follow up.

Findings

Thirty-two participants (65%) out of a possible forty-nine completed the DCE questionnaire at follow up. Twenty-two (69%) participants were female, and ten participants (31%) were male. The pilot DCE demonstrated the method could be used with participants with a mean age of 79 years (SD 7.54). Logit regression in STATA indicated participants preferences for increased time with the healthcare professional who delivers rehabilitation (β -coefficient = 0.005, 95% CI: 0.000 to 0.010), and a physiotherapy/occupational therapy assistant to deliver the rehabilitation sessions (β -coefficient = -0.596, 95% CI: -0.862 to -0.445).

Interpretation

Previous DCEs conducted in Australia focused on patient outcomes such as pain and dose of therapy. The design of the DCE described here used attributes associated with service configuration, which could have the potential to inform service implementation and assist service design, incorporating the preferences of patients. It could also provide lessons for future DCEs conducted with similar populations.

Funding

This work was funded by the National Institute for Health Research Health Technology Assessment (NIHR HTA) Programme (Funder Reference: 11/33-03).

Contributors

NHW is PI of the larger feasibility study. NHW, JLR and NUD conducted the focus groups and professional survey. NUD and JMC conducted the systematic review. JMC and RTE developed the DCE. JMC designed the DCE questionnaire. JMC and STY conducted data analysis and interpreted the results. JMC wrote the abstract with input from JLR, NUD, NHW, STY and RTE. All authors have seen and approved the final version of the abstract. JMC is not an early career researcher.

Conflicts of interest

JMC, JLR, NUD, NHW, STY and RTE authors report grants from NIHR HTA, during the conduct of the study. STY did not receive any direct salary funding from the NIHR HTA. JMC, RTE and NHW report grants from Public Health Wales, outside the submitted work.

Acknowledgments

This work was funded by the National Institute for Health Research Health Technology Assessment (NIHR HTA) Programme (Funder Reference: 11/33-03) and will be published in full in Health Technology Assessment. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the HTA programme, NIHR, NHS or the Department of Health.

Word count

348 words