



PRIFYSGOL  
**BANGOR**  
UNIVERSITY

## **Second-to-fourth digit ratio (2D:4D) is unrelated to measures of somatic reproductive effort among young men from Cebu, the Philippines**

Georgiev, Alexander V.; Ryan, Calen P.; Gettler, Lee T.; McDade, Thomas W.; Kuzawa, Christopher W.

**American Journal of Physical Anthropology**

DOI:

[10.1002/ajpa.23215](https://doi.org/10.1002/ajpa.23215)

Published: 01/07/2017

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

*Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*

Georgiev, A. V., Ryan, C. P., Gettler, L. T., McDade, T. W., & Kuzawa, C. W. (2017). Second-to-fourth digit ratio (2D:4D) is unrelated to measures of somatic reproductive effort among young men from Cebu, the Philippines. *American Journal of Physical Anthropology*, 163(3), 437-445. <https://doi.org/10.1002/ajpa.23215>

### **Hawliau Cyffredinol / General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Published in: American Journal of Physical Anthropology, DOI: 10.1002/ajpa.23215*

*(Version of record online: 27 March 2017)*

**Second-to-fourth digit ratio (2D:4D) is unrelated to measures of somatic reproductive effort among young men from Cebu, The Philippines**

Alexander V. Georgiev<sup>1\*</sup>, Calen P. Ryan<sup>1</sup>, Lee T. Gettler<sup>2,3</sup>, Thomas W. McDade<sup>1,4</sup> & Christopher W. Kuzawa<sup>1,4</sup>

<sup>1</sup>Department of Anthropology, Northwestern University, Evanston, IL

<sup>2</sup>Department of Anthropology, University of Notre Dame, Notre Dame, IN

<sup>3</sup>The Eck Institute for Global Health, University of Notre Dame, Notre Dame, IN

<sup>4</sup>Institute for Policy Research, Northwestern University, Evanston, IL

**\*Correspondence:** Alexander V. Georgiev, School of Biological Sciences, Bangor University, Deiniol Road, Bangor, Gwynedd, LL57 2UW, UK. Email: [a.georgiev@bangor.ac.uk](mailto:a.georgiev@bangor.ac.uk)

**Abbreviated title:** Digit ratios and male somatic reproductive effort

**Key words:** prenatal androgen exposure, muscularity, adiposity, testosterone

**Grant sponsorship:** This work was supported by the Wenner-Gren Foundation (Grants 7356 and 8186), National Science Foundation (Grants BCS-0542182 and BCS- 0962212), Interdisciplinary Obesity Center (Grant RR20649), and Center for Environmental Health and Susceptibility (Grant ES10126; Project 7-2004-E).

## **ABSTRACT**

**Objectives:** A low second-to-fourth (2D:4D) digit ratio, a retrospective marker of high prenatal androgens, predicts increased investment in costly sexually dimorphic traits in men in some studies, though results are mixed. Here we test the hypothesis that the relationship of low 2D:4D ratios with increased muscularity and decreased adiposity depends on current testosterone (T) levels, such that digit ratio will be a particularly strong predictor of outcomes among men exhibiting a mating-effort-oriented endocrinological profile (high T). We also test the association between 2D:4D and somatic traits independently of T.

**Materials and methods:** We related 2D:4D digit ratios, and their interaction with T, to handgrip strength, lean mass, arm muscle area, and skinfold thickness in a sample of young, childless men (20-22 y) from Cebu, Philippines (N = 623).

**Results:** Digit ratio did not significantly predict men's T-dependent somatic traits. Interactions between 2D:4D and morning T, similarly, did not predict male muscularity or adiposity. Although two of the interactions were significant or marginally significant ( $P < 0.1$ ), after adjusting for multiple testing the evidence in support of our hypothesis was weak.

**Discussion:** We found no evidence that 2D:4D predicted measures of somatic reproductive effort in this sample of young men from Cebu, who as a group could be considered mostly mating-oriented. These relationships were also not contingent upon, or stronger, when considering the moderating effect of concurrent T levels. In this sample, 2D:4D was therefore either a poor proxy of prenatal androgen exposure or prenatal androgens had limited influence on adult somatic outcomes.

## Introduction

Prenatal exposure to testosterone (T) or other androgens is known to have organizational effects on a range of sexually dimorphic traits in vertebrates (Phoenix et al., 1959; Smith et al., 2013). It can influence neural development, as well as the expression of social and sexual behavior in later life (Forger et al., 1996; Thornton et al., 2009; Hines et al., 2015). Increased prenatal androgen exposure has also been linked to greater fetal and postnatal growth in some study systems, particularly among male offspring (Gill et al., 1998; Helle et al., 2012).

Androgens play a role in the processes leading to differentiation of mesenchymal pluripotent stem cells into the myogenic lineage and the inhibition of their differentiation into the adipogenic lineage (Bhasin et al., 2003; Singh et al., 2003; De Gendt and Verhoeven, 2012). Such insights gleaned from *in vitro* studies and knock-out models of mice likely go some ways towards explaining the generally greater muscle mass and lower adiposity of men compared to women – differences that are initiated *in utero*, given known sex differences in weight and body composition at birth (Catalano et al., 1995). Indeed, experimental and comparative data suggest that changes in the role of prenatal testosterone exposure on growth might be an important mechanism underlying the rapid evolution of pronounced sexual size dimorphism in some lineages (Cox et al., 2009). Intra-specifically, therefore, differential exposure to prenatal androgens may lead to inter-individual variation in male growth and physiology with possible downstream effects on reproductive performance and success.

Because of the challenges of studying the effects of prenatal androgen in humans, the ratio of the second to the fourth digit of the hand (2D:4D ratio) has been widely used as a non-invasive, retrospective marker of intra-uterine androgen exposure (Manning et al., 1998; McIntyre, 2006; Voracek and Loibl, 2009; Voracek, 2011). There is both direct and indirect

evidence from animal model experiments and human studies that, under the influence of prenatal androgens, the embryonic fourth digit grows longer than the second digit, thus resulting in a lower 2D:4D ratio (Brown et al., 2002; Okten et al., 2002; Lutchmaya et al., 2004; Berenbaum et al., 2009; Manning, 2011; Zheng and Cohn, 2011; Rivas et al., 2014). Consistent with these effects, men on average tend to have lower 2D:4D ratios than women in most human populations (Peters et al., 2002; Hönekopp and Watson, 2010). Additional support for the role of prenatal androgen exposure as an influence on the 2D:4D ratio comes from studies of opposite-sex dizygotic twins (van Anders et al., 2006), birth order and the sex of older siblings (Saino et al., 2006), and individuals with congenital conditions affecting pre-natal androgen exposure, such as congenital adrenal hyperplasia (Brown et al., 2002; Okten et al., 2002; Rivas et al., 2014) and complete androgen insensitivity syndrome (Berenbaum et al., 2009). Finally, a direct measure of prenatal hormones via amniocentesis showed that infants that experienced higher T to estrogen ratios in utero had lower 2D:4D ratios at 2 years of age (Lutchmaya et al., 2004). Although these results generally align with the premise that 2D:4D reflects prenatal androgen exposure, inconsistencies across studies have fueled continuing skepticism about the ratio and its interpretation (Berenbaum et al., 2009; Constantinescu and Hines, 2012; Hines et al., 2015). For instance, in at least one population women have lower 2D:4D ratios than men (Apicella et al., 2016). Some studies of individuals with congenital conditions and/or direct measures of prenatal or perinatal hormones also do not provide unequivocal support for the prediction that higher prenatal androgen exposure leads to lower 2D:4D ratios (Buck, 2003; Hickey et al., 2010; Ventura et al., 2013; Hollier et al., 2015).

Despite this lack of consensus on the utility of the 2D:4D ratio as a marker of inter-individual differences in prenatal androgen exposure, many studies have documented

biologically meaningful relationships between 2D:4D ratios and a range of behavioral (Coco, 2013; Butovskaya et al., 2015; Wlodarski et al., 2015; Del Giudice and Angeleri, 2016), endocrinological (Kilduff et al., 2013; Crewther et al., 2015) and reproductive outcomes (Manning et al., 2000; Honekopp et al., 2006; Manning et al., 2009; Klimek et al., 2014), which are interpreted as evidence for effects of prenatal androgen action on longer-term outcomes (but see Putz et al., 2004; Hönekopp et al., 2007).

In terms of male traits related to reproductive effort, lower 2D:4D has been proposed to be a correlate of male competitiveness, fertility and attractiveness, in part through the long-term effects of prenatal androgen exposure on sexually selected male traits that underlie athletic performance, such as strength and endurance (Manning and Taylor, 2001). With some exceptions (Gallup et al., 2007; Voracek et al., 2010; Folland et al., 2012; Muller et al., 2013; Ranson et al., 2015) studies have found that men with lower 2D:4D ratios are stronger, more muscular, and larger (Fink et al., 2006; Hone and McCullough, 2012; Zhao et al., 2012; Halil et al., 2013; Klimek et al., 2014). They also exhibit superior performance in competitive endurance sports and tests of overall physical fitness (Manning and Taylor, 2001; Hönekopp et al., 2006; Bennett et al., 2010; Tamiya et al., 2012; Ranson et al., 2015), and have higher maximal lung oxygen uptake (Hill et al., 2012). A meta-analysis of more than 2500 participants from 24 populations concluded that a lower 2D:4D ratio generally predicted numerous measures of athletic prowess, although findings were relatively modest and also heterogeneous across studies (Hönekopp and Schuster, 2010).

According to the organizational – activational hypothesis, the intrauterine effects of T are presumed to permanently alter (organize) tissues during development, which then experience the activational effects of circulating T, production of which begins at puberty (Phoenix, 2009;

Wallen, 2009). While the scope of the organizational – activational hypothesis was initially restricted specifically to the role of T in organizing tissues related to adult mating behavior (Phoenix et al., 1959), this framework has been extended to account for sex differences in overall phenotypic development (Arnold, 2009). Following from this, one issue that has rarely been considered in the digit ratio literature is the possible contingency of relationships between digit ratios and T-dependent somatic traits on the concurrent activating effect of T. In humans, testosterone production has been shown to decline markedly as adult males shift from mating to pairbonding (Mazur and Michalek, 1998; Burnham et al., 2003) and parenting (Gray et al., 2002; Gettler et al., 2011; Gray et al., 2016). Although functional explanations for these changes in T primarily focus on the hormone’s behavioral role, a similar logic could apply to somatic mating effort. Maintaining costly androgen-dependent somatic traits (e.g., muscularity and strength) that may affect male reproductive success, either through providing advantages in male-male competition or via female mate choice (Puts, 2010; Apicella, 2014), would be particularly important while men are engaging in mating (as opposed to paternal) effort. It has been proposed that high T in men of reproductive age is key to maintaining muscle and strength, which is achieved in part by diverting energy away from maintenance functions like immunity and energy reserves (Bribiescas, 2001; Kuzawa et al., 2016; but see Alvarado et al., 2015). If prenatal androgen exposure affects somatic traits that influence mating success, these relationships should be especially salient among men who have high T and are invested in mating effort. Whether this is the case is presently unclear, as analyses of 2D:4D ratios and somatic traits have not considered the role of study participants’ T in conjunction with life history stage.

Here we examine the relationship between adult 2D:4D ratios and measures of investment in somatic mating effort (lean mass, grip strength, and arm muscle area) and energy stores (adiposity) in a large, prospectively recruited cohort of men (20-22 years old) living in Cebu, The Philippines. Previous cross-sectional analyses conducted on these men at the same age have shown that T levels were only significantly lower among fathers and that single and pair-bonded age mates had similarly high T (Kuzawa et al., 2009). We thus focus our analyses on a subset of men who have maintained higher T levels throughout their adult lives (i.e. have not become fathers yet). We expect a strong relationship between somatic traits and 2D:4D among men, because their T has likely been consistently high, on average, in support of traits associated with mating effort. To further assess the role of T in affecting the possible relationships between digit ratios and somatic traits, we also tested for interactions between morning T levels and digit ratios, with the prediction that men with high T and low 2D:4D would exhibit greater grip strength, arm muscle area and lean mass, and lower adiposity, in comparison to men with low T and high 2D:4D. We focused in particular on waking T levels, which are higher than measures from the rest of the day (Kuzawa et al., 2016) and predictive of muscularity among the physically active men in this population (Gettler et al., 2010).

## **Materials and Methods**

### **Study site and subjects**

Data come from the Cebu Longitudinal Health and Nutrition Survey (CLHNS), a population-based birth cohort study of mothers and their infants born in 1983-84 from randomly selected urban and rural neighborhoods in Metropolitan Cebu, Philippines (Adair et al., 2011). This research was conducted under conditions of informed consent with human subjects



clearance from the Institutional Review Boards of the University of North Carolina, Chapel Hill and Northwestern University.

### **Data collection**

Adult somatic measures (height, grip strength, arm muscle area, lean mass, and skinfold thickness) were measured in 2005 using standard procedures, as described in detail elsewhere (Gettler et al., 2010). Briefly, grip strength was measured to the nearest kg in triplicate with a dynamometer, with the mean used in analyses. Arm muscle area was estimated from arm circumference (measured with a flexible tape) and triceps skinfolds (measured in triplicate with a skinfold caliper), adjusted for the estimated area of the humerus. Lean mass was calculated from estimated % body composition calculated from triceps, suprailiac, and subscapular skinfold thickness using a body composition predictive formula validated for Asian populations. Information on socioeconomic status, relationship status (co-habitation), parenthood and other factors were collected during in-home interviews (Kuzawa et al., 2009). Digit ratios were measured directly in triplicate with calipers in 2009. For each digit, we calculated the mean of three repeat measures of the 2D:4D ratio for each hand. Intraclass correlation coefficients for the repeated digit measures were high for all digits measured on both hands ( $>0.99$ ) but we caution that these measures were undertaken in close succession by the same observer and are thus not statistically independent.

Testosterone was measured from saliva samples collected by study participants immediately upon waking and assayed subsequently at the Laboratory for Human Biology at Northwestern University (Kuzawa et al., 2009). T concentrations were determined in duplicate using an enzyme immunoassay protocol developed and validated for use with saliva samples (Salimetrics #1-2402, State College, PA). The inter-assay coefficients of variation were 11.5%

and 13.7 % for high and low controls, respectively (Gettler et al., 2010). T values were adjusted for time of saliva collection prior to analysis (Kuzawa et al. 2010).

We limit analyses to the sample of men who did not have any children (to rule out those with reduced T levels due to parenthood), who had a complete set of morphological and hormone measures collected in 2005, and who were not shift workers due to unusual wake/sleep patterns that affected T production (Lee et al., 2014). In addition, we excluded 4 and 9 outliers from the measures of the L and R-hand 2D:4D ratios, respectively (values > 3SD of the mean) retaining a final analysis sample of 623 men.

### **Data analysis**

To assess the relationship between digit ratios and somatic traits of interest (grip strength of the stronger hand, arm muscle area, lean mass, and skinfold thickness) we first report results from simple correlational tests (without controlling for potentially confounding factors), to allow a direct comparison with previously published studies. Second, for both hands, we constructed two models for each trait: first, using only 2D:4D as a predictor, and second, including the interaction between 2D:4D and morning T levels. Previous studies have examined this ratio both for the left and the right hand, using both a continuous ratio and a binary metric (low/masculinized v. high/feminized; low being  $< 1.0$ ; high  $\geq 1.0$ ). While we focused our analyses and interpretation on continuously measured digit ratios, for the purposes of comparison with the existing literature we also report findings from models that consider a binary measure of high/low 2D:4D. To address the issue of multiple testing we first report unadjusted P-values for explorative purposes (because some previous studies only examined one of these 4 alternative ways of assessing digit masculinization) and then consider the adjusted critical P-value after correcting for running 32 separate models ( $P = 0.002$ ). All models controlled for additional

variables that could confound associations between digit ratio and somatic outcomes, including adult household income, age, and height, and binary measures of whether the men engaged regularly in physically demanding work, basketball playing, and weightlifting (see Gettler et al., 2010). All continuous variables were converted to z-scores prior to analysis to allow direct unitless comparison of coefficients across different outcomes. Testosterone values were log-transformed first and adjusted for time of saliva collection, after which the residuals were converted to z-scores. Digit ratio values were modeled as residuals from models regressing measured 2D:4D ratio on observer identity to account for small but systematic differences in measurement across 5 observers. In presenting descriptive statistics and plots, and carrying out simple correlational tests, we use 2D:4D ratios adjusted for observer. Tests are two-tailed throughout with  $\alpha = 0.05$  (and  $p < 0.1$  considered a statistical trend), unless noted otherwise, and performed in Stata 13.1 (Stata Corporation, College Station, TX). Power analysis was performed in G\*Power 3.1 (Faul et al., 2009).

## RESULTS

In this sample of young adult men (Table 1), 2D:4D digit ratios were slightly lower on the left hand (L2D:4D: mean  $\pm$  SE =  $0.961 \pm 0.001$ ) than on the right (R2D:4D:  $0.965 \pm 0.001$ ;  $t = -4.59$ ,  $N = 623$ ,  $P < 0.0001$ ) with 56.98% of men exhibiting a lower (relatively more masculinized) 2D:4D ratio on their L hand, compared to the right. The 2D:4D ratio in the sample was categorized as low/masculine ( $< 1.00$ ) for 91.3% of the men based upon the L hand and 90.4% for the R hand.

==== TABLE 1 =====

Men with greater adiposity (sum of skinfolds) tended to have higher R2D:4D (all correlational tests performed on log-transformed values;  $r = 0.077$ ,  $N = 623$ ,  $P = 0.056$ ) but this

relationship was not significant for the left hand ( $r = 0.004$ ,  $P = 0.921$ ). No significant relationships were found between 2D:4D and grip strength (L:  $r = -0.044$ ,  $P = 0.270$ ; R:  $r = -0.035$ ,  $P = 0.384$ ), arm muscle area (L:  $r = -0.023$ ,  $P = 0.561$ ; R:  $r = -0.055$ ,  $P = 0.169$ ), or lean mass (L:  $r = -0.010$ ,  $P = 0.801$ ; R:  $r = -0.008$ ,  $P = 0.849$ ) (Fig. 1).

In multiple regression models accounting for potentially confounding factors (height, age, regular involvement in physically demanding activity and sports, and household income) none of the four somatic measures we tested (grip strength, arm muscle area, lean mass, and sum of skinfolds) were significantly associated with 2D:4D on either hand. The only relationship that approached statistical significance (before correcting for multiple testing) was the positive association between R2D:4D and adiposity (sum of skinfolds;  $P = 0.078$ , Table 2). There was no evidence for non-linear relationships between 2D:4D ratios and somatic traits (Fig. 1).

===== FIGURE 1 =====

In simple correlation tests, waking T levels were not associated with 2D:4D on either hand (L:  $r = -0.014$ ,  $P = 0.721$ ; R:  $r = -0.045$ ,  $P = 0.264$ ). To evaluate the potential role of T as a moderating influence on relationships between 2D:4D ratios and somatic traits we included a 2D:4D x waking T interaction term. One of the multiple regression models showed a significant positive interaction between R2D:4D and morning T in predicting grip strength ( $P = 0.043$ ). In a separate model, R2D:4D and morning T interacted positively and almost significantly ( $P = 0.077$ ) to predict arm muscle area (Table 2). With the positive slope steepest among men with the highest T, these interactions suggested that men with higher T and more feminized (higher) 2D:4D tended to have greater grip strength and more arm musculature. Re-running the models described above with categorical measures of left and right 2D:4D (high vs. low) yielded consistently non-significant findings (results not shown). Neither the significant interaction ( $P =$

0.043) or the two marginally significant main effects ( $P < 0.1$ ) noted above, remained significant after adjusting for multiple testing (32 separate models), which lowered the significance threshold to  $P = 0.002$ .

==== TABLE 2 ====

## **DISCUSSION**

The aim of this study was to examine the relationship between 2D:4D ratios, a putative marker of prenatal androgen exposure, and somatic traits associated with mating effort in a large sample of young Filipino men. We focused our analyses on young adults who had never been fathers and who were thus likely to have maintained higher average T levels throughout their adult lives (Kuzawa et al., 2009). Although the relationship between 2D:4D and T-dependent somatic traits would be expected to be strongest in these “mating-oriented” men, we found little evidence that digit ratios predicted somatic traits, with or without considering the potentially moderating role of concurrent T levels. These findings suggest that digit ratio is a poor predictor of adult somatic mating effort in this young adult sample of men from Cebu.

Although previous studies examining the relationship between digit ratios and adult somatic traits in humans have produced conflicting results, a meta-analysis of published findings found support for a relationship between low 2D:4D ratios and athletic prowess (including grip strength) in both men and women (Hönekopp and Schuster, 2010). The putative effect of prenatal androgen exposure on athleticism, however, appears inconsistent across studies, and with low to moderate effect sizes (Hönekopp and Schuster, 2010). Studies evaluating the relationship between 2D:4D and adult somatic development (specifically musculature and

adiposity) are less common and have thus far not been the subject of meta-analytical review. In older adults 2D:4D has been shown to be negatively correlated with calf circumference and skeletal muscle mass index in both men and women (Halil et al., 2013). Men and women with higher 2D:4D ratios also have more feminine body fat distribution, although there does not appear to be a relationship between total fatness and digit ratios (Ertuğrul, 2013). A lack of a significant relationship between 2D:4D and either fat free mass or adiposity was also reported in at least one other study (Muller et al., 2013). In the context of these findings, our study contributes to increasing awareness of the low predictive power of adult 2D:4D ratios, as a proxy of prenatal androgen exposure (Berenbaum et al., 2009; Constantinescu and Hines, 2012; Hines et al., 2015).

Given that we focused analyses on a subgroup of young men who were most likely to exhibit peak androgen-dependent somatic trait development, and also examined the potential interaction between concurrent T levels and digit ratios in predicting these traits, our lack of significant findings is particularly striking. Not only were these relationships largely not statistically significant but the single significant effect before accounting for multiple testing (R2D:4D x AMT, Table 2) suggested a relationship between digit ratios and grip strength that was in the opposite direction to the one predicted – i.e. men with higher T levels but with more feminized (higher) digit ratios (rather than with more masculinized/lower ratios) had greater grip strength. Nevertheless, the fact that two of the interactions involving morning T levels were the closest to being significant of all predictors tested here (Table 2) suggests that future work examining the role of prenatal and adult androgen action in concert might provide more insight into male somatic development than studies focusing on either in isolation.

Although, it is possible in theory that the lack of significant findings in our study reflects low power, our sample size suggests otherwise. With 623 men in our dataset, our study was among the larger studies of the adult correlates of digit ratios, and we estimate that we were powered to detect a relationship between 2D:4D and somatic traits as low as  $|r| = 0.14$ . This value is lower than the average effect size across studies included in a meta-analysis of 2D:4D and athletic prowess (including grip strength) with  $r \approx -0.26$  (Hönekopp and Schuster, 2010). For comparison, in our dataset the relationship between digit ratios and grip strength was  $r \approx -0.04$ .

Set against the larger body of work on this topic our findings are not particularly unexpected. The literature on the putative relationship between 2D:4D measures and a range of adult somatic and behavioral phenotypes is characterized by inconsistent results and frequent null findings. The use of many operational definitions of 2D:4D ratios (left hand, right hand, average of the two hands, continuous or categorical), the frequent lack of multiple-testing corrections (especially when more than one type of operational definition is used), and the inherently noisy nature of this variable (given that most studies measure ratios based on finger length externally rather than the finger bones underneath) contribute to uncertainty in interpreting findings and are likely also compounded by reporting bias (Apicella et al., 2015). Our findings similarly suggest that caution is needed when using digit ratios as proxies of inter-individual differences in prenatal androgen exposure. Alternatively, if 2D:4D ratios do reflect such exposures well, the role of these early influences in shaping investment in somatic reproductive effort in adult men may be limited when placed in the context of other developmental factors, including the early post-natal period when there is evidence for continued organizational effects of T on somatic development (Kuzawa et al., 2010; Kiviranta et al., 2016).

Longitudinal research on pre- and early post-natal organizational effects of androgens on adult reproductive traits should incorporate direct hormonal measures to help distinguish between these two explanations.

### **Acknowledgements**

We thank the many researchers at the Office of Population Studies, University of San Carlos, Cebu, the Philippines, for their central role in study design and data collection, and the Filipino participants, who generously provided their time for this study. Elizabeth Quinn, Katy Sharrock, Jeffrey Huang, Iram Azam, Divya Mallampati, Brian Dubin, and Laura Rogers helped with lab work. We thank the editors and two anonymous reviewers for their constructive comments on a previous version of this manuscript.

### **Literature cited**

- Adair LS, Popkin BM, Akin JS, Guilkey DK, Gultiano S, Borja J, Perez L, Kuzawa CW, McDade T, Hindin MJ. 2011. Cohort profile: the Cebu longitudinal health and nutrition survey. *International Journal of Epidemiology* 40:619–625.
- Alvarado LC, Muller MN, Emery Thompson M, Klimek M, Nenko I, Jasienska G. 2015. The Paternal Provisioning Hypothesis: effects of workload and testosterone production on men's musculature. *Behaviour* 158:19–35.
- Apicella CL, Carré JM, Dreber A. 2015. Testosterone and economic risk taking: a review. *Adaptive Human Behavior and Physiology* 1:358–385.
- Apicella CL, Tobolsky VA, Marlowe FW, Miller KW. 2016. Hadza hunter-gatherer men do not have more masculine digit ratios (2D:4D). *Behaviour* 159:223–232.
- Apicella CL. 2014. Upper-body strength predicts hunting reputation and reproductive success in Hadza hunter-gatherers. *Evolution and Human Behavior* 35:508–518.
- Arnold AP. 2009. The organizational–activational hypothesis as the foundation for a unified theory of sexual differentiation of all mammalian tissues. *Hormones and Behavior* 55:570–578.



- Bennett M, Manning JT, Cook CJ, Kilduff LP. 2010. Digit ratio (2D:4D) and performance in elite rugby players. *Journal of Sports Sciences* 28:1415–1421.
- Berenbaum SA, Bryk KK, Nowak N, Quigley CA, Moffat S. 2009. Fingers as a Marker of Prenatal Androgen Exposure. *Endocrinology* 150:5119–5124.
- Bhasin S, Taylor WE, Singh R, Artaza J, Sinha-Hikim I, Jasuja R, Choi H, Gonzalez-Cadavid NF. 2003. The mechanisms of androgen effects on body composition: mesenchymal pluripotent cell as the target of androgen action. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences* 58:M1103–10.
- Bribiescas RG. 2001. Reproductive ecology and life history of the human male. *Yearbook of Physical Anthropology* 44:148–176.
- Brown WM, Hines M, Fane BA, Breedlove SM. 2002. Masculinized Finger Length Patterns in Human Males and Females with Congenital Adrenal Hyperplasia. *Hormones and Behavior* 42:380–386.
- Buck JJ. 2003. In-utero androgen exposure and 2nd to 4th digit length ratio--comparisons between healthy controls and females with classical congenital adrenal hyperplasia. *Human Reproduction* 18:976–979.
- Burnham TC, Chapman JF, Gray PB, McIntyre MH, Lipson SF, Ellison PT. 2003. Men in committed, romantic relationships have lower testosterone. *Hormones and Behavior* 44:119–122.
- Butovskaya M, Burkova V, Karelin D, Fink B. 2015. Digit ratio (2D:4D), aggression, and dominance in the Hadza and the Datoga of Tanzania. *Am J Hum Biol* 27:620–627.
- Catalano PM, Drago NM, Amini SB. 1995. Factors affecting fetal growth and body composition. *Am J Obstet Gynecol* 172:1459–1463.
- Coco M. 2013. The second-to-fourth digit ratio correlates with aggressive behavior in professional soccer players. *Mol Med Report* 7:1733–1738.
- Constantinescu M, Hines M. 2012. Relating Prenatal Testosterone Exposure to Postnatal Behavior in Typically Developing Children: Methods and Findings. *Child Dev Perspect*:n/a–n/a.
- Cox RM, Stenquist DS, Calsbeek R. 2009. Testosterone, growth and the evolution of sexual size dimorphism. *Journal of Evolutionary Biology* 22:1586–1598.
- Crewther B, Cook C, Kilduff L, Manning J. 2015. Digit ratio (2D:4D) and salivary testosterone, oestradiol and cortisol levels under challenge: Evidence for prenatal effects on adult endocrine responses. *Early Human Development* 91:451–456.
- De Gendt K, Verhoeven G. 2012. Tissue- and cell-specific functions of the androgen receptor revealed through conditional knockout models in mice. *Molecular and Cellular*

- Endocrinology 352:13–25.
- Del Giudice M, Angeleri R. 2016. Digit Ratio (2D:4D) and Attachment Styles in Middle Childhood: Indirect Evidence for an Organizational Effect of Sex Hormones. *Adaptive Human Behavior and Physiology* 2:1–10.
- Ertuğrul B. 2013. Sexually dimorphic human body fat distribution and second-to-fourth digit ratio. *Eurasian Journal of Anthropology* 3:56–62.
- Faul F, Erdfelder E, Buchner A, Lang A-G. 2009. Statistical power analyses using G\*Power 3.1: Tests for correlation and regression analyses. *Behavior Research Methods* 41:1149–1160.
- Fink B, Thanzami V, Seydel H, Manning JT. 2006. Digit ratio and hand-grip strength in German and Mizos men: Cross-cultural evidence for an organizing effect of prenatal testosterone on strength. *Am J Hum Biol* 18:776–782.
- Folland JP, Mc Cauley TM, Phipers C, Hanson B, Mastana SS. 2012. Relationship of 2D:4D finger ratio with muscle strength, testosterone, and androgen receptor CAG repeat genotype. *Behaviour* 148:81–87.
- Forger NG, Galef BG, Clark MM. 1996. Intrauterine position affects motoneuron number and muscle size in a sexually dimorphic neuromuscular system. *Brain Research* 735:119–124.
- Gallup AC, White DD, Gallup GG Jr. 2007. Handgrip strength predicts sexual behavior, body morphology, and aggression in male college students. *Evolution and Human Behavior* 28:423–429.
- Gettler LT, Agustin SS, Kuzawa CW. 2010. Testosterone, physical activity, and somatic outcomes among Filipino males. *Yearbook of Physical Anthropology* 142:590–599.
- Gettler LT, McDade TW, Feranil AB, Kuzawa CW. 2011. Longitudinal evidence that fatherhood decreases testosterone in human males. *Proceedings of the National Academy of Sciences* 108:16194–16199.
- Gill JW, Hosking BJ, Egan AR. 1998. Prenatal programming of mammalian growth—a review of the role of steroids. *Livestock production science*.
- Gray PB, Kahlenberg SM, Barrett ES, Lipson SF, Ellison PT. 2002. Marriage and fatherhood are associated with lower testosterone in males. *Evolution and Human Behavior* 23:193–201.
- Gray PB, McHale TS, Carré JM. 2016. A review of human male field studies of hormones and behavioral reproductive effort. *Hormones and Behavior*:1–16.
- Halil M, Gurel EI, Kuyumcu ME, Karaismailoglu S, Yesil Y, Ozturk ZA, Yavuz BB, Cankurtaran M, Ariogul S. 2013. Digit (2D:4D) ratio is associated with muscle mass (MM) and strength (MS) in older adults: Possible effect of in utero androgen exposure. *Archives of Gerontology and Geriatrics* 56:358–363.

- Helle S, Laaksonen T, Huitu O. 2012. Sex-specific offspring growth according to maternal testosterone, corticosterone, and glucose levels. *Behavioral Ecology* 24:205–212.
- Hickey M, Doherty DA, Hart R, Norman RJ, Mattes E, Atkinson HC, Sloboda DM. 2010. Maternal and umbilical cord androgen concentrations do not predict digit ratio (2D:4D) in girls: A prospective cohort study. *Psychoneuroendocrinology* 35:1235–1244.
- Hill R, Simpson B, Manning J, Kilduff L. 2012. Right–left digit ratio (2D:4D) and maximal oxygen uptake. *Journal of Sports Sciences* 30:129–134.
- Hines M, Constantinescu M, Spencer D. 2015. Early androgen exposure and human gender development. *Biol Sex Differ* 6:3.
- Hollier LP, Keelan JA, Jamnadass ESL, Maybery MT, Hickey M, Whitehouse AJO. 2015. Adult digit ratio (2D:4D) is not related to umbilical cord androgen or estrogen concentrations, their ratios or net bioactivity. *Early Human Development* 91:111–117.
- Hone LSE, McCullough ME. 2012. 2D:4D ratios predict hand grip strength (but not hand grip endurance) in men (but not in women). *Evolution and Human Behavior* 33:780–789.
- Honekopp J, Voracek M, Manning J. 2006. 2nd to 4th digit ratio (2D:4D) and number of sex partners: Evidence for effects of prenatal testosterone in men. *Psychoneuroendocrinology* 31:30–37.
- Hönekopp J, Bartholdt L, Beier L, Liebert A. 2007. Second to fourth digit length ratio (2D:4D) and adult sex hormone levels: New data and a meta-analytic review. *Psychoneuroendocrinology* 32:313–321.
- Hönekopp J, Schuster M. 2010. A meta-analysis on 2D:4D and athletic prowess: Substantial relationships but neither hand out-predicts the other. *PERSONALITY AND INDIVIDUAL DIFFERENCES* 48:4–10.
- Hönekopp J, T Manning J, Müller C. 2006. Digit ratio (2D:4D) and physical fitness in males and females: Evidence for effects of prenatal androgens on sexually selected traits. *Hormones and Behavior* 49:545–549.
- Hönekopp J, Watson S. 2010. Meta-analysis of digit ratio 2D:4D shows greater sex difference in the right hand. *Am J Hum Biol* 22:619–630.
- Kilduff L, Cook CJ, Bennett M, Crewther B, Bracken RM, Manning J. 2013. Right–left digit ratio (2D:4D) predicts free testosterone levels associated with a physical challenge. *Journal of Sports Sciences* 31:677–683.
- Kiviranta P, Kuiri-Hanninen T, Saari A, Lamidi ML, Dunkel L, Sankilampi U. 2016. Transient Postnatal Gonadal Activation and Growth Velocity in Infancy. *Pediatrics* 138:e20153561–e20153561.
- Klimek M, Galbarczyk A, Nenko I, Alvarado LC, Jasienska G. 2014. Digit ratio (2D:4D) as an

- indicator of body size, testosterone concentration and number of children in human males. *Ann Hum Biol* 41:518–523.
- Kuzawa CW, Georgiev AV, McDade TW, Bechayda SA, Gettler LT. 2016. Is there a testosterone awakening response in humans? *Adaptive Human Behavior and Physiology* 2:166–183.
- Kuzawa CW, Gettler LT, Muller MN, McDade TW, Feranil AB. 2009. Fatherhood, pairbonding and testosterone in the Philippines. *Hormones and Behavior* 56:429–435.
- Kuzawa CW, McDade TW, Adair LS, Lee N. 2010. Rapid weight gain after birth predicts life history and reproductive strategy in Filipino males. *Proceedings of the National Academy of Sciences* 107:16800–16805.
- Lee J, Fried R, Thayer Z, Kuzawa CW. 2014. Preterm delivery as a predictor of diurnal cortisol profiles in adulthood: Evidence from Cebu, Philippines. *Am J Hum Biol* 26:598–602.
- Lutchmaya S, Baron-Cohen S, Raggatt P, Knickmeyer R, Manning JT. 2004. 2nd to 4th digit ratios, fetal testosterone and estradiol. *Early Human Development* 77:23–28.
- Manning J, Barley L, Walton J, Lewis-Jones D, Trivers R, Singh D, Thornhill R, Rohde P, Bereczkei T, Henzi P, Soler M, Szwed A. 2000. The 2nd:4th digit ratio, sexual dimorphism, population differences, and reproductive success: evidence for sexually antagonistic genes? *Evolution and Human Behavior* 21:163–183.
- Manning JT, Henzi P, Venkatramana P, Martin S, Singh D. 2009. Second to fourth digit ratio: ethnic differences and family size in English, Indian and South African populations. *Ann Hum Biol* 30:579–588.
- Manning JT, Scutt D, Wilson J, Lewis-Jones DI. 1998. The ratio of 2nd to 4th digit length: a predictor of sperm numbers and concentrations of testosterone, luteinizing hormone and oestrogen. *Hum Reprod* 13:3000–3004.
- Manning JT, Taylor RP. 2001. Second to fourth digit ratio and male ability in sport: implications for sexual selection in humans. *Evolution and Human Behavior* 22:61–69.
- Manning JT. 2011. Resolving the role of prenatal sex steroids in the development of digit ratio. *Proceedings of the National Academy of Sciences* 108:16143–16144.
- Mazur A, Michalek J. 1998. Marriage, Divorce, and Male Testosterone. *Social Forces* 77:315–330.
- McIntyre MH. 2006. The use of digit ratios as markers for perinatal androgen action. *Reprod Biol Endocrinol* 4:10–9.
- Muller DC, Manning JT, Hopper JL, English DR, Giles GG, Severi G. 2013. No strong association between second to fourth digit ratio (2D:4D) and adult anthropometric measures with emphasis on adiposity. *Ann Hum Biol* 40:201–204.

- Okten A, Kalyoncu M, Yariş N. 2002. The ratio of second- and fourth-digit lengths and congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *Early Human Development* 70:47–54.
- Peters M, Mackenzie K, Bryden P. 2002. Finger length and distal finger extent patterns in humans. *Behaviour* 117:209–217.
- Phoenix CH, Goy RW, Gerall AA, Young WC. 1959. Organizing action of prenatally administered testosterone propionate on the tissues mediating mating behavior in the female guinea pig. *Endocrinology* 65:369–382.
- Phoenix CH. 2009. Organizing action of prenatally administered testosterone propionate on the tissues mediating mating behavior in the female guinea pig. *Hormones and Behavior* 55:566.
- Puts DA. 2010. Beauty and the beast: mechanisms of sexual selection in humans. *Evolution and Human Behavior* 31:157–175.
- Putz DA, Gaulin SJC, Sporter RJ, McBurney DH. 2004. Sex hormones and finger length: What does 2D:4D indicate? *Evolution and Human Behavior* 25:182–199.
- Ranson R, Stratton G, Taylor SR. 2015. Digit ratio (2D:4D) and physical fitness (Eurofit test battery) in school children. *Early Human Development* 91:327–331.
- Rivas MP, Moreira LMA, Santo LDE, Marques ACSS, El-Hani CN, Toralles MBP. 2014. New studies of second and fourth digit ratio as a morphogenetic trait in subjects with congenital adrenal hyperplasia. *Am J Hum Biol* 26:559–561.
- Saino N, Leoni B, Romano M. 2006. Human digit ratios depend on birth order and sex of older siblings and predict maternal fecundity. *Behav Ecol Sociobiol* 60:34–45.
- Singh R, Artaza JN, Taylor WE, Gonzalez-Cadavid NF, Bhasin S. 2003. Androgens Stimulate Myogenic Differentiation and Inhibit Adipogenesis in C3H 10T1/2 Pluripotent Cells through an Androgen Receptor-Mediated Pathway. *Endocrinology* 144:5081–5088.
- Smith AS, Birnie AK, French JA. 2013. Prenatal Androgens Affect Development and Behavior in Primates. In: Clancy KBH, Hinde K, Rutherford JN, editors. *Building Babies: Primate Development in Proximate and Ultimate Perspective*. New York, NY: Springer New York. p 103–131.
- Tamiya R, Lee SY, Ohtake F. 2012. Second to fourth digit ratio and the sporting success of sumo wrestlers. *Evolution and Human Behavior* 33:130–136.
- Thornton J, Zehr JL, Loose MD. 2009. Effects of prenatal androgens on rhesus monkeys: A model system to explore the organizational hypothesis in primates. *Hormones and Behavior* 55:633–644.
- van Anders SM, Vernon PA, Wilbur CJ. 2006. Finger-length ratios show evidence of prenatal hormone-transfer between opposite-sex twins. *Hormones and Behavior* 49:315–319.

- Ventura T, Gomes MC, Pita A, Neto MT, Taylor A. 2013. Digit ratio (2D:4D) in newborns: Influences of prenatal testosterone and maternal environment. *Early Human Development* 89:107–112.
- Voracek M, Loibl LM. 2009. Scientometric analysis and bibliography of digit ratio (2D:4D) research, 1998-2008. *Psychological Reports* 104:922–956.
- Voracek M, PUM U, DRESSLER SG. 2010. Investigating digit ratio (2D:4D) in a highly male-dominated occupation: The case of firefighters. *Scandinavian Journal of Psychology* 51:146–156.
- Voracek M. 2011. Special issue preamble: Digit ratio (2D:4D) and individual differences research. *PERSONALITY AND INDIVIDUAL DIFFERENCES* 51:367–370.
- Wallen K. 2009. The Organizational Hypothesis: Reflections on the 50th anniversary of the publication of Phoenix, Goy, Gerall, and Young (1959). *Hormones and Behavior* 55:561–565.
- Wlodarski R, Manning J, Dunbar RIM. 2015. Stay or stray? Evidence for alternative mating strategy phenotypes in both men and women. *Biology Letters* 11:20140977–20140977.
- Zhao D, Li B, Yu K, Zheng L. 2012. Digit ratio (2D:4D) and handgrip strength in subjects of Han ethnicity: Impact of sex and age. *Behaviour* 149:266–271.
- Zheng Z, Cohn MJ. 2011. Developmental basis of sexually dimorphic digit ratios. *Proceedings of the National Academy of Sciences*.




Figure 1. Relationships between 2D:4D ratio and adult somatic traits. Digit ratios for the L hand are shown as they were on average more masculinized (lower) than those for the R hand (plots for the R hand are similar). Raw data shown. See text for statistical details.

**Table 1 Characteristics of Cebu men in the analytical sample (N = 623). Digit ratios adjusted for collection of data by multiple observers (see text for details).**

<b>Variable</b>	<b>Mean</b>	<b>SD</b>	<b>Min</b>	<b>Max</b>
Age (years)	21.47	0.29	20.79	22.60
L2D:4D (adjusted)	0.961	0.029	0.840	1.057
R2D:4D (adjusted)	0.965	0.027	0.888	1.058
Morning T (pg/ml)	196.90	77.82	26.12	588.43
Grip strength of stronger hand (kg)	72.98	22.03	19.67	129.00
Arm muscle area (cm <sup>2</sup> )	34.87	7.81	14.17	60.41
Lean mass (kg)	46.65	5.78	32.58	68.15
Sum of skinfolds (mm)	37.44	17.78	11.00	137.00
Height (cm)	162.78	5.95	144.50	181.20



**Table 2.** Multiple linear regression models linking adult somatic traits to left and right 2D:4D ratios with and without considering interactions with waking testosterone (N = 623)\*.

<b>GRIP STRENGTH</b>						
<b>Predictor</b>	<b>Predictor statistics</b>			<b>Full model statistics</b>		
	<b>Estimate</b>	<b>SE</b>	<b>P</b>	<b>F</b>	<b>P</b>	<b>Adj. R2</b>
L2D:4D	-0.038	0.038	0.323	F (7, 615) = 2.42	0.019	0.016
L2D:4D x AMT**	0.071	0.044	0.108	F (9, 613) = 2.40	0.011	0.020
R2D:4D	-0.031	0.040	0.436	F (7, 615) = 2.37	0.022	0.015
R2D:4D x AMT	0.097	0.048	0.043	F (9, 613) = 2.52	0.008	0.022

<b>ARM MUSCLE AREA</b>						
<b>Predictor</b>	<b>Predictor statistics</b>			<b>Full model statistics</b>		
	<b>Estimate</b>	<b>SE</b>	<b>P</b>	<b>F</b>	<b>P</b>	<b>Adj. R2</b>
L2D:4D	-0.025	0.037	0.498	F (7, 615) = 7.24	<0.0001	0.065
L2D:4D x AMT	0.050	0.043	0.243	F (9, 613) = 5.91	<0.0001	0.066
R2D:4D	-0.055	0.039	0.155	F (7, 615) = 7.48	<0.0001	0.068
R2D:4D x AMT	0.082	0.046	0.077	F (9, 613) = 6.3	<0.0001	0.071

<b>LEAN MASS</b>						
<b>Predictor</b>	<b>Predictor statistics</b>			<b>Full model statistics</b>		
	<b>Estimate</b>	<b>SE</b>	<b>P</b>	<b>F</b>	<b>P</b>	<b>Adj. R2</b>
L2D:4D	-0.019	0.031	0.529	F (7, 615) = 50.34	<0.0001	0.357
L2D:4D x AMT	-0.003	0.035	0.934	F (9, 613) = 39.94	<0.0001	0.355
R2D:4D	-0.021	0.032	0.515	F (7, 615) = 50.35	<0.0001	0.357
R2D:4D x AMT	0.032	0.038	0.405	F (9, 613) = 39.16	<0.0001	0.356

<b>SUM OF SKINFOLDS</b>						
<b>Predictor</b>	<b>Predictor statistics</b>			<b>Full model statistics</b>		
	<b>Estimate</b>	<b>SE</b>	<b>P</b>	<b>F</b>	<b>P</b>	<b>Adj. R2</b>
L2D:4D	0.008	0.038	0.828	F (7, 615) = 5.18	<0.0001	0.045
L2D:4D x AMT	-0.025	0.044	0.567	F (9, 613) = 4.05	<0.0001	0.042
R2D:4D	0.070	0.039	0.078	F (7, 615) = 5.64	<0.0001	0.050
R2D:4D x AMT	-0.004	0.047	0.934	F (9, 613) = 4.37	<0.0001	0.047

\* All models control for age, height, household income, and engagement in strenuous physical activity, basketball playing and weightlifting (results not shown). 2D:4D entered as residuals of raw 2D:4D on observer ID (See text for details).

\*\* Waking salivary testosterone.