

DR. MARTINA H STIASNY (Orcid ID : 0000-0001-6365-2610)

DR. CATRIONA CLEMMESSEN (Orcid ID : 0000-0003-4102-7341)

Article type : Primary Research Articles

## **Divergent responses of Atlantic cod to ocean acidification and food limitation**

Running title: Acidification affects growth of cod larvae

**Authors: Martina H. Stiasny<sup>1,2</sup>, Michael Sswat<sup>3</sup>, Felix H. Mittermayer<sup>1</sup>, Inger-Britt Falk-Petersen<sup>4</sup>, Nalani K. Schnell<sup>5</sup>, Velmurugu Puvanendran<sup>6</sup>, Atle Mortensen<sup>6</sup>, Thorsten B.H. Reusch<sup>1</sup> and Catriona Clemmesen<sup>1\*</sup>**

<sup>1</sup> GEOMAR Helmholtz Centre for Ocean Research, Evolutionary Ecology of Marine Fishes, Düsternbrooker Weg 20, 24105 Kiel, Germany

<sup>2</sup> Department of Economics, Sustainable Fisheries, University of Kiel, Wilhelm-Seelig-Platz 1, 24118 Kiel, Germany

<sup>3</sup> GEOMAR Helmholtz Centre for Ocean Research, Biological Oceanography, Düsternbrooker Weg 20, 24105 Kiel, Germany

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/gcb.14554

This article is protected by copyright. All rights reserved.

<sup>4</sup> University of Tromsø, Faculty of Biosciences, Fisheries and Economics, Breivika, N-9037 Tromsø, Norway

<sup>5</sup> Institut de Systématique, Évolution, Biodiversité, ISYEB–UMR 7205–CNRS, MNHN, UPMC, EPHE, Muséum national d’Histoire naturelle, Sorbonne Universities, 57 rue Cuvier, CP 30, 75005 Paris, France

<sup>6</sup> Nofima AS, Postboks 6122, NO-9291 Tromsø, Norway

\* Corresponding author: e-mail: clemmesen@geomar.de Tel: +494316004558, Fax: +494316004553

Keywords: early life-stages, energy limitation, food regimes, *Gadus morhua*, gill development, histology, laboratory experiment, lipid content, ossification, RNA/DNA ratios.

Paper type: Primary Research Article

### Abstract

**In order to understand the effect of global change on marine fishes, it is imperative to quantify the effects on fundamental parameters such as survival and growth. Larval survival and recruitment of the Atlantic cod (*Gadus morhua*) was found to be heavily impaired by end-of-century levels of ocean acidification. Here, we analysed larval growth among 35-36 days old surviving larvae, along with organ development and ossification of the skeleton. We combined CO<sub>2</sub>-treatments (ambient: 503  $\mu$ atm, elevated: 1179  $\mu$ atm) with food availability in order to evaluate the effect of energy limitation in addition to the ocean acidification stressor. As expected, larval size (as a proxy for growth) and skeletogenesis were positively affected by high food availability. We found**

significant interactions between acidification and food availability. Larvae fed ad libitum showed little difference in growth and skeletogenesis due to the CO<sub>2</sub> treatment. Larvae under energy limitation were significantly larger and had further developed skeletal structures in the elevated CO<sub>2</sub> treatment compared to the ambient CO<sub>2</sub> treatment. However, the elevated CO<sub>2</sub> group revealed impairments in critically important organs, such as the liver, and had comparatively smaller functional gills indicating a mismatch between size and function. It is therefore likely that individual larvae that had survived acidification treatments, will suffer from impairments later during ontogeny. Our study highlights important allocation trade-off between growth and organ development, which is critically important to interpret acidification effects on early life-stages of fish.

## Introduction

Over the last 150 years anthropogenic activities have led to an increase in atmospheric concentrations of carbon dioxide, which may increase globally to 750-1000 ppm CO<sub>2</sub> by 2100 under current climate projections (i.e. RCP 8.5 (IPCC, 2013)). Significant effects on marine ecosystems are to be expected through changes in ocean acidification levels (direct effects) and increases in temperature via the indirect effects of CO<sub>2</sub> as a potent greenhouse gas (Doney, Fabry, Feely, & Kleypas, 2009; Orr et al., 2005). Many commercially and ecologically important fish populations are progressively subjected to these environmental changes with effects at all biological levels already visible now. Owing to the higher solubility of CO<sub>2</sub> in colder waters an even sharper increase in ocean acidification levels in these high latitude regions is predicted (Steinacher, Joos, Frölicher, Plattner, & Doney, 2009). A thorough understanding of these effects on individuals and fish populations becomes

increasingly important, particularly for species that are commercially exploited and where management may need adjustment under consideration of global change effects (Denman, Christian, Steiner, Pörtner, & Nojiri, 2011; Kjesbu et al., 2014; Lam, Cheung, & Sumaila, 2014).

Studies on the effects of ocean acidification on marine fishes, especially the early life stages that are most vulnerable to ocean acidification, have steadily increased in recent years and revealed different effects. Such a pronounced interspecies variability is possibly due to different experimental setups, different analyzed life stages (often limited to very early life stages, e.g. embryos, non-feeding larvae), and short-term versus long-term effects of the applied treatments. For example, no effect on growth rate was observed for walleye pollack (*Theragra chalcogramma*) (Hurst, Fernandez, & Mathis, 2013), juvenile scup (*Stenotomus chrysops*) (Perry et al., 2015) and the larvae of yellowtail kingfish (*Seriola lalandi*) (Watson et al., 2018). For other species a decrease in growth rate under elevated CO<sub>2</sub> was observed, e.g. in the inland silverside (*Menidia beryllina*) (Baumann, Talmage, & Gobler, 2012), for yellowfin tuna (*Thunnus albacares*) (Bromhead et al., 2015), and Atlantic herring (*Clupea harengus*) (Frommel et al., 2014). However, another study found no effect on Atlantic herring (Sswat, Stiasny, Jutfelt, Riebesell, & Clemmesen, 2018) . Still others showed an increase in growth rate e.g. in orange clown fish (*Amphiprion percula*) (Munday, Donelson, Dixon, & Endo, 2009), in cod (Frommel et al., 2012) in summer flounder (*Paralichthys dentatus*) (Chambers et al., 2013, 2014) and in sand smelt larvae (*Atherina presbyter*) (Silva et al., 2016) under elevated pCO<sub>2</sub> levels.

Most of the studies on the effect of ocean acidification on larval fishes have been using ad libitum or generally high food densities that are unrealistic in nature. Only recently studies have taken the possible interaction of CO<sub>2</sub> level and food supply into consideration, assuming that ad libitum food availability could potentially compensate for any direct effect of CO<sub>2</sub> on survival, growth and development (Gobler, Merlo, Morrell, & Griffith, 2018; Hurst, Laurel, Hanneman, Haines, & Ottmar, 2017; Sswat, Stiasny, Taucher, et al., 2018; Stiasny et al., 2018). Kreiss *et al.* (2015) have hypothesized that ocean acidification increases costs for osmoregulation and baseline energy demands in adult cod. A limited energy uptake due to lower food availability may therefore be required for ocean acidification effects to become experimentally visible. For example, in juvenile blue mussels, food availability outweighed the negative effects of ocean acidification (Thomsen, Casties, Pansch, Körtzinger, & Melzner, 2013) and an increase in feeding rate in corals was able to mitigate the negative effects of ocean acidification on calcification (Towle, Enochs, & Langdon, 2015). We therefore wanted to test ocean acidification effects in combination with food limitation.

Knowledge on the bioenergetics of development under ocean acidification and how organisms prioritize endogenous energy flow toward development and physiological function is mostly unknown (Matson, Yu, Sewell, & Hofmann, 2012). It is increasingly important to understand the underlying mechanisms of CO<sub>2</sub> effects on growth and how energy limitation may lead to important yet poorly understood possible metabolic trade-offs. We have therefore chosen to include RNA/DNA ratio analyses in this study to determine larval condition. RNA/DNA ratios, indicators of protein biosynthesis machinery, can reflect the nutritional condition of larval fish (Clemmesen, 1994; Buckley *et al.*, 2008;) and have been used to reflect changes in energy availability in relation to ocean acidification (Franke & Clemmesen, 2011; Frommel et al., 2014) and other environmental stressors (Chícharo & Chícharo, 2008).

Frommel *et al.* (2012) suggested that the increase in body mass of cod larvae under experimental ocean acidification was mostly caused by an increase in lipid content observed under elevated CO<sub>2</sub> levels and indicated an altered pathway in lipid and protein metabolisms. Díaz-Gil *et al.* (2015) observed shifts in the relative proportions of fatty acids in the tissues of red drum and Silva *et al.* (2016) showed a higher cellular energy consumption rate at elevated CO<sub>2</sub> levels in sand smelt measured by changes in enzyme levels. A possible reason for the sensitivity of these larval stages to ocean acidification might be the functional development of organs such as the gastro-intestinal tract (Zambonino Infante & Cahu, 2001), the gills and the urinary system with their osmoregulatory functions (Varsamos, Nebel, & Charmantier, 2005). Yet, research on the functional development under the influence of ocean acidification in fish larvae is still limited. Hence one of the key objectives of this study was to determine both, growth parameters, including lipid content, as well as developmental organ and skeletal differences. So far, only limited information is available how ocean acidification effects these areas of development (Frommel *et al.* 2012, Pimentel *et al.*, 2014a; Crespel *et al.*, 2017).

We studied how the trade-offs among growth and development under combined ocean acidification and food limitation played out in Atlantic cod (*Gadus morhua*), one of the most important commercial species in the North Atlantic (FAO, 2016). Larvae of the North-East Arctic cod stock showed a doubling of mortality rates after hatching under end-of-the-century acidification levels with marked negative effects when modelling subsequent population recruitment (Koenigstein *et al.*, 2018; Stiasny *et al.*, 2016). Here we assessed whether the surviving larvae (35 to 36 dph) subjected to ambient and elevated CO<sub>2</sub> treatments (503 +/- 89 and 1179 +/- 87  $\mu$ atm *p*CO<sub>2</sub>, 7.9 and 7.61 pH, respectively) were affected in their growth and development, by explicitly considering the nutritional status of the larvae in a fully crossed

design. The chosen level of ocean acidification falls within the possible scenarios of climate change in the Arctic (AMAP, 2013). We compared responses of larvae fed different amounts of prey in a 2 x 2 factorial design to include the effect of energy limitation. Single and interactive effects were estimated on various growth and condition parameters, as well as skeletal and organ development. We discuss the potential trade-offs between growth and development to these different acidification levels and feeding regimes.

### **Materials and methods**

The experiment was performed in 2014 at the Norwegian National Cod Breeding Centre (recently renamed to Centre for Marine Aquaculture), in Tromsø, Norway. In order to obtain eggs and larvae, adult cod were caught alive in the Barents Sea at roughly 70°15'N, 19°00'E in March 2014 and transferred to the Centre in Tromsø. They were kept in large breeding tanks (25 m<sup>3</sup>) at ambient temperature, light and CO<sub>2</sub> conditions. Spawning occurred naturally two weeks after introduction into the breeding tanks and all eggs were collected from the outflow. The fertilized eggs were transferred to incubators with either ambient (503 ± 89 μatm CO<sub>2</sub>) or increased CO<sub>2</sub> (1179 ± 87 μatm) concentrations. Egg developmental stages were examined under a stereomicroscope to ensure that several females participated in these spawning events (Fridgeirsson, 1978). Three different egg developmental stages indicating different spawning times during the night, which cannot be produced by the spawning event of one female, were observed. This result indicates that at least three different females had participated at the spawning event, since a single female will not spawn multiple times within one day (Kjesbu, 1989). At 100% hatch, occurring at 112 degree-days, 11,000 larvae were transferred into each of the twelve 190 L rearing tanks with a constant flow-through of water from two header tanks. Six tanks each were supplied from a header tank with ambient water

and from a header tank containing water with increased CO<sub>2</sub> concentrations. For the egg incubation and the start of the experiment the temperature was set to 6°C and was later, in the larval tanks, raised to 10°C simultaneously in all tanks, while kept at constant light conditions (24h light). Larvae were fed with *Nannochloropsis* (until day 12 days post-hatching (dph)) and *Brachionus* until 24 dph, after which feeding was gradually shifted to *Artemia* nauplii. Prey concentrations fed at each feeding were the same for both treatments, but number of feedings and therefore total amount fed per day differed (Supplement S Table 1). Though prey availability in this set-up is hard to compare to natural conditions, since larvae were pulse-fed with high concentrations of prey, it is likely that the energy limitation, even in the low food treatment, is still above most conditions encountered by larvae in the field.

Larvae in one tank in the ambient CO<sub>2</sub> treatment were abruptly lost over night, due to an unknown factor, resulting in six replicates for the elevated CO<sub>2</sub> treatment and five for the ambient treatment. This study was carried out in strict accordance with the laboratory regulations applicable in Norway. The application was approved by the National Regulatory Committee on the Ethics of Animal Experiments (Permit FOTS id 6382). All conditions and samplings were conducted to minimize suffering.

The values in the ambient CO<sub>2</sub> treatment were  $p\text{CO}_2$ :  $503 \pm 89 \mu\text{atm}$ ;  $\text{pH } 7.9 \pm 0.15$  and  $p\text{CO}_2$ :  $1179 \pm 87 \mu\text{atm}$ ;  $\text{pH } 7.61 \pm 0.03$  levels in the elevated CO<sub>2</sub> rearing tanks were achieved by controlling the pH values in a header tank with pH probes connected to an IKS computer system. The volume of the header tank ensured a thorough mixing and equilibration of CO<sub>2</sub> before the water entered the rearing tank thereby assuring constant conditions in the rearing tanks. The pH and temperature were furthermore manually checked



every day in all the rearing tanks with a separate pH/temperature probe (WTW pH/Cond 340i/3320). Water chemistry, including DIC and alkalinity, was measured weekly. For further details please consult Stiasny *et al.* (2016) or Stiasny *et al.* (2018).

For the growth measurements larvae at 36 dph were sampled alive, euthanized with MS222 (Ethyl 3-aminobenzoate methanesulfonate) and frozen at -20°C (N= 11-27 per treatment).

The sampled larvae were later photographed under a stereomicroscope next to a micrometer scaling bar. The photographs were then used to measure the larval sizes using the software ImageJ. In order to measure dry weight, larvae were freeze-dried (Christ Alpha 1-4 freeze dryer, Martin Christ Gefriertrocknungsanlagen GmbH, Osterrode, Germany) before being weighed (Sartorius SC2 microbalance, Sartorius AG, Göttingen, Germany, precision 0.1µg).

For the ossification analysis, larvae at 36 dph were fixed in 70% ethanol (Schnell, Konstantinidis, & Johnson, 2016). For the investigation of the skeletal development, the larvae were cleared and double stained (c&s) in an acid-free c&s method, following a modified protocol of Walker & Kimmel (2007). After fixation specimens were stained for cartilage in 6 ml of an acid-free alcian blue staining solution corresponding to Part A of Walker & Kimmel (2007). After 24 h the specimens were washed in a 70% ethanol solution to remove excessive alcian blue and transferred for another 24 h into 6 ml of a 0.5% KOH solution containing four droplets of a 3% H<sub>2</sub>O<sub>2</sub> solution and alizarin red powder. Tissues of the larvae were then cleared by KOH, bleached by H<sub>2</sub>O<sub>2</sub> and stained for calcified structures by alizarin red. Finally specimen were transferred to a 70% glycerin solution for dissection and digital documentation. All specimen were digitally photographed with an Axiocam microscope camera attached to a ZEISS Discovery V20 stereomicroscope and processed with

the Zeiss ZEN software. The number of ossified vertebrae of between seven to 19 larvae per treatment was counted on these photographs. To measure the gill structures seven to twelve larvae per treatment were used. The branchial basket of each larva was removed and the third ceratobranchial was dissected out and photographed in order to measure the ossified structure (the pink stained structure in Figure 1) in length (horizontal line in Figure 1a), the length of the longest gill filament (vertical line in Figure 1a) and the area of gill filaments above the ossified part of the ceratobranchial (as shown in red in Figure 1b).

For the histological analysis, four to nine larvae per treatment were fixed in 4% buffered formaldehyde at 35 dph, embedded in *Technovit*<sup>®</sup> or paraffin, sectioned transversely or longitudinally at 3µm, followed by staining with methylene blue or haematoxylin and eosin respectively. *Technovit*-sections from head region (with eyes, gills and heart), front part of gut (with liver, pancreatic tissue, kidney tissue) as well as paraffin sections were studied and photographed with the microscope (Leitz Aristoplan with a Leica DFC295 camera). Moderate or numerous amounts of vacuoles in the pigment layer of the retina were noted and given a 4-category, qualitative score (normal 0/ some +/- several ++/ many +++). A similar score was used for registration of lipid vacuoles in the cod larvae livers.

Prior to RNA/DNA analyses and lipid determination all larvae previously stored at -80°C were freeze dried for 16 h and weighed (precision 0.1 µg). Measurements of RNA and DNA were made using the fluorimetric method described by Clemmesen (1993) and modified by Malzahn *et al.* (2007). Five larvae from three replicate tanks each were freeze dried, weighed and the tissue was homogenized in 400 µl or 800 µl 0.01% sodiumdodecyl sulfate Tris buffer

(TE SDS), depending on the individual larval dry mass. When necessary, the homogenate was diluted up to 10-fold with 0.01% TE-SDS prior to fluorimetric determination. Ethidium bromide was used as a specific nucleic acid fluorescent dye for both RNA and DNA, and the total fluorescence was measured (Fluoroskan Ascent, Thermo Scientific, Waltham, Massachusetts, USA). RNase was then used to digest all RNA enzymatically. The RNA fluorescence was calculated by subtracting the DNA fluorescence from the total fluorescence. By using the calibration curve fitted to the standard measurements (23 s r-RNA Boehringer, Boehringer Ingelheim GmbH, Ingelheim am Rhein, Germany) the amount of RNA was calculated. Following Le Pecq & Paoletti (1966), the DNA concentration was calculated using the relationship between RNA and DNA fluorescence with a slope ratio of standard DNA to standard RNA of 2.2, which adjusts for the relative fluorescence intensity difference of RNA and DNA.

Total lipids were extracted from individual freeze-dried and weighted cod larvae (five additional larvae from each tank) using a modification of the Folch method (Folch, Lees, & Stanley, 1957) with dichloromethane/methanol/chloroform (1:1:1 v/v/v). Freeze dried larvae were individually placed in 1.5 ml of the solution in a glass vial, securely capped with Teflon lined screwcaps and stored at -80°C for 72 h. The defatted carcasses were transferred to Eppendorf vials and placed with open lids in a desiccator for 48 h to allow for evaporation of the remaining lipid solvents before determining the defatted dry weight on a microscale (Sartorius SC2 microbalance). By subtracting the dry weight of the defatted carcass from the total larval dry weight, the amount of lipids could be gravimetrically determined and presented as lipids in % of dry weight.

For the analyses linear mixed-effects models were used to test for interactions between the CO<sub>2</sub> and the food treatment (as fixed effects) and their effect on larval size, while using the tank ID as a random effect. In case of the gill and the vertebrae measurements, standard length was included as a covariate to account for differences in gill size or stage of ossification that could be due to larval size rather than CO<sub>2</sub> treatment. The model always followed the following formula: `model <- lmer (Parameter ~ CO2 treatment * Food treatment [*standard length] + (1 | Tank ID))`. No assumptions of this analysis were violated. Lipid content was logit transformed as recommended for proportional data (Warton & Hui, 2011). No other parameters needed to be transformed. P-values were later obtained using the `anova(model)` function.

All statistical analyses were run in the programs R (Version 3.3.2) and RStudio (Version 1.0.136). Graphics were done in the R package `ggplot2` (Wickham, 2009) and the linear mixed-effects model was run in the R package “`lme4`” (Bates, Mächler, Bolker, & Walker, 2015).

## **Results**

The CO<sub>2</sub> and food treatment showed a significant interaction effect on larval size, both in terms of standard length (SL,  $t=-3.49$ ,  $p<0.001$ ) as well as dry weight (DW,  $t=-2.021$ ,  $p=0.043$ ) (All results of the statistical analyses are shown in detail in Supplement S Table 2).

Larvae in the high-food treatment were significantly longer ( $t=-3.97$ ,  $p<0.0001$ ) and heavier ( $t=-8.521$ ,  $p<0.0001$ ) than those in the low-food treatment. While larval size at the high-food regime was not affected by the acidification, larvae at the low-food regime were heavier

(mean: 0.49mg vs. 0.29mg) and longer (mean: 8.9mm vs. 7.8mm) under elevated CO<sub>2</sub> compared to ambient CO<sub>2</sub> levels (Figure 2). The variance due to the tank ID was consistently small in all parameter's analyses.

The number of ossified vertebrae increased linearly with the standard length in all treatments (Supplement S Table 2, Figure 3). After taking this relationship into account, both CO<sub>2</sub> and food treatment had a strong interactive effect on the number of vertebrae that were fully ossified ( $t=2.054$ ,  $p<0.0001$ ). Under ambient CO<sub>2</sub>, larvae in the high-food treatment had about twice as many fully ossified vertebrae than those in the low-food treatment, both normalized for standard length. However, larvae in the low-food, elevated CO<sub>2</sub> treatment had more fully ossified vertebrae than those in the low-food, ambient CO<sub>2</sub> treatment, closely resembling the relationship between standard length and number of ossified vertebrae of the high food/ambient CO<sub>2</sub> treatment. In the elevated CO<sub>2</sub> treatment, the relationship between the number of fully ossified vertebrae and standard length was similar between high and low-food treatment larvae (Figure 3).

The RNA/DNA ratios were most strongly affected by the food treatment ( $t=-10.39$ ,  $p<0.0001$ ) with much higher ratios in the high food treatment. Additionally, there was an interactive effect between the CO<sub>2</sub> and food treatments ( $t=-2.32$ ,  $p=0.02$ ). In the high-food regime, RNA/DNA ratios were similar between the CO<sub>2</sub> treatments. However, in the low-food regime RNA/DNA ratios did differ between CO<sub>2</sub> regimes with 25% lower RNA/DNA ratios in the ambient treatment (Figure 4a).

The lipid content (Figure 4b) showed a strong interactive effect of the CO<sub>2</sub> and food treatment ( $t=3.38$ ,  $p<0.001$ ). It was on average between 18 and 19% in all larvae from the elevated CO<sub>2</sub> treatment and in those from the ambient CO<sub>2</sub>, high-food treatment, but significantly higher in the larvae from the ambient CO<sub>2</sub>, low-food treatment with a mean above 22%.

The length of the ossified gill arches was most strongly linked to the larval size ( $t=1.905$ ,  $p<0.0001$ , Supplement S Table 2). Nevertheless, there was also a strong interactive effect of the two treatments ( $t=0.931$ ,  $p=0.006$ ). (Figure 5a). Under ambient CO<sub>2</sub> concentrations larvae in the low food treatment had shorter ossified gill arches than those in the high food treatment, even after applying standard length as a covariate. Larvae under elevated CO<sub>2</sub> showed no interaction with feeding level, with gill arch lengths increasing linearly with larval size.

The size of the gill filaments showed a different picture. Both filament length and gill area were strongly positively correlated to the standard length of the larvae ( $t=8.75$ ,  $p=0.02$ ;  $t=1.764$ ,  $p<0.0001$  respectively, Figure 5 b, c). The length of the longest gill filament was additionally affected by the CO<sub>2</sub> treatment ( $t=-2.232$ ,  $p<0.0001$ ) showing longer filaments under ambient CO<sub>2</sub> compared to elevated CO<sub>2</sub> independent of larval size or food treatment (Figure 5b). Gill area was significantly affected by the interaction of both treatments ( $t=1.181$ ,  $p<0.001$ ). In the ambient CO<sub>2</sub> treatment, larger gill areas were found in the high-food regime than in the low-food regime. In the elevated CO<sub>2</sub> treatment larvae showed a

similar gill area to standard length relationship independent of their own food treatment. Overall the larvae in the elevated CO<sub>2</sub> treatment had gill areas close to those in the ambient, low food treatment, and were smaller than those in the ambient, high food treatment (Figure 5c).

The severity of ocean acidification and food limitation induced organ damage depended on the particular organ. While vacuoles in the eyes were found across all treatments, vacuoles in the pancreas, the kidneys and particularly in the liver were most common in larvae in the elevated CO<sub>2</sub>, low-food regime. Damage in the pancreas and the kidneys was mild, but more severe vacuolization was observed in the liver of larvae from the elevated CO<sub>2</sub> low-food treatment (Figure 6).

## **Discussion**

Both prey availability and ocean acidification produced strong interactive effects on growth and development of Arcto-Norwegian cod larvae that would have remained undetected when manipulating both factors in isolation. This result is particularly relevant, since we chose a realistic end-of-the-century CO<sub>2</sub> concentration for the ocean acidification level (AMAP, 2013; Denman et al., 2011). Specifically, only larvae in the low-food treatment showed significant differences according to the CO<sub>2</sub> treatment. Here, larvae were heavier, longer, had higher RNA/DNA ratios, lower lipid contents and longer ossified gill arches compared to larvae from the ambient CO<sub>2</sub> treatment. This was unexpected since osmoregulation in fish is an energetically costly process, facilitated by ATPase (Melzner et al., 2009), which would suggest that the larvae from the elevated CO<sub>2</sub> treatment should be energetically more limited.

Accepted Article

It is noteworthy that growth and skeletal development is similar in larvae from the high CO<sub>2</sub>, low-food treatment and larvae from the high-food, ambient CO<sub>2</sub> treatment. For a quick recapitulation: larvae in the low-food, elevated CO<sub>2</sub> treatment had more fully ossified vertebrae than those in the low-food, ambient CO<sub>2</sub> treatment, and with that closely resemble the relationship between standard length and number of ossified vertebrae of the high food/ambient CO<sub>2</sub> treatment. This result in combination with the liver histology (see discussion below) indicates a compensatory effect in larvae, facing stressors like high CO<sub>2</sub> and low food, in order to maintain their growth and development. This compensatory effect might be achieved by altered energy allocation in the liver (i.e. highly vacuolized and less storage in the liver) and the lower body lipid content resulting in changed organ development/histomorphology and ossification. As a consequence particularly the liver may possibly be impaired by the vacuolization found in the low-food, elevated CO<sub>2</sub> treatment, which was not found in the high-food, elevated CO<sub>2</sub> treatment. Since our sample size for histological analysis was very low and statistical analyses were not possible, the conclusion drawn from the histological data have to be taken with caution. But similar organ impairments of liver and gills have been already documented in cod, herring, tuna and summer flounder in response to ocean acidification (Chambers et al., 2013; Frommel et al., 2012, 2014, 2016; Stiasny et al., 2018). The observed lower body lipid content in the larvae from the low food elevated CO<sub>2</sub> treatment might be associated with the increase in ossification, since dietary lipids are important energy stores for skeletal development in marine fishes and might have been allocated for the ossification process (Lall & Lewis-McCrea, 2007; Kjørsvik *et al.*, 2009). Additionally, high lipid levels of larvae in the low-food, ambient CO<sub>2</sub> treatment suggest that less energy was used for ossification and developmental processes compared to the other treatments. This hypothesis is supported by the smaller larval size and fewer ossified vertebrae.



The gill size was always smaller in the elevated CO<sub>2</sub> treatment, independent of food regime and larval size. Considering that the active surface of the gill is a complex three dimensional structure and therefore several fold larger than the dimensional area that was measured (Lefevre, McKenzie, & Nilsson, 2017), these differences suggest a marked impairment of gill functionality, with potentially severe effects on individual performance, including osmoregulation, oxygen uptake and swimming performance. For example, Watson et al. (2018) found that juvenile kingfish had reduced swimming performance when reared at elevated CO<sub>2</sub>, which could be a consequence of reduced gill area as observed here. Considering the gill is also the primary organ for acid-base regulation, it poses the question whether ocean acidification effects may lead to a vicious circle in larval development. Ocean acidification may hamper gill development, leaving the young fish with even less potential to react to pH changes. There is significant need for further research looking into the effects of ocean acidification on physiology and development.

Increased calcification of the otoliths due to acidification has been shown for several species (Bignami, Enochs, Manzello, Sponaugle, & Cowen, 2013; Hurst et al., 2012; Maneja et al., 2013; Pimentel et al., 2014; Réveillac et al., 2015). This is likely due to the increased concentration of bicarbonate ions in the blood, which is elevated as a buffering mechanism during hypercapnia or pH stress (Melzner et al., 2009). The skeleton of marine fishes on the other hand is composed of calcium phosphate rather than calcium carbonate as are otoliths. Nevertheless, it appears that the ossification of skeletal elements is also affected by pH stress. Crespel *et al.* (2017) found an increase in bone calcification and a lower occurrence of bone deformities in 45 days old sea bass larvae reared under elevated CO<sub>2</sub> levels (1520 ppm). This suggests that the observed faster skeleton development might be an indirect effect of internal

pH homeostasis and not caused by changes in the regulatory pathways. Nonetheless, the mechanism relating water CO<sub>2</sub> content to skeleton mineralisation remains unknown (Crespel et al., 2017). This highlights a major lack in our physiological understanding of ocean acidification effects in fish.

Previous data on increased larval mortality are important to interpret the ocean acidification effects on growth and development presented here (Stiasny et al., 2016). Some of the effects observed here may actually be a population-level rather than a physiological response, namely when particular phenotypic traits were correlated with susceptibility to ocean acidification, leaving, for example, more larvae in the population that showed increased growth under enhanced *p*CO<sub>2</sub>. Unfavourable prey availability selects for phenotypes with increased growth (Folkvord, Rukan, Johannessen, & Moksness, 1997; Meekan & Fortier, 1996). It is well established that enhanced growth results in higher survival during early-life-stages, since larvae outgrow the predator field faster (Bailey & Houde, 1989). As another explanation, increased CO<sub>2</sub> concentrations may have led to increased foraging behaviour and food consumption (e.g. (*Amphiprion melanopus*) (Nowicki, Miller, & Munday, 2012) which may also apply to Atlantic cod larvae in our experiment and explain larger sizes under ocean acidification. Foraging behaviour or actual food consumption should be considered in future experiments.

A recent meta-analysis revealed that the data basis is still too limited to allow for generalizations with regard to ocean acidification effects on growth among fish species (Cattano et al., 2018). The strong interaction between food supply and CO<sub>2</sub> treatment on growth and development observed in this study highlights that food regime is important in

assessing the impacts of ocean acidification on marine fishes. Therefore, some of the laboratory studies may reflect an incomplete or inaccurate picture of the likely responses especially for wild populations, since the majority are using ad libitum feeding conditions.

In conclusion it appears the cod larvae make the trade-off to spend more energy on growth and ossification of skeletal elements in the elevated CO<sub>2</sub> treatment compared to larvae in the ambient treatment, while putting less effort into organ development, when they are energy limited. While our study confirms increased larval size under ocean acidification - but only in an energy limited situation – it suggests metabolic trade-offs associated with near future CO<sub>2</sub> levels. This information is critically needed for the evaluation of ocean acidification effects.

An increase in size may not translate into increased Darwinian fitness of larvae, i.e. enhanced reproductive contribution to the next generation. Increased larval size may result in developmental patterns that can possibly decrease fitness through decreased functionality of the organs or even impairments. To date, it remains unclear which factors drive these divergent responses and cause the larvae to invest so heavily into growth. Taken together, our study suggests that the responses of organisms and populations should be assessed as comprehensively as possible to not miss one of the critical biological effects.

### **Acknowledgements**

The authors thank three anonymous reviewers, whose comments have significantly improved the manuscript. Additionally, we thank Richard Nash for his editing eye during the revision process, significantly improving the language of this manuscript. Funding was provided through the BIOAcid project (Biological Impacts of Ocean ACIDification) funded by the

German Ministry for Education and Research (BMBF) (Grant No. 03F0655A), the AquaExcel transnational access grant for aquaculture infrastructures (Grant No. 0104/06/08/21) and the Bonus Baltic Sea research and development programme (Art 185) BIO-C3 project, funded jointly by the EU and the BMBF (Grant No. 03F0682A). The experiment was conducted at the Norwegian Cod Breeding Centre, NOFIMA, Tromsø, Norway. The authors are grateful to all scientists and staff at the station and all scientists from BIOAcid, who supported the experiment.

## References

- AMAP. (2013). *AMAP Assessment 2013: Arctic Ocean Acidification*. Oslo, Norway.
- Bailey, K. M., & Houde, E. D. (1989). Predation on Eggs and Larvae of Marine Fishes and the Recruitment Problem. *Advances in Marine Biology*, 25. [https://doi.org/10.1016/S0065-2881\(08\)60187-X](https://doi.org/10.1016/S0065-2881(08)60187-X)
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, 67(1), 1–48. <https://doi.org/10.18637/jss.v067.i01>
- Baumann, H., Talmage, S. C., & Gobler, C. J. (2012). Reduced early life growth and survival in a fish in direct response to increased carbon dioxide. *Nature Climate Change*, 2(1), 38–41. <https://doi.org/10.1038/nclimate1291>
- Bignami, S., Enochs, I. C., Manzello, D. P., Sponaugle, S., & Cowen, R. K. (2013). Ocean acidification alters the otoliths of a pantropical fish species with implications for sensory function. *Proceedings of the National Academy of Sciences of the United States of*

*America*, 110(18), 7366–7370. <https://doi.org/10.1073/pnas.1301365110>

Boglione, C., Gisbert, E., Gavaia, P., Witten, P. E., Moren, M., Fontagné, S., & Koumoundouros, G. (2013). Skeletal anomalies in reared European fish larvae and juveniles. Part 2: Main typologies, occurrences and causative factors. *Reviews in Aquaculture*, 5(SUPPL.1), 121–167. <https://doi.org/10.1111/raq.12016>

Bromhead, D., Scholey, V., Nicol, S., Margulies, D., Wexler, J., Stein, M., ... Lehodey, P. (2015). The potential impact of ocean acidification upon eggs and larvae of yellowfin tuna (*Thunnus albacares*). *Deep Sea Research Part II: Topical Studies in Oceanography*, 113, 268–279. <https://doi.org/10.1016/j.dsr2.2014.03.019>

Buckley, L. J., Caldarone, E. M., & Clemmesen, C. (2008). Multi-species larval fish growth model based on temperature and fluorometrically derived RNA/DNA ratios: Results from a meta-analysis. *Marine Ecology Progress Series*, 371, 221–232. <https://doi.org/10.3354/meps07648>

Cattano, C., Claudet, J., Domenici, P., & Milazzo, M. (2018). Living in a high CO<sub>2</sub> world: a global meta-analysis shows multiple trait-mediated fish responses to ocean acidification. *Ecological Monographs*, 88(3), 320–335. <https://doi.org/10.1002/ecm.1297>

Chambers, R. C., Candelmo, A. C., Habeck, E. A., Poach, M. E., Wieczorek, D., Cooper, K. R., ... Phelan, B. A. (2013). Ocean acidification effects in the early life-stages of summer flounder, *Paralichthys dentatus*. *Biogeosciences Discussions*, 10(8), 13897–13929. <https://doi.org/10.5194/bgd-10-13897-2013>

Chambers, R. C., Candelmo, A. C., Habeck, E. A., Poach, M. E., Wieczorek, D., Cooper, K. R., ... Phelan, B. A. (2014). Effects of elevated CO<sub>2</sub> in the early life stages of summer flounder, *Paralichthys dentatus*, and potential consequences of ocean acidification. *Biogeosciences*, 11, 1613–1626. <https://doi.org/10.5194/bg-11-1613-2014>

- Chícharo, M. a, & Chícharo, L. (2008). RNA:DNA ratio and other nucleic acid derived indices in marine ecology. *International Journal of Molecular Sciences*, 9(8), 1453–1471. <https://doi.org/10.3390/ijms9081453>
- Clemmesen, C. (1993). Improvements in the fluorimetric determination of the RNA and DNA content of individual marine fish larvae. *Marine Ecology Progress Series*, 100, 177–183. <https://doi.org/10.3354/meps100177>
- Clemmesen, C. M. (1994). The effect of food availability, age or size on the RNA/DNA ratio of individually measured herring larvae : Laboratory calibration. *Marine Biology*, 118, 377–382.
- Crespel, A., Zambonino-Infante, J. L., Mazurais, D., Koumoundouros, G., Fragkoulis, S., Quazuguel, P., ... Claireaux, G. (2017). The development of contemporary European sea bass larvae (*Dicentrarchus labrax*) is not affected by projected ocean acidification scenarios. *Marine Biology*, 164(7). <https://doi.org/10.1007/s00227-017-3178-x>
- Denman, K., Christian, J. R., Steiner, N., Pörtner, H.-O., & Nojiri, Y. (2011). Potential impacts of future ocean acidification on marine ecosystems and fisheries : current knowledge and recommendations for future research. *ICES Journal of Marine Science*, 68(6), 1019–1029. <https://doi.org/10.1093/icesjms/fsr074>
- Díaz-Gil, C., Catalán, I. A., Palmer, M., Faulk, C. K., & Fuiman, L. A. (2015). Ocean acidification increases fatty acids levels of larval fish. *Biology Letters*, 11(7). <https://doi.org/10.1098/rsbl.2015.0331>
- Doney, S. C., Fabry, V. J., Feely, R. A., & Kleypas, J. A. (2009). Ocean Acidification: The Other CO<sub>2</sub> Problem. *Annual Review of Marine Science*, 1(1), 169–192. <https://doi.org/10.1146/annurev.marine.010908.163834>
- FAO. (2016). *The State of World Fisheries in 2015. Contributing to food security and*

*nutrition for all*. Rome.

Folch, J., Lees, M., & Stanley, G. H. S. (1957). A simple method for the isolation and purification of total lipids from animal tissues. *Journal of Biological Chemistry*, 226, 497–509.

Folkvord, A., Rukan, K., Johannessen, A., & Moksness, E. (1997). Early life history of herring larvae in contrasting feeding environments determined by otolith microstructure analysis. *Journal of Fish Biology*, 51(sa), 250–263. <https://doi.org/10.1111/j.1095-8649.1997.tb06102.x>

Franke, A., & Clemmesen, C. (2011). Effect of ocean acidification on early life stages of Atlantic herring (*Clupea harengus* L.). *Biogeosciences*, 8(12), 3697–3707. <https://doi.org/10.5194/bg-8-3697-2011>

Fridgeirsson, E. (1978). Embryonic development of five species of gadoid fishes in Icelandic waters. *Rit Fiskideildar*, 5(6), 1–71.

Frommel, A. Y., Maneja, R., Lowe, D., Malzahn, A. M., Geffen, A. J., Folkvord, A., ... Clemmesen, C. (2012). Severe tissue damage in Atlantic cod larvae under increasing ocean acidification. *Nature Climate Change*, 2(1), 42–46. <https://doi.org/10.1038/nclimate1324>

Frommel, A. Y., Maneja, R., Lowe, D., Pascoe, C. K., Geffen, A. J., Folkvord, A., ... Clemmesen, C. (2014). Organ damage in Atlantic herring larvae as a result of ocean acidification. *Ecological Applications*, 24(5), 1131–1143. <https://doi.org/10.1890/13-0297.1>

Frommel, A. Y., Margulies, D., Wexler, J. B., Stein, M. S., Scholey, V. P., Williamson, J. E., ... Havenhand, J. (2016). Ocean acidification has lethal and sub-lethal effects on larval development of yellowfin tuna, *Thunnus albacares*. *Journal of Experimental Marine*

*Biology and Ecology*, 482, 18–24. <https://doi.org/10.1016/j.jembe.2016.04.008>

Gobler, C. J., Merlo, L. R., Morrell, B. K., & Griffith, A. W. (2018). Temperature, Acidification, and Food Supply Interact to Negatively Affect the Growth and Survival of the Forage Fish, *Menidia beryllina* (Inland Silverside), and *Cyprinodon variegatus* (Sheepshead Minnow). *Frontiers in Marine Science*, 5(March), 1–12. <https://doi.org/10.3389/fmars.2018.00086>

Hurst, T. P., Fernandez, E. R., & Mathis, J. T. (2013). Effects of ocean acidification on hatch size and larval growth of walleye pollock (*Theragra chalcogramma*). *ICES Journal of Marine Science*, 70(4), 812–822. <https://doi.org/10.1093/icesjms/fst053>

Hurst, T. P., Fernandez, E. R., Mathis, J. T., Miller, J. A., Stinson, C. M., & Ahgeak, E. F. (2012). Resiliency of juvenile walleye pollock to projected levels of ocean acidification. *Aquatic Biology*, 17, 247–259. <https://doi.org/10.3354/ab00483>

Hurst, T. P., Laurel, B. J., Hanneman, E., Haines, S. A., & Ottmar, M. L. (2017). Elevated CO<sub>2</sub> does not exacerbate nutritional stress in larvae of a Pacific flatfish. *Fisheries Oceanography*, 26(3), 336–349. <https://doi.org/10.1111/fog.12195>

IPCC. (2013). *Climate Change 2013 - The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel of Climate Change*. (T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, ... P. M. Midgley, Eds.). Cambridge: Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. <https://doi.org/10.1017/CBO9781107415324>

Kjesbu, O. S. (1989). The spawning activity of cod, *Gadus morhua* L. *Journal of Fish Biology*, 34(2), 195–206. <https://doi.org/10.1111/j.1095-8649.1989.tb03302.x>

Kjesbu, O. S., Bogstad, B., Devine, J. A., Gjørseter, H., Howell, D., Ingvaldsen, R. B., ... Skjæraasen, J. E. (2014). Synergies between climate and management for Atlantic cod



fisheries at high latitudes. *Proceedings of the National Academy of Sciences*, 111(9), 3478–3483. <https://doi.org/10.1073/pnas.1316342111>

Kjørsvik, E., Olsen, C., Wold, P. A., Hoehne-Reitan, K., Cahu, C. L., Rainuzzo, J., ... Olsen, Y. (2009). Comparison of dietary phospholipids and neutral lipids on skeletal development and fatty acid composition in Atlantic cod (*Gadus morhua*). *Aquaculture*, 294(3–4), 246–255. <https://doi.org/10.1016/j.aquaculture.2009.06.012>

Koenigstein, S., Dahlke, F. T., Stiasny, M. H., Storch, D., Clemmesen, C., & Pörtner, H.-O. (2018). Forecasting future recruitment success for Atlantic cod in the warming and acidifying Barents Sea. *Global Change Biology*, 24(1), 526–535. <https://doi.org/10.1111/gcb.13848>

Kreiss, C. M., Michael, K., Lucassen, M., Jutfelt, F., Motyka, R., Dupont, S., & Pörtner, H. O. (2015). Ocean warming and acidification modulate energy budget and gill ion regulatory mechanisms in Atlantic cod (*Gadus morhua*). *Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology*, 185(7), 767–781. <https://doi.org/10.1007/s00360-015-0923-7>

Lall, S. P., & Lewis-McCrea, L. M. (2007). Role of nutrients in skeletal metabolism and pathology in fish - An overview. *Aquaculture*, 267(1–4), 3–19. <https://doi.org/10.1016/j.aquaculture.2007.02.053>

Lam, V. W. Y., Cheung, W. W. L., & Sumaila, U. R. (2014). Marine capture fisheries in the Arctic: winners or losers under climate change and ocean acidification? *Fish and Fisheries*, 1–23. <https://doi.org/10.1111/faf.12106>

Le Pecq, J. B., & Paoletti, C. (1966). A new fluorometric method for RNA and DNA determination. *Annual Review of Biochemistry*, 17, 100–107.

Lefevre, S., McKenzie, D. J., & Nilsson, G. E. (2017). Models projecting the fate of fish

populations under climate change need to be based on valid physiological mechanisms.

*Global Change Biology*, 1–11. <https://doi.org/10.1111/gcb.13652>

Malzahn, A. M., Aberle, N., Clemmesen, C., & Boersma, M. (2007). Nutrient limitation of primary producers affects planktivorous fish condition. *Limnology and Oceanography*, 52(5), 2062–2071. <https://doi.org/10.4319/lo.2007.52.5.2062>

Maneja, R., Frommel, A., Geffen, A., Folkvord, A., Piatkowski, U., Chang, M., & Clemmesen, C. (2013). Effects of ocean acidification on the calcification of otoliths of larval Atlantic cod *Gadus morhua*. *Marine Ecology Progress Series*, 477(May 2010), 251–258. <https://doi.org/10.3354/meps10146>

Matson, P. G., Yu, P. C., Sewell, M. A., & Hofmann, G. E. (2012). Development Under Elevated  $p\text{CO}_2$  Conditions Does Not Affect Lipid Utilization and Protein Content in Early Life-History Stages of the Purple Sea Urchin, *Strongylocentrotus purpuratus*. *The Biological Bulletin*, 223(3), 312–327. <https://doi.org/10.1086/BBLv223n3p312>

Meekan, M. G., & Fortier, L. (1996). Selection for fast growth during the larval life of Atlantic cod *Gadus morhua* on the Scotian Shelf. *Marine Ecology Progress Series*, 137, 25–37. <https://doi.org/10.3354/meps137025>

Melzner, F., Gutowska, M. A., Langenbuch, M., Dupont, S., Lucassen, M., Thorndyke, M. C., ... Pörtner, H.-O. (2009). Physiological basis for high  $\text{CO}_2$  tolerance in marine ectothermic animals: pre-adaptation through lifestyle and ontogeny? *Biogeosciences Discussions*, 6(3), 4693–4738. <https://doi.org/10.5194/bgd-6-4693-2009>

Munday, P. L., Donelson, J. M., Dixson, D. L., & Endo, G. G. K. (2009). Effects of ocean acidification on the early life history of a tropical marine fish. *Proceedings of The Royal Society B: Biological Sciences*, 276(1671), 3275–3283. <https://doi.org/10.1098/rspb.2009.0784>

- Nowicki, J. P., Miller, G. M., & Munday, P. L. (2012). Interactive effects of elevated temperature and CO<sub>2</sub> on foraging behavior of juvenile coral reef fish. *Journal of Experimental Marine Biology and Ecology*, 412, 46–51. <https://doi.org/10.1016/j.jembe.2011.10.020>
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. a, ... Yool, A. (2005). Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature*, 437(7059), 681–686. <https://doi.org/10.1038/nature04095>
- Perry, A. L., Low, P. J., Ellis, J. R., & Reynolds, J. D. (2005). Climate change and distribution shifts in marine fishes. *Science (New York, N.Y.)*, 308(5730), 1912–1915. <https://doi.org/10.1126/science.1111322>
- Perry, D. M., Redman, D. H., Widman, J. C., Meseck, S., King, A., & Pereira, J. J. (2015). Effect of ocean acidification on growth and otolith condition of juvenile scup, *Stenotomus chrysops*. *Ecology and Evolution*, 5(18), 4187–4196. <https://doi.org/10.1002/ece3.1678>
- Pimentel, M., Pegado, M., Repolho, T., & Rosa, R. (2014). Impact of ocean acidification in the metabolism and swimming behavior of the dolphinfish (*Coryphaena hippurus*) early larvae. *Marine Biology*, 161(3), 725–729. <https://doi.org/10.1007/s00227-013-2365-7>
- Pimentel, M. S., Faleiro, F., Dionisio, G., Repolho, T., Pousao-Ferreira, P., Machado, J., & Rosa, R. (2014). Defective skeletogenesis and oversized otoliths in fish early stages in a changing ocean. *J Exp Biol*, 217(Pt 12), 2062–2070. <https://doi.org/10.1242/jeb.092635>
- Poloczanska, E. S., Brown, C. J., Sydeman, W. J., Kiessling, W., Schoeman, D. S., Moore, P. J., ... Richardson, A. J. (2013). Global imprint of climate change on marine life. *Nature Climate Change*, 3(10), 919–925. <https://doi.org/10.1038/nclimate1958>
- Réveillac, E., Lacoue-Labarthe, T., Oberhänsli, F., Teyslié, J. L., Jeffree, R., Gattuso, J. P., &

Martin, S. (2015). Ocean acidification reshapes the otolith-body allometry of growth in juvenile sea bream. *Journal of Experimental Marine Biology and Ecology*, 463, 87–94. <https://doi.org/10.1016/j.jembe.2014.11.007>

Schnell, N. K., Konstantinidis, P., & Johnson, G. D. (2016). High-proof Ethanol Fixation of Larval and Juvenile Fishes for Clearing and Double Staining. *Copeia*, 104(3), 617–622. <https://doi.org/10.1643/CI-15-382>

Silva, C. S. E., Novais, S. C., Lemos, M. F. L., Mendes, S., Oliveira, A. P., Gonçalves, E. J., & Faria, A. M. (2016). Effects of ocean acidification on the swimming ability, development and biochemical responses of sand smelt larvae. *Science of The Total Environment*, 563–564(September), 89–98. <https://doi.org/10.1016/j.scitotenv.2016.04.091>

Sswat, M., Stiasny, M. H., Jutfelt, F., Riebesell, U., & Clemmesen, C. (2018). Growth performance and survival of larval Atlantic herring , under the combined effects of elevated temperatures and CO<sub>2</sub>. *PloS One*, 13(1), 1–19. <https://doi.org/10.1371/journal.pone.0191947>

Sswat, M., Stiasny, M. H., Taucher, J., Algueró-Muñiz, M., Bach, L. T., Jutfelt, F., ... Clemmesen, C. (2018). Food web changes under ocean acidification promote herring larvae survival. *Nature Ecology and Evolution*, 2(May), 1–5. <https://doi.org/10.1038/s41559-018-0514-6>

Steinacher, M., Joos, F., Frölicher, T. L., Plattner, G.-K., & Doney, S. C. (2009). Imminent ocean acidification projected with the NCAR global coupled carbon cycle-climate model. *Biogeosciences Discussions*, 5(6), 4353–4393. <https://doi.org/10.5194/bgd-5-4353-2008>

Stiasny, M. H., Mittermayer, F. H., Göttler, G., Bridges, C. R., Falk-Petersen, I.-B.,

Puvanendran, V., ... Clemmesen, C. (2018). Effects of parental acclimation and energy limitation in response to high CO<sub>2</sub> exposure in Atlantic cod. *Scientific Reports*, 8(1). <https://doi.org/10.1038/s41598-018-26711-y>

Stiasny, M. H., Mittermayer, F. H., Sswat, M., Voss, R., Jutfelt, F., Chierici, M., ... Clemmesen, C. (2016). Ocean Acidification Effects on Atlantic Cod Larval Survival and Recruitment to the Fished Population. *PLoS ONE*, 11(8), e0155448. <https://doi.org/10.1371/journal.pone.0155448>

Thomsen, J., Casties, I., Pansch, C., Körtzinger, A., & Melzner, F. (2013). Food availability outweighs ocean acidification effects in juvenile *Mytilus edulis*: Laboratory and field experiments. *Global Change Biology*, 19(4), 1017–1027. <https://doi.org/10.1111/gcb.12109>

Towle, E. K., Enochs, I. C., & Langdon, C. (2015). Threatened Caribbean coral is able to mitigate the adverse effects of ocean acidification on calcification by increasing feeding rate. *PLoS ONE*, 10(4). <https://doi.org/10.1371/journal.pone.0123394>

Vanderplancke, G., Claireaux, G., Quazuguel, P., Madec, L., Ferraresso, S., Sévère, A., ... Mazurais, D. (2015). Hypoxic episode during the larval period has long-term effects on European sea bass juveniles (*Dicentrarchus labrax*). *Marine Biology*, 162(2), 367–376. <https://doi.org/10.1007/s00227-014-2601-9>

Varsamos, S., Nebel, C., & Charmantier, G. (2005). Ontogeny of osmoregulation in postembryonic fish: A review. *Comparative Biochemistry and Physiology - A Molecular and Integrative Physiology*, 141(4 SPEC. ISS.), 401–429. <https://doi.org/10.1016/j.cbpb.2005.01.013>

Walker, M., & Kimmel, C. (2007). A two-color acid-free cartilage and bone stain for zebrafish larvae. *Biotechnic & Histochemistry*, 82(1), 23–28.

<https://doi.org/10.1080/10520290701333558>

Warton, D. I., & Hui, F. K. C. (2011). The arcsine is asinine: the analysis of proportions in ecology. *Ecology*, 92(1), 3–10. <https://doi.org/10.1890/10-0340.1>

Watson, S. A., Allan, B. J. M., McQueen, D. E., Nicol, S., Parsons, D. M., Pether, S. M. J., ... Munday, P. L. (2018). Ocean warming has a greater effect than acidification on the early life history development and swimming performance of a large circumglobal pelagic fish. *Global Change Biology*, 24(9), 4368–4385. <https://doi.org/10.1111/gcb.14290>

Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis*. New York City, US: Springer International Publishing.

Zambonino Infante, J. L., & Cahu, C. L. (2001). Ontogeny of the gastrointestinal tract of marine fish larvae. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology December*, 130(4), 477–487. [https://doi.org/10.1016/S1532-0456\(01\)00274-5](https://doi.org/10.1016/S1532-0456(01)00274-5)

### Figure Captions

Figure 1. Third ceratobranchial with gill filaments of 36 days old cod larvae showing (a) the measured lengths of the ossified gill arch (horizontal line), which is stained in pink, and the longest gill filament (vertical line) and (b) the measured gill filament area (shown encircled in red).

Figure 2. (a) Standard length of 36 days old cod larvae in mm and (b) Dry weight in mg (mean +/- SD) (N=11-26 per treatment).

Figure 3. Image of cleared and double stained specimens showing number of ossified vertebrae (coloured in red) in 36 days old cod larvae in (a) ambient CO<sub>2</sub>, high food (b) acidification, high food (c) ambient CO<sub>2</sub>, low food, (d) acidification, low food. (e) Number of fully ossified vertebrae in relation to standard length, lines show the linear relationship per treatment.

Figure 4. (a) RNA/DNA content of the 36 days old cod larvae (N=10-18 per treatment) and (b) Lipid content of the larvae (N=11-19 per treatment; mean +/- SD).

Figure 5. (a) Length of the ossified gill arch (mm) of 36 days old cod larvae (b) length of the longest gill filament (mm) (c) gill area (mm<sup>2</sup>) all in relation to standard length (in mm), lines show the linear relationships per treatment.

Figure 6. Frequency of organ impairments in (a) liver, (b) eyes, (c) pancreas, and (d) kidney of the 35 days old cod larvae in a subjective scale from 0 (no impairments) to +++ (very severe/many impairments). (N=6,5,3,9 (from left in each panel))

#### **Author contributions**

C.C., M.H.S. and M.S. designed the experiment; M.H.S., M.S., F.H.M. and C.C. performed the experiment; V.P. and A.M. supported the experiment and provided the facilities; N.K.S. performed the ossification analysis; I.B.F.P. performed the histological analysis; M.H.S., M.S., C.C. analyzed data; M.H.S., T.B.H.R., and C.C. wrote the main paper; All authors discussed the results and implications and commented on the manuscript at all stages.

### Additional information

The authors declare no competing financial interests. Correspondence and requests for materials should be addressed to C.C.













