

Title	SNC log symplectic structures on Fano products
Author(s)	Okumura, Katsuhiko
Citation	代数幾何学シンポジウム記録 (2018), 2018: 149-149
Issue Date	2018
URL	http://hdl.handle.net/2433/236417
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

SNC log symplectic structures on Fano products

Katsuhiko Okumura
Waseda University

Introduction to Poisson structures

Definition (Poisson bracket).

The holomorphic **Poisson bracket** on X :

- (bilinear form) $\{-, -\} : \mathcal{O}_X \times \mathcal{O}_X \rightarrow \mathcal{O}_X$
- (skew-symmetric) $\{f, g\} = -\{g, f\}$,
- (Jacobi identity) $\{f, \{g, h\}\} + \{g, \{h, f\}\} + \{h, \{f, g\}\} = 0$,
- (Leibniz rule) $\{f, g \cdot h\} = \{f, g\}h + \{f, h\}g$.

Definition (Poisson structure).

The holomorphic **Poisson structure** on the smooth variety X :

$\Pi \in \Gamma(X, \wedge^2 T_X)$ s.t. $[\Pi, \Pi] = 0 \in \wedge^3 T_X$, where $[-, -]$ is the Schouten bracket.

$$[\frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}, \frac{\partial}{\partial x_k} \wedge \frac{\partial}{\partial y_l}] = \frac{\partial^2 \Pi}{\partial x_i \partial y_j} \wedge \frac{\partial}{\partial x_k} \wedge \frac{\partial}{\partial y_l} + \frac{\partial^2 \Pi}{\partial y_j \partial x_k} \wedge \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_l} + \frac{\partial^2 \Pi}{\partial x_i \partial x_k} \wedge \frac{\partial}{\partial y_j} \wedge \frac{\partial}{\partial y_l} + \frac{\partial^2 \Pi}{\partial y_j \partial y_l} \wedge \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_k}$$

Remark.

- {Poisson structures on X } \leftrightarrow {Poisson brackets on X }
- (\cdot):
 $\rightarrow : \Pi : \mathcal{O}_X \times \mathcal{O}_X \rightarrow \mathcal{O}_X; (f, g) \mapsto \Pi^\#(df, dg)$
 $\leftarrow : \Pi = \sum_{i,j} \{x_i, x_j\} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}$
- $[\Pi, \Pi] = 0 \iff$ Jacobi identity

Definition (Degeneracy divisor).

- Poisson structure (X, Π) is **generically symplectic**
 $\iff \dim X = 2n$ and $\Pi^n \neq 0$.

Suppose that (X, Π) is a generically symplectic.

- $D(\Pi) := \{x \in X \mid \Pi^n(x) = 0\}$ forms a divisor called the **degeneracy divisor**.
- (X, Π) is a **log symplectic structure**
 $\iff D(\Pi)$ is a reduced divisor,
- (X, Π) is a **SNC log symplectic structure**
 $\iff (X, \Pi)$ is a log symplectic structure and $D(\Pi)$ is a simple normal crossing divisor.
 $\Pi^n \in \Gamma(X, \wedge^{2n} T_X) \rightsquigarrow D(\Pi) \sim -\mathcal{K}_X$.

Motivations & Main Result

Main Theorem

X_i : Fano variety over \mathbb{C} , $\text{Pic}(X_i) = \mathbb{Z}$,
 $\dim X_i = n_i \geq 3$,
 $X = \prod_{i=1}^m X_i$, $\dim X = 2n$,
 Π : SNC log symplectic structure.

- $X_i = \mathbb{P}^{n_i}$
- Π : **diagonal Poisson structure**

Background

Question

How many (X, Π) with conditions

- X : smooth projective variety
- $D(\Pi)$: reduced SNC

Theorem (Lima, Pereira).

X : Fano variety over \mathbb{C} , $\text{Pic}(X) = \mathbb{Z}$, $\dim X = 2n \geq 4$,
 Π : SNC log symplectic structure on X .

- \Rightarrow • $X = \mathbb{P}^n$
- Π : **diagonal Poisson structure on \mathbb{P}^n**

How about if $\rho(X) \geq 2$

Corollary (O).

X : Fano variety over \mathbb{C} , $\text{Pic}(X) = \mathbb{Z}$,
 $\dim X \geq 3$,
 $X = \mathbb{P}^n \iff \exists \Pi$: SNC log symplectic structure on $X \times X$

Diagonal Poisson structures and form as bivector fields

Definition (Diagonal Poisson structure).

$X : \mathbb{A}^{2n}$ or (product of) \mathbb{P}^n , $\dim X = 2n$
 (X, Π) is a **diagonal Poisson structure**
 $\iff D(\Pi)$ is composed of all coordinate hyperplanes.

Theorem (Polishchuk).

There is a surjective map of bivector fields:

$$\varphi_n : \{(\mathbb{A}^{n+1}, \bar{\Pi}) \mid \bar{\Pi} : \text{quadratic}\} \rightarrow \{(\mathbb{P}^n, \Pi)\}$$

Furthermore, $\bar{\Pi}$ is Poisson on $\mathbb{A}^{n+1} \Rightarrow \varphi_n(\bar{\Pi})$ is also Poisson on \mathbb{P}^n .

(\cdot):

$$\left\{ \frac{x_i}{x_k}, \frac{x_j}{x_k} \right\} = \frac{1}{x_k^2} \{x_i, x_j\} - \frac{x_i}{x_k^3} \{x_k, x_j\} - \frac{x_j}{x_k^3} \{x_i, x_k\}$$

$\{x_i, x_j\}$ is quadratic, $\rightsquigarrow \left\{ \frac{x_i}{x_k}, \frac{x_j}{x_k} \right\} \in k \left[\frac{x_i}{x_k}, \dots, \frac{x_j}{x_k} \right]$ \square

Fact (Pym).

$X := \prod_{i=1}^m \mathbb{P}^{n_i}$, $\dim X = \sum_{i=1}^m n_i = 2n$, Π : diagonal Poisson str. on X ,
coordinate: $[x_{10}, \dots, x_{1n_1}; x_{20}, \dots, x_{2n_2}; \dots; x_{m0}, \dots, x_{mn_m}]$,
 $\Rightarrow \exists \sigma = \sum_{1 \leq i, k \leq m, 1 \leq j \leq n_i, 1 \leq l \leq n_k} \Delta_{ijkl}$: diagonal Poisson str. on $\mathbb{A}^{2n-m} \simeq \prod_{i=1}^m \mathbb{A}^{n_i-1}$, where $\Delta_{ij} = c_{ijk} x_{ij} x_{kl} \frac{\partial}{\partial x_{ij}} \wedge \frac{\partial}{\partial x_{kl}}$
s.t. σ induces Π on X .

(\cdot):

Definition. (r-matrix construction)

Π is constructed by **r-matrix construction** w.r.t a Lie group G :

Π is a image of r along $\mathfrak{g} \rightarrow \Gamma(X, T_X)$,

where \mathfrak{g} : Lie algebra of G , r : r-matrix for G , i.e. $[r, r] = 0$.

The Fact comes from r-matrix construction for $G = (\mathbb{C}^*)^n \times (\mathbb{C}^*)^{n_2} \times \dots \times (\mathbb{C}^*)^{n_m}$. \square

Key lemma (Pym).

(X, Π) : SNC log symplectic structure

$D(\Pi) = \sum_{j=1}^k D_j$: irreducible decomposition of the degeneracy divisor.

$$\Rightarrow \text{ch}(T_X) - \text{ch}(T_X^\vee) = 2 \sinh[D_j]$$

(\cdot): We have 2 exact sequences;

$$0 \rightarrow \Omega_X^1 \rightarrow \Omega_X^1(\log D) \rightarrow \bigoplus_{j=1}^m \mathcal{O}_{D_j} \rightarrow 0,$$

$$0 \rightarrow \mathcal{O}_X(-D_j) \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_{D_j} \rightarrow 0.$$