



Australian  
National  
University

# CRE Theme 1: Nutrition and Food security

How effective are family based and institutional nutrition interventions in improving children's diet and health? A systematic review

Andrew P Black, Katina D'Onise, Robyn McDermott, Hassan Vally, Kerin O'Dea



University of South Australia



## ACKNOWLEDGEMENTS

This research is a project of the Australian Primary Health Care Research Institute, which is supported by a grant from the Australian Government Department of Health. The information and opinions contained in it do not necessarily reflect the views or policy of the Australian Primary Health Care Research Institute or the Australian Government Department of Health.

## CITATION

Andrew P Black<sup>1,2</sup>, Katina D'Onise<sup>1</sup>, Robyn McDermott<sup>1,3</sup>, Hassan Vally<sup>4</sup>, Kerin O'Dea<sup>1</sup>. (2017). How effective are family based and institutional nutrition interventions in improving children's diet and health? A systematic review. Report prepared for the Australian Primary Health Care Research Institute (APHCRI), Canberra, ACT.

1. School of Health Sciences, University of South Australia, Adelaide, South Australia, Australia
2. Bulgarr Ngaru Medical Aboriginal Corporation, Grafton, NSW, Australia
3. College of Public Health, Medical and Veterinary Sciences, James Cook University, Cairns, Queensland, Australia
4. School of Psychology & Public Health, La Trobe University, Melbourne, Victoria, Australia

Professor Robyn McDermott  
James Cook University  
Cairns Qld 4870 Australia  
T 61 7 4232 1604  
E [robyn.mcdermott@jcu.edu.au](mailto:robyn.mcdermott@jcu.edu.au)  
<http://www.ccdp.jcu.edu.au>

# CONTENTS

## Contents

Background .....	4
Methods .....	5
CRITERIA FOR INCLUSION IN THE REVIEW.....	5
Types of study.....	5
Participants .....	5
Interventions and specific comparisons to be made .....	5
Outcomes of interest .....	5
Search Methods.....	6
Results .....	8
DESCRIPTION AND SCOPE OF THE INCLUDED STUDIES .....	8
OVERALL METHODOLOGICAL ASSESSMENT .....	9
OUTCOMES.....	10
Impact on Fruit and Vegetable (F&V) intake.....	10
Impact on Fat intake.....	11
Impact on Energy Dense, Nutrient Poor (EDNP) foods intake.....	12
Other Health Outcomes .....	12
SUSTAINABILITY OF NUTRITION AND HEALTH OUTCOMES .....	13
Discussion .....	14
Conclusion .....	16
References .....	17
Appendix 1: Figure 1, Systematic literature review search strategy and results.....	23
Appendix 2: Table 1, Characteristics of included studies.....	24
Appendix 3: Table 2, Impact of nutrition interventions on children's fruit and vegetable intake .....	32
Appendix 4: Table 3, Impact of nutrition interventions on children's fat intake .....	34

## Background

Despite increased interest in food and cooking, high levels of sub-optimal dietary behaviour have been documented in many countries. Nutrition surveys report low levels of fruit and vegetable (F&V) intake, inadequate intake of important nutrients and high intake of energy-dense nutrient poor (EDNP) foods in all age-groups.<sup>1-8</sup> Less healthy dietary behaviour is an important factor underlying the high prevalence of chronic non-communicable diseases, including cardiovascular disease and diabetes mellitus, and increasing obesity rates.<sup>9-11</sup> The early signs of these chronic diseases and risk factors are increasingly being documented in health literature relating to children and adolescents.<sup>12, 13</sup> This emphasises the importance of promoting and supporting the development of healthy eating habits from an early age.

In this context, a key challenge is how to support and engage families to make sustainable changes to healthier eating, despite the challenges posed within contemporary society. Preschool and childcare are increasingly central in family life, in addition to the importance of schools. Given the ready access to children, these settings have been widely used in programs which aim to improve the dietary intake of young children.<sup>14</sup> These nutrition programs have included the reintroduction and expansion of school meal programs, healthy lifestyle promotion programs, economic incentives, enhancement of school canteens and restrictions on the availability of less healthy foods within schools. Recent reviews have found that school-based nutrition programs are moderately effective at improving diet quality, reducing adiposity and improving fruit intake in the short-term.<sup>15-17</sup>

In addition, many of these institutional programs have also incorporated strategies to engage with families to strengthen their impact, as this is seen as vital to improving the sustainability of healthier eating patterns in the community.<sup>15, 18</sup> The emphasis on the role of the family in promoting healthy eating is underpinned by an understanding of the social and cultural context which shapes the preparation, sharing and eating of food.<sup>19</sup> Healthy changes to an individual child's eating patterns will be reinforced if the family also shares the same influences and models the changes in behaviour.<sup>20</sup> The question that this poses is how to influence the dietary patterns in the family. Notwithstanding the major role that women have traditionally had in food preparation, fathers and children have also been shown to have an important impact on family food choices.<sup>21</sup> Current research has been increasingly focused on reducing the increasing prevalence of obesity and thus nutrition interventions have frequently been part of multi-component healthy lifestyle programs.<sup>15</sup> Golley et al. have identified features associated with more effective engagement of parents in child obesity prevention programs including greater parent involvement in implementation, use of self-monitoring and goal-setting behaviour change techniques.<sup>22, 23</sup> However, the evidence of the optimal strategies to involve parents in the promotion of healthier eating in the family remains inconclusive, particularly in school settings.<sup>24</sup>

The objective of this systematic review was to document the potential for family-based and institutional nutrition programs to improve the nutrition and health of young children in high income countries. Another aim was to determine how sustainable the impacts of these nutrition programs are on children's nutrition and health. Finally, lower socio-economic status is associated with both higher risk of chronic non-communicable disease and lower uptake of health promoting behaviours, including healthy eating. Thus, the impact of these nutrition programs that may help to reduce this social inequality was also reviewed.

## Methods

The review was undertaken using the principles outlined in the Cochrane Handbook<sup>25</sup> and the Cochrane Public Health and Health Promotion guidelines<sup>26</sup> with the exception that only one reviewer scanned the title list and/or abstract and that only studies published in English were included. The following definitions were used to identify relevant studies:

- > Family-Based programs – all programs which involved one or more adults with or without their children in any setting
- > Institutional programs – programs involving groups of children in organised settings such as schools, pre-schools, childcare including out of school hours care, youth and church groups
- > Nutrition program – all interventions where one major aim was to improve the quality of dietary intake.

## CRITERIA FOR INCLUSION IN THE REVIEW

### Types of study

Randomised controlled trials (RCTs) including cluster randomised trials were eligible for inclusion in the review.

### Participants

Eligible participants were adults and children from families in high income countries as defined on the World Bank List of High Income Countries ([worldbank.org.au](http://worldbank.org.au)). Low income countries were excluded as the availability of financial resources and the extent of malnutrition in these countries means that the issues are likely to be very different. The age range for inclusion of studies was children from birth through to 12 years (early childhood and primary school). The participants were 'well' children from the local population.

Socio-economic disadvantage was defined as families from areas that are described as disadvantaged (e.g. low income area, ghetto, social housing projects); of low socio-economic status (e.g. low income as defined by the researchers); and disadvantaged minorities (e.g. Indigenous peoples). Socio-economic disadvantage was not an inclusion/exclusion criterion, but was used to describe the impact of interventions on this population.

### Interventions and specific comparisons to be made

Eligible interventions were those that aimed to improve nutritional intake undertaken for at least 12 weeks. In the RCTs, at least one group participated in a nutrition program and another group received a control intervention. Controls may have received no intervention, delayed intervention, or attention control. As an important aspect of school nutrition programs, school meals programs were included, updating the evidence from a 2007 review of school feeding programs.<sup>27</sup> Obesity treatment interventions for children were excluded as these programs have been reviewed recently.<sup>28, 29</sup> Obesity prevention interventions for children have also been reviewed<sup>14, 15</sup> and were only included if the studies reported on nutritional outcomes in addition to changes in anthropometry.

### Outcomes of interest

To be eligible for the review, a study had to report valid measures of at least one primary outcome.

## Primary outcomes

- > Nutritional intake (measured by validated dietary assessment techniques, food purchasing, or biomarkers).

## Health status

- > Any measure of physical health-Mortality, Morbidity, Hospital and Emergency department admissions
- > Child growth and development outcomes- using standardised measures
- > Longer term effects following program completion (e.g. in the year(s) following to look at longer term outcomes)
- > Adverse outcomes – Stigmatisation, dependency, decreased total family expenditure on food (including subsidy), increase in high fat/high sugar foods (including takeaway food), and obesity or excessive weight loss.

## Search Methods

The following electronic databases from 1980 to December 2014 were searched: Medline, Central (Cochrane)/DARE, Embase and CINAHL, with a search strategy that incorporated terms including both Medical Subject headings and keywords for:

1. Food
2. Family-based programs and Institutional nutrition programs
3. Nutrition and health outcomes.

In addition, filters for high income countries and study design were applied. A Medline search strategy was developed and adapted to the other databases as required. The complete search strategy is available from the authors. In addition, the reference lists of included studies were assessed to identify other eligible studies. Only articles published in English were included in the review.

## Data synthesis and analysis

The search results were downloaded into an Endnote library and titles and/or abstracts assessed by one of two authors for eligibility. The full-text manuscripts of those deemed potentially eligible were assessed by either author and any discrepancies resolved by consensus with the other author. Data of eligible studies were entered into a standard template and data entry checked after data extraction completed. The primary health and nutrition outcomes, any adverse outcomes, together with details of the intervention, including length of follow-up, demographics of participants and theoretical basis were extracted. The study authors were contacted to try to obtain additional data if necessary.

The interpretation of the results was facilitated by converting outcomes presented as a mean with a measure of variance to a Cohen's d effect size estimate using an online calculator.<sup>30</sup> The impact of the interventions was compared using effect size (Cohen's d) as a standard measure, which was interpreted as shown: Small 0.1-0.2, moderate 0.3-0.5, large  $\geq 0.6$ . Due to the heterogeneity of the interventions and study outcomes, a narrative synthesis of the results is presented.<sup>25</sup>

## Quality appraisal

The included studies were assessed systematically for methodological quality and risk of bias using the Effective Public Health Practice Project (EPHPP) critical appraisal tool.<sup>31</sup> This tool assesses the risk of bias based on potential selection bias, study design, likelihood of confounding, blinding of outcome assessors and participants, appropriateness of data collection and completeness of follow-up. Studies were classified as high risk of potential

bias if two or more of the above categories were assessed as weak (Weak), moderate risk of potential bias if one category was assessed as weak (Moderate) and low risk of potential bias if none of the above categories were assessed as weak (Strong).

## Results

The systematic literature search identified 6,122 articles of which 786 were retrieved for abstract review. There were 43 studies that met the inclusion criteria (53 articles) and were included in the systematic review (see Figure 1, Appendix 1). A further four studies were excluded for inadequate reporting which prevented assessment of methods<sup>32, 33</sup> or outcomes,<sup>34</sup> ineligible design<sup>35</sup> or no reporting of relevant outcomes.<sup>36</sup> Thus, this review reports on the results of 39 studies (Table 1, Appendix 2).

## DESCRIPTION AND SCOPE OF THE INCLUDED STUDIES

The included studies were all RCTs. Almost half the studies (n=18) were conducted in the United States with 16 studies in a range of European countries, including five in the United Kingdom, as well as three studies in Australia and two in New Zealand. The dates of publication ranged from 1996-2014; however, 31 (79.5%) of the studies were published in the last 10 years.

The majority of the studies (82%) were undertaken in schools (n=25) or preschools (n=6) with the most common approach incorporating multiple components to promote healthy eating and/or other aspects of healthy lifestyle particularly increased physical activity (Table 1). These multiple component nutrition programs included combinations of class curriculum activities, school food service modifications, home activities, enhanced physical education/activities and strategies to engage parents/families (see Box 1). There were also four studies of school meal programs,<sup>37-40</sup> one school gardening study,<sup>41</sup> a school fruit program study<sup>42</sup> and two school canteen studies.<sup>43, 44</sup> The duration of the studies ranged from three months to the ongoing Special Turku Coronary Risk Factor Intervention Project for Children (STRIP) studies<sup>45, 46</sup> which reported follow-up

*Box 1. Approaches used in various combinations in the multi-component school/ preschool programs and family-based programs.*

### School/Preschool

- Fruit and or vegetable snacks (Free or Paid)
- School lunch program
- School garden
- Improvements in school meals/ tuckshop/ canteen facilities
- School breakfast program
- Nutrition education- Classroom
- Meal preparation sessions child (and/or parent)
- Physical activity sessions
- Change agent to support healthy nutrition/physical activity
- School food policy changes
- Observation/Rewards eating healthy foods at lunch/in class
- School-wide promotion messages
- Multimedia activities promoting healthy lifestyles
- Teacher modelling healthy eating
- Homework activities/Newsletter

### Family

- Individual and/or Group Nutrition education
- DVDs/Newsletters/Workbooks
- Non-residential camp
- Practical nutrition and/or physical activity sessions
- Internet education/activities
- Phone call from program staff



of children recruited as infants at 9 and 11 years. Twenty-three studies (59%) were between six months and two years in duration (Table 1).

Fourteen of the 25 school studies included efforts to involve families;<sup>47-60</sup> however, the uptake of this was limited in most studies where it was assessed.<sup>48-50, 52, 57</sup> There was only one school study predominantly focused on family activities; however, it was less intensive as it involved monthly nutrition or physical activity challenges.<sup>53</sup> Five of the six preschool studies involved parent-focused activities,<sup>61-65</sup> with three of these including parents in nutrition education sessions.<sup>62-64</sup>

The participants in the primary school based programs were predominantly older children aged 8-12 with many of the studies at multiple schools involving hundreds of students. There were two smaller scale programs at single schools with 60-100 participants.<sup>59, 66</sup> There were three particularly large studies: a trial of universal free school breakfast program in the USA (compared with the existing means-tested school breakfast program) assessing 4,358 students (153 schools);<sup>39</sup> an evaluation of a free school breakfast program in Wales involving 4,350 students at 111 schools (1,975 with follow-up at all time-points),<sup>38</sup> and the Child and Adolescent Trial for Cardiovascular Health (CATCH), a multi-component school program in the USA with 5,106 students (96 schools) at baseline.<sup>57</sup> The majority of the school and preschool studies involved populations that were representative of the community and thus included children from families with varying socio-economic status (SES)/educational attainment. Six studies included predominantly disadvantaged children, either by selecting schools fulfilling criteria for low SES,<sup>38, 54, 67</sup> an intervention for Native American children,<sup>50</sup> or by undertaking the study in low income rural communities.<sup>51, 53</sup>

The majority of the studies in schools and preschools involved a mixture of support for school staff, either teachers or cafeteria staff, and/or research staff to deliver enhanced nutrition education including opportunities for practical food experiences.<sup>41, 44, 47-50, 54, 56, 58, 62-69</sup>

These were usually at one or more geographically clustered schools. In addition, two studies had a major focus on teachers and/or parents as role models of healthy eating,<sup>63, 69</sup> while another study involved high school students as facilitators of the nutrition education.<sup>59</sup> Six studies instituted large scale programs- either developed directly with the input of the education or health department,<sup>55, 60, 70</sup> or instituted with low-cost resources or requiring low intensity input by school staff that was consistent with the local school curriculum.<sup>52, 57, 61</sup>

The three publicly funded programs included Food Dudes in United Kingdom primary schools,<sup>55</sup> Project Energise in primary schools in Waikato, New Zealand<sup>60</sup> and Munch and Move in preschools in New South Wales, Australia.<sup>70</sup> In addition, there were studies evaluating publicly funded national school meals programs- breakfast programs in the USA<sup>39</sup> and Wales<sup>38</sup> and a school fruit program in Norway.<sup>42</sup> There were also smaller research trials of a school breakfast program in New Zealand<sup>40</sup> and a school lunch program in Denmark.<sup>37</sup>

There were only eight family-based studies (Table 1) and these predominantly targeted the parents or parents together with their children.<sup>20, 46, 71-76</sup> Four of the five studies aimed at parents involved monthly (or less frequent) nutrition education for individuals or groups together with individual assessment.<sup>46, 72, 73, 76</sup> However, the two studies involving parents with their children incorporated practical nutrition or physical activity sessions one to three times a week.<sup>74, 75</sup> The only study to directly target children involved a 4-week summer camp program in the USA with internet follow-up for the children and their parents.<sup>71</sup> These family-based studies had relatively small numbers of participants except for an Australian study involving parent groups for those with infants<sup>72</sup> and the STRIP.<sup>46</sup> The recruitment strategies used in these studies meant that unless a disadvantaged population was explicitly targeted,<sup>74, 75</sup> then participants were predominantly not disadvantaged.<sup>71-73, 76</sup>

## OVERALL METHODOLOGICAL ASSESSMENT

Of the 39 RCTs, 15 (38%) were assessed to be at low risk of bias, 15 were assessed to have a moderate risk of bias and nine to have a high risk of bias (Table 1). Underlying

theories and/or clear rationale for interventions were common among the included studies. The development of 26 of the studies was based on one or more specific theories of behaviour most commonly social cognitive theory (n=13) and social learning theory (n=7). In addition, the five school meals programs were developed with a clear rationale. There was no clear relationship between a theoretical basis or rationale and effectiveness of the intervention.

Nutritional intake was assessed using standardised self-reported (or parent-report) measures in 29 of the 38 studies which reported these data (76%) with two of these incorporating assistance by research staff and one using measured self-reporting. Five studies (13%) used measurement/ observation by research staff and four (11%) used a combination of self-report and observation. In addition, most of the studies reported standard categories of macro- and micro-nutrient intake alongside other outcomes, which increases the likelihood of finding statistically significant differences by chance alone due to multiple comparisons.<sup>77</sup> Finally, only 12 studies (31%) reported on follow-up at a time after the intervention, and of these only three reported long-term follow-up greater than one year (Table1).

Despite these potential methodological limitations, the consistent results from these RCTs and the observation that most appeared adequately powered to detect relevant improvements in health outcomes for individuals and populations provides support for the findings of this review.

## OUTCOMES

The focus of this review and hence the most frequent outcome reported was dietary intake, with all but one study reporting on dietary intake through at least one measure. The most commonly reported outcomes were fruit and vegetable intake and fat intake. The intake of energy-dense nutrient-poor (EDNP) foods was another focus analysed in fewer studies. These are reported in detail below.

Many of these programs also aimed to promote healthy lifestyles. Hence in addition to dietary intake, physical activity levels and/or sedentary behaviour was reported by 15 studies (12 school studies and three family studies), screen-time in six studies and body mass index was reported by 19 studies (12 school studies and all seven family studies). Analysis of these outcomes was not undertaken. Other biomedical outcomes were reported in only four studies, which were generally the longer-term studies and/or more intensive studies aimed at reduction of fat intake and adiposity. The determinants of improved dietary intake were reported by a limited number of studies- attitudes and knowledge in five school studies, self-efficacy in two studies and fruit and vegetable (F&V) access in one study. Adverse effects were not reported in these studies.

There were 93 effect size estimates extracted or calculated in the 39 included studies. Thirty-one (33%) of these effect sizes (Cohen's d) were greater than 0.2 and had a 95% confidence interval that did not include the null value. However, there were also five studies which reported no impact on intake of energy-dense, nutrient-poor foods and three studies which reported no effect on at least one biomedical outcome, and no effect size estimates were calculated. The 31 positive effects were on increasing F&V intake and reducing fat intake; however there was no obvious relationship between study components and effectiveness. The impact of involving parents in school/preschool studies was only systematically assessed in the CATCH study,<sup>57</sup> which did not find a difference in outcomes between the school-based intervention with or without parental involvement.

### Impact on Fruit and Vegetable (F&V) intake

Twenty-five studies reported on F&V intake (Table 2). Of these studies, five reported a null effect on fruit and/or vegetable intake.<sup>41, 47, 49, 52, 72</sup> There were eleven studies which found a null to small effect on F&V intake.<sup>42, 44, 48, 51, 54, 61, 65, 67, 70, 76, 78</sup> Five studies reported a moderate

effect on F&V intake<sup>43, 55, 59, 62, 66</sup> and four studies which found a large effect on fruit and/or vegetable intake.<sup>20, 56, 69, 71</sup> In addition, two studies which reported a null to small effect on vegetable or overall F&V intake, also reported a moderate to large effect on fruit intake.<sup>47, 54</sup> Ten of the 25 studies were assessed to be at low risk of bias (Table 1) with eight of these reporting null to small effect on F&V intake.

There were two family-based studies and five school studies that reported a medium or large effect on F&V intake and were at low or moderate risk of bias. Five of these studies were based on social cognitive theory or social learning theory and involved high intensity interventions and/or innovative strategies to engage children. The family-based studies included the The Girls health Enrichment Multi-site Studies (GEMS) study involving eight year-old girls at a four week summer camp in the USA with weekly internet sessions for girls and their parents afterwards for eight weeks.<sup>71</sup> This study incorporated activities to promote F&V intake and physical activity into the regular camp program.

The other family-based study involved obese parents and their non-obese children in a comprehensive weight-control program for the parents comparing the impact of promoting F&V or low fat/sugar diets.<sup>20</sup> Both of these studies were delivered with high fidelity and had high follow-up albeit in small sample size. The five school studies included one which compared exposure to a teacher modelling eating fruit (or other healthy snacks) daily or healthy lifestyle curriculum with a control group.<sup>69</sup> Both of the intervention groups increased fruit intake at the end of the one year intervention. However, only the exposure group maintained this increased fruit intake one year later. There was also a lunch rewards program with observation three days/week.<sup>43</sup> Another school curriculum based program was of low intensity but involved high school students to deliver the program to younger children.<sup>59</sup> All three of these were delivered with high fidelity and achieved high follow-up. There were two other school programs, one involving daily videos and rewards together with F&V provision<sup>55</sup> and the other involving four physical activity and two nutrition education sessions weekly.<sup>66</sup>

However, the fidelity of these two interventions and the follow-up achieved was not clear. Three of these five school studies<sup>55, 59, 66</sup> involved parents in some way while the other two had no parental involvement;<sup>43, 69</sup> however, there was no systematic assessment of the impact of involving parents in any of these studies. (Table 2)

Overall, there is some evidence that both school-based studies and family-based studies, if designed and implemented well, have a positive impact on F&V intake, particularly fruit. These successful programs incorporated engaging and innovative strategies including multimedia programs to motivate the children. This impact was demonstrated for the duration of the intervention and for up to 12 months post-intervention follow-up. For example, the one study of a school garden program found no impact on F&V intake overall, but did report a significant increase in F&V intake in students at schools where a successful school garden was achieved.<sup>41</sup> However, the majority of the studies had a null or small effect on F&V intake.

### Impact on Fat intake

Fifteen studies reported fat outcomes (Table 3, Appendix 4). Of these studies, six had a null effect,<sup>37, 41, 47, 51, 52, 56</sup> four had a null to small effect,<sup>39, 50, 57, 66</sup> three had a moderate effect<sup>53, 75, 78</sup> and two studies a large effect on dietary fat intake.<sup>45, 73</sup> Five of the 15 studies were assessed to have a low risk of bias (Table 1) with one of these reporting a null effect, three a small effect and one a moderate effect on fat intake.

The two studies with a large effect had reduction in fat as a main aim of the study, whereas the majority of the other studies focused on fruit and vegetable intake. Hendrie et al.<sup>73</sup> targeted parents of school-aged children to promote low fat dairy products. This study involved a low intensity intervention in a relatively small sample size; however it was delivered with high fidelity, based on social learning theory and a high follow-up was achieved. In addition to the reduction in fat intake, there was a small change in LDL cholesterol, but no change in BMI. The STRIP study<sup>45, 46</sup> is an ongoing, relatively low

intensity program in which families with infants were recruited to attend twice yearly nutrition education and assessment promoting a low saturated fat diet, predominately *via* parents. This study also had high fidelity, but was not based on any explicit theory. These STRIP studies reported on the follow-up of a small sub-sample, aged 9–11 years at follow up. In addition to these two family-based studies, four of the eight school studies involving parents<sup>50, 53, 57, 66</sup> (all assessed to have a low risk of bias) reported smaller reductions in fat intake of children, although the only study which tested the impact of parental involvement systematically reported no additional benefit compared to no parental component.<sup>57</sup> (Table 3).

Overall, there is some evidence of benefit for studies that specifically target fat intake and are set in the home/parent based intervention, although the majority of the studies had a null or very small effect.

### Impact on Energy Dense, Nutrient Poor (EDNP) foods intake

There were nine studies which reported intake of EDNP foods. Of these, three found no effect,<sup>59, 63, 70</sup> two studies reported a small effect for one outcome measure but not the other related to EDNP foods,<sup>61, 79</sup> two showed a moderate effect (although findings not statistically significant).<sup>78, 20</sup> and two studies had limited reporting of results.<sup>64, 67</sup> The studies reporting a moderate effect were family-based studies. Epstein et al. recruited obese parents, with non-obese children, to a weight control program, with a reduction in EDNP foods among the children ( $d=-0.51$  95%CI -1.30, 0.27) for those in the arm that focused on reducing intake of high fat/sugar foods (compared with the F&V arm).<sup>20</sup> This study had a small sample size, moderate intensity over six months and then six months post intervention follow-up and was at moderate risk of bias. The Bounce program involved healthy lifestyle activities after school for mother-daughter pairs, with reductions in high fat foods (Cohen's  $d=0.40$   $p=0.26$ ) and sweetened beverages (Cohen's  $d=0.36$ ,  $p=0.31$ ) among girls in the intervention group.<sup>78</sup> This was also a small study with high intensity over three months and a moderate risk of bias. Given the nature of these studies, the results should be interpreted cautiously. Overall, the studies demonstrated limited evidence for benefits in reducing EDNP food intake.

### Other Health Outcomes

Five of the studies reported biomedical outcomes, although as the duration/follow-up of the studies was <3years, except for the STRIP study, the focus was on proxy measures. There was a null effect on blood pressure and lipids in the STRIP and CATCH studies.<sup>45, 57</sup> The STRIP studies found a small to moderate effect on reducing insulin resistance in 9-year old children [(Cohen's  $d=-0.16$  (95%CI -0.58, 0.27) in girls and -0.58 (95%CI -1.02, -0.13) in boys] and null to small effects on lipids.<sup>45</sup> In a later follow-up, in 11-year olds, there was a small to moderate effect on improving endothelial function [Cohen's  $d=0.11$  (95%CI -0.18, 0.39) in girls and 0.34 (95%CI 0.05, 0.63) in boys] with reduced total and LDL cholesterol in boys only.<sup>46</sup>

There was decreased blood pressure (BP) in the 10-12-year old children [systolic BP mean difference=-0.23 (95%CI -0.43, -0.02); diastolic BP mean difference=-0.14 (95%CI -0.30, 0.04)] but not the 5-7-year old children in Project Energise – a large school-based healthy lifestyle program in New Zealand.<sup>60</sup> Project Energise involved a change agent to champion the promotion of healthy nutrition and increased physical activity in curricular and extra-curricular activities in primary schools. The study also found small decreases in adiposity in 5-7-year old children only. In addition to the reductions in fat intake, Hendrie et al. found reduced LDL cholesterol mean difference=-0.15 mmol/L (95%CI -0.30, -0.01, Cohen's  $d=0.52$ ) and total cholesterol mean difference=-0.12 (95%CI -0.28, 0.05) but no changes in other plasma lipids three months after the program.<sup>73</sup> In the school lunch study in Denmark, Damsgaard et al. found no change in the metabolic syndrome score with reductions in mean blood pressure, total cholesterol, TAG and insulin resistance offset by increased waist circumference and reduced HDL cholesterol.<sup>80</sup> Of these five studies, only the CATCH study was assessed to have a low risk of bias. Thus, the limited evidence of improvements in cardiovascular disease risk factors, should also be interpreted cautiously.

## **Dietary and Health Outcomes and socio-economic status/disadvantaged populations**

The five school-based studies and two family-based studies which were targeted at disadvantaged populations (Table 1) did not clearly demonstrate that these programs were more or less effective in these populations compared to the overall assessments described above. Four of the studies reported on F&V intake with one reporting a large effect,<sup>54</sup> one a moderate effect<sup>78</sup> and two a small effect<sup>51, 67</sup> (Table 2). Four of the five studies which reported on fat intake in these populations showed moderate effect<sup>50, 53, 75, 78</sup> and one showed no effect<sup>51</sup> (Table 3). The Bounce study, described above, involved disadvantaged African-American mother-daughter pairs in an urban setting and reported reduced fat intake, increased F&V and reductions in EDNP food, as noted previously.<sup>78</sup> This is consistent with the sub-group analysis reported in the three year follow-up CATCH study, which showed that overall dietary intake after this program in rural USA was the same in African-American and Hispanic-American children as in white children.<sup>81</sup>

## **SUSTAINABILITY OF NUTRITION AND HEALTH OUTCOMES**

There were four studies of three or more years duration included in the review.<sup>39, 46, 50, 57</sup> Three were school-based programs. Two were multi-component programs – the Pathways program which aimed to reduce adiposity in American Indian school children and the CATCH program which aimed to reduce risk factors for cardiovascular disease in primary school students (Table 1). Both reported small to moderate effect on reducing total fat and/or saturated fat intake after three years: Pathways Total fat intake mean difference-2.5% and CATCH Total fat intake mean difference-2% (Table 3). The other school-based program was the Universal School Breakfast Pilot Program in the USA.<sup>39</sup> This program operated successfully for three years; however, follow-up dietary intake data were only collected at 12 months. This showed that intake of a nutritionally substantive breakfast increased from 76% to 80% in intervention schools, but that breakfast skipping was 4% in both intervention and control schools. The other long-term study was the family-based STRIP study in Finland. The children participating in this program had lower intake of total and saturated fat at nine years and saturated fat at eleven years than control children, although results varied by gender (Table 3).

Long-term follow-up of programs (>1 year post-intervention) to assess the ongoing impact was reported by two studies in addition to the ongoing STRIP study described above. A three year follow-up of the CATCH study reported that these children still had significantly lower intake of total and saturated fat compared to controls although overall dietary intake was not significantly different between the two groups (based on the Healthy Eating Index).<sup>81</sup> In a three-year follow-up of the Free School Fruit program in Norway, increases in F&V intake, noted at the conclusion of the program and one year after the program, persisted although reduced from the conclusion of the program.<sup>82</sup> It was estimated that boys ate an extra 0.38 serves of F&V / day and girls 0.44 serves of F&V / day as a result of prior participation in this school fruit program compared to an extra 0.6 serves/day at the conclusion and 0.5 serves/day at one year.<sup>42</sup> Thus, there is consistent although limited evidence that ongoing multicomponent nutrition programs can improve dietary patterns sufficiently to impact on population health (if sustained) and that these effects persist, although attenuated for up to three years.

## Discussion

The majority of the 39 RCTs included in this review were in schools or preschools with only eight undertaken in family settings. There were substantial (and statistically significant) improvements in dietary intake in 31% of outcomes assessed. The positive impacts were found in increased F&V intake and reduced fat intake. Although there were effective family-based and institutional nutrition programs, there is insufficient evidence to determine the impact of involving parents in preschool/school nutrition programs. The findings do strongly support the importance of social context to healthy eating. The two studies (both directed at parents) which reported a large decrease in fat intake indicate that the promotion of simple dietary information which is well understood and engages parents is effective and enables them to have a major impact on their children's dietary intake.<sup>45, 73</sup> Similarly, school nutrition programs can improve the intake of healthy foods, particularly fruit. Based on successful outcomes observed in this review, the most effective strategies included the use of innovative strategies to engage and motivate the children including rewards, cartoon characters promoting healthy foods, modelling by teachers and the use of older peer educators. A theoretical framework, particularly the use of social cognitive theory or social learning theory, also appeared to support positive impacts of these programs. None of these features were sufficient to achieve positive outcomes; rather, it supports that the effective implementation of an innovative and well-designed intervention is more likely to improve dietary intake.

In a 2010 review, Hingle et al.<sup>83</sup> concluded that there was insufficient evidence of the impact of parental involvement in dietary interventions to improve children's dietary intake. This is consistent with our finding that although parental/family involvement was an element of most preschool/school studies, it was not central to most interventions or consistently implemented. Hingle et al. did find that more direct approaches to engage parents were more likely to have a positive impact on children's diet. The positive impacts demonstrated in the family studies in this review provide further evidence to support this finding. The use of social cognitive theory or social learning theory to enhance the effectiveness of nutrition interventions has been previously identified.<sup>83, 84</sup> Interestingly, Segal et al.<sup>85</sup> have highlighted that it is the extent to which the theory is consistent with the needs of the target population and the program components are consistent with the nominated theory which may enhance effectiveness. Thus, identifying and explicitly assessing elements of social cognitive theory (or other relevant theory) will enhance the understanding of how to design and implement an effective nutrition program.

The sustainability of the dietary improvements of successful nutrition programs is vital to achieving long-term health benefits. The three studies reporting on longer term follow-up reinforced that simple dietary interventions - promoting a lower fat diet to families (STRIP study) and a school fruit program - can produce sustained improvements in dietary intake. Ongoing large scale nutrition programs in schools/preschools are easier to implement and maintain provided funding and support is available. This was demonstrated by three school studies of three years duration including the Pathways program in Native American schools,<sup>50, 57</sup> and the US free school breakfast program.<sup>39</sup> There is also the large scale implementation of other preschool and school-based programs, such as Food Dudes in the UK,<sup>86</sup> Project Energise in New Zealand<sup>60</sup> and Munch and Move in Australia.<sup>70</sup> It is relevant to consider how practical it is to implement complex nutrition programs in other settings. Published articles cannot include the details of educational resources or activities undertaken and there are resource implications of making this material available online. It is also difficult to convey or replicate the interpersonal interactions within program activities that may contribute to the intervention's effectiveness. The Food Dudes program demonstrates the sustained commitment and resources required to implement a successful nutrition intervention in other locations.<sup>86, 87</sup>

From a population perspective, the sustainability of nutrition programs must also be weighed against the opportunity cost implicit in funding such an ongoing program. Overall, these programs have only a modest impact on dietary intake, and alone are unlikely to mitigate the

many changes within contemporary society which have undermined healthy dietary intake and lifestyles. Other strategies which help to create an environment which supports people's health and nutrition, including restricting junk food advertising or enhancing the availability of healthier food choices, need to be compared with the impact of both school and family nutrition programs. Considered assessment of these strategies, including economic analysis, will help to develop a range of effective programs that together can support healthy eating in the population.

The potential to improve dietary intake is relevant to most of the population. This review found that school and community nutrition programs undertaken in disadvantaged communities were as effective as interventions in other settings and thus should impact dietary intake sufficiently to improve health outcomes in these populations. However, there was limited reporting of results by socio-economic status (SES) in the larger school studies, unless it was relevant to the rationale for the intervention, such as school meal programs. It was also observed that family-based programs recruited predominantly families with above average SES, unless undertaken explicitly targeting a disadvantaged community. Thus, the evidence suggests that child nutrition programs are beneficial in all children irrespective of SES, although it is important to consider strategies to maximise disadvantaged families' participation in effective family-based nutrition programs.

The most important limitation in this review was the validity and reliability of the self-reported dietary intake data. Although the studies used standardised methodologies predominantly 24-hour recalls and FFQs, the potential for bias and the challenge of precision is well recognised.<sup>88</sup> It is particularly challenging for young children to estimate quantities of foods consumed and the role of parents is limited when children have consumed food separately from their parents. A smaller number of studies used direct observation and measurement by researchers, which are also limited in the capacity to assess all foods eaten throughout the day. The use of more objective measures to assess dietary intake, including the use of biomarkers and electronic shopping data can supplement and strengthen the conclusions that can be drawn from dietary self-report measures.<sup>89, 90</sup> These dietary self-report measures remain central to studies of nutritional interventions; hence attention to standardised reporting of these results facilitates the synthesis required in a systematic review. A further limitation was the inclusion of English language articles only, although only eight potentially eligible studies were excluded on the basis of language.

The evidence from this review indicates that schools have been shown to be an effective setting to improve primary school aged children's dietary intake. Although the programs have been shown to be feasible to conduct on an ongoing basis, there is still limited evidence of the long-term benefits. Thus, careful evaluation of both the effectiveness and cost-effectiveness of such programs remains important. Given the moderate impact, it is relevant to consider how direct family-based strategies may be incorporated into these programs or operate concurrently in the same communities. Effective family-based programs have generally operated on a small scale and the feasibility of implementing these on a larger scale or multiple settings need further investigation.

## Conclusion

Family-based nutrition programs which support parents with simple nutrition information have demonstrated potential to substantially reduce dietary fat and increase F&V, while school-based nutrition programs have shown the potential to moderately increase F&V intake, particularly fruit. Effective components of the family-based programs have been simple dietary messages directed towards parents with regular follow-up, while the effective school-based programs have incorporated role-models including peers, teachers and heroic figures, rewards and increased access to healthy foods. However, there is limited evidence of the sustainability of effective family and school nutrition programs, particularly their impact on biomedical health outcomes. Given the limited impact of individual programs, complementary nutrition interventions are needed which build a supportive environment and provide the opportunities for everyone to eat healthily.



## References

1. Rangan AM, Schindeler S, Hector DJ, et al. Consumption of 'extra' foods by Australian adults: types, quantities and contribution to energy and nutrient intakes. *Eur J Clin Nutr.* 2009 Jul;63(7):865-871.
2. CSIRO Preventative Health National Research Flagship, University of South Australia. *2007 Australian National Children's Nutr and Phys Act Survey.* Main Findings Canberra: Department of Health and Ageing;2007.
3. Hall JN, Moore S, Harper SB, et al.. Global Variability in Fruit and Vegetable Consumption. *Am J Prev Med.* 2009;36(5):402-409.e405.
4. Rehm CD, Monsivais P, Drewnowski A. The quality and monetary value of diets consumed by adults in the United States. *The Am J Clin Nutr* 2011;94(5):1333-1339.
5. Hamer M, Mishra GD. Dietary patterns and cardiovascular risk markers in the UK Low Income Diet and Nutrition Survey. *Nutr Metab Cardiovascular Diseases.* 2010;20(7):491-497.
6. Kant AK. Reported consumption of low-nutrient-density foods by American children and adolescents: nutritional and health correlates, NHANES III, 1988 to 1994. *Arch Pediatr Adolesc Med.* 2003;157(8):789-796.
7. Diethelm K, Jankovic N, Moreno LA, et al. Food intake of European adolescents in the light of different food-based dietary guidelines: results of the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) Study. *Pub Health Nutr.* 2012;15(03):386-398.
8. Moore LV, Thompson FE. Adults Meeting Fruit and Vegetable Intake Recommendations—United States, 2013. *MMWR Morbidity and mortality weekly report.* 2015;64(26):709-713.
9. Bazzano LA, Li TY, Joshipura KJ, et al. Intake of fruit, vegetables, and fruit juices and risk of diabetes in women. *Diabetes Care.* 2008;31(7):1311-1317.
10. World Health Organisation. *Global Health Risks: Mortality and burden of disease attributable to selected major risks.* Geneva: World Health Organisation, Department of Health Statistics and Informatics;2009.
11. Australian Institute of Health and Welfare. *Australia's Health 2010.* Australia's health series no. 12. Cat. no. AUS 122. Canberra: Australian Institute of Health and Welfare;;2010.
12. Huang RC, Mori TA, Burke V, et al. Synergy between adiposity, insulin resistance, metabolic risk factors, and inflammation in adolescents. *Diabetes Care.* 2009 Apr;32(4):695-701.
13. Strong JP, Malcom GT, McMahan C, et al. Prevalence and extent of atherosclerosis in adolescents and young adults: Implications for prevention from the pathobiological determinants of atherosclerosis in youth study. *JAMA.* 1999;281(8):727-735.
14. Foltz JL, May AL, Belay B, et al.. Population-Level Intervention Strategies and Examples for Obesity Prevention in Children. *Annual Review of Nutr.* 2012;32:391-415.
15. Waters E, de Silva-Sanigorski A, Burford BJ, et al. Interventions for preventing obesity in children. *Cochrane Database of Syst Reviews.* 2011(12).
16. Wang D, Stewart D. The implementation and effectiveness of school-based nutrition promotion programmes using a health-promoting schools approach: a systematic review. *Pub Health Nutr.* 2013;16(6):1082-1100.
17. Evans CEL, Christian MS, Cleghorn CL, et al. Systematic review and meta-analysis of school-based interventions to improve daily fruit and vegetable intake in children aged 5 to 12 y. *Am J Clin Nutr.* 2012;96(4):889-901.
18. Hendrie GA, Brindal E, Corsini N, et al. Combined Home and School Obesity Prevention Interventions for Children: What Behavior Change Strategies and Intervention Characteristics Are Associated With Effectiveness? *Health Ed Behav.* 2012;39(2):159-171.

19. Mann J, Truswell AS. *Essentials of Human Nutrition*. 3rd ed. Oxford: Oxford University Press; 2007.
20. Epstein LH, Gordy CC, Raynor HA, et al. Increasing fruit and vegetable intake and decreasing fat and sugar intake in families at risk for childhood obesity. *Obesity Research* [serial on the Internet]. 2001; 9(3): Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/569/CN-00347569/frame.html>.
21. Hall L, Collins CE, Morgan PJ, et al. Children's intake of fruit and selected energy-dense nutrient-poor foods is associated with fathers' intake. *J Am Diet Assoc*. 2011 Jul;111(7):1039-1044.
22. Golley RK, Hendrie GA, Slater A, et al. Interventions that involve parents to improve children's weight-related nutrition intake and activity patterns - what nutrition and activity targets and behaviour change techniques are associated with intervention effectiveness? *Obes Rev*. 2011 Feb;12(2):114-130.
23. Hendrie GA, Brindal E, Corsini N, et al. Combined home and school obesity prevention interventions for children: what behavior change strategies and intervention characteristics are associated with effectiveness? *Health Ed Behav*. 2012 Apr;39(2):159-171.
24. Van Lippevelde W, Verloigne M, De Bourdeaudhuij I, et al. Does parental involvement make a difference in school-based nutrition and physical activity interventions? A systematic review of randomized controlled trials. *Int J Pub Health*. 2012/08/01;57(4):673-678.
25. Higgins JPT, Green S, (eds). *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0* [updated March 2011]. The Cochrane Collaboration; 2011. Available from: [www.cochrane-handbook.org](http://www.cochrane-handbook.org).
26. Armstrong R, Waters E, Jackson N, et al. *Guidelines for Systematic reviews of health promotion and public health interventions*. Version 2. Australia: Melbourne University; 2007.
27. Kristjansson B, Petticrew M, MacDonald B, et al. School feeding for improving the physical and psychosocial health of disadvantaged students. *Cochrane Database of Syst Reviews* [serial on the Internet]. 2007; (1): Available from: <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD004676.pub2/abstract>.
28. Kothandan SK. School based interventions versus family based interventions in the treatment of childhood obesity- a systematic review. *Arch Pub Health*. 2014;72(1):3.
29. Collins CE, Burrows TL, Bray J, et al. Effectiveness of parent-centred interventions for the prevention and treatment of childhood overweight and obesity in community settings: A systematic review. *JBI Database of Syst Reviews and Implem Reports*. 2013;11 (9):180-257.
30. Wilson DB. *Practical Meta-Analysis Effect Size Calculator*. Oslo, Norway: The Campbell Collaboration; 2001 [20/6/2015]; Available from: [http://www.campbellcollaboration.org/resources/effect\\_size\\_input.php](http://www.campbellcollaboration.org/resources/effect_size_input.php).
31. National Collaborating Centre for Methods and Tools. *Quality Assessment Tool for Quantitative Studies*. Hamilton, Ontario: McMaster University; 2008 [updated 13 April, 2010; 14/2/2015]; Available from: <http://www.nccmt.ca/registry/view/eng/14.html>
32. Reynolds KD, Raczynski JM, Binkley D, et al. Design of "High 5": a school-based study to promote fruit and vegetable consumption for reduction of cancer risk. *J Cancer Ed*. 1998;13(3):169-177.
33. Reynolds KD, Franklin FA, Binkley D, et al. Increasing the fruit and vegetable consumption of fourth-graders: results from the high 5 project. *Preventive Med* [serial on the Internet]. 2000; 30(4): Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/941/CN-00371941/frame.html>.
34. Nader PR, Sallis JF, Patterson TL, et al. A family approach to cardiovascular risk reduction: results from the San Diego Family Health Project. *Health Ed Q*. 1989 1989;16 (2):229-244.

35. Moore L, Tapper K. The impact of school fruit tuck shops and school food policies on children's fruit consumption: a cluster randomised trial of schools in deprived areas. *J Epidem & Community Health*. 2008 Oct;62(10):926-931.
36. Zask A, Adams JK, Brooks LO, et al. Tooty Fruity Veggie: an obesity prevention intervention evaluation in Australian preschools. *Health promotion J Austr* [serial on the Internet]. 2012; 23(1): Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1440-1788.2012.02903.x>
37. Andersen R, Biloft-Jensen A, Christensen T, et al. Dietary effects of introducing school meals based on the New Nordic Diet – a randomised controlled trial in Danish children. The OPUS School Meal Study. *Br J Nutr*. 2014;111(11):1967-1976.
38. Moore L, Moore GF, Tapper K, et al. Free breakfasts in schools: design and conduct of a cluster randomised controlled trial of the Primary School Free Breakfast Initiative in Wales [ISRCTN18336527]. *BMC Pub Health*. 2007;7:258.
39. Crepinsek MK, Singh A, Bernstein LS, et al. Dietary Effects of Universal-Free School Breakfast: Findings from the Evaluation of the School Breakfast Program Pilot Project. *J Am Dietet Assoc*. 2006 November;106 (11):1796-1803.
40. Ni Mhurchu C, Gorton D, Turley M, et al. *Effects of a free school breakfast programme on children's attendance, academic achievement, and short-term hunger: A stepped-wedge, cluster randomised controlled trial*. Australasian Medical Journal. 2011;Conference: 35th Annual Scientific Meeting Joint Annual Scientific Meeting of the Nutrition Society of New Zealand and the Nutrition Society of Australia Queenstown New Zealand. Conference Start: 20111129 Conference End: 20111202. Conference Publication: (var.pagings). 4 (12):805.
41. Christian MS, Evans CEL, Nykjaer C, et al. Evaluation of the impact of a school gardening intervention on children's fruit and vegetable intake: A randomised controlled trial. *Int J Behav Nutr Phys Act*. 2014 August 16;11 (1)(99).
42. Bere E, Veierød MB, Bjelland M, et al.. Free school fruit--sustained effect 1 year later. *Health Ed Research*. 2006;21(2):268-275.
43. Hendy HM, Williams KE, Camise TS. Kid's Choice Program improves weight management behaviors and weight status in school children. *Appetite*. 2011 Apr;56(2):484-494.
44. Perry CL, Bishop DB, Taylor GL, et al.. A randomized school trial of environmental strategies to encourage fruit and vegetable consumption among children. *Health Ed Behav: the official publication of the Society for Pub Health Ed*. 2004 Feb;31 (1):65-76.
45. Kaitosaari T, Rönnemaa T, Viikari J, et al. Low-saturated fat dietary counseling starting in infancy improves insulin sensitivity in 9-year-old healthy children: the Special Turku Coronary Risk Factor Intervention Project for Children (STRIP) study. *Diabetes Care*. 2006;29(4):781-785.
46. Raitakari OT, Rönnemaa T, Järvisalo MJ, et al. Endothelial function in healthy 11-year-old children after dietary intervention with onset in infancy: the Special Turku Coronary Risk Factor Intervention Project for children (STRIP). *Circulation*. 2005;112(24):3786-3794.
47. Anderson AS, Porteous LEG, Foster E, et al. The impact of a school-based nutrition education intervention on dietary intake and cognitive and attitudinal variables relating to fruits and vegetables. *Pub Health Nutr*. 2005;8(6):650-656.
48. Baranowski T, Davis M, Resnicow K, et al. Gimme 5 fruit, juice, and vegetables for fun and health: outcome evaluation.[Erratum appears in Health Educ Behav 2000 Jun;27(3):390]. *Health Ed Behav*. 2000 Feb;27(1):96-111.
49. Bere E, Veierød MB, Bjelland M, et al. Outcome and process evaluation of a Norwegian school-randomized fruit and vegetable intervention: Fruits and Vegetables Make the Marks (FVMM). *Health Ed Research*. 2006;21(2):258-267.
50. Caballero B, Clay T, Davis SM, et al. Pathways: a school-based, randomized controlled trial for the prevention of obesity in American Indian schoolchildren. *Am J Clin Nutr*. 2003;78(5):1030-1038.

51. Cohen JFW, Kraak VI, Choumenkovitch SF, et al. The CHANGE study: a healthy-lifestyles intervention to improve rural children's diet quality. *J Academy Nutr Dietet*. 2014 Jan;114(1):48-53.
52. Evans CE, Ransley JK, Christian MS, et al. A cluster-randomised controlled trial of a school-based fruit and vegetable intervention: Project Tomato. *Pub Health Nutr* [serial on the Internet]. 2013; 16(6): Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/416/CN-00966416/frame.html>.
53. Greening L, Harrell KT, Low AK, et al. Efficacy of a school-based childhood obesity intervention program in a rural southern community: TEAM Mississippi Project. *Obesity* (Silver Spring, Md) [serial on the Internet]. 2011; 19(6): Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/952/CN-00801952/frame.html>.
54. Hoffman JA, Franko DL, Thompson DR, et al. Longitudinal behavioral effects of a school-based fruit and vegetable promotion program. *J Pediatr Psych*. 2010;35 (1):61-71.
55. Horne PJ, Hardman CA, Lowe CF, et al. Increasing parental provision and children's consumption of lunchbox fruit and vegetables in Ireland: the Food Dudes intervention. *Eur J Clin Nutr*. 2009 May;63(5):613-618.
56. Kristjansdottir AG, Johannsson E, Thorsdottir I. Effects of a school-based intervention on adherence of 7-9-year-olds to food-based dietary guidelines and intake of nutrients. *Pub Health Nutr*. 2010;13(8):1151-1161.
57. Luepker RV, Perry CL, McKinlay SM, et al. Outcomes of a field trial to improve children's dietary patterns and physical activity. The Child and Adolescent Trial for Cardiovascular Health. CATCH collaborative group. *JAMA* [serial on the Internet]. 1996; 275(10): Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/560/CN-00123560/frame.html>.
58. Llargues E, Franco R, Recasens A, et al. Assessment of a school-based intervention in eating habits and physical activity in school children: The AVall study. *J Epidem Community Health*. 2011 October;65 (10):896-901.
59. Muth ND, Chatterjee A, Williams D, et al. Making an IMPACT: effect of a school-based pilot intervention. *North Carolina Med J*. 2008 Nov-Dec;69(6):432-440.
60. Rush E, Reed P, McLennan S, et al.. A school-based obesity control programme: Project Energize. Two-year outcomes. *Br J Nutr*. 2012;107(4):581-587.
61. Bayer O, von Kries R, Strauss A, et al. Short- and mid-term effects of a setting based prevention program to reduce obesity risk factors in children: a cluster-randomized trial. *Clin Nutr*. 2009;28(2):122-128.
62. De Bock F, Breitenstein L, Fischer JE. Positive impact of a pre-school-based nutritional intervention on children's fruit and vegetable intake: results of a cluster-randomized trial. *Pub Health Nutr*. 2012 Mar;15(3):466-475.
63. Natale R, Scott SH, Messiah SE, et al. Design and methods for evaluating an early childhood obesity prevention program in the childcare center setting. *BMC Pub Health* [serial on the Internet]. 2013; 13: Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/201/CN-00864201/frame.html>.
64. Natale RA, Lopez-Mitnik G, Uhlhorn SB, et al. Effect of a child care center-based obesity prevention program on body mass index and nutrition practices among preschool-aged children. *Health Promot Pract*. 2014 Sep;15(5):695-705.
65. Vereecken C, Huybrechts I, van Houte H, et al. Results from a dietary intervention study in preschools "Beastly Healthy at School". *Int J Pub Health*. 2009;54(3):142-149.
66. Hopper CA, Munoz KD, Gruber MB, et al.. A school-based cardiovascular exercise and nutrition program with parent participation: an evaluation study. *Children's Health Care*. 1996;25(3):221-235.

67. Breslin G, Brennan D, Rafferty R, et al. The effect of a healthy lifestyle programme on 8-9 year olds from social disadvantage. *Arch Disease in Childhood*. 2012;97(7):618-624.
68. Kain J, Uauy R, Taibo M. Chile's school feeding programme: Targeting experience. *Nutr Research*. 2002;22 (5):599-608.
69. Perikkou A, Gavrieli A, Kougioufa MM, et al.. A Novel Approach for Increasing Fruit Consumption in Children. *J Acad Nutr Dietet*. 2013 September;113 (9):1188-1193.
70. Hardy LL, King L, Kelly B, et al. Munch and Move: Evaluation of a preschool healthy eating and movement skill program. *Int J Behav Nutr Phys Act*. 2010 03 Nov;7(80).
71. Baranowski T, Baranowski JC, Cullen KW, et al. The Fun, Food, and Fitness Project (FFFP): the Baylor GEMS pilot study. *Ethnicity & Disease* [serial on the Internet]. 2003; 13(1 Suppl 1): Available from:  
<http://onlinelibrary.wiley.com/doi/cochrane/central/articles/929/CN-00436929/frame.html>.
72. Cameron AJ, Ball K, Hesketh KD, et al. Variation in outcomes of the Melbourne Infant, Feeding, Activity and Nutrition Trial (InFANT) Program according to maternal education and age. *Prevent Med*. 2014 January;58 (1):58-63.
73. Hendrie GA, Golley RK. Changing from regular-fat to low-fat dairy foods reduces saturated fat intake but not energy intake in 4-13-y-old children. *Am J Clin Nutr*. 2011;93(5):1117-1127.
74. Olvera NN, Knox B, Scherer R, et al. A Healthy Lifestyle Program for Latino Daughters and Mothers. *Am J Health Ed*. 2008 2008/09/01;39(5):283-295.
75. Stolley MR, Fitzgibbon ML. Effects of an obesity prevention program on the eating behavior of African American mothers and daughters. *Health Ed Behav* [serial on the Internet]. 1997; 24(2): Available from:  
<http://onlinelibrary.wiley.com/doi/cochrane/central/articles/938/CN-00137938/frame.html>.
76. Tabak RG, Tate DF, Stevens J, et al. Family ties to health program: a randomized intervention to improve vegetable intake in children. *J Nutr Ed Behav* [serial on the Internet]. 2012; 44(2): Available from:  
<http://onlinelibrary.wiley.com/doi/cochrane/central/articles/983/CN-00832983/frame.html>.
77. Sterne JA, Davey Smith G. Sifting the evidence-what's wrong with significance tests? *BMJ*. 2001 Jan 27;322(7280):226-231.
78. Olvera N, Bush JA, Sharma SV, et al. BOUNCE: a community-based mother-daughter healthy lifestyle intervention for low-income Latino families. *Obesity* (Silver Spring, Md) [serial on the Internet]. 2010; 18 Suppl 1: Available from:  
<http://onlinelibrary.wiley.com/doi/cochrane/central/articles/806/CN-00729806/frame.html>.
79. Murphy S, Moore G, Tapper K, et al. Free healthy breakfasts in primary schools: a cluster randomised controlled trial of a policy intervention in Wales, UK. *Pub Health Nutr*. 2011;14(2):219-226.
80. Damsgaard CT, Dalskov SM, Laursen RP, et al. Provision of healthy school meals does not affect the metabolic syndrome score in 8-11-year-old children, but reduces cardiometabolic risk markers despite increasing waist circumference. *Br J Nutr*. 2014 Dec 14;112(11):1826-1836.
81. Dwyer J, Cosentino C, Li D, et al. Evaluating school-based interventions using the Healthy Eating Index. *J Am Dietetic Assoc*. 2002 Feb;102(2):257-259.
82. Bere E, Veierod MB, Skare O, et al. Free School Fruit--sustained effect three years later. *Int J Behav Nutr Phys Act*. 2007;4:5.
83. Hingle MD, O'Connor TM, Dave JM, et al. Parental Involvement in Interventions to Improve Child Dietary Intake: A Systematic Review. *Preventive Med*. 2010 05/10;51(2):103-111.
84. Nyberg G, Sundblom E, Norman A, et al. Effectiveness of a universal parental support programme to promote healthy dietary habits and physical activity and to prevent

- overweight and obesity in 6-year-old children: the Healthy School Start Study, a cluster-randomised controlled trial. *PLoS One*. 2015;10(2):e0116876.
85. Segal L, Sara Opie R, Dalziel K. Theory! The missing link in understanding the performance of neonate/infant home-visiting programs to prevent child maltreatment: a systematic review. *Milbank Q*. 2012 Mar;90(1):47-106.
  86. Erjavec M, Viktor S, Horne P, et al. Implementing a healthy eating programme: changing children's eating habits for life. *Community Practitioner*. 2012;85(4):39-42.
  87. National Social Marketing Centre. *Food Dudes Overview*. National Social Marketing Centre; 2010 [updated 04/04/2011 Accessed 6/02/2016]; Available from: <http://www.thensmc.com/resources/showcase/food-dudes>.
  88. Gibson RS. *Principles of nutritional assessment*. 2nd ed. Oxford: Oxford University Press; 2005.
  89. Ni Mhurchu C, Blakely T, Jiang Y, et al. Effects of price discounts and tailored nutrition education on supermarket purchases: a randomized controlled trial. *Am J Clin Nutr*. 2010 Mar;91(3):736-747.
  90. Black AP, Vally H, Morris P, et al.. Nutritional impacts of a fruit and vegetable subsidy programme for disadvantaged Australian Aboriginal children. *Br J Nutr*. 2013;110(12):2309-2317.
  91. Bere E, Hilsen M, Klepp KI. Effect of the nationwide free school fruit scheme in Norway. *Br J Nutr*. 2010;104(4):589-594.
  92. Davis SM, Clay T, Smyth M, et al. Pathways curriculum and family interventions to promote healthful eating and physical activity in American Indian schoolchildren. *Preventive Med*. 2003;37(6 part 2):S24-34.
  93. Himes JH, Ring K, Gittelsohn J, et al. Impact of the Pathways intervention on dietary intakes of American Indian schoolchildren. *Preventive Med* [serial on the Internet]. 2003; 37(6 Pt 2): Available from: <http://onlinelibrary.wiley.com/doi/10.1016/j.ypmed.2003.08.004>.
  94. Campbell KJ, Lioret S, McNaughton SA, et al. A Parent-Focused Intervention to Reduce Infant Obesity Risk Behaviors: A Randomized Trial. *Pediatr*. 2013 April 1, 2013;131(4):652-660.
  95. Hendrie G, Sohonpal G, Lange K, et al. Change in the family food environment is associated with positive dietary change in children. *Int J Behav Nutr Phys Act*. 2013 07 Jan;10(4).
  96. Moore GF, Murphy S, Chaplin K, et al. Impacts of the Primary School Free Breakfast Initiative on socio-economic inequalities in breakfast consumption among 9-11-year-old schoolchildren in Wales. *Pub Health Nutr*. 2014 Jun;17(6):1280-1289.
  97. Natale RA, Messiah SE, Asfour L, et al. Role modeling as an early childhood obesity prevention strategy: effect of parents and teachers on preschool children's healthy lifestyle habits. *J Dev Behav Pediatr*. 2014 Jul-Aug;35(6):378-387.

## Appendix 1: Figure 1, Systematic literature review search strategy and results




Figure 1: Literature search strategy and results

## Appendix 2: Table 1, Characteristics of included studies

Study, Location	Setting, Disadvantaged population (if indicated)	Interventions	Participants: Age during intervention	Sample size †	Sample size at Follow-up †	Duration of intervention	Follow-up duration post-intervention	Health & Nutrition Outcomes	Potential risk of bias rating
Andersen 201437 OPUS school meal study  Denmark	School	School meals (free)	Years 3 & 4 Mean age 10 years	834 no breakdown by I/C	663	6 month crossover trial	0	Dietary intake (self-report)	M
Anderson 200547  Scotland	School	School multicomponent	Year 2 & Year 6 classes 6-7 years & 10-11 years	I=158 C=136	I=64 C=65	9 months	0	Dietary intake (self-report) Attitudes & knowledge of healthy eating	W
Baranowski 200048 Gimme 5  USA	School	School multicomponent-comprehensive	Years 3-5	1864 no breakdown by I/C	1253	2 years	0	Dietary intake (self-report) F&V preferences, Self-efficacy	S
Baranowski 200371 GEMS Fun, Food and Fitness  USA	Family	Summer camp/ Internet F/U for family	8 years	I=19 C=16	I=17, C=14	3 months	<1mth	Dietary intake (self-report), BMI, PA levels (accelerometry, self-report)	M
Bayer 200961 Tigerkids Germany	Preschool	Preschool - multicomponent	Preschool Mean age 6 years	I=1049 C=560, I2=1040 C2=565	I= 850, C=469; I2=872 C2=468	2 years	0	Dietary intake (Parent-report), BMI	S
Bere 2006a,2007, 201042, 81, 90 Free School Fruit Norway	School	School-Free Fruit & multicomponent	Year 6/7 Mean age 11.3 years	577 no breakdown by IC	I=286 C=231	2 years (only 1 year free fruit)	1 year, 3years	Dietary intake (self-report), F&V preferences	S



Study, Location	Setting, Disadvantaged population (if indicated)	Interventions	Participants: Age during intervention	Sample size †	Sample size at Follow-up †	Duration of intervention	Follow-up duration post-intervention	Health & Nutrition Outcomes	Potential risk of bias rating
Bere 2006b49 F&V Makes the Marks Norway	School	School- Paid fruit & multicomponent	Years 6/7 Mean age 11.3 years	450 no breakdown by I/C	I=190 C=179	1 year	1 year	Dietary intake (self-report), F&V accessibility, preferences, modelling	S
Breslin 201267 Sport For Life Northern Ireland	School Low SES schools	School multicomponent	Year 3 8-9 years	I=209 C=207	?	3 months	1 month	Dietary intake (self-report), BMI, PA levels & Screen-time (accelerometry, self-report)	S
Caballero 200350 Davis 200391, Himes 200392 Pathways study USA	School Native American community schools	School multicomponent	Year 3 Mean age 7.6 years	I=879 C=825	I=727, C=682	3 years	0	Dietary intake (self-report, direct observation), BMI, PA levels (accelerometry, self-report)	S
Cameron 201472 Campbell 201393 Melbourne InFANT  Australia	Family	Parent groups	Mother-infant pairs Infants 4-15 months	I=271 C=271	I= 241 C=239	18 months	0	Dietary intake (parent report), BMI, PA levels & Screen-time (accelerometry)	S
Christian 201441 England	School	School gardening	Primary school ages 7-11	I=529 C=727	I=312 C=329	2 years	0	Dietary intake (observer & parent-report)	W
Cohen 201451 CHANGE study  USA	School Low income rural community	School multicomponent	Primary school Years 1-6	1302 no breakdown by I/C	I=251 C=181	2 years	0	Dietary intake (Assisted self-report)	M

Study, Location	Setting, Disadvantaged population (if indicated)	Interventions	Participants: Age during intervention	Sample size †	Sample size at Follow-up †	Duration of intervention	Follow-up duration post-intervention	Health & Nutrition Outcomes	Potential risk of bias rating
Crepinsek 200639 Universal Free School Breakfast Program USA	School	Free school breakfast program	Primary School Years 2-6	4358 no breakdown by I/C	I=2212 C=2066	3 years- Evaluation done at 12 months	0	Dietary intake (parent & self-report)	S
De Bock 201262 Germany	Preschool	Preschool - multicomponent	Preschool Mean age 4.3 years	I=194 C=183	202 No breakdown by IC	6 months	6 months	Dietary intake (parent report), BMI	W
Epstein 200120 USA	Family	Family - Parent sessions	30 families- 1 parent & 1 child Mean age 8.7 years	I=15 I2=15	27 No breakdown by IC	6 months	6 months	Dietary intake (parent & self-report), BMI	M
Evans 201352 Project Tomato England	School	School multicomponent	Year 3 (7-8yrs)	I=530 C=550	I=331 C=347	10 months	10 months	Dietary intake (observer & parent-report)	S
Greening 201153 TEAM Mississippi Project USA	School Low SES rural community	School/Family	6-10 years Mean age 8.3 years	I=224 C=283	I=204 C=246	8 months	0	Dietary intake (parent-report), Fitness & PA levels	S
Hardy 201070 Munch & Move Australia	Preschool	Preschool - multicomponent	Mean age 4.5yrs	I=263 C=167	I=218 C=141	5 months	0	Lunch foods (Measurement), Movement skills	M
Hendrie 201173 Hendrie 201394 Australia	Family	Family - Parent sessions	4-13 year old children Mean age 8.6yrs	I=76 C=69	I=76 C=64	3 months	3 months	Dietary intake (parent & self-report), BMI, Plasma lipids	M

Study, Location	Setting, Disadvantaged population (if indicated)	Interventions	Participants: Age during intervention	Sample size †	Sample size at Follow-up †	Duration of intervention	Follow-up duration post-intervention	Health & Nutrition Outcomes	Potential risk of bias rating
Hendy 201143 Kid's Choice Program USA	School	School rewards program (Lunch)	Years 1-4	457 Both groups	382 both groups	3 months	0	Dietary intake (observation), BMI, PA levels (Pedometer)	S
Hoffman 201054 USA	School Low SES urban community	School multicomponent	Kinder, Year 1 Mean age 6.2 years	I=149 C=148	I=113 C=97	6 months	1 year	Dietary intake (observation), BMI, F&V preferences	W
Hopper 199666 USA	School	School multicomponent	Year 2 & 4 Mean age 8.9 years	I=48 C=49	I=48 C=49	12 weeks	0	Dietary intake (Assisted self & parent-report), BMI	M
Horne 200955 Food Dudes Ireland	School	School multicomponent	4-11 years	I=228 C=207	?	12 months	0	Dietary intake (observation)	M
Kristjansdottir 201056 Iceland	School	School multicomponent	7-9 years	265 No breakdown by I/C	I=58 C=48	2 years	0	Dietary intake (Parent report & measurement)	W
Luepker 199657, Dwyer 200280 CATCH USA	School	School food service changes, curriculum and home program	Year 3 initially 8 to 11 years	5106	4019 (1532 at 3 years post intervention)	3 years	3 years	BP, BMI/skin fold thickness, serum lipids and Apolipoprotein B, dietary intake (self-report), PA/screen time (self-report), fitness (9 minute run)	S
Llargues 201158 Avall Study Spain	School	School based curriculum multicomponent, plus home	Year 1 5-7 years	598, I=272 C=237	I=272, C=236	2 years	0	BMI, dietary intake (parent-report), PA/screen time (parent-report)	S

Study, Location	Setting, Disadvantaged population (if indicated)	Interventions	Participants: Age during intervention	Sample size †	Sample size at Follow-up †	Duration of intervention	Follow-up duration post-intervention	Health & Nutrition Outcomes	Potential risk of bias rating
Moore 2007, 201438, 95 Murphy 201179 Welsh Primary School Free Breakfast Initiative Wales	School	Free school breakfast program	9 to 11 years	RCT trial with repeated X sectional design	1975 had at least baseline and one follow up	12months	0	Dietary intake (Self-report), Classroom cognitive tests, Attitudes towards breakfast (Self-report), Strengths & Difficulties questionnaire (teacher-report)	W
Muth 200859 IMPACT USA	School	School curriculum based	Year 4 Mean age 9.9 years	I=38 C=37	I=37 C=36	3 months plus student leader training	0	Dietary intake (Self-report), PA/sedentary behaviour/screen time (Self-report)	S
Natale 201363, 96 Healthy Caregivers, Healthy children USA	Preschool	Pre-school multicomponent, environmental and family	2 to 5 years	I=726 C=379	I=238 C=69	2 years	0	Dietary intake (parent-report), sedentary behaviour (parent-report)	W
Natale 201464 Healthy Inside-Healthy Outside (HI-HO) USA	Preschool	Pre-school multicomponent plus home	2 to 5 years	I=238, C=69 dyads	185 dyads	6 months	6 months	Dietary intake (parent-report), BMI, PA levels and screen time (parent-report)	M

Study, Location	Setting, Disadvantaged population (if indicated)	Interventions	Participants: Age during intervention	Sample size †	Sample size at Follow-up †	Duration of intervention	Follow-up duration post-intervention	Health & Nutrition Outcomes	Potential risk of bias rating
Ni Mhurchu 201140 Free school breakfast New Zealand	School	Free school breakfast program	Primary school children Mean age 9.4 years	424	375	1 year for longest	0	Breakfast intake/hunger (self-report), food security (family-report) Academic achievement, school attendance (records), Strengths & Difficulties (teacher-report)	M
Olvera 2008, 201074, 78 Bounce program USA	Family Latino mothers/daughters	Community-based PA and nutrition intervention	Years 3 to 6	46 dyads, I=26 C=20	I=18 C=17	12 weeks	0	Dietary intake (self & parent report), BMI, PA (accelerometer, shuttle run test) exercise heart rate, peak oxygen consumption	M
Perrikou 201369 Cyprus	School	Teacher modelling (EXPO), curriculum (EDUC) two IV streams	9 years	I1=68, I2=79, C=71	I1=59, I2=67, C=58	29 weeks	1 year	Dietary intake (observation)	M
Perry 200444 Cafeteria Power Plus Project USA	School	Cafeteria-based intervention	Years 1 & 3	1668, no breakdown by I/C	1168	2 years	0	Dietary intake (parent-report), BMI	M

Study, Location	Setting, Disadvantaged population (if indicated)	Interventions	Participants: Age during intervention	Sample size †	Sample size at Follow-up †	Duration of intervention	Follow-up duration post-intervention	Health & Nutrition Outcomes	Potential risk of bias rating
Raitakari 200546, Kaitosaari 200645 Special Turku Coronary Risk Factor Intervention Project for Children (STRIP) Finland	Family	Family-based low saturated fat diet/nutrition intervention	Infants 7 months old at commencement	I=540 C=522 (families as unit)	I=179 C=190	Ongoing FU visits	9 years & 11 years	Dietary intake (self & parent report), PA (self-report), endothelial function (brachial artery ultrasound), BP, BMI, lipids, glucose, insulin, HOMA-IR, Apolipoprotein A, Apolipoprotein B	W
Rush, 201260 Project Energize New Zealand	School	School based nutrition and PA change agent mediated	Primary school children Years 1 -6	I=692 C=660	1352 (~30% lost to follow up, analysis restricted to matched groups)	2 years	0	BP, BMI, % body fat (bioelectrical impedance)	M
Stolley 199775 USA	Family Low income African American families	Community-based PA and nutrition IV	7-12 year olds	Pairs I=32, C=33	I=23 daughters & 20 mums C=27 daughters & 18 mums	12 weeks	0	BMI, dietary intake (self-report)	M
Tabak 201276 Family Ties to Health Program USA	Family	Family - Phone calls and newsletters	Preschool-aged children 2-5 years	I=25 C=23	I=22 C=21	4 months	0	Dietary intake (parent report)	W

<b>Study, Location</b>	<b>Setting, Disadvantaged population (if indicated)</b>	<b>Interventions</b>	<b>Participants: Age during intervention</b>	<b>Sample size †</b>	<b>Sample size at Follow-up †</b>	<b>Duration of intervention</b>	<b>Follow-up duration post-intervention</b>	<b>Health &amp; Nutrition Outcomes</b>	<b>Potential risk of bias rating</b>
Vereecken 200965 Beastly Healthy at School Belgium	Preschool	Pre-school multicomponent	Preschool children 2.5 to 5 years	I=742 kids C=480	I=618 kids C=445	12 weeks	6 months	Dietary intake (observation and parent-report)	S

*Risk of bias: S = Strong; M = Moderate; W = Weak*

*† I = Intervention group; C = Control group*

### Appendix 3: Table 2, Impact of nutrition interventions on children's fruit and vegetable intake

Study	F&V results* (95% CI)	Outcome method	Direction of association Intervention vs Control
<b>Family studies</b>			
Baranowski 2003	Fruit & Veg (including juice) intake d=1.3 (0.55, 2.11)	24-hour recall x2	+
Cameron 2014	Fruit intake d=0.06 Veg intake d=0.05	24-hour recall x3	-
Epstein 2001	F&V intake d=1.05 (0.23-1.87) (↑ F&V group cf. ↓Fat/Sugar group)	FFQ	+/-
Olvera 2008, 2010	F&V d=0.34	SPAN questionnaire	+
Tabak 2012	V intake d=0.24 (-0.35, 0.84)	FFQ	-
<b>Preschool/School studies with no parent component</b>			
Bere 2006a	F&V intake (1 yr) d=0.21 (0.04, 0.39) F&V intake (2 yrs) d=0.19 (0.02, 0.37)	24-hour recall	+
Breslin 2012	Veg intake d=0.17 (-0.03, 0.36) Fruit intake d=0.0 (-0.19, 0.19)	FFQ	-
Christian 2014	F&V intake d=-0.2 (-0.3, 0.0) Veg intake d=-0.1 (-0.2, 0.1) Fruit intake d=-0.1 (-0.3, 0.0)	24-hour recall	-
Hardy 2010	Fruit in lunch d=-0.11 (-0.40, 0.18) Veg in lunch d=0.20 (-0.09, 0.48)	Lunchbox audit	-
Hendy 2011	F or V eaten first d=0.61 (0.36, 0.81)	Direct observation	+
Perrikou 2013	Fruit intake d=1.43 (1.05, 1.79) Exposure d=1.41 (1.05, 1.77) Education at 1 year	2 day food record (parent)	+
Perry 2004	Fruit intake d=0.09 Fruit (no juice) d=0.12	Direct observation	+



Study	F&V results* (95% CI)	Outcome method	Direction of association Intervention vs Control
<b>Preschool/School studies with parent component</b>			
Anderson 2005	Fruit intake d=0.48 (0.13, 0.83) F&V d=0.04 (-0.30, 0.39)	3-day food record (self-report)	+/-
Baranowski 2000	F&V intake Mean difference 0.2 serves/day (1 yr), 0.2 Serves/day (2 yrs)	7 day food record	+
Bayer 2009	Fruit intake d=0.14, Veg. intake d=0.13	FFQ	+
Bere 2006b	F&V intake (1 yr) d=0.03(-0.17, 0.23) F&V intake (2 yrs) d=-0.1(- 0.29, 0.12)	24-hour recall	-
Cohen 2014	F&V intake/1000kcal d=0.1 F intake/1000kcal d=0.1 V intake/1000kcal d=0.1	24-hour recall/FFQ	-
De Bock 2012	Change in Fruit intake d=0.38 (0.18, 0.59) Change in Veg intake 0.33 (0.12, 0.53)	Short questions	+
Evans 2012	F&V intake d=0.01 (-0.14, 0.16)	24-hour recall	-
Hoffman 2010	Fruit intake 1 yr d=0.86, 2 yrs d=0.55 Veg intake 1yr d=0.34, 2 yrs No difference (at school)	Direct measurement	+/-
Hopper 1996	Change in F&V serves d=0.40 (0.00, 0.80) (at school)	Direct observation & measurement	+
Horne 2009	F&V consumed at lunch at 1 yr d=0.35 (0.16, 0.54)	Direct observation & measurement	+
Kristjansdottir 2010	F&V intake d=0.92 (0.52, 1.32)	3 day weighed food diary	+
Muth 2008	F&V Mean difference +0.85 serves/day	SPAN questionnaire (child)	+
Vereecken 2009	Fruit intake d=0.19 (0.00, 0.38) Veg intake d=0.08 (-0.11, 0.27) Fruit intake (at school) d=0.10 (-0.02, 0.22)	FFQ (parent) Audit (teacher)	+/-

\*d= Cohen's d where data available otherwise mean difference presented.

## Appendix 4: Table 3, Impact of nutrition interventions on children's fat intake

Study	Fat results (Cohens' d or mean difference* (95% CI)	Outcome method	Direction of association Intervention vs Control
<b>Family studies</b>			
Hendrie 2011	Total fat Mean diff=-10.9g/day (-19.3, -2.5), Saturated fat Mean diff=-8.1g/day (-11.9 to -4.3)	24-hour recall x3	+
Kaitosaari 2006 STRIP study	Total fat Boys d=-0.24 (-0.68, 0.20); Girls d=-0.78 (-1.22,-0.34) Saturated fat Boys d=-0.79 (-1.22,-0.34) Girls d=-1.17 (-1.63, -0.71) Polyunsaturated fat Boys d=0.55 (0.10, 0.99) Girls d=0.62 (0.19, 1.06)	4-day food record	+
Raitakari 2005	Saturated fat Girls age 11 d=-0.46 (-0.76, -0.17) Boys d=-0.95 (-1.25, -0.64)	4-day food record	+
Olvera 2008, 2010	Reduced intake of high fat foods d=0.39 (-0.28, 1.06)	FFQ	+
Stolley 1997	Saturated fat d= 0.39 (-0.17, 0.95) Fat % of Total energy d= 0.54 (-0.02, 1.11)	FFQ	+
<b>School studies with no parent component</b>			
Andersen 2014	Total fat d=0.00 (-0.11,0.10) Saturated fat d=0.00 (-0.11, 0.10) Monounsaturated fat d= 0.00 (-0.11,0.10) Polyunsaturated fat d= 0.00 (-0.10,0.10) Trans fatty acid d=0.00 (-0.10,0.10)	7-day food diary	-
Christian 2014	Total fat intake d=0.02 (-0.13, 0.18)	24-hour recall	-
Crepinsek 2006	Total fat d=-0.11(-0.17, -0.04) Saturated fat Mean diff=-0.20% (% of total energy)	24-hour recall	+

Study	Fat results (Cohens' d or mean difference* (95% CI)	Outcome method	Direction of association Intervention vs Control
<b>School studies with parent component</b>			
Anderson 2005	Fat as % energy Mean diff=-0.1%	3-day food diary	-
Caballero 2003 Pathways study	Total Fat Mean diff=-2.5% (-3.9,-1.1) Total fat at lunch Mean diff= -4.2% (-7.1,-1.3) (% of total energy)	Direct observation, 24-hour recall	+
Cohen 2014	Energy from saturated fat d=-0.02 (-0.21, 0.17)	24-hour recall, Modified FFQ	-
Evans 2012	Total fat Mean diff=1.2g/day (-2.8,5.1), Saturated fat Mean diff=0.0g/day (-1.5,1.5)	24-hour recall	-
Greening 2011	Reduced dietary fat d=0.33 (0.15, 0.52)	Child dietary fat questionnaire	+
Hopper 1996	Saturated fat Mean diff=-0.63g/day (p>0.05)	24-hour recall	+
Kristjansdottir 2010	Total fat (g/day) d=0.18 (-0.2,0.56) Saturated Fat d= 0.04 (-0.34, 0.42) Monounsaturated fat d=0.35 (-0.04, 0.73) PUFA d=0.03 (-0.35, 0.41)	3-day weighed food record	+/-
Leupker 1996	Fat as % energy d=-0.23 (-0.35, -0.11) Monounsaturated fat d= -0.15 (-0.26, 0.03) Saturated fat d=-0.20 (-0.32, -0.09) Polyunsaturated fat d=- 0.16, (-0.28, 0.045) (All % Total energy)	24-hour recall	+

\*d= Cohen's d where data available otherwise mean difference presented