

Research Article

The Quark-Gluon Plasma Equation of State and the Generalized Uncertainty Principle

L. I. Abou-Salem, N. M. El Naggar, and I. A. Elmashad

Physics Department, Faculty of Science, Benha University, Benha 13518, Egypt

Correspondence should be addressed to L. I. Abou-Salem; loutfy.abousalem@fsc.bu.edu.eg

Received 13 July 2015; Accepted 19 September 2015

Academic Editor: Elias C. Vagenas

Copyright © 2015 L. I. Abou-Salem et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The publication of this article was funded by SCOAP³.

The quark-gluon plasma (QGP) equation of state within a minimal length scenario or Generalized Uncertainty Principle (GUP) is studied. The Generalized Uncertainty Principle is implemented on deriving the thermodynamics of ideal QGP at a vanishing chemical potential. We find a significant effect for the GUP term. The main features of QCD lattice results were quantitatively achieved in case of $n_f = 0$, $n_f = 2$, and $n_f = 2 + 1$ flavors for the energy density, the pressure, and the interaction measure. The exciting point is the large value of bag pressure especially in case of $n_f = 2 + 1$ flavor which reflects the strong correlation between quarks in this bag which is already expected. One can notice that the asymptotic behavior which is characterized by Stephan-Boltzmann limit would be satisfied.

1. Introduction

Essential modifications in Heisenberg's uncertainty principle are predicted near Planck scale which is called Generalized Uncertainty Principle (GUP). One of the most exciting predictions of some approaches related to quantum gravity [1], perturbative string theory [2], and black holes [3] is the minimal length concept existence. For a recent review, see [4]. These approaches seem to modify almost all mechanical Hamiltonians. Thus, quantum mechanics can be studied in the presence of a minimal length [5–8]. Also, in quantum optics, the GUP implications can be measured directly which confirm the theoretical predictions [9–11]. In particular, exact solutions of various relativistic [12–15] and nonrelativistic problems [16–22] have been obtained in the presence of a minimal length $\Delta x_0 = \hbar(\beta)^{1/2}$. The Generalized Uncertainty Principle [23, 24] was implemented on deriving the thermodynamics of ideal quark-gluon plasma of massless quark flavor [25, 26].

In other minimal length formalism [5, 6], the Heisenberg algebra associated with the momentum \hat{p} and the position coordinates \hat{x}_i is given by

$$[\hat{x}_i, \hat{p}_j] = i\hbar\delta_{ij}(1 + \beta p^2), \quad (1)$$

where $\beta > 0$ is the minimal length parameter. The Generalized Uncertainty Principle (GUP) corresponding to it reads

$$\Delta x_i \Delta p_j \geq \delta_{ij} [1 + \beta (\Delta p)^2 + \beta \langle p \rangle^2] \quad (2)$$

which yields a minimal observable length $\Delta x_0 = \hbar(\beta)^{1/2}$. Hence, the momentum would be subject of a modification and becomes

$$p_i = p_{oi} (1 + \beta p_0^2), \quad (3)$$

where $x_i = x_{0i}$ and p_{0i} satisfy the canonical commutation relations $[x_{0i}, p_{0j}] = i\hbar\delta_{ij}$. Also, p_{0i} can be interpreted as the momentum at low energies and p_i at high energies. Since the GUP modifies the Hamiltonian, it is important to study these effects quantitatively. These effects were investigated on condensed matter, atomic systems [10–12, 27], the Liouville theorem (LT) in statistical mechanics [28], and the weak equivalence principle (WKP). Recently, another approach based on super gravity was implemented to QCD and to QGP especially [29–31].

In this paper, the effect of the GUP on QGP equation of state of massless quark flavors at a vanishing chemical potential μ is studied. We calculate the corrections to various

thermodynamic quantities, like the energy density and the pressure. Then, these corrections with bag model are used to describe the quark-gluon plasma equation of state and compare it with QCD lattice results. This paper is organized as follows. In Section 2, we derive thermodynamics of QGP consisting of a noninteracting massless bosons and fermions with impact of GUP approach. The results, discussions, and conclusions are given in Sections 3 and 4.

2. Thermodynamics of Quark-Gluon Plasma with GUP Effect

In this section, we derive the thermodynamics of QGP in case of bosons and fermions taking into account the GUP impact. Then, the thermodynamical equations such as pressure and energy density of quark-gluon plasma are obtained.

2.1. In Case of Bosons. At finite temperature T and chemical potential μ , the grand canonical partition function z_B for noninteracting massive bosons with g internal degrees of freedom is given by [32]

$$z_B = \prod_k \left[\sum_{l=0}^{\infty} \exp\left(-l \frac{E(k) - \mu}{T}\right) \right]^g \quad (4)$$

$$= \prod_k \left[1 - \exp\left(\frac{E(k) - \mu}{T}\right) \right]^{-g}, \quad (5)$$

where l is the occupation number for each quantum state with energy $E(k) = \sqrt{k^2 + m^2}$ with mass m and k is the momentum of the particle. Here the infinite product is taken for all possible momentum states.

For simplicity we consider chiral limit (i.e., vanishing mass) and a vanishing chemical potential, which experiments ensure at high energy. Then the dominant excitation in the hadronic phase is a massless pion, while that in the quark-gluon plasma is massless quarks and gluons. For a particle of mass M having a distant origin and an energy comparable to the Planck scale, the momentum would be a subject of a tiny modification so that the dispersion relation would too. According to GUP approach, the dispersion relation reads

$$E^2(k) = k^2 c^2 (1 + \beta k^2)^2 + M^2 c^4, \quad (6)$$

where M and c are the mass of the particle and the speed of light as introduced by Lorentz and implemented in special relativity, respectively. For simplicity we use natural units in which $\hbar = c = 1$. Hence, we have

$$E(k) = k(1 + \beta k^2). \quad (7)$$

For large volume, the sum over all states of single particle can be rewritten in terms of an integral [33]

$$\sum_k \rightarrow \frac{V}{(2\pi)^3} \int_0^{\infty} d^3k \rightarrow \frac{V}{2\pi^2} \int_0^{\infty} \frac{k^2 dk}{(1 + \beta k^2)^4}. \quad (8)$$

Thus, the partition function, (4), becomes

$$\begin{aligned} \ln z_B &= \frac{-Vg}{2\pi^2} \int_0^{\infty} k^2 \frac{\ln[1 - \exp(-E(k)/T)]}{(1 + \beta k^2)^4} dk = \frac{-Vg}{2\pi^2} \\ &\cdot \int_0^{\infty} k^2 \frac{\ln[1 - \exp(-(k/T)(1 + \beta k^2))]}{(1 + \beta k^2)^4} dk \\ &= \frac{Vg}{2\pi^2} \left[\frac{k}{6\beta(1 + \beta k^2)^3} \right. \\ &\cdot \ln \left[1 - \exp\left(-\frac{k}{T}(1 + \beta k^2)\right) \right] \Bigg|_0^{\infty} - \frac{Vg}{2\pi^2} \\ &\cdot \int_0^{\infty} \frac{1}{6\beta(1 + \beta k^2)^3} \left[\ln \left[1 - \exp\left(-\frac{k(1 + \beta k^2)}{T}\right) \right] + \frac{k(1 + 3\beta k^2)}{T} \right. \\ &\cdot \left. \frac{1}{\exp(k(1 + \beta k^2)/T) - 1} \right] dk, \end{aligned} \quad (9)$$

where ‘‘Integration by Parts’’ was used for solving the above equation. It is obvious that the first term in (9) vanishes. Thus,

$$\begin{aligned} \ln z_B &= \frac{-Vg}{2\pi^2} \\ &\cdot \int_0^{\infty} \frac{1}{6\beta(1 + \beta k^2)^3} \left[\ln \left[1 - \exp\left(-\frac{k(1 + \beta k^2)}{T}\right) \right] \right. \\ &\cdot \left. + \frac{k(1 + 3\beta k^2)}{T} \frac{1}{\exp(k(1 + \beta k^2)/T) - 1} \right] dk. \end{aligned} \quad (10)$$

For solving (10) let $x = (k/T)(1 + \beta k^2)$ so that $dx = (1 + 3\beta k^2)dk/T$ and

$$\begin{aligned} \ln z_B &= \frac{-Vg}{2\pi^2} \left[\int_0^{\infty} \frac{\ln[1 - e^{-x}] T dx}{6\beta(1 + \beta k^2)^3 (1 + 3\beta k^2)} \right] \\ &- \frac{Vg}{2\pi^2} \left[\int_0^{\infty} \frac{k dx}{6\beta(1 + \beta k^2)^3 (e^x - 1)} \right]. \end{aligned} \quad (11)$$

The momentum k as a function of x variable can be approximated to the first order of β as follows:

$$\begin{aligned} k &= xT - \beta k^3 \\ &= xT - \beta(xT - \beta k^3)^3 = xT - \beta x^3 T^3 \left(1 - \frac{\beta k^3}{xT}\right)^3 \\ &\approx xT - \beta x^3 T^3. \end{aligned} \quad (12)$$

Employing the value of k into integrand terms of (11) and approximating them to the first order of β are as follows.

For the first term,

$$\frac{T}{6\beta(1+\beta k^2)^3(1+3\beta k^2)} = \frac{T(1-6\beta k^2+9\beta^2 k^4)}{6\beta} \quad (13)$$

$$\approx \frac{T}{6\beta} - x^2 T^3 + \frac{7\beta}{2} x^4 T^5.$$

For the second term,

$$\frac{k}{6\beta(1+\beta k^2)^3} = \frac{k-3\beta k^3}{6\beta} \quad (14)$$

$$\approx \frac{xT}{6\beta} - \frac{2x^3 T^3}{3} + \frac{3}{2}\beta x^5 T^5.$$

Substituting the values of terms in (13) and (14) into (11) and solving them analytically, we have

$$\ln z_B = \frac{\pi^2}{90} VgT^3 - \frac{16\pi^4}{315} g\beta T^5. \quad (15)$$

In case of $\beta \rightarrow 0$, the above equation is reduced to the partition function for bosons without GUP effect [32].

2.2. In Case of Fermions. At finite temperature T and chemical potential μ , the grand canonical partition function z_f for noninteracting massive bosons with g internal degrees of freedom is given by [32]

$$z_f = \prod_k \left[\sum_{l=0,1}^{\infty} \exp\left(-l \frac{E(k)-\mu}{T}\right) \right]^g \quad (16)$$

$$= \prod_k \left[1 + \exp\left(-\frac{E(k)-\mu}{T}\right) \right]^g, \quad (17)$$

where l is the occupation number for each quantum state with energy $E(k) = \sqrt{k^2 + m^2}$ with mass m and k is the momentum of the particle. Here the infinite product is taken for all possible momentum states.

For simplicity we consider chiral limit (i.e., vanishing mass) and a vanishing chemical potential, which experiments ensure at high energy. Then the dominant excitation in the hadronic phase is a massless pion, while that in the quark-gluon plasma is massless quarks and gluons. For a particle of mass M having a distant origin and an energy comparable to the Planck scale, the momentum would be a subject of a tiny modification. According to GUP approach, the dispersion relation reads

$$E^2(k) = k^2 c^2 (1 + \beta k^2)^2 + M^2 c^4, \quad (18)$$

where M and c are the mass of the particle and the speed of light as introduced by Lorentz and implemented in special relativity, respectively. For simplicity we use natural units in which $\hbar = c = 1$. Hence, we have

$$E(k) = k(1 + \beta k^2). \quad (19)$$

For large volume, the sum over all states of single particle can be rewritten in terms of an integral [33]

$$\sum_k \rightarrow \frac{V}{(2\pi)^3} \int_0^\infty d^3 k \rightarrow \frac{V}{2\pi^2} \int_0^\infty \frac{k^2 dk}{(1 + \beta k^2)^4}. \quad (20)$$

Thus, the partition function, (16), becomes

$$\begin{aligned} \ln z_f &= \frac{Vg}{2\pi^2} \int_0^\infty \frac{k^2 \ln(1 + \exp(-E(k)/T))}{(1 + \beta k^2)^4} dk = \frac{Vg}{2\pi^2} \\ &\cdot \int_0^\infty \frac{k^2 \ln(1 + \exp(-k(1 + \beta k^2)/T))}{(1 + \beta k^2)^4} dk \\ &= \frac{-Vg}{2\pi^2} \frac{k \ln(1 + \exp(-k(1 + \beta k^2)/T))}{6\beta(1 + \beta k^2)^3} \Big|_0^\infty \quad (21) \\ &+ \frac{Vg}{2\pi^2} \int_0^\infty \frac{\ln(1 + \exp(-k(1 + \beta k^2)/T)) dk}{6\beta(1 + \beta k^2)^3} \\ &- \frac{Vg}{2\pi^2} \\ &\cdot \int_0^\infty \frac{k(1 + 3\beta k^2) dk}{6\beta T(1 + \beta k^2)^3(1 + \exp(k(1 + \beta k^2)/T))}, \end{aligned}$$

where ‘‘Integration by Parts’’ was used for solving the above equation. It is obvious that the first term in (21) vanishes. Thus,

$$\begin{aligned} \ln z_f &= \frac{Vg}{2\pi^2} \left[\int_0^\infty \frac{\ln(1 + \exp(-k(1 + \beta k^2)/T)) dk}{6\beta(1 + \beta k^2)^3} \right. \\ &\left. - \int_0^\infty \frac{k(1 + 3\beta k^2) dk}{6\beta T(1 + \beta k^2)^3(1 + \exp(k(1 + \beta k^2)/T))} \right]. \quad (22) \end{aligned}$$

For solving (22) let $x = (k/T)(1 + \beta k^2)$ so that $dx = (1 + 3\beta k^2)dk/T$ and

$$\begin{aligned} \ln z_f &= \frac{Vg}{2\pi^2} \left[\int_0^\infty \frac{\ln[1 + e^{-x}] T dx}{6\beta(1 + \beta k^2)^3(1 + 3\beta k^2)} \right. \\ &\left. - \int_0^\infty \frac{k(1 + 3\beta k^2) dx}{6\beta(1 + 3\beta k^2)(e^x + 1)} \right]. \quad (23) \end{aligned}$$

Employing the value of k into integrand terms of (23) and approximating them to the first order of β are as follows.

For the first term,

$$\begin{aligned} \frac{T}{6\beta(1 + \beta k^2)^3(1 + 3\beta k^2)} &= \frac{(1 - 3\beta k^2)(1 - 3\beta k^2)}{6\beta} \\ &\approx \frac{T}{6\beta} - x^2 T^3 + \frac{7\beta}{2} x^4 T^5. \quad (24) \end{aligned}$$

For the second term,

$$\begin{aligned} \frac{k}{6\beta(1+3\beta k^2)} &= k(1-3\beta k^2) \\ &\simeq \frac{xT}{6\beta} - \frac{2}{3}x^3T^3 + \frac{3}{2}\beta x^5T^5. \end{aligned} \quad (25)$$

Substituting the values of terms in (24) and (25) into (23) and solving them analytically, we have

$$\ln z_f = \frac{7}{8} \frac{\pi^2}{90} VgT^3 - \frac{31\pi^4}{630} Vg\beta T^5. \quad (26)$$

In case of $\beta \rightarrow 0$, the above equation is reduced to the partition function for fermions without GUP effect [32].

Now, the QGP equation of state of free massless quarks and gluons can be derived from the above equations. The total grand canonical partition function of QGP state can be given by adding the grand partition functions coming from the contribution of bosons (gluons), fermions (quarks), and vacuum [39] as follows:

$$\ln z_{\text{QGP}} = \ln z_F + \ln z_B + \ln z_V, \quad (27)$$

where $\ln z_B$, $\ln z_F$, and $\ln z_V$ are the grand canonical functions of gluons, quarks, and vacuum, respectively.

In our case, the vacuum pressure can be represented with the bag constant B in case of Bag model [40].

Since the value of vacuum partition function equals $\ln z_V = -VB/T$ and $\ln z = -\Omega/T$, using these values (27) reads

$$\frac{\Omega}{V}\Big|_{\text{QGP}} = \frac{\Omega}{V}\Big|_{\text{quarks}} + \frac{\Omega}{V}\Big|_{\text{gluons}} + \frac{\Omega}{V}\Big|_{\text{vacuum}}, \quad (28)$$

where Ω is the grand canonical potential. Substituting (15) and (26) into (28), we have

$$\begin{aligned} \frac{\Omega}{V}\Big|_{\text{QGP}} &= -\left(g_g + \frac{7}{8}g_q\right) \frac{\pi^2}{90} T^4 \\ &\quad + \left(g_g \frac{16\pi^4}{315} + g_q \frac{31\pi^4}{630}\right) \beta T^6 + B. \end{aligned} \quad (29)$$

Hence, the QGP equation of state reads

$$\begin{aligned} P_{\text{QGP}} &= -\frac{\Omega}{V}\Big|_{\text{QGP}} \\ &= \left(g_g + \frac{7}{8}g_q\right) \frac{\pi^2}{90} T^4 \\ &\quad - \left(g_g \frac{16\pi^4}{315} + g_q \frac{31\pi^4}{630}\right) \beta T^6 - B \\ &= \frac{\sigma_{\text{SB}}}{3} T^4 - \left(g_g \frac{16\pi^4}{315} + g_q \frac{31\pi^4}{630}\right) \beta T^6 - B, \end{aligned} \quad (30)$$

where σ_{SB} is SB constant and is given by $(g_g + (7/8)g_q)\pi^2/30$. Hence, the energy density of QGP state of matter is given by

$$\varepsilon_{\text{QGP}} = \sigma_{\text{SB}} T^4 - 3 \left(g_g \frac{16\pi^4}{315} + g_q \frac{31\pi^4}{630}\right) \beta T^6 + B. \quad (31)$$

3. Results and Discussions

The main features of QCD lattice results show a clear N_f -dependence for both energy density and pressure (i.e., they become larger as the number of degrees of freedom increases). As it is clear from the Monte Carlo lattice results [34, 35], the pressure $P(T)$ rapidly increases at $T \simeq T_c$ which may be agreed with our predictions after adding the GUP effect on QGP equation of state. The bag pressure B can be determined by comparing the obtained equation with that of QCD lattice results.

The main problem appears when one starts to adjust the behavior of the energy density, through varying the value of the bag pressure; with the QCD lattice results, one can not obtain a qualitative agreement in case of the pressure using the same bag parameter value [41].

For overcoming this problem, the bag model was modified [42]. In this technique, the thermodynamical relation between the energy density and the pressure [33] was used

$$T \frac{dP}{dT} - P(T) = \varepsilon(T). \quad (32)$$

Solving the above first-order differential equation, we obtain the QGP equation of state as follows:

$$P = \frac{\sigma_{\text{SB}}}{3} T^4 - \left(g_g \frac{16}{525} + g_q \frac{31}{1050}\right) \pi^4 \beta T^6 - B + AT, \quad (33)$$

where A is a constant coming from the partial differential equation solution and can be determined from comparing the calculated QGP equation of state with the QCD lattice results.

To adjust the high temperature behavior for $P(T)$ and $\varepsilon(T)$, we will consider the suppression factor of the Stefan Boltzmann constant used in quasi particle approach [43]. In this approach, the system of interacting gluons may be treated as a noninteracting quasi particles gas with gluon quantum numbers, but with thermal mass (i.e., $m(T)$). The modified SB constant σ reads [41]

$$\sigma = \kappa(a) \sigma_{\text{SB}}, \quad (34)$$

where $\kappa(a)$ is a suppression factor. For $a \rightarrow 0$, it follows $\kappa \rightarrow 1$. Also, the function $\kappa(a)$ decreases monotonously and goes to zero at $a \rightarrow \infty$. Thus, the final form of the bag model equation of state (33) with incorporating the GUP modification is given by

$$P = \frac{\sigma}{3} T^4 - \left(g_g \frac{16}{525} + g_q \frac{31}{1050}\right) \pi^4 \beta T^6 - B + AT. \quad (35)$$

With σ and B being free model parameters

$$\varepsilon = \sigma T^4 - 3 \left(g_g \frac{16}{315} + g_q \frac{31}{630}\right) \pi^4 \beta T^6 + B. \quad (36)$$

In case of $\beta \rightarrow 0$, the above equations (35) and (36) are reduced to the pressure and the energy density equations obtained in the modified bag model [42] (i.e., without incorporating the GUP modification).

The theoretical calculation of the pressure and the energy density of the QGP using (35) and (36) compared to the lattice




FIGURE 1: The symbols show the MC LR for the pressure and the energy density in the SU(3) gluodynamics [34, 35]; the line corresponds to the equation of state (35) for the pressure in the left panel and (36) for the energy density in the right panel with $\sigma = 4.20719$, $A = -2.0019T_c^3$, and $B = -1.465358T_c^4$. The arrows in the above figures correspond to $P/\varepsilon = 1/3$ (i.e., the SB limit).




FIGURE 2: The symbols show the MC LR for the pressure and the energy density in case of $n_f = 2$ QCD equation of state [36]; the line corresponds to the equation of state (35) for the pressure in the left panel and (36) for the energy density in the right panel with $\sigma = 10.403$, $A = -8.72895T_c^3$, and $B = -4.9T_c^4$. The arrows in the above figures correspond to $P/\varepsilon = 1/3$ (i.e., the SB limit).

results in case of $n_f = 0$ are given in Figure 1. In this case we used the following parameter values: $g_{\text{QGP}} = 16$, $g_q = 0$, $g_g = 16$, $T_c = 0.200$ GeV, and $\beta = 0.0001$ GeV $^{-1}$ as taken in [11, 32, 43].

The theoretical calculation of the pressure and the energy density of the QGP using (35) and (36) compared to the lattice results in case of $n_f = 2$ are given in Figure 2. In this case we used the following parameter values: $g_{\text{QGP}} = 37$, $g_q = 24$, $g_g = 16$, $T_c = 0.152$ GeV, and $\beta = 0.0001$ GeV $^{-1}$ as taken in [11, 32, 43].

The theoretical calculation of the pressure and the energy density of the QGP using (35) and (36) compared to the lattice results in case of $n_f = 2 + 1$ are given in Figure 3. In this case we used the following parameter values: $g_{\text{QGP}} = 47.5$, $g_q = 36$, $g_g = 16$, $T_c = 0.152$ GeV, and $\beta = 0.0001$ GeV $^{-1}$ as taken in [11, 32, 43].

Also, we can calculate the interaction measure $(\varepsilon - 3P)/T^4$ for both $n_f = 0$ and $n_f = 2 + 1$ and compare them with lattice results for the interaction measure in the SU(3) gluodynamics [34, 35] and in case of $n_f = 2 + 1$ QCD equation of state [37, 38], respectively. These results can be shown in Figure 4.

It can be shown that the inflection point of $(\varepsilon - 3P)/T^4$ is at $T = 0.1846$ GeV for $n_f = 0$ and $T = 0.202$ GeV for $n_f = 2 + 1$ which are different from that obtained in [42].

4. Conclusion

In the present work, the effect of the GUP on QGP of massless quark flavors at a vanishing chemical potential, μ , is studied. The equation of state and the energy density are derived for the QGP state consisting of noninteracting massless bosons and fermions with impact of GUP approach. Also, the total grand canonical partition function of QGP state is given. One can conclude that a significant effect for the GUP term exists in case of studying the thermal properties of the QGP state. The main features of QCD lattice results were quantitatively achieved in case of $n_f = 0$, $n_f = 2$, and $n_f = 2 + 1$ for the equation of state, the energy density, and the interaction measure. The interesting point in our results is the large value of bag pressure especially in case of $n_f = 2 + 1$ flavor. It nearly equals 4.46 times the value obtained in [42] without taking into account the negative sign of it, which reflects the




FIGURE 3: The symbols show the MC LR for the pressure and the energy density in case of $n_f = 2 + 1$ QCD equation of state [37, 38]; the line corresponds to the equation of state (35) for the pressure in the left panel and (36) for the energy density in the right panel with $\sigma = 12.22017$, $A = -13.8577T_c^3$, and $B = -10.4353T_c^4$. The arrows in the above figures correspond to $P/\varepsilon = 1/3$ (i.e., the SB limit).




FIGURE 4: The symbols show the lattice results for the interaction measure in the SU(3) gluodynamics (left panel) [34, 35] and in case of $n_f = 2 + 1$ QCD equation of state (right panel) [37, 38].

strong correlation between quarks inside the bag which is already expected. The negative sign may be regarded as the tendency of the bag to reduce its volume. One can conclude that the modification of the QGP bag model equation of state using the GUP effect can reproduce the QCD lattice results which stands for the real data of QGP. Finally, with this study, one may be encouraged to implement GUP effect in other thermodynamic properties of QGP.



Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1] L. J. Garay, "Quantum gravity and minimum length," *International Journal of Modern Physics A*, vol. 10, no. 2, pp. 145–166, 1995.
- [2] D. J. Gross and P. F. Mende, "String theory beyond the Planck scale," *Nuclear Physics B*, vol. 303, no. 3, pp. 407–454, 1988.
- [3] M. Maggiore, "A generalized uncertainty principle in quantum gravity," *Physics Letters B*, vol. 304, no. 1-2, pp. 65–69, 1993.
- [4] S. Hossenfelder, "Minimal length scale scenarios for quantum gravity," *Living Reviews in Relativity*, vol. 16, no. 2, 2013.
- [5] A. Kempf, "Uncertainty relation in quantum mechanics with quantum group symmetry," *Journal of Mathematical Physics*, vol. 35, no. 9, pp. 4483–4496, 1994.
- [6] A. Kempf, G. Mangano, and R. B. Mann, "Hilbert space representation of the minimal length uncertainty relation," *Physical Review D*, vol. 52, no. 2, pp. 1108–1118, 1995.
- [7] A. Kempf, "Non-pointlike particles in harmonic oscillators," *Journal of Physics A: Mathematical and General*, vol. 30, no. 6, pp. 2093–2101, 1997.
- [8] H. Hinrichsen and A. Kempf, "Maximal localization in the presence of minimal uncertainties in positions and in momenta," *Journal of Mathematical Physics*, vol. 37, no. 5, pp. 2121–2137, 1996.
- [9] I. Pikovski, M. R. Vanner, M. Aspelmeyer, M. S. Kim, and Č. Brukner, "Probing planck-scale physics with quantum optics," *Nature Physics*, vol. 8, no. 5, pp. 393–397, 2012.
- [10] S. Das and E. C. Vagenas, "Universality of quantum gravity corrections," *Physical Review Letters*, vol. 101, no. 22, Article ID 221301, 4 pages, 2008.

- [11] A. F. Ali, S. Das, and E. C. Vagenas, "Proposal for testing quantum gravity in the lab," *Physical Review D*, vol. 84, no. 4, Article ID 044013, 10 pages, 2011.
- [12] C. Quesne and V. M. Tkachuk, "Dirac oscillator with nonzero minimal uncertainty in position," *Journal of Physics A: Mathematical and General*, vol. 38, no. 8, pp. 1747–1765, 2005.
- [13] K. Nouicer, "An exact solution of the one-dimensional Dirac oscillator in the presence of minimal lengths," *Journal of Physics A*, vol. 39, no. 18, pp. 5125–5134, 2006.
- [14] C. Quesne and V. M. Tkachuk, "Generalized deformed commutation relations with nonzero minimal uncertainties in position and/or momentum and applications to quantum mechanics," *Symmetry, Integrability and Geometry: Methods and Applications*, vol. 3, article 016, 18 pages, 2007.
- [15] T. K. Jana and P. Roy, "Exact solution of the Klein-Gordon equation in the presence of a minimal length," *Physics Letters A*, vol. 373, no. 14, pp. 1239–1241, 2009.
- [16] L. N. Chang, D. Minic, N. Okamura, and T. Takeuchi, "Exact solution of the harmonic oscillator in arbitrary dimensions with minimal length uncertainty relations," *Physical Review D*, vol. 65, no. 12, Article ID 125027, 2002.
- [17] I. Dadić, L. Jonke, and S. Meljanac, "Harmonic oscillator with minimal length uncertainty relations and ladder operators," *Physical Review D*, vol. 67, no. 8, Article ID 087701, 4 pages, 2003.
- [18] K. Gemba, Z. T. Hlousek, and Z. Papp, "Algebraic solution of the harmonic oscillator with minimal length uncertainty relations," <http://arxiv.org/abs/0712.2078>.
- [19] F. Brau, "Minimal length uncertainty relation and the hydrogen atom," *Journal of Physics A: Mathematical and General*, vol. 32, no. 44, pp. 7691–7696, 1999.
- [20] T. V. Fityo, I. O. Vakarchuk, and V. M. Tkachuk, "One-dimensional Coulomb-like problem in deformed space with minimal length," *Journal of Physics A: Mathematical and General*, vol. 39, no. 9, p. 2143, 2006.
- [21] R. Akhoury and Y.-P. Yao, "Minimal length uncertainty relation and the hydrogen spectrum," *Physics Letters B*, vol. 572, no. 1-2, pp. 37–42, 2003.
- [22] S. Benczik, L. N. Chang, D. Minic, and T. Takeuchi, "Hydrogen-atom spectrum under a minimal-length hypothesis," *Physical Review A—Atomic, Molecular, and Optical Physics*, vol. 72, no. 1, Article ID 012104, 2005.
- [23] S. Das and E. C. Vagenas, "Phenomenological implications of the generalized uncertainty principle," *Canadian Journal of Physics*, vol. 87, no. 3, pp. 233–240, 2009.
- [24] S. Das, E. C. Vagenas, and A. F. Ali, "Discreteness of space from GUP II: relativistic wave equations," *Physics Letters B*, vol. 690, no. 4, pp. 407–412, 2010.
- [25] N. M. El Naggar, L. I. Abou-Salem, I. A. Elmashad, and A. F. Ali, "A study on Quark-Gluon plasma equation of state using generalized uncertainty principle," *Journal of Modern Physics*, vol. 4, pp. 13–20, 2013.
- [26] A. Tawfik, L. I. Abou-Salem, I. Elmashad, J.-U. Nabi, and A. F. Ali, "Quantum gravity effect on the Quark-Gluon plasma," *SOP Transactions on Theoretical Physics*, vol. 1, no. 1, pp. 1–6, 2014.
- [27] A. F. Ali, S. Das, and E. C. Vagenas, "The generalized uncertainty principle and quantum gravity phenomenology," in *Proceedings of the 12th Marcel Grossmann Meeting*, pp. 2407–2409, Paris, France, July 2009.
- [28] A. F. Ali, "Minimal length in quantum gravity, equivalence principle and holographic entropy bound," *Classical and Quantum Gravity*, vol. 28, no. 6, Article ID 065013, 2011.
- [29] J. Sadeghi, B. Pourhassan, and S. Heshmatian, "Application of AdS/CFT in quark-gluon plasma," *Advances in High Energy Physics*, vol. 2013, Article ID 759804, 13 pages, 2013.
- [30] J. Sadeghi and B. Pourhassan, "STU–QCD correspondence," *Canadian Journal of Physics*, vol. 91, no. 12, pp. 995–1019, 2013.
- [31] J. Sadeghi and B. Pourhassan, "Drag force of moving quark at $\mathcal{N} = 2$ supergravity," *Journal of High Energy Physics*, vol. 2008, no. 12, article 026, 2008.
- [32] K. Yagi, T. Hatsuda, and Y. Miake, *Quark-Gluon Plasma From Big Bang to Little Bang*, Cambridge University Press, Cambridge, UK, 2005.
- [33] W. Greiner, L. Neise, and H. Stöcker, *Thermodynamics and Statistical Mechanics*, Springer, New York, NY, USA, 1997.
- [34] G. Boyd, J. Engels, F. Karsch et al., "Equation of state for the SU(3) gauge theory," *Physical Review Letters*, vol. 75, no. 23, pp. 4169–4172, 1995.
- [35] G. Boyd, J. Engels, F. Karsch et al., "Thermodynamics of SU(3) lattice gauge theory," *Nuclear Physics B*, vol. 469, no. 3, pp. 419–444, 1996.
- [36] F. Karsch, E. Laermann, and A. Peikert, "The pressure in 2, 2+1 and 3 flavour QCD," *Physics Letters, Section B: Nuclear, Elementary Particle and High-Energy Physics*, vol. 478, no. 4, pp. 447–455, 2000.
- [37] S. Borsányi, Z. Fodor, C. Hoelbling et al., "Is there still any T_c mystery in lattice QCD? Results with physical masses in the continuum limit III," *Journal of High Energy Physics*, vol. 2010, no. 9, article 073, 2010.
- [38] S. Borsanyi, G. Endrődi, Z. Fodor et al., "The QCD equation of state with dynamical quarks," *Journal of High Energy Physics*, vol. 2010, article 77, 2010.
- [39] J. Letessier and J. Rafelski, *Hadrons and Quark-Gluon Plasma*, Cambridge University Press, Cambridge, UK, 2004.
- [40] A. Chodos, R. L. Jaffe, K. Johnson, C. B. Thorn, and V. F. Weisskopf, "New extended model of hadrons," *Physical Review D: Particles and Fields*, vol. 9, no. 12, pp. 3471–3495, 1974.
- [41] V. V. Begun, M. I. Gorenstein, and O. A. Mogilevsky, "Modified bag models for the Quark-Gluon plasma equation of state," *International Journal of Modern Physics E*, vol. 20, no. 8, pp. 1805–1815, 2011.
- [42] V. V. Begun, M. I. Gorenstein, and O. A. Mogilevsky, "Equation of state for the quark gluon plasma with the negative bag constant," *Ukrainian Journal of Physics*, vol. 55, no. 9, pp. 1049–1052, 2010.
- [43] M. I. Gorenstein and S. N. Yang, "Gluon plasma with a medium-dependent dispersion relation," *Physical Review D*, vol. 52, no. 9, pp. 5206–5212, 1995.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

