

## Sexual Orientation, Disclosure, and Cardiovascular Stress Reactivity

Robert-Paul Juster<sup>1\*</sup>, David Matthew Doyle<sup>2</sup>, Mark L. Hatzenbuehler<sup>3</sup>,  
Bethany G. Everett<sup>4</sup>, L. Zachary DuBois<sup>5</sup>, & Jennifer J. McGrath<sup>6</sup>

1–Department of Psychiatry and Addiction, University of Montreal, Montreal, Canada

2–Department of Psychology, University of Exeter, Exeter, United Kingdom

3–Department of Sociomedical Sciences, Columbia University, New York, United States

4–Department of Sociology, University of Utah, Salt Lake City, United States

5–Department of Anthropology, University of Oregon, Eugene, United States

6–Department of Psychology, Concordia University, Montreal, Canada

### Corresponding Author:

Robert-Paul Juster, PhD

Centre d'études sur le sexe/genre, l'allostasie, et la résilience (CESAR)

Centre de recherche de l'Institut universitaire en santé mentale de Montréal

7331 Hochelaga, Pavillon Fernand-Seguin, FS-145-12

Montreal, Quebec, Canada H1N 3V2

Telephone : 514-251-4000 Ext 3415 ou Ext 3253

Email: [robert-paul.juster@umontreal.ca](mailto:robert-paul.juster@umontreal.ca)

1  
2 **Abstract** Stigma may strain the heart health of lesbian, gay, and bisexual (LGB) individuals. To date,  
3  
4 however, LGB-related differences in cardiovascular diagnosis, risk factors, and basal biomarkers are  
5  
6 inconsistently reported. Using a laboratory-based stress paradigm, the current study assessed whether  
7  
8 cardiovascular stress reactivity differs as a function of sexual orientation and disclosure status ('coming  
9  
10 out') in a sample of healthy young LGB and heterosexual adults. Eighty-seven participants aged 18 to 45  
11  
12 ( $M = 24.61 \pm 0.61 SE$ ) identifying as LGB and heterosexual (47%) were exposed to the Trier Social Stress  
13  
14 Test, a well-validated laboratory stressor involving public speaking and mental arithmetic. Throughout a  
15  
16 two-hour session, ambulatory recordings for heart rate and blood pressure were collected. Self-report  
17  
18 questionnaires were also administered to assess psychosocial and demographic variables. Gay/bisexual  
19  
20 men showed higher heart rate and lesbian/bisexual women showed marginally higher mean arterial blood  
21  
22 pressure in response to a stressor, compared to sex- and age-matched heterosexuals. No significant  
23  
24 differences emerged when comparing LGB individuals who had completely disclosed and those that had  
25  
26 not completely disclosed their sexual orientation to family and friends. Compared to heterosexuals, heart  
27  
28 rate is higher among gay/bisexual men and blood pressure is marginally higher among lesbian/bisexual  
29  
30 women, when exposed to a laboratory-based stressor. These preliminary findings contribute to a small  
31  
32 literature on sexual orientation differences in stress reactive biomarkers that requires further exploration.  
33  
34  
35  
36  
37  
38  
39  
40

41 **Lay Abstract** In response to stress exposure in a laboratory, gay/bisexual men showed higher heart rate  
42  
43 than heterosexual men. By contrast, lesbian/bisexual showed a non-significant tendency towards higher  
44  
45 blood pressure than heterosexual women. These preliminary findings suggest that the heart health of LGB  
46  
47 individuals might be strained by stigma exposure.  
48  
49  
50  
51

52 **Keywords:** sexual orientation; cardiovascular stress reactivity; blood pressure; heart rate; Trier Social  
53  
54 Stress Test  
55  
56  
57  
58  
59  
60

## **Introduction**

Cardiovascular disease (CVD) is the leading cause of death in the United States (Xu, Murphy, Kochanek, & Bastian, 2016) and worldwide (Mozaffarian et al., 2015). Numerous studies have indicated that socially disadvantaged groups, such as racial/ethnic minorities, are at heightened risk for CVD relative to socially advantaged majority groups. Recently, researchers have begun to examine whether sexual minorities – that is, individuals who identify as lesbian, gay, or bisexual (LGB), who engage in same-sex behaviors, and/or who report same-sex attractions – are at increased risk for CVD outcomes compared to heterosexuals due in part to stigma and stress processes (IOM, 2011; Lick, Durso, & Johnson, 2013).

To date, studies have provided mixed evidence for LGB health disparities in CVD. Inconsistencies are due in part to LGB subgroup differences in CVD risk profiles (Caceres, Brody, & Chyun, 2016). Among sexual minority men, some studies have reported no differences in CVD risk factors (e.g., smoking, obesity, hypertension) relative to heterosexual men (Clark et al., 2015; Conron, Mimiaga, & Landers, 2010; Farmer, Bucholz, Flick, Burroughs, & Bowen, 2013), while others have reported increased risk factors among certain subgroups, such as bisexual men (Conron, et al., 2010; Farmer, Bucholz, et al., 2013). In a nationally representative study of blood pressure by sexual orientation, gay men showed double the risk of hypertension compared to heterosexual men, while no differences were found among women (Everett and Mollborn, 2013). By contrast, other nationally representative studies have identified increased CVD risk among sexual minority women compared to heterosexual women (Clark, et al., 2015; Conron, et al., 2010; Farmer, Jabson, Bucholz, & Bowen, 2013; Simoni, Smith, Oost, Lehavot, & Fredriksen-Goldsen, 2017). A systematic review concluded that there is an elevated risk for CVD in both sexual minority men and women that should be further assessed by combining subjective and objective measures of CVD risk (Caceres et al., 2017).

One approach to further study susceptibilities to CVD among LGB individuals is stress reactivity paradigms. When physical and psychosocial threats are detected by the brain, stress

1  
2 response systems (e.g., neuroendocrine, cardiovascular) are activated (McEwen, 1998). This  
3  
4 involves the *sympathetic-adrenal-medullary* (SAM)-axis release of catecholamines (e.g., adrenalin)  
5  
6 within seconds that increases heart rate and blood pressure, followed by the *hypothalamic-*  
7  
8 *pituitary-adrenal* (HPA)-axis production of glucocorticoids (e.g., cortisol) within minutes to sustain  
9  
10 the stress response. Chronic activation of the stress response may damage physiological integrity.  
11  
12 It has been proposed that cumulative activation of stress responses may be compounded by  
13  
14 stigma faced by LGB individuals who are consequently more vulnerable to stress-related diseases  
15  
16 (Juster, bVencill, & Johnson, 2017; Lick, et al., 2013).  
17  
18  
19  
20  
21  
22

23 The *reactivity hypothesis* (Manuck, 1994) states that exaggerated physiological reactivity to  
24  
25 stressors is linked to stress-related conditions like CVD (Lovallo, 2010; Lovallo and Gerin, 2003;  
26  
27 Schwartz et al., 2003). Cardiovascular stress reactivity refers to the magnitude or pattern of an  
28  
29 individual's hemodynamic responses to such stressors. It is believed that cardiovascular reactivity is  
30  
31 amplified among individuals exposed to social adversities (Treiber et al., 2003). In addition to the  
32  
33 magnitude of reactivity, the concept of recovery plays an important role in understanding stress-  
34  
35 induced disease risk. *Recovery* represents the duration and prolongation of stress responses after  
36  
37 the stressor ceases (Brosschot, 2010; Brosschot, Pieper, & Thayer, 2005). Stress reactivity/recovery  
38  
39 dynamics are characterized by much individual variability both in terms of exposure to stress and  
40  
41 in terms of recovery (Earle, Linden, & Weinberg, 1999; Linden, Earle, Gerin, & Christenfeld, 1997;  
42  
43 Rutledge, Linden, & Paul, 2000).  
44  
45  
46  
47  
48  
49  
50  
51

52 The Trier Social Stress Test (TSST) is a classic stress reactivity paradigm used extensively to assess  
53  
54 stress reactivity/recovery linked to stress-related disease (Allen et al., 2017). The TSST involves a 10-min  
55  
56 anticipation phase, 5-min performing a mock job interview, and 5-min calculating mental arithmetic  
57  
58  
59  
60

1  
2 (Kirschbaum, Pirke, & Hellhammer, 1993). This paradigm elicits social-evaluative threat (Dickerson and  
3  
4 Kemeny, 2002) that robustly activates the SAM- and HPA-axes. Stress reactivity and recovery can then be  
5  
6 discerned by assessing objective biomarkers (e.g., cortisol, blood pressure) pre- and post-TSST.  
7

8  
9 Recently, three international TSST studies have assessed HPA-axis functioning of LGB  
10  
11 individuals. First, American LGB individuals ( $N = 74$ ) growing up in less socially tolerant states  
12  
13 (compared to more socially tolerant states) evidenced blunted HPA-axis production of cortisol in  
14  
15 response to a modified TSST in which participants described an event where they experienced  
16  
17 rejection based on their sexual orientation (Hatzenbuehler and McLaughlin, 2013). The authors  
18  
19 suggest that this hypo-reactive cortisol pattern might indicate a pathophysiological profile  
20  
21 associated with chronic stress, trauma, and/or fatigue (Fries, Hesse, Hellhammer, & Hellhammer,  
22  
23 2005). Second, a Canadian study of young adults ( $N = 87$ ) showed that sexual orientation  
24  
25 modulates cortisol reactivity to the TSST (Juster et al., 2015). Specifically, lesbian/bisexual women  
26  
27 showed higher cortisol during recovery than heterosexual women, while gay/bisexual men  
28  
29 demonstrated lower cortisol throughout testing compared to heterosexual men. Third, an Israeli  
30  
31 study of gay and heterosexual men exposed to a modified TSST ( $N = 36$ ) showed that gender atypical  
32  
33 behavior and heterosexual sexual orientation predicted higher levels of social interaction anxiety; however,  
34  
35 these variables were not associated with cortisol dynamics (Jacobson, Cohen, & Diamond, 2016).  
36  
37  
38  
39  
40  
41  
42  
43  
44

45  
46 To summarize, sexual minority men may show dampened HPA-axis reactivity, while sexual  
47  
48 minority women appear to show both HPA-axis hypo-reactivity (Hatzenbuehler and McLaughlin, 2013)  
49  
50 and hyper-reactivity (Juster, et al., 2015). This underlines the need to conduct sex/gender-specific  
51  
52 analyses when assessing LGB populations. In addition, other biological systems need to be  
53  
54 assessed. Of pertinence to the present study, cardiovascular stress reactivity has yet to be reported in a  
55  
56  
57  
58  
59  
60

1  
2 TSST study of sexual orientation. To the best of our knowledge, only one American study assessed  
3  
4 cardiovascular reactivity by exposing 27 healthy adult gay men to a psychosocial stressor that  
5  
6 involved discussing difficulties associated with concealing one's sexual orientation in day-to-day  
7  
8 life (Pérez-Benitez, O'Brien, Carels, Gordon, & Chiros, 2007). Men with high concealment of their  
9  
10 sexual orientation but who engaged in more disclosure during the laboratory task exhibited  
11  
12 greater cardiovascular recovery (assessed via heart rate and stroke volume) than those men who  
13  
14 engaged in less disclosure during the task (Pérez-Benitez, et al., 2007). Despite low power and no  
15  
16 sexual minority women, this study highlights the role that concealment and disclosure may have  
17  
18 on cardiovascular reactivity and the heart health of sexual minority men. Given that the existing  
19  
20 literature on health disparities in cardiovascular disease is mixed, further investigation of  
21  
22 cardiovascular stress processes is warranted.  
23  
24  
25  
26  
27  
28  
29  
30

31 The current study assessed whether TSST-induced cardiovascular stress reactivity differs as a  
32  
33 function of sexual orientation and disclosure status in a sample of healthy young LGB and heterosexual  
34  
35 adults. Due to heightened exposure to stigma, we expected LGB individuals to show amplified  
36  
37 cardiovascular stress reactivity. First, we hypothesized that sexual minority women and men would show  
38  
39 higher cardiovascular stress reactivity to the TSST than sex- and age-matched heterosexual individuals.  
40  
41 Second, among the LGB participants only, we hypothesized that LGB individuals who had fully disclosed  
42  
43 their sexual orientation would evidence lower cardiovascular stress reactivity than those LGB individuals  
44  
45 who had not yet fully disclosed and may therefore experience additional distress.  
46  
47  
48  
49  
50  
51

## 52 **Methods**

### 53 *Participants*

54  
55  
56  
57  
58  
59  
60

1  
2 Eighty-seven participants aged 18 to 45 ( $M = 24.61 \pm 0.61 SE$ ) identifying as lesbian or gay (8  
3  
4 women and 20 men), bisexual (13 women and 5 men) or heterosexual (20 women and 21 men) were  
5  
6 recruited from Montreal as part of a broader study. The main exclusionary criteria were medicinal use of  
7  
8 synthetic steroid hormones, major health problems (e.g., HIV/AIDS, cancer), and severe mental illness  
9  
10 (e.g., schizophrenia). No participant was taking medications related to cardiovascular problems (e.g., anti-  
11  
12 hypertensives) or that could interfere with cardiovascular functioning (e.g., stimulants).  
13  
14

15  
16 According to a recent systemic review (Simoni, et al., 2017), two demographic variables of  
17  
18 importance in research on physical health conditions among sexual minorities are race/ethnicity and  
19  
20 educational attainment. Race/ethnicity was coded as “0” for white individuals (74.5%) and “1” for non-  
21  
22 white individuals (25.5%). Education was coded as “0” for ‘no college degree attained’ (53.5%) and “1”  
23  
24 for ‘college degree attained’ (46.5%). Both race/ethnicity and education were controlled for in our  
25  
26 statistical analyses.  
27  
28

### 29 *Sexual Orientation*

30  
31 Sexual orientation was assessed and cross-validated using three methods: (a) response to separate  
32  
33 advertisements recruiting either lesbian/gay/bisexual or heterosexual participants; (b) asking participants  
34  
35 their identified sexual orientation in an open-ended manner during a telephone screening interview; and (c)  
36  
37 administration of a modified 5-item Klein Sexual Orientation Scale (Klein, Sepekoff, & Wolf, 1990). This  
38  
39 instrument uses a 7-point Likert scale to assess sexual attractions, sexual behavior, sexual fantasies,  
40  
41 lifestyle preference, and sexual identity along a continuum of sexual experiences “in your life up to now,”  
42  
43 with low scores representing greater heterosexuality and high scores representing greater homosexuality.  
44  
45 The sample’s responses showed near perfect internal consistency ( $\alpha = 0.98$ ). Based on correspondence  
46  
47 among these three methods, sexual orientation was coded as “LGB” ( $n = 46$ ) or “heterosexual” ( $n = 41$ ).  
48  
49  
50  
51

### 52 *Disclosure Status*

53  
54 Disclosure was measured using a four-item inventory created by our group that asked the LGB  
55  
56 participants ( $n = 46$ ) their age for four key milestones regarding same-sex sexual attractions that included  
57  
58  
59  
60

1 self-recognition, self-identification, disclosure to friends, and disclosure to family. Participants were coded  
2 as “disclosed” ( $n = 31$ ) if they provided ages for all four items or as “non-disclosed” ( $n = 14$ ) if one or  
3  
4  
5  
6  
7 more of the items were not answered completely. One LGB participant did not provide this information  
8  
9 and was dropped from analyses assessing disclosure status among LGB participants.

### 11 *Mastery*

12  
13 The 7-item Pearlin & Schooler Personal Mastery Scale (Pearlin and Schooler, 1978) was used to  
14 assess perceived mastery and control. This widely employed instrument uses a 4-point Likert scale  
15 (strongly disagree to strongly agree) to items such as “there is little I can do to change many of the  
16 important things in my life.” Reliability was good in the current sample (Cronbach’s  $\alpha = .82$ ). Individuals  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
with higher mastery are believed to appraise themselves as better able to manage stressors, which may  
buffer their physiological arousal to threat (Roepke et al., 2008). Mastery was therefore used as a covariate  
based on previous literature linking its role in modulating stress processes (Chida and Hamer, 2008).

### 30 *General Protocol*

31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
This study was approved by the research ethics board of Louis-H. Lafontaine Hospital and therefore  
accords with proper ethical conduct. Upon a 15-minute study explanation and screening interview by  
telephone, eligible participants were scheduled for two laboratory visits. The current analysis  
focuses on an afternoon/evening visit scheduled between 12:00 and 19:00 ( $M = 14.34$ ,  $SE = 0.11$ )  
that lasted two hours in which participants were exposed to the standardized psychosocial  
stressor (e.g., TSST). Upon debriefing, participants were compensated with \$50 CAD.

### 47 *Visit Order*

48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
The order of visits was counterbalanced randomly to manipulate experienced novelty of the testing  
environment (Sindi, Fiocco, Juster, Pruessner, & Lupien, 2013). In the first group (morning/afternoon;  $n =$   
49), participants received a blood draw (findings reported elsewhere; (Juster, Almeida, et al., 2016; Juster,  
Ouellet, et al., 2016; Juster, Smith, Ouellet, Sindi, & Lupien, 2013)) in the morning during their first visit



1 and were exposed to the TSST in the afternoon during their second visit about a week later; this was  
2 reversed for the second group (afternoon/morning;  $n = 37$ ). Because the second group arrived for the first  
3 time to our laboratory when exposed to the TSST, we expected that they would be more distressed than the  
4 first group who had already familiarized themselves with the laboratory setting. Given that novelty to  
5 testing environments can be appraised as stressful (Sindi, et al., 2013), preliminary analyses assessed  
6 whether visit order modulated cardiovascular functioning. This manipulation does not represent a circadian  
7 dissimilarity since participants were all exposed to the TSST in the afternoon (arrival time:  $M = 14.34$ ,  $SE$   
8  $= 0.11$ ; departure time:  $M = 16.17$ ,  $SE = 0.11$ ).

### 20 *Stress Reactivity Paradigm*

21  
22 The two-hour afternoon visit involved exposure to a modified version (Andrews et al., 2007;  
23 Wadiwalla et al., 2010) of the TSST (Kirschbaum, et al., 1993). Upon arrival, participants acclimated to the  
24 laboratory environment for approximately 40 minutes while we recorded their cardiovascular parameters  
25 every 10 mins. They were then given instructions for the TSST and asked to prepare their mock job  
26 interview speech. After this ten-minute anticipation phase, participants walked for approximately 15  
27 seconds to another room where they were asked to deliver a five-minute mock job interview, followed by  
28 five minutes of mental arithmetic in front of an unseen “behavioral expert” (who was actually a trained  
29 confederate) seated behind a one-way mirror. The participant and the “behavioral expert” communicated  
30 via an intercommunication device and the participant’s performance was recorded by a video camera.  
31 Upon completion of the TSST, participants were returned to the original room and cardiovascular  
32 recordings resumed with recordings every 10 mins. until the study concluded.

### 48 *Cardiovascular Measures*

49  
50 Auscultatory cardiovascular functioning was recorded throughout the two-hour visit before and  
51 after the TSST. No recordings were taken during the TSST. We employed an electronic  
52 sphygmomanometer (A&D Medical: Model UA-631 V) on participants’ non-dominant arm (selection of  
53 three cuff-sizes based on size). Participants were seated and asked to refrain from moving or talking while  
54  
55  
56  
57  
58  
59

1  
2 the cuff inflated. In total, we recorded heart rate (HR), systolic blood pressure (SBP), and diastolic blood  
3  
4 pressure (DBP) at ten-minute intervals at the following time-points relative to the TSST (**Figure 1**): -40  
5  
6 min, -30 min, -20 min, -10 min, 0 (TSST), +10 min, +20 min, +30 min, +40 min, and +50 min.  
7  
8

9 Cardiovascular recordings of HR, SBP, and DBP were averaged according to three phases (**Figure**  
10  
11 **1**): *acclimation* (-40 min, -30 min, -20 min, and -10 min pre-TSST), *reactivity* (immediately before and  
12  
13 after the TSST), and *recovery* (+10 min, +20 min, +30 min, and +40 min post-TSST). This aggregated  
14  
15 approach enhances reliability and reduces measurement error (Kamarck, Debski, & Manuck,  
16  
17 2000). SBP and DBP were transformed into mean arterial pressure (MAP) using a standard formula:  $\sum$   
18  
19 (DBP + 1/3 (SBP – DBP)).  
20  
21  
22

#### 23 24 *Statistical Analysis*

25  
26 To optimize power due to fewer lesbians and bisexual men, we combined groups of lesbian/gay and  
27  
28 bisexual individuals within sex (20 women and 26 men) and contrasted them to sex-matched heterosexual  
29  
30 individuals (20 women and 21 men). The direction of results was similar for lesbian and bisexual women  
31  
32 as well as gay and bisexual men, albeit of reduced statistical strength as indicated by lower p-value when  
33  
34 assessing differences with heterosexual individuals.  
35  
36

37  
38 To assess group differences in descriptive information reported in **Table 1**, we employed univariate  
39  
40 analysis of variance (ANOVA), Tukey's post-hoc analysis, and chi-square tests. Our main analyses were  
41  
42 stratified by sex given our focus on sexual orientation differences in cardiovascular stress reactivity. Sex-  
43  
44 disaggregation of analyses is also recommended by the National Institute of Health when assessing sex-  
45  
46 specific phenomena (NIH, 2014), and sexual orientation may modulate stress reactivity in a sex-specific  
47  
48 manner (Allen, et al., 2017).  
49  
50

51  
52 Mixed-design repeated-measures analysis of covariance (ANCOVA) were run with sexual  
53  
54 orientation (LGB and heterosexual) entered as the between-subject factor and time (repeated measures of  
55  
56 HR and MAP for acclimation, reactivity, and recovery) entered as the within-subjects factor while  
57  
58  
59

1  
2 adjusting for covariates (race/ethnicity, education, mastery, and visit order). Greenhouse-Geisser  
3  
4 corrections are reported whenever Mauchly's tests denoted violations in sphericity. Significant time-by-  
5  
6 group interaction effects were assessed using post-hoc one-way ANOVAs, and significant time effects  
7  
8 were assessed using paired-sample t-tests. Supplemental analyses assessed SBP and DBP separately. Effect  
9  
10 sizes are interpreted as follows: small ( $\eta^2_p \cong 0.01$ ), medium ( $\eta^2_p \cong 0.06$ ), or large ( $\eta^2_p \cong 0.14$ ) effects (Fritz,  
11  
12 Morris, & Richler, 2012).  
13  
14

## 15 16 17 18 Results 19

### 20 21 *Descriptive Statistics*

22  
23 Descriptive analyses assessed sex differences by sexual orientation. As seen in **Table 1**, LGB  
24  
25 respondents did not differ from heterosexual individuals with respect to a broad array of sample  
26  
27 characteristics. Two exceptions involved significant group differences for sexual orientation and  
28  
29 contraception. First, LGB participants reported greater propensity towards same-sex only responses on the  
30  
31 Klein Sexual Orientation Scale ( $ps < .001$ ) than heterosexuals. Second, heterosexual women were more  
32  
33 likely to be using oral contraceptives than lesbian/bisexual women ( $p = .017$ ). Note that this refers to orally  
34  
35 administered hormonal contraceptives and not those delivered by other modes. **Table 1** also shows that  
36  
37 risk factors for CVD like tobacco smoking, alcohol (over)consumption, and illicit drug use were not  
38  
39 different between groups by sexual orientation.  
40  
41  
42

### 43 44 *Preliminary Analysis*

45  
46 Potential confounders of heart rate (HR) and mean arterial pressure (MAP) were scrutinized in  
47  
48 preliminary analyses using repeated-measures ANOVA as a function of visit order (morning/afternoon:  $n =$   
49  
50 49; afternoon/morning group:  $n = 37$ ), self-reported menstrual cycle status (follicular:  $n = 20$ ; luteal:  $n =$   
51  
52 20), and oral contraceptive use (users:  $n = 14$ ; non-users:  $n = 26$ ). For visit order, no between-subject  
53  
54 differences were detected for HR ( $p = .45$ ) and MAP ( $p = 0.30$ ). By contrast, time-by-visit order interaction  
55  
56  
57  
58  
59  
60

1 effects were significant for HR ( $F_{(6.88,578.13)} = 2.73, p = .009, \eta^2_p = .032$ ) and marginal for MAP ( $F_{(6.15,516.4)} =$   
2 1.891,  $p = .079, \eta^2_p = .022$ ). Those coming to the laboratory for the first time showed greater cardiovascular  
3 reactivity than those coming for the second time. Among women, no between-group differences in HR and  
4 MAP were found as a function of menstrual cycle (respectively,  $p = .998$  and  $p = .952$ ) or oral contraceptive  
5 use (respectively,  $p = .415$  and  $p = .804$ ). Therefore, only visit order is controlled for in the main analyses.

### 13 *Sexual Orientation Differences in Cardiovascular Stress Reactivity*

14  
15 The following mixed-design repeated-measures ANCOVAs assessed HR and MAP as a function of  
16 sexual orientation while adjusting for race/ethnicity, education, mastery, and visit order split by sex.

17  
18 For HR among women (**Figure 2A**), only a within-subjects time effect was detected ( $F_{(1.56,53.13)} =$   
19 31.72,  $p < .001, \eta^2_p = .483$ ): HR increased between acclimation and reactivity ( $p < .001$ ) and decreased  
20 from reactivity to recovery ( $p < .001$ ). No group difference was detected as a function of sexual orientation  
21 ( $p = .494, \eta^2_p = .014$ ).

22  
23 For HR among men (**Figure 2B**), a within-subjects time effect was also detected ( $F_{(1.41,56.47)} = 9.01$  ,  
24  $p = .001, \eta^2_p = .184$ ): HR increased between acclimation and reactivity ( $p < .001$ ) and decreased from  
25 reactivity to recovery ( $p < .001$ ). In addition, a time-by-sexual orientation interaction effect was detected  
26 ( $F_{(1.41,56.47)} = 3.56$  ,  $p = .050, \eta^2_p = .082$ ). Post-hoc analysis revealed that gay/bisexual men had higher HR  
27 during acclimation ( $p = .029, \eta^2_p = .114$ ) and reactivity ( $p = .034, \eta^2_p = .108$ ), and marginally during  
28 recovery ( $p = .055, \eta^2_p = .089$ ), compared to heterosexual men. Between-subjects effects were significant  
29 for sexual orientation ( $F_{(1,40)} = 4.85, p = .033, \eta^2_p = .108$ ), whereby gay/bisexual men showed higher overall  
30 HR across the entire visit than heterosexual men.

31  
32 For MAP among women (**Figure 2C**), a within-subjects time effect was significant ( $F_{(1.56,53.94)} =$   
33 20.47,  $p < .001, \eta^2_p = .376$ ): MAP increased from acclimation to reactivity ( $p < .001$ ) and decreased from  
34 reactivity to recovery ( $p < .001$ ). A marginally significant time-by-sexual orientation interaction effect was  
35 detected ( $F_{(1.56,53.16)} = 3.30, p = .056, \eta^2_p = .089$ ). Post-hoc analysis revealed that lesbian/bisexual women  
36 had marginally higher MAP than heterosexual women during the reactivity phase ( $p = .054, \eta^2_p = .105$ ), but  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

not during acclimation or recovery. The between-subjects effect for sexual orientation was marginally significant ( $F_{(1,34)} = 3.12, p = .086, \eta^2_p = .084$ ), whereby lesbian/bisexual women showed higher overall MAP than heterosexual women. In supplemental analyses assessing blood pressure separately, between-groups differences were marginally higher among lesbian/bisexual women in contrast to heterosexual women for DBP ( $p = .076, \eta^2_p = .090$ ) and were non-significant for SBP ( $p = .195, \eta^2_p = .049$ ).

For MAP among men (**Figure 2D**), a within-subjects time effect was significant ( $F_{(1.42,56.77)} = 25.93, p < .001, \eta^2_p = .393$ ): MAP increased from acclimation to reactivity ( $p < .001$ ) and decreased from reactivity to recovery ( $p < .001$ ). No group difference was found between gay/bisexual men and heterosexual men in MAP ( $p = .978, \eta^2_p < .001$ ) nor in supplemental analyses assessing SBP and DBP separately.

#### *Disclosure Status in Relation to Cardiovascular Stress Reactivity among Sexual Minorities*

The following mixed-design repeated-measures ANCOVAs re-assessed HR and MAP as a function of disclosure status and sex while adjusting for race/ethnicity, education, mastery, and visit order in the full sample. Here, we did not disaggregate analyses by sex, but rather entered sex as a covariate/moderator to preserve power and to ascertain if sex differences were present among the LGB individuals. The groups were divided as follows for those who had completely disclosed (men:  $n = 19$ , women:  $n = 12$ ) and those who had not fully disclosed (men:  $n = 7$ , women:  $n = 7$ ) their sexual orientation to family and friends.

For HR, no between-groups differences were found for HR by disclosure status ( $p = .239, \eta^2_p = .037$ ), sex ( $p = .656, \eta^2_p = .005$ ), or disclosure status X sex interaction ( $p = .931, \eta^2_p < .001$ ).

For MAP, we detected a time X disclosure status X sex interaction effect ( $F_{(1.42,52.42)} = 4.13, p = .034, \eta^2_p = .100$ ); however, follow up analyses did not attain significance. While non-significant, sexual minority women who had not disclosed showed a tendency towards higher MAP during reactivity. No between-group differences were found for MAP by disclosure status ( $p = .401, \eta^2_p = .019$ ), sex ( $p = .973, \eta^2_p < .001$ ), or disclosure status X sex interaction ( $p = .651, \eta^2_p = .006$ ). Results remained non-significant when assessing SBP and DBP separately.

## **Discussion**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The current study provides preliminary evidence that cardiovascular stress reactivity may differ by sexual orientation. Using a well-established psychosocial stress paradigm (Allen, et al., 2017), we found that gay/bisexual men showed higher HR compared to heterosexual men. By contrast, lesbian/bisexual women showed marginally higher MAP compared to heterosexual women; however, this result was only at trend level. Those LGB individuals who had not completely disclosed their sexual orientation to family and friends did not show significant differences in cardiovascular parameters during the acclimation, reactivity, and recovery phases. This study contributes to a small literature on stress reactive biomarkers among LGB individuals, a health disparities population. All the same, we must be cautious in our interpretations given marginal group differences and the number of non-significant findings which may alternatively suggest that there are in fact no sexual orientation differences in cardiovascular stress reactivity.

Compared to heterosexual men, sexual minority men evidenced higher HR not only throughout the TSST reactivity phases but also during the acclimation phase of the 2 hour laboratory visit. The magnitude of this effect is medium to large and this provides the first report of cardiovascular stress reactivity differences as a function of sexual orientation in men. Increased HR via the autonomic nervous system among gay/bisexual men is a pattern consistent with a large body of literature among racial/ethnic minority populations that likewise report an increased cardiovascular reactivity/recovery due to discrimination (Lick, et al., 2013; Williams and Mohammed, 2009). Future research using more sophisticated measures of autonomic nervous system functioning (e.g., heart rate variability) are warranted. Moreover, continuous measures of high-frequency cardiovascular activity during specific tasks would allow a more nuanced assessment of stress experienced during the job talk versus mathematics phases of the TSST. In contrast to HR, we found no evidence for differences between men for MAP, which is inconsistent with a report of a 2-fold increased risk for hypertension among gay men (Everett and Mollborn, 2013). It is noteworthy that past research assessing CVD risk among LGB individuals used measures like diagnosis, self-reports, and

1  
2 basal biomarkers that cannot be easily compared to our use of biomarkers representing acclimation,  
3  
4 reactivity, and recovery.  
5

6 In contrast to sexual minority men, sexual minority women appear to exhibit marginally amplified  
7  
8 MAP reactivity relative to heterosexuals, but otherwise no differences in HR. Specifically, lesbian/bisexual  
9  
10 women showed marginally elevated overall and reactive MAP compared to heterosexual women, a finding  
11  
12 of medium effect magnitude and at trend level. As our study is the first to report on cardiovascular stress  
13  
14 reactivity in sexual minority women, we can only compare our findings to previous research on basal  
15  
16 biomarkers related to CVD, which showed no sexual orientation differences in hypertension for women  
17  
18 (Everett and Mollborn, 2013). The current findings provide preliminary evidence that sexual minority  
19  
20 women may show marginally higher blood pressure in a stress reactivity paradigm than heterosexual  
21  
22 women. As this result was at trend level, we reserve caution in concluding that blood pressure differs by  
23  
24 sexual orientation for women. This will need to be replicated and compared with other physiological systems.  
25  
26  
27  
28  
29

30 Our analyses did not identify differences in cardiovascular stress reactivity according to  
31  
32 disclosure. We must be cautious about this conclusion, however, given our limited power and  
33  
34 sample heterogeneity. Because LGB individuals who conceal their sexual orientation are inhibiting  
35  
36 a wide range of complex behaviors (Pachankis, 2007), they are likely allocating resources to this  
37  
38 inhibition at a potential physiological price (Juster, et al., 2013; Pérez-Benitez, et al., 2007). On the  
39  
40 other hand, past research indicates that disclosure may not always be associated with better  
41  
42 health outcomes for sexual minorities (Doyle and Molix, 2016). For example, increased  
43  
44 harassment, victimization, distress, HPA-axis production, and suicidality can occur as part of  
45  
46 disclosing (D'Augelli, 2002; Huebner, Rebchook, & Kegeles, 2004; Igartua, Gill, & Montoro, 2003;  
47  
48 McGregor et al., 2001; Oetjen and Rothblum, 2000; Waldo, 1999). Therefore, it is important to  
49  
50 consider broader contextual factors and individual differences that may modify the salutogenic  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 and/or pathogenic effects of disclosure processes. Lastly, sexual identity formation and disclosure  
3  
4 are complex processes that were crudely captured in our binary classification that would benefit  
5  
6 from more refined measurement in future work.  
7  
8

### 9 10 *Limitations*

11  
12 Despite medium to large effect sizes, our results are derived from a small convenience sample  
13  
14 measured at one time point. Additional psychosocial variables (e.g., relationship factors, health behaviors)  
15  
16 should be considered to fully understand CVD risk among LGB individuals (Frech, Lynch, & Barr, 2016).  
17  
18 CVD is influenced by numerous health behaviors like diet, smoking, alcohol consumption, exercise, and  
19  
20 sleep that may function as moderators. In addition, variables like mastery that we included as a covariate  
21  
22 might be involved in causal pathways that link sexual orientation (and minority stigma) to cardiovascular  
23  
24 reactivity, since it is possible that having a history of being stigmatized might result in increased reactivity  
25  
26 to social stressors in a laboratory setting. Larger sample sizes would allow future researcher to test  
27  
28 mediation or moderation effects of such health behaviors and psychosocial profiles.  
29  
30  
31

32  
33 In addition, CVD risk factors may cluster differently among subgroups of the LGB community  
34  
35 (IOM, 2011). Moreover, inconsistencies in how CVD risk factors, biomarkers, and disorders are measured  
36  
37 in the literature might lead to discrepancies that future research should consider (Caceres, et al., 2016).  
38  
39 Lastly, our study did not include transgender people that evidence stress biomarker profiles that appear to  
40  
41 increase CVD risk (Dubois, 2012; DuBois, Powers, Everett, & Juster, 2017). Finally, the sex of participants  
42  
43 interacts with the sex of TSST behavioral judges (Goodman, Janson, & Wolf, 2017); however, 87% of our  
44  
45 participants interacted with a female TSST behavioral judge, a distribution that should be better balanced in  
46  
47 future studies. Despite these limitations, our preliminary study contributes to a small literature on sexual  
48  
49 orientation differences in cardiovascular health and points to several areas for future research on stress  
50  
51 reactivity within the context of sexual orientation.  
52  
53  
54

### 55 56 *Future Directions*



1  
2 This preliminary study provides several directions for future research. First, the current  
3  
4 cardiovascular stress reactivity findings among men (i.e., gay/bisexual men show amplified HR throughout  
5  
6 the TSST session compared to heterosexual men) complement our previous report that showed low HPA-  
7  
8 axis reactivity among gay/bisexual men in this same sample (Juster, et al., 2015). These findings  
9  
10 collectively suggest a counter-regulation of HPA-axis and cardiovascular functioning among sexual  
11  
12 minority men. It is not, however, clear whether these profiles can be considered maladaptive. Noteworthy  
13  
14 as well in this sample is that other risk factors for CVD (BMI, triglycerides) were actually lower among  
15  
16 sexual minority men (Juster, et al., 2013). The compiled findings using this sample suggest that changes in  
17  
18 one physiological system examined individually may draw an incomplete picture of the underlying stress  
19  
20 processes involved. Instead, attending to coordinated change across multiple systems and their dynamic  
21  
22 alignment using multi-systemic approaches is ideally suited to evaluating the correlates of social  
23  
24 adversities (Lucas et al., 2017). We suggest that future studies employ more rigorous measurement of these  
25  
26 systems with more frequent sampling (e.g., heart rate variability).  
27  
28  
29  
30

31  
32 Second, lifecourse perspectives must be considered when assessing stigma-related determinants of  
33  
34 CVD trajectories among LGB populations. However, the current study did not measure sources of stigma  
35  
36 experienced throughout the lifecourse specifically related to one's sexual orientation other than our crude  
37  
38 measure of disclosure. As these sources of sexual minority stigma can have long-term effects on CVD  
39  
40 (Lick, et al., 2013), future prospective research (Katz-Wise et al., 2017a) is required to determine how  
41  
42 stigma 'gets under the skin' of LGB individuals across life and across generations. It has been  
43  
44 recommended that measures of victimization should use standard timeframes (e.g., lifetime reports, past  
45  
46 year reports, before/after 'coming out' reports) to better understand developmental variation and age trends  
47  
48 (Katz-Wise and Hyde, 2012). Timing of sexual orientation development is also important to consider  
49  
50 (Katz-Wise et al., 2017b). For example, those with earlier timing of same-sex/gender sexual experiences  
51  
52 are at greater risk of mental health problems than those at experience such experiences at a later age when  
53  
54 more emotionally and cognitively mature (Katz-Wise, et al., 2017a).  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Third, stress reactivity paradigms are inherently limited in their lab-to-life generalizability. It has been argued that cardiovascular reactivity in the laboratory is at best only moderately generalizable to non-laboratory situations; however, psychosocial stress paradigms (e.g., TSST) may be more representative of daily life stressors than cognitive or physical tasks routinely used in the cardiovascular reactivity literature (Schwartz, et al., 2003). Lastly, future research should assess genetic, environmental, and behavioral risk factors together when assessing cardiovascular endophenotypes (Collaboration, 2017; Schwartz, et al., 2003) to broaden our understanding of key mechanisms. In addition, we encourage future studies to consider measuring both sex-based and gender-based variables to refine understanding of subgroup differences that cannot be detected by focusing solely on male/female distinctions.

### *Conclusions*

Due to increased exposure to stigma, it has been proposed that sexual minorities may be at increased risk of cardiovascular disease. To date, the literature supporting this proposal has been mixed. Using a stress reactivity paradigm among a small convenience sample, we conclude that gay/bisexual men evidence higher heart rate during acclimation and reactivity phases compared to heterosexual men. Marginal differences were found for lesbian/bisexual women showing higher blood pressure than heterosexual women. These preliminary findings will need to be replicated in future studies that use more comprehensive approaches to assess the heart health of sexual and gender minority populations.

### **Disclosure statement**

The authors have no conflicts of interest to disclose.

### **Acknowledgements**

This study was funded by a grant (222055) from the Canadian Institutes of Health Research. Robert Paul Juster held a Doctoral scholarship from the Institute of Aging of the Canadian Institutes of Health Research (SIA 95402) during data collection of this manuscript. Current support during manuscript

preparation is thanks to the Canadian Institutes of Health Research Banting postdoctoral fellowship. We wish to thank our participants for their commitment to this study.

## References

- Allen, A. P., Kennedy, P. J., Dockray, S., Cryan, J. F., Dinan, T. G., & Clarke, G. (2017). The Trier Social Stress Test: Principles and practice. *Neurobiol Stress*, 6, pp. 113-126. doi:10.1016/j.ynstr.2016.11.001 Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/28229114>
- Andrews, J., Wadiwalla, M., Juster, R. P., Lord, C., Lupien, S. J., & Pruessner, J. C. (2007). Effects of manipulating the amount of social-evaluative threat on the cortisol stress response in young healthy men. *Behav Neurosci*, 121(5), pp. 871-876. doi:2007-13974-006 [pii] 10.1037/0735-7044.121.5.871 Retrieved from [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=17907819](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17907819)
- Brosschot, J. F. (2010). Markers of chronic stress: Prolonged physiological activation and (un)conscious perseverative cognition. *Neurosci Biobehav Rev* doi:S0149-7634(10)00005-9 [pii] 10.1016/j.neubiorev.2010.01.004 Retrieved from [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=20096302](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20096302)
- Brosschot, J. F., Pieper, S., & Thayer, J. F. (2005). Expanding stress theory: prolonged activation and perseverative cognition. *Psychoneuroendocrinology*, 30(10), pp. 1043-1049. doi:10.1016/j.psyneuen.2005.04.008 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15939546>
- Caceres, B. A., Brody, A., & Chyun, D. (2016). Recommendations for cardiovascular disease research with lesbian, gay and bisexual adults. *J Clin Nurs*, 25(23-24), pp. 3728-3742. doi:10.1111/jocn.13415 Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/27239792>
- Caceres, B. A., Brody, A., Luscombe, R. E., Primiano, J. E., Marusca, P., Sitts, E. M., & Chyun, D. (2017). A Systematic Review of Cardiovascular Disease in Sexual Minorities. *Am J Public Health*, 107(4), p 570. doi:10.2105/AJPH.2016.303630a Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/28272943>
- Chida, Y., & Hamer, M. (2008). Chronic psychosocial factors and acute physiological responses to laboratory-induced stress in healthy populations: a quantitative review of 30 years of investigations. *Psychol Bull*, 134(6), pp. 829-885. doi:2008-14745-003 [pii] 10.1037/a0013342 Retrieved from [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18954159](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18954159)
- Clark, C. J., Borowsky, I. W., Salisbury, J., Usher, J., Spencer, R. A., Przedworski, J. M., . . . Everson-Rose, S. A. (2015). Disparities in long-term cardiovascular disease risk by sexual identity: The National Longitudinal Study of Adolescent to Adult Health. *Prev Med*, 76, pp. 26-30. doi:10.1016/j.ypmed.2015.03.022 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/25849883>

- 1  
2 Collaboration, N. C. D. R. F. (2017). Worldwide trends in blood pressure from 1975 to 2015: a pooled  
3 analysis of 1479 population-based measurement studies with 19.1 million participants. *Lancet*,  
4 389(10064), pp. 37-55. doi:10.1016/S0140-6736(16)31919-5 Retrieved from  
5 <http://www.ncbi.nlm.nih.gov/pubmed/27863813>  
6
- 7  
8 Conron, K. J., Mimiaga, M. J., & Landers, S. J. (2010). A population-based study of sexual orientation  
9 identity and gender differences in adult health. *Am J Public Health*, 100(10), pp. 1953-1960.  
10 doi:10.2105/AJPH.2009.174169 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/20516373>  
11
- 12  
13 D'Augelli, A. R. (2002). Mental health problems among lesbian, gay, and bisexual youths ages 14 to 21.  
14 *Clinical Child Psychology and Psychiatry*, 7(3), pp. 433-456.
- 15  
16 Dickerson, S. S., & Kemeny, M. E. (2002). Acute stressors and cortisol reactivity: a meta-analytic review.  
17 *Psychosomatic Medicine*, 54, pp. 105-123.
- 18  
19 Doyle, D. M., & Molix, L. (2016). Minority stress and inflammatory mediators: covering moderates  
20 associations between perceived discrimination and salivary interleukin-6 in gay men. *J Behav Med*,  
21 39(5), pp. 782-792. doi:10.1007/s10865-016-9784-0 Retrieved from  
22 <https://www.ncbi.nlm.nih.gov/pubmed/27534538>  
23
- 24  
25 Dubois, L. Z. (2012). Associations between transition-specific stress experience, nocturnal decline in  
26 ambulatory blood pressure, and C-reactive protein levels among transgender men. *Am J Hum Biol*,  
27 24(1), pp. 52-61. doi:10.1002/ajhb.22203 Retrieved from  
28 <http://www.ncbi.nlm.nih.gov/pubmed/22120883>  
29
- 30  
31 DuBois, L. Z., Powers, S., Everett, B. G., & Juster, R. P. (2017). Stigma and diurnal cortisol among  
32 transitioning transgender men. *Psychoneuroendocrinology*, 82, pp. 59-66.  
33 doi:10.1016/j.psyneuen.2017.05.008 Retrieved from  
34 <https://www.ncbi.nlm.nih.gov/pubmed/28511045>
- 35  
36 Earle, T. L., Linden, W., & Weinberg, J. (1999). Differential effects of harassment on cardiovascular and  
37 salivary cortisol stress reactivity and recovery in women and men. *J Psychosom Res*, 46(2), pp.  
38 125-141. doi:S0022399998000750 [pii] Retrieved from  
39 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_u](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_u)  
40 [ids=10098822](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=10098822)  
41
- 42  
43 Everett, B., & Mollborn, S. (2013). Differences in hypertension by sexual orientation among U.S. young  
44 adults. *J Community Health*, 38(3), pp. 588-596. doi:10.1007/s10900-013-9655-3 Retrieved from  
45 <https://www.ncbi.nlm.nih.gov/pubmed/23397511>  
46
- 47  
48 Farmer, G. W., Bucholz, K. K., Flick, L. H., Burroughs, T. E., & Bowen, D. J. (2013). CVD risk among  
49 men participating in the National Health and Nutrition Examination Survey (NHANES) from 2001  
50 to 2010: differences by sexual minority status. *J Epidemiol Community Health*, 67(9), pp. 772-778.  
51 doi:10.1136/jech-2013-202658 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/23766523>
- 52  
53 Farmer, G. W., Jabson, J. M., Bucholz, K. K., & Bowen, D. J. (2013). A population-based study of  
54 cardiovascular disease risk in sexual-minority women. *Am J Public Health*, 103(10), pp. 1845-  
55 1850. doi:10.2105/AJPH.2013.301258 Retrieved from  
56 <http://www.ncbi.nlm.nih.gov/pubmed/23948018>  
57  
58  
59  
60

- 1  
2 Frech, A., Lynch, J. L., & Barr, P. (2016). Health consequences of same and opposite-sex unions:  
3 partnership, parenthood, and cardiovascular risk among young adults. *J Behav Med*, 39(1), pp. 13-  
4 27. doi:10.1007/s10865-015-9673-y Retrieved from  
5 <http://www.ncbi.nlm.nih.gov/pubmed/26323506>  
6
- 7 Fries, E., Hesse, J., Hellhammer, J., & Hellhammer, D. H. (2005). A new view on hypocortisolism.  
8 *Psychoneuroendocrinology*, 30(10), pp. 1010-1016. doi:S0306-4530(05)00089-2 [pii]  
9 10.1016/j.psyneuen.2005.04.006 Retrieved from  
10 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15950390)  
11 [950390](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15950390)  
12  
13
- 14 Fritz, C. O., Morris, P. E., & Richler, J. J. (2012). Effect size estimates: current use, calculations, and  
15 interpretation. *J Exp Psychol Gen*, 141(1), pp. 2-18. doi:10.1037/a0024338 Retrieved from  
16 <http://www.ncbi.nlm.nih.gov/pubmed/21823805>  
17
- 18 Goodman, W. K., Janson, J., & Wolf, J. M. (2017). Meta-analytical assessment of the effects of protocol  
19 variations on cortisol responses to the Trier Social Stress Test. *Psychoneuroendocrinology*, 80, pp.  
20 26-35. doi:10.1016/j.psyneuen.2017.02.030 Retrieved from  
21 <https://www.ncbi.nlm.nih.gov/pubmed/28292684>  
22  
23
- 24 Hatzenbuehler, M. L., & McLaughlin, K. A. (2013). Structural Stigma and Hypothalamic-Pituitary-  
25 Adrenocortical Axis Reactivity in Lesbian, Gay, and Bisexual Young Adults. *Ann Behav Med*, 47,  
26 pp. 39-47. doi:10.1007/s12160-013-9556-9 Retrieved from  
27 <http://www.ncbi.nlm.nih.gov/pubmed/24154988>  
28  
29
- 30 Huebner, D. M., Rebchook, G. M., & Kegeles, S. M. (2004). Experiences of harassment, discrimination,  
31 and physical violence among young gay and bisexual men. *American Journal of Public Health*,  
32 94(7), pp. 1200-1203. Retrieved from  
33 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_u](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15226143)  
34 [ids=15226143](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15226143)  
35  
36
- 37 Igartua, K. J., Gill, K., & Montoro, R. (2003). Internalized homophobia: A factor in depression, anxiety,  
38 and suicide in the gay and lesbian population. *Canadian Journal of Community Mental Health*,  
39 22(2), pp. 15-30.  
40
- 41 IOM. (2011). *The Health of Lesbian, Gay, Bisexual, and Transgender People: Building a Foundation for*  
42 *Better Understanding* Washinton, DC: The National Academies Press.  
43
- 44 Jacobson, R., Cohen, H., & Diamond, G. M. (2016). Gender Atypicality and Anxiety Response to Social  
45 Interaction Stress in Homosexual and Heterosexual Men. *Arch Sex Behav*, 45(3), pp. 713-723.  
46 doi:10.1007/s10508-015-0528-y Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/25946903>  
47  
48
- 49 Juster, R. P., Almeida, D., Cardoso, C., Raymond, C., Johnson, P. J., Pfaus, J. G., . . . Lupien, S. J. (2016).  
50 Gonads and strife: Sex hormones vary according to sexual orientation for women and stress indices  
51 for both sexes. *Psychoneuroendocrinology*, 72, pp. 119-130. doi:10.1016/j.psyneuen.2016.06.011  
52 Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/27398882>  
53  
54
- 55 Juster, R. P., bVencill, J. A., & Johnson, P. J. (2017). Impact of stress and strain on current LGBT health  
56 disparities. In K. Eckstrand & J. Potter (Eds.), *Trauma, Resilience, and Health Promotion for LGBT*  
57  
58  
59  
60

1  
2 *Patients: What Every Healthcare Provide Should Know* (Vol. 1, pp. 35-48). New York: Springer  
3 Press.  
4

- 5 Juster, R. P., Hatzenbuehler, M. L., Mendrek, A., Pfaus, J. G., Smith, N. G., Johnson, P. J., . . . Pruessner,  
6 J. C. (2015). Sexual orientation modulates endocrine stress reactivity. *Biological Psychiatry*, *77*(7),  
7 pp. 668-676. doi:10.1016/j.biopsycho.2014.08.013 Retrieved from  
8 <http://www.ncbi.nlm.nih.gov/pubmed/25444167>  
9
- 10 Juster, R. P., Ouellet, E., Lefebvre-Louis, J. P., Sindi, S., Johnson, P. J., Smith, N. G., & Lupien, S. J.  
11 (2016). Retrospective coping strategies during sexual identity formation and current  
12 biopsychosocial stress. *Anxiety Stress Coping*, *29*, pp. 119-138.  
13 doi:10.1080/10615806.2015.1004324 Retrieved from  
14 <http://www.ncbi.nlm.nih.gov/pubmed/25599254>  
15
- 16 Juster, R. P., Smith, N. G., Ouellet, E., Sindi, S., & Lupien, S. J. (2013). Sexual orientation and disclosure  
17 in relation to psychiatric symptoms, diurnal cortisol, and allostatic load. *Psychosomatic Medicine*,  
18 *75*(2), pp. 103-116. doi:10.1097/PSY.0b013e3182826881 Retrieved from  
19 <http://www.ncbi.nlm.nih.gov/pubmed/23362500>  
20
- 21 Kamarck, T. W., Debski, T. T., & Manuck, S. B. (2000). Enhancing the laboratory-to-life generalizability  
22 of cardiovascular reactivity using multiple occasions of measurement. *Psychophysiology*, *37*(4), pp.  
23 533-542. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10934912>  
24
- 25 Katz-Wise, S. L., & Hyde, J. S. (2012). Victimization experiences of lesbian, gay, and bisexual individuals:  
26 a meta-analysis. *J Sex Res*, *49*(2-3), pp. 142-167. doi:10.1080/00224499.2011.637247 Retrieved  
27 from <https://www.ncbi.nlm.nih.gov/pubmed/22380586>  
28
- 29 Katz-Wise, S. L., Rosario, M., Calzo, J. P., Scherer, E. A., Sarda, V., & Austin, S. B. (2017a). Associations  
30 of Timing of Sexual Orientation Developmental Milestones and Other Sexual Minority Stressors  
31 with Internalizing Mental Health Symptoms Among Sexual Minority Young Adults. *Arch Sex*  
32 *Behav*, *46*(5), pp. 1441-1452. doi:10.1007/s10508-017-0964-y Retrieved from  
33 <https://www.ncbi.nlm.nih.gov/pubmed/28271349>  
34
- 35 Katz-Wise, S. L., Rosario, M., Calzo, J. P., Scherer, E. A., Sarda, V., & Austin, S. B. (2017b).  
36 Endorsement and Timing of Sexual Orientation Developmental Milestones Among Sexual Minority  
37 Young Adults in the Growing Up Today Study. *J Sex Res*, *54*(2), pp. 172-185.  
38 doi:10.1080/00224499.2016.1170757 Retrieved from  
39 <https://www.ncbi.nlm.nih.gov/pubmed/27148762>  
40
- 41 Kirschbaum, C., Pirke, K. M., & Hellhammer, D. H. (1993). The 'Trier Social Stress Test'--a tool for  
42 investigating psychobiological stress responses in a laboratory setting. *Neuropsychobiology*, *28*(1-  
43 2), pp. 76-81. doi:119004 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8255414>  
44
- 45 Klein, F., Sepekoff, B., & Wolf, T. J. (1990). Sexual orientation: A multivariable dynamic process. In T.  
46 Geller (Ed.), *Bisexuality: A Reader and Sourcebook*: Times Change Press.  
47
- 48 Lick, D. J., Durso, L. E., & Johnson, K. L. (2013). Minority stress and physical health among sexual  
49 minorities. *Perspect Psychol Sci*, *8*, pp. 521-548.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2 Linden, W., Earle, T. L., Gerin, W., & Christenfeld, N. (1997). Physiological stress reactivity and recovery:  
3 conceptual siblings separated at birth? *J Psychosom Res*, 42(2), pp. 117-135.  
4 doi:S0022399996002401 [pii] Retrieved from  
5 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_u](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9076640)  
6 [ids=9076640](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9076640)  
7
- 8  
9 Lovallo, W. R. (2010). Cardiovascular responses to stress and disease outcomes: a test of the reactivity  
10 hypothesis. *Hypertension*, 55(4), pp. 842-843. doi:HYPERTENSIONAHA.110.149773 [pii]  
11 10.1161/HYPERTENSIONAHA.110.149773 Retrieved from  
12 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_u](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20194293)  
13 [ids=20](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20194293)  
14 [194293](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20194293)  
15
- 16 Lovallo, W. R., & Gerin, W. (2003). Psychophysiological reactivity: mechanisms and pathways to  
17 cardiovascular disease. *Psychosom Med*, 65(1), pp. 36-45. Retrieved from  
18 <http://www.ncbi.nlm.nih.gov/pubmed/12554814>  
19
- 20 Lucas, T., Wegner, R., Pierce, J., Lumley, M. A., Laurent, H. K., & Granger, D. A. (2017). Perceived  
21 Discrimination, Racial Identity, and Multisystem Stress Response to Social Evaluative Threat  
22 Among African American Men and Women. *Psychosom Med*, 79(3), pp. 293-305.  
23 doi:10.1097/PSY.0000000000000406 Retrieved from  
24 <http://www.ncbi.nlm.nih.gov/pubmed/27806018>  
25
- 26  
27 Manuck, S. B. (1994). Cardiovascular reactivity in cardiovascular disease: "once more unto the breach". *Int*  
28 *J Behav Med*, 1(1), pp. 4-31. doi:10.1207/s15327558ijbm0101\_2 Retrieved from  
29 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_u](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16250803)  
30 [ids=16250803](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16250803)  
31
- 32  
33 McEwen, B. S. (1998). Stress, adaptation, and disease. Allostasis and allostatic load. *Ann N Y Acad Sci*,  
34 840, pp. 33-44. Retrieved from  
35 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_u](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9629234)  
36 [ids=9629234](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9629234)  
37
- 38  
39 McGregor, B. A., Carver, C. S., Antoni, M. H., Weiss, S., Yount, S. E., & Ironson, G. (2001). Distress and  
40 internalized homophobia among lesbian women treated for early stage breast cancer. *Psychology of*  
41 *Women Quarterly*, 25, pp. 1-8.  
42
- 43  
44 Mozaffarian, D., Benjamin, E. J., Go, A. S., Arnett, D. K., Blaha, M. J., Cushman, M., . . . Stroke Statistics,  
45 S. (2015). Heart disease and stroke statistics--2015 update: a report from the American Heart  
46 Association. *Circulation*, 131(4), pp. e29-322. doi:10.1161/CIR.000000000000152 Retrieved from  
47 <https://www.ncbi.nlm.nih.gov/pubmed/25520374>  
48
- 49  
50 NIH. (2014). *Monitoring adherence to the NIH policy on the inclusion of women and minorities as subjects*  
51 *in clinical research: Tracking on human subjects research as reported in the fiscal year 2009 and*  
52 *fiscal year 2010*. Retrieved from <http://orwh.od.nih.gov/resources/policyreports/index.asp>.  
53
- 54  
55 Oetjen, H., & Rothblum, E. D. (2000). When lesbians aren't gay: factors affecting depression among  
56 lesbians. *Journal of Homosexuality*, 39(49-73)  
57
- 58  
59 Pachankis, J. E. (2007). The psychological implications of concealing a stigma: a cognitive-affective-  
60 behavioral model. *Psychol Bull*, 133(2), pp. 328-345. doi:2007-02367-008 [pii]

- 1  
2 10.1037/0033-2909.133.2.328 Retrieved from  
3 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=17](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17338603)  
4 [338603](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17338603)  
5
- 6 Pearlin, L. I., & Schooler, C. (1978). The structure of coping. *J Health Soc Behav*, 19(1), pp. 2-21.  
7 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/649936>  
8
- 9 Pérez-Benitez, C. O., O'Brien, W. H., Carels, R. A., Gordon, A. K., & Chiros, C. E. (2007). Cardiovascular  
10 correlates of disclosing homosexual orientation. *Stress and Health*, 23, pp. 141-152.  
11
- 12 Roepke, S. K., Mausbach, B. T., Aschbacher, K., Ziegler, M. G., Dimsdale, J. E., Mills, P. J., . . . Grant, I.  
13 (2008). Personal mastery is associated with reduced sympathetic arousal in stressed Alzheimer  
14 caregivers. *Am J Geriatr Psychiatry*, 16(4), pp. 310-317. doi:10.1097/JGP.0b013e3181662a80  
15 Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/18378556>  
16
- 17 Rutledge, T., Linden, W., & Paul, D. (2000). Cardiovascular recovery from acute laboratory stress:  
18 reliability and concurrent validity. *Psychosom Med*, 62(5), pp. 648-654. Retrieved from  
19 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_u](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11020094)  
20 [ids=11020094](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11020094)  
21
- 22 Schwartz, A. R., Gerin, W., Davidson, K. W., Pickering, T. G., Brosschot, J. F., Thayer, J. F., . . . Linden,  
23 W. (2003). Toward a causal model of cardiovascular responses to stress and the development of  
24 cardiovascular disease. *Psychosom Med*, 65(1), pp. 22-35. Retrieved from  
25 <http://www.ncbi.nlm.nih.gov/pubmed/12554813>  
26
- 27 Simoni, J. M., Smith, L., Oost, K. M., Lehavot, K., & Fredriksen-Goldsen, K. (2017). Disparities in  
28 Physical Health Conditions Among Lesbian and Bisexual Women: A Systematic Review of  
29 Population-Based Studies. *J Homosex*, 64(1), pp. 32-44. doi:10.1080/00918369.2016.1174021  
30 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/27074088>  
31
- 32 Sindi, S., Fiocco, A. J., Juster, R. P., Pruessner, J., & Lupien, S. J. (2013). When we test, do we stress?  
33 Impact of the testing environment on cortisol secretion and memory performance in older adults.  
34 *Psychoneuroendocrinology*, 38(8), pp. 1388-1396. doi:10.1016/j.psyneuen.2012.12.004 Retrieved  
35 from <http://www.ncbi.nlm.nih.gov/pubmed/23352228>  
36
- 37 Treiber, F. A., Kamarck, T., Schneiderman, N., Sheffield, D., Kapuku, G., & Taylor, T. (2003).  
38 Cardiovascular reactivity and development of preclinical and clinical disease states. *Psychosom*  
39 *Med*, 65(1), pp. 46-62. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12554815>  
40
- 41 Wadiwalla, M., Andrews, J., Lai, B., Buss, C., Lupien, S. J., & Pruessner, J. C. (2010). Effects of  
42 manipulating the amount of social-evaluative threat on the cortisol stress response in young healthy  
43 women. *Stress*, 13(3), pp. 214-220. doi:10.3109/10253890903277561 Retrieved from  
44 <http://www.ncbi.nlm.nih.gov/pubmed/20392193>  
45
- 46 Waldo, C. R. (1999). Working in a majority context: A structural model of heterosexism as minority stress  
47 in the workplace. *Journal of Counseling Psychology*, 46, pp. 218-232.  
48
- 49 Williams, D. R., & Mohammed, S. A. (2009). Discrimination and racial disparities in health: evidence and  
50 needed research. *J Behav Med*, 32(1), pp. 20-47. doi:10.1007/s10865-008-9185-0 Retrieved from  
51 <https://www.ncbi.nlm.nih.gov/pubmed/19030981>  
52  
53  
54  
55  
56  
57  
58  
59  
60



Xu, J. Q., Murphy, S. L., Kochanek, K. D., & Bastian, B. A. (2016). *Deaths: Final data for the 2013*. Hyattsville, MD

### Table and Figure Captions

**Figure 1.** Study timeline. Ten cardiovascular recordings were taken at ten-minute intervals and aggregated into phases representing acclimation, reactivity, and recovery from the Trier Social Stress Test (TSST).


**Figure 2.** Estimated mean ( $\pm$ SE) heart rate (A–B) and mean arterial pressure (C–D) in response to the Trier Social Stress Test as a function of sexual orientation among women ( $n = 40$ ) and men ( $n = 46$ ) adjusted for race/ethnicity, education, mastery, and visit order. Note: \* =  $p < 0.05$ ; † =  $p < 0.10$ .

1  
2 **Table 1.** Sample descriptive statistics according to sexual orientation and sex ( $N = 87$ ). Group differences  
3  
4 were identified for sexual orientation and contraception variables.  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review Only



Information	Sample	Lesbian/Bisexual Women	Heterosexual Women	Gay/Bisexual Men	Heterosexual Men	<i>p</i>
<i>N</i>	87	20	20	26	21	-
<b>Demographic</b>						
Age, <i>M (SE)</i>	24.61 (0.61)	24.10 (1.34)	25.45 (1.13)	23.77 (0.98)	25.33 (1.47)	0.685
<b>Race/ethnicity</b>						
White, %	70.1	75.0	55.0	73.1	76.2	0.147
Black, %	5.7	15.0	10.0	0.0	0.0	0.147
Asian, %	12.6	0.0	10.0	23.1	14.3	0.147
Hispanic, %	6.9	0.0	20.0	3.8	4.8	0.147
Arab, %	4.6	10.0	5.0	0.0	4.8	0.147
<b>Occupation</b>						
Workers, %	34.5	40.0	35.0	23.1	42.9	0.490
Students, %	65.5	60.0	65.0	76.9	57.1	0.490
Working/studying hours/week, <i>M (SE)</i>	28.02 (1.82)	28.40 (3.65)	28.5 (4.38)	29.96 (3.24)	24.91 (3.54)	0.789
<b>Sexual orientation</b>						
Sexual attractions, <i>M (SE)</i>	3.51 (0.26)	4.75 (0.33)	1.40 (0.15)	5.96 (0.30)	1.29 (0.10)	< 0.001
Sexual behaviors, <i>M (SE)</i>	3.34 (0.26)	4.25 (0.44)	1.30 (0.11)	6.00 (0.29)	1.114 (0.8)	< 0.001
Sexual fantasies, <i>M (SE)</i>	3.61 (0.26)	5.15 (0.25)	1.65 (0.25)	5.81 (0.32)	1.29 (0.12)	< 0.001
Lifestyle preferences, <i>M (SE)</i>	3.37 (0.27)	5.15 (0.44)	1.15 (0.08)	5.54 (0.33)	1.10 (0.07)	< 0.001
Sexual identity, <i>M (SE)</i>	3.52 (0.27)	5.05 (0.43)	1.15 (0.08)	6.04 (0.26)	1.19 (0.09)	< 0.001
<b>Socio-economic</b>						
Post-secondary education, %	95.3	95.0	90.0	100.0	95.0	0.575
Personal annual income, \$ CAD, <i>M (SE)</i>	16,000 (0.17)	14,500 (0.34)	19,000 (0.53)	14,000 (0.19)	16,800 (0.33)	0.737
Household annual income, \$ CAD, <i>M (SE)</i>	32,100 (0.32)	37,000 (0.68)	25,000 (0.54)	27,100 (0.52)	39,000 (0.75)	0.311
<b>Health and well-being</b>						
Medication use, %	18.4	25.0	15.0	15.4	14.3	0.105
Oral contraceptive use, %	16.1	20.0	50.0	50.0	-	0.017
Minor physical condition, %	34.5	50.0	40.0	23.1	28.6	0.258
<b>Psychiatric history</b>						
None, %	28.7	15.0	40.0	26.9	33.3	0.522
Past history, %	8.0	15.0	5.0	11.5	33.3	0.522
Family history, %	35.6	35.0	45.0	30.8	33.3	0.522
Both past and family history, %	27.6	35.0	10.0	30.8	33.3	0.522
<b>Subjective dimensions</b>						
Self-rated health, <i>M (SE)</i>	3.72 (0.09)	3.50 (0.21)	3.95 (0.14)	3.69 (0.17)	3.75 (0.20)	0.419
Self-rated physique, <i>M (SE)</i>	3.57 (0.10)	3.15 (0.22)	3.70 (0.18)	3.27 (0.14)	3.40 (0.28)	0.284
Self-rated diet, <i>M (SE)</i>	3.30 (0.11)	3.25 (0.19)	3.40 (0.25)	3.34 (0.19)	3.15 (0.25)	0.832
<b>Behavioral</b>						
<b>Tobacco smoking</b>						
Smokers, %	11.5	5.0	10.0	11.5	19.0	0.376
Social smokers, %	14.9	10.0	5.0	19.2	23.8	0.376
Non-smokers, %	73.6	85.0	85.0	69.2	57.1	0.376
<b>Alcohol consumption (weekly)</b>						
0 or infrequently, %	25.3	25.0	50.0	18.1	4.8	0.134
1 to 5, %	40.2	45.0	40.0	26.9	52.4	0.134
6 to 10, %	26.4	25.0	10.0	38.5	28.6	0.134
11 or more, %	8.0	5.0	0.0	11.5	14.3	0.134
<b>Illicit drug use</b>						
None, %	66.7	60.0	85.0	61.5	61.9	0.334
Occasional (monthly or annually), %	24.1	35.0	5.0	30.8	23.8	0.334
Regular (daily or weekly), %	9.2	5.0	10.0	7.7	14.3	0.334
<b>Interpersonal</b>						
Single, %	72.1	60.0	80.0	80.8	65.0	0.315
Children, %	4.7	10.0	5.0	0.0	5.0	0.463
Siblings, %	86.0	95.0	85.0	80.8	85.0	0.578
Parents alive, %	94.2	95.0	90.0	92.3	100.0	0.559
Infrequent family gatherings, %	36.1	30.0	30.0	57.7	20.0	0.169
Non-religious/spiritual, %	78.8	76.5	84.2	84.0	68.4	0.569

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11





Stress **B**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29



**D**



Information	Sample	Lesbian/Bisexual Women	Heterosexual Women	Gay/Bisexual Men	Heterosexual Men	<i>p</i>
<i>N</i>	87	20	20	26	21	-
<b>Demographic</b>						
Age, <i>M (SE)</i>	24.61 (0.61)	24.10 (1.34)	25.45 (1.13)	23.77 (0.98)	25.33 (1.47)	0.685
<b>Race/ethnicity</b>						
White, %	70.1	75.0	55.0	73.1	76.2	0.147
Black, %	5.7	15.0	10.0	0.0	0.0	0.147
Asian, %	12.6	0.0	10.0	23.1	14.3	0.147
Hispanic, %	6.9	0.0	20.0	3.8	4.8	0.147
Arab, %	4.6	10.0	5.0	0.0	4.8	0.147
<b>Occupation</b>						
Workers, %	34.5	40.0	35.0	23.1	42.9	0.490
Students, %	65.5	60.0	65.0	76.9	57.1	0.490
Working/studying hours/week, <i>M (SE)</i>	28.02 (1.82)	28.40 (3.65)	28.5 (4.38)	29.96 (3.24)	24.91 (3.54)	0.789
<b>Sexual orientation</b>						
Sexual attractions, <i>M (SE)</i>	3.51 (0.26)	4.75 (0.33)	1.40 (0.15)	5.96 (0.30)	1.29 (0.10)	< 0.001
Sexual behaviors, <i>M (SE)</i>	3.34 (0.26)	4.25 (0.44)	1.30 (0.11)	6.00 (0.29)	1.114 (0.8)	< 0.001
Sexual fantasies, <i>M (SE)</i>	3.61 (0.26)	5.15 (0.25)	1.65 (0.25)	5.81 (0.32)	1.29 (0.12)	< 0.001
Lifestyle preferences, <i>M (SE)</i>	3.37 (0.27)	5.15 (0.44)	1.15 (0.08)	5.54 (0.33)	1.10 (0.07)	< 0.001
Sexual identity, <i>M (SE)</i>	3.52 (0.27)	5.05 (0.43)	1.15 (0.08)	6.04 (0.26)	1.19 (0.09)	< 0.001
<b>Socio-economic</b>						
Post-secondary education, %	95.3	95.0	90.0	100.0	95.0	0.575
Personal annual income, \$ CAD, <i>M (SE)</i>	16,000 (0.17)	14,500 (0.34)	19,000 (0.53)	14,000 (0.19)	16,800 (0.33)	0.737
Household annual income, \$ CAD, <i>M (SE)</i>	32,100 (0.32)	37,000 (0.68)	25,000 (0.54)	27,100 (0.52)	39,000 (0.75)	0.311
<b>Health and well-being</b>						
Medication use, %	18.4	25.0	15.0	15.4	14.3	0.105
Oral contraceptive use, %	16.1	20.0	50.0	23.1	-	0.017
Minor physical condition, %	34.5	50.0	40.0	26.9	28.6	0.258
<b>Psychiatric history</b>						
None, %	28.7	15.0	40.0	11.5	33.3	0.522
Past history, %	8.0	15.0	45.0	30.8	33.3	0.522
Family history, %	35.6	35.0	10.0	30.8	33.3	0.522
Both past and family history, %	27.6	35.0	10.0	30.8	33.3	0.522
<b>Subjective dimensions</b>						
Self-rated health, <i>M (SE)</i>	3.72 (0.09)	3.50 (0.21)	3.95 (0.14)	3.69 (0.17)	3.75 (0.20)	0.419
Self-rated physique, <i>M (SE)</i>	3.57 (0.10)	3.15 (0.22)	3.70 (0.18)	3.27 (0.14)	3.40 (0.28)	0.284
Self-rated diet, <i>M (SE)</i>	3.30 (0.11)	3.25 (0.19)	3.40 (0.25)	3.34 (0.19)	3.15 (0.25)	0.832
<b>Behavioral</b>						
<b>Tobacco smoking</b>						
Smokers, %	11.5	5.0	10.0	11.5	19.0	0.376
Social smokers, %	14.9	10.0	5.0	19.2	23.8	0.376
Non-smokers, %	73.6	85.0	85.0	69.2	57.1	0.376
<b>Alcohol consumption (weekly)</b>						
0 or infrequently, %	25.3	25.0	50.0	18.1	4.8	0.134
1 to 5, %	40.2	45.0	40.0	26.9	52.4	0.134
6 to 10, %	26.4	25.0	10.0	38.5	28.6	0.134
11 or more, %	8.0	5.0	0.0	11.5	14.3	0.134
<b>Illicit drug use</b>						
None, %	66.7	60.0	85.0	61.5	61.9	0.334
Occasional (monthly or annually), %	24.1	35.0	5.0	30.8	23.8	0.334
Regular (daily or weekly), %	9.2	5.0	10.0	7.7	14.3	0.334
<b>Interpersonal</b>						
Single, %	72.1	60.0	80.0	80.8	65.0	0.315
Children, %	4.7	10.0	5.0	0.0	5.0	0.463
Siblings, %	86.0	95.0	85.0	80.8	85.0	0.578
Parents alive, %	94.2	95.0	90.0	92.3	100.0	0.559
Infrequent family gatherings, %	36.1	30.0	30.0	57.7	20.0	0.169
Non-religious/spiritual, %	78.8	76.5	84.2	84.0	68.4	0.569