




Texas Agricultural Extension Service

# Feline Immunodeficiency Virus



This publication was produced jointly  
by the  
Texas A&M University College of  
Veterinary Medicine  
and the  
Texas Agricultural Extension Service

Alice M. Wolf  
and Bruce Lawhorn\*

Feline immunodeficiency virus (FIV) causes chronic disease in cats (acquired immunodeficiency syndrome, or AIDS) similar to the disease caused by human immunodeficiency virus (HIV) in human beings. Although the two viruses belong to the same family, the feline virus is not transmissible to human beings and the human virus is not transmissible to cats.

FIV infection has been found in cats worldwide. The disease occurs more commonly in male cats than in females; the highest incidence is in cats older than 5 years of age. Free-roaming cats and mixed breed cats are at highest risk for contracting FIV infection. The disease is rare in cats housed strictly indoors or in catteries.

FIV is transmitted from cat to cat very efficiently by cat bite wounds and blood transfusions. Infection by casual contact (e.g., using common food bowls or litter pans) does not occur. To date, it does not appear that an infected female can pass the virus to her kittens before they are born or through her milk. Again, FIV is very specifically adapted to infect cats; human beings are not at risk from this disease.

Several weeks following infection with FIV the cat will have a short, mild illness with fever and lymph node enlargement. Often the signs of illness during this stage are so subtle that the owner will not notice that the cat is sick. These early signs will disappear without treatment after a few days or weeks. The virus then undergoes a period of dormancy. Although FIV is still present in the cat, the animal will be apparently healthy and show no signs of illness for months or years.

Eventually the virus will become reactivated and upset the balance of the cat's immune system. At this time the cat will begin to show signs of chronic infections of the mouth (gingivitis), nasal cavity (rhinitis and sinusitis) or skin. Chronic intractable diarrhea occurs in some cats; tumors appear in others. Neurologic signs such as behavioral changes, dementia, aggression, inappropriate elimination behavior and seizures have been seen in some FIV-infected cats.

---

\* Associate professor of small animal medicine and surgery and Extension veterinarian, The Texas A&M University System.

The diagnosis of FIV infection is confirmed by finding antibodies to the FIV virus in the cat's blood. This test can be performed by your veterinarian. Any sick cat should be tested because of the association of FIV with chronic illness in the cat. Because apparently healthy cats may harbor FIV in a dormant state for many years, it is wise to also test new adult cats coming into a household and cats entering closed populations, especially catteries.

There is no specific treatment for FIV infection. Drugs currently used against HIV infection (e.g., AZT) are toxic to cats. Veterinarians treat the symptoms, such as infection or cancer, and give supportive care. Unfortunately, the outlook for cats in the advanced stage of this disease is poor. Even if the signs of illness can be temporarily controlled, FIV infection is ultimately fatal to the cat.

At this time there is no vaccine for preventing FIV infection. You can reduce the risk by neutering your cat to reduce roaming and territorial fighting behavior, and by keeping your cat indoors.

*Educational programs conducted by the Texas Agricultural Extension Service serve people of all ages regardless of socioeconomic level, race, color, sex, religion, handicap or national origin.*

---

Issued in furtherance of Cooperative Extension Work in Agriculture and Home Economics, Acts of Congress of May 8, 1914, as amended, and June 30, 1914, in cooperation with the United States Department of Agriculture. Zerle L. Carpenter, Director, Texas Agricultural Extension Service, The Texas A&M University System.