

Towards evidence based m-health application design in cancer patient healthy lifestyle interventions.

Panagiotis E. Antoniou

Medical Physics Laboratory, Medical School
Aristotle University of Thessaloniki
Thessaloniki, Greece
pantonio@otenet.gr

Maria Karagianni

Medical Physics Laboratory, Medical School
Aristotle University of Thessaloniki
Thessaloniki, Greece
mkaragiannipsy@gmail.com

Octavio Rivera-Romero

Departamento de Tecnología Electrónica
Universidad de Sevilla
Seville, Spain
orivera@us.es

Panagiotis D. Bamidis

Medical Physics Laboratory, Medical School
Aristotle University of Thessaloniki
Thessaloniki, Greece
bamidis@med.auth.gr

Abstract— Cancer is one of the most prevalent diseases in Europe and the world. Significant correlations between dietary habits and cancer incidence and mortality have been confirmed by the literature. Physical activity habits are also directly implicated in the incidence of cancer. Lifestyle behaviour change may be benefited by using mobile technology to deliver health behaviour interventions. M-Health offers a promising cost-efficient approach to deliver en-masse interventions. Smartphone apps with constructs such as gamification and personalized have shown potential for helping individuals lose weight and maintain healthy lifestyle habits. However, evidence-based content and theory-based strategies have not been incorporated by those apps systematically yet. The aim of the current work is to put the foundations for a methodologically rigorous exploration of wellness/health intervention literature/app landscape towards detailed design specifications for connected health m-apps. In this context, both the overall work plan is described as well as the details for the significant steps of application space and literature space review. Both strategies for research and initial outcomes of it are presented. The expected evidence based design process for patient centered health and wellness interventions is going to be the primary input in the implementation process of upcoming patient centered health/wellness m-health interventions.

Keywords: *m-health; cancer; health apps; application design; evidence centered design*

I. INTRODUCTION

In Europe in 2012, 3.45 million new cases of cancer and 1.75 million deaths from cancer were estimated [1]. Breast cancer (464,000 cases), colorectal (447,000), prostate (417,000) and lung (410,000) were the most common types of cancer, representing half of the overall burden of cancer in Europe. Significant correlations between dietary habits and cancer incidence and mortality have been confirmed by

ecological studies [2][3]. Physical activity habits are directly implicated in the incidence of cancer [4].

Ehberman et al. 2013 reported cancers associated with excess weight and lack of sufficient physical activity. An increased risk of adenocarcinoma of esophagus, colon and rectal cancer, pancreas cancer, kidney cancer, and endometrial and postmenopausal breast cancer has been associated with excess weight [5], [6], [7], [8]. Also, there are evidences suggesting an association between an increased risk of gallbladder, liver, thyroid, and hematopoietic cancers and excess weight [5]. On the other hand, an increased risk of colon, endometrial, and breast cancers may be associated with lack of physical activity [5], [9], [10], [11], [12].

Andersson et al. [13] conducted a review of the current evidence related to excess body fat and cancer. The incident cancer burden attributable to excess body mass in 30 Europe countries was estimated by Renehan et al. [14] finding a risk of 3.2% in men and 8.6% in women. Evidences from prospective studies have confirmed associations between excess weight and thyroid, liver, leukaemia, malignant melanoma, multiple myeloma, and non- Hodgkin lymphoma [15], gastric cardia cancer [16], and advanced prostate cancer [17], [18].

Healthy lifestyle recommendations are not only suggested for cancer prevention but also for patients and survivors. Cancer prognosis and quality of life among cancer survivors may be negatively affected by excess weight and lack of physical activity. There are studies reporting excess weight is associated with poorer survival among patients with breast cancer [19], [20], [21] and colorectal cancer [22], [23], [24], [25]. Obesity is associated with mortality and incidence of late-stage disease but not with incidence of early-stage disease among prostate cancer patients [25]. Physical activity after diagnosis of breast or colon cancer may have reduced cancer-specific mortality. Physical activity and physical exercise interventions may reduce

disease and treatment-related symptoms of prostate cancer patients during and after treatments, improving their quality of life [26], [27], [28], [29], [30].

Co-morbid chronic diseases are common among cancer patients/survivors. Cancer and its treatment are associated with an increased risk for co-morbid conditions including heart disease, diabetes, and stroke [31]. Type 2 diabetes has been associated with colon, breast, and pancreas cancer [1]. Patients with type 2 diabetes are at increased risk of liver, pancreas, endometrium, colorectal, breast, and bladder cancers [32], [33], [34]. Risk reduction of type 2 diabetes and cardiovascular disease may be achieved by body weight reduction [1]. Lifestyle modification programmes involving diet, exercise, and behavioural techniques has been effective to reduce weight and to avoid weight gain [1].

Finally, cancer patients and survivors' mental wellbeing, psychological and psychosocial factors are also affected by diet and physical activity. Depression is common among cancer patients/survivors [35], [36], [37], [38], (Honda & Goodwin 2004) affecting negatively the physical and psychological health of cancer survivors. Depression is also associated with obesity, diabetes, and cardiovascular disease [39], [40], [41]. Exercise has been identified as a treatment to mitigate depression symptoms in cancer survivors [42], [43].

Lifestyle behaviour change may be benefited by using mobile technology to deliver health behaviour interventions [44], [45], [46]. mHealth offers a promising cost-efficient approach to deliver them enabling in masse interventions [47], [48], [49]. Smartphone apps have shown potential for helping some individuals lose weight and maintain healthy lifestyle habits [44], [47]. Behavior change techniques, gamification and personalization features should be included in apps to promote healthy lifestyle. However, evidence-based content [48], [49], [50] and theory-based strategies [46], [51] could not be incorporated by those apps as was revealed in Breton et al. 2011 [50].

Cowan et al. performed a content analysis of apps primarily aimed at increasing exercise [51]. They found that overall the apps contained few features based on behavioral change theory.

Azar et al. conducted a comparative, descriptive assessment of the top-rated free apps to assist with weight management in iTunes App Store [52]. The objective of this review was to evaluate diet/nutrition and anthropometric tracking apps based on incorporation of features consistent with behaviour change theories. 23 apps were analysed. They found low overall scores for inclusion of behavioural theory-based strategies.

Helf & Hlavacs performed a critical review of motivational tools (gamification) to increase user engagement included in health-related apps [53]. They found low scores in dimensions requiring interdisciplinary work, such as designing interventions based on scientific theories or content personalization based on personality types or traits. From the previous discourse it is clear that there is a significant body of literature that explores health and wellness mobile applications. With this fact in mind this work aims to put the foundations for exploring this literature

towards evidence based m-health application design for prevalent chronic diseases such as breast and prostate cancer.

II. METHODS

Towards the aforementioned goal, a survey of publications was conducted. This survey assess the efficacy of mobile/wearables applications for maintaining healthy lifestyles or affecting healthy behavioral change. In this context, an initial tentative research question was formulated:

What are the most effective app features in the whole wearables/mobiles ecosystem (including gamified and AR/VR apps) for maintaining healthy lifestyles or affecting health behavioral change?

From answering this research question it would be possible to refine a list of effective application features that would be used as the pool from which to draw when designing applications for specific topics, diseases and comorbidities. For that, a literature and application review was organized for mobile, tablet, and wearables features for maintaining healthy lifestyle and affecting healthy behavioral change. Initial constraints were:

- A Focus on multilingual applications
 - A Focus on features that are relevant to exercise motivation and nutritional improvement
 - Inclusion of applications that are available to both mobile devices and wearables.
- Within these constraints a detailed work plan was devised consisting of the following specific tasks:
- Definition of the conceptual review field.
 - Definition of the application review field.
 - Definition of Search engines.
 - Definition of keywords subcategories
 - Behavioral change theories, behavioral change constructs
 - Specific diseases and demographics
 - mobile/wearables app ecosystem features
 - Exclusion criteria
 - Focus on excluding publications that are not providing evidence based direct or indirect links between relevant topics
 - Focus on app exclusion criteria based on non-international market reach, language barrier, popularity threshold, non-proper feature documentation.
 - Conducting the Literature Survey.
 - Initial Keyword search in the literature; archiving of results.
 - Application of exclusion criteria.
 - Between observer consistency check
 - Conducting the app stores' keyword search
 - Application of exclusion criteria
 - Archiving of final results.
 - Analysis of discovered data.
 - Aggregation of reported efficacy links between behavioral change theories and specific diseases and target groups. Distilling of most impactful behavioral change theories within the scope of the selected diseases and target groups

- Aggregation of reported links between behavioral theory constructs and behavioral change theories. Distilling of behavioral change constructs in scope with previously determined theories
- Aggregation of literature reported links between behavioral change constructs or theories and mobile/wearable applications' features. Distilling of application features relevant to previously determined constructs and theories.
- Aggregation of literature reported efficacy of application features in motivating behavioral change for healthy exercise and nutrition habits (from strand 1 described above)
- Aggregation of feature proliferation in the existing mobiles/wearables application ecosystem.
- Conceptual, evidence supported, design guidelines formulation
 - Critical revision of aggregated material.
 - Hierarchical classification of shortlisted app features based on implicit impact to specific demographic and disease
 - Hierarchical classification of shortlisted app features based on proliferation in the overall application ecosystem.
 - Formulation of feature guidelines regarding application design for the specific demographics and diseases.
- Presentation and finalization of the work.
 - Drafting and publication of a research paper with the outcomes of this work.

For the review part of this endeavor it was determined that the research would follow a scoping review method. This is an appropriate method to systematically scan and evaluate which studies are within or out of the scope of the research area that is explored for evidence [54]. Different types of literature review methods were considered, but a scoping review study best fit our research purpose with the emphasis placed on the scoping technique to “map” relevant literature in the field of interest rather than collecting similar evidence for a highly focused research question. The method is effective in identifying gaps in the evidence base where no research has been conducted and identifying emerging results in new fields of research; The methodological framework of Arksey and O'Malley [55] is adhered to as it aligns with the proposed work plan. This consists of five stages of scoping and review:

- Identify the research question,
- Identify relevant studies,
- Select studies,
- Chart the data, and
- Collate, summarize, and report the results.

III. RESULTS - OUTCOMES

Having established the previous methodologic foundation the final research questions for this work were formulated:

- *What are the literature validated behavioral change constructs that support the efficacy of the various mobile and tablet application features in healthy lifestyle change and maintenance apps?*

• *What is the minimum specific set of behavioral change theories that are the literature validated to be effective in motivating breast cancer patients for healthy lifestyle?*

• *What is the minimal ideal feature set for maximum impact of mobile apps for healthy lifestyle maintenance/change on breast cancer patients based on existing behavioral theory research?*

To move into the process of identifying relevant studies the keywords that were considered based on BCT experience and topical research (cf. introduction) are summarized in Table I.

TABLE I Keywords (per broad subject)

Behavioral change theories	Behavioral change techniques	Mobile application Topics and features
Theory of reasoned action/planned behavior (TPB)	Stress management, General communication skills training, Model/demonstrate the behavior, Relapse prevention/coping planning, Facilitate social comparison, Goal setting (behavior), Action planning, Provide feedback on performance, Barrier identification/problem solving, Provide instruction, Teach to use prompts/cues, Provide normative information about others' behavior, Plan social support/social change, Provide rewards for behavior, Prompt self-monitoring of behavior, Provide information on the consequences in general, Provide information on the consequences for individual,	Mobile OR Ubiquitous OR Tablet
Transtheoretical model (TTM)		
Social cognitive theory (SCT)		
Elaboration likelihood model (ELM)		Phone
Extended parallel process model (EPPM)		Physical activity
Self-regulation theory (SRT)		Exercise
Precaution adoption process model (PAPM)		Health OR Healthy
Diffusion of innovations model (DIM)		Diet
Health belief model (HBM)		Food
Social		

norms theory (SNT)	approximation, response differentiation, avoidance, sensitization, desensitization, flooding, aversive conditioning/punishment, contingency management, convert control, Contingency management for behavior elimination : time-out, response differentiation, token economy, response cost, negative practice, flooding- response prevention	
--------------------	---	--

TABLE II Search engines to include in the keyword search.

Literature search engines included for the scoping review keyword search.
PubMed
IEEE xplora
ACM digital library
Web of Science
Cochrane Library
ERIC
Embase
SCOPUS
Google scholar
BASE

Practically these keywords were organized in formal queries in several search engines (Table 2). An example query for each research question is presented for PubMed in Table 3 but shall be implemented to all the aforementioned search engines where there is capacity for advanced searches. It must be noted that the behavioral change techniques were deemed too numerous to include explicitly, thus only the theories behind them were used in the queries.

TABLE III Search engines to include in the keyword search.

PubMed search argument:		
Q1) Cancer and physical activity mobile apps.	("Mobile" [TIAB] OR "Ubiquitous" [TIAB] OR "Tablet" [TIAB] OR "Phone" [TIAB]) AND ("Physical activity" [TIAB] OR "Exercise" [TIAB] OR "Diet" [TIAB] OR "Dietary" [TIAB] OR "Food" [TIAB] OR "Healthy Lifestyle"[TIAB]) AND ("Cancer"[All Fields] OR "Oncology"[All Fields] OR "Neoplasm" [All Fields] OR "Neoplastic" [All Fields])	
Q2) Behavioral change theories and mobile apps.	("Mobile" [TIAB] OR "Ubiquitous" [TIAB] OR "Tablet" [TIAB] OR "Phone" [TIAB]) AND ("Behavioral Change Theory"[All Fields] OR "Theory of reasoned action/planned behavior" [All Fields] OR "Transtheoretical model" [All Fields] OR "Social cognitive theory" [All Fields] OR "Elaboration likelihood model" [All Fields] OR "Extended parallel process model" [All Fields] OR "Self-regulation theory" [All Fields] OR "Precaution adoption process model" [All Fields] OR "Diffusion of	

	innovations model" OR "Health belief model" [All Fields] OR "Social norms theory" [All Fields] OR "Learning Theory" [All Fields])
--	---

Regarding the search strategy in the app stores (Google Play & App store), due to the fact that a manual search in the Google play store as well as the Apple App store would be an encyclopedic task it was decided for the 42matters (<https://42matters.com/app-market-data>) search API to be utilized in order to perform this endeavor.

IV. DISCUSSION

The overall aim of this work is to create a conceptual, evidence based design recommendation towards developing mobile and wearables applications to support patients suffering from the specific chronic diseases and belonging in the specific demographic. A mapping of the conceptual framework behind this strand is presented in Figure 1.




Figure 1 Conceptual framework of implicit link between specific app features and specific disease and demographics.

To proceed towards this endeavor we aim to

- Critically explore research linking app features with behavioral constructs
- Critically explore research linking specific behavioral constructs with overall behavioral change theories.
- Critically explore research about the efficacy of the aforementioned theories and constructs on our joint endeavor's specific diseases and comorbidities (Breast and Prostate cancer and their comorbidities especially in the elderly) so that
- A literature based, evidence supported, efficacy link can be formulated between specific mobile/wearable application features and the specific diseases and comorbidities of interest to the joint research group.

Previous work from members of this research team in this strand has a demonstrated experience in gamified engagement endeavors for the elderly. It has developed and piloted its own elderly exergaming platform with published results [60],[61]. Furthermore it has experience in ecologic validity of living lab environments having established in its premises a prototype Living Lab. This living lab consists of an environment equipped with monitoring sensors and analytics software for performing technology assessment in

ecologically valid environments and stealth assessment of cognitive and physical capacities of the elderly [62]. Furthermore, this research team has extensive experience in integrating mobile and wearable devices in elderly ambient assisted living environments, with EU funded research providing published results in these endeavors [63]. These expected outcomes of this work are the necessary refinement effort that focuses on describing the conceptual basis and initial methodological findings that align behavioral change theories and the means to apply them. The expected evidence based design process for patient centered health and wellness interventions is going to be the primary input in the implementation process of upcoming patient centered health/wellness m-health interventions.

ACKNOWLEDGMENT

This work has, in part been supported by the ENJECT COST action (COST Action TD1405 - Reference code: COST-STSM-ECOST-STSM-TD1405-220216-070451) (DISCOVER) through the Short Term Scientific Missions instrument (STSM) that P.A. and O.R. collaborated in.

REFERENCES

- [1] Ferlay J, Steliarova-Foucher E, Lortet-Tieulent J, Rosso S, Coebergh JW, Comber H, Forman D, Bray F. Cancer incidence and mortality patterns in Europe: estimates for 40 countries in 2012. *Eur J Cancer*. 2013 Apr;49(6):1374-403. doi: 10.1016/j.ejca.2012.12.027. PubMed PMID: 23485231.sfgdfg
- [2] Norat, T., Scoccianti, C., Boutron-Ruault, M.-C., Anderson, A., Berrino, F., Cecchini, M., ... Romieu, I. (2015). European Code against Cancer 4th Edition: Diet and cancer. *Cancer Epidemiology*, 39 Suppl 1, S56-66. <http://doi.org/10.1016/j.canep.2014.12.016>
- [3] Scocciantia C, Cecchinib M, Anderson AS, Berrinod F, Boutron-Ruaulte M, Espinaa C, et al. European Code Against Cancer. In: Alcohol drinking and Cancer; 4th ed. 2015. <http://dx.doi.org/10.1016/j.canep.2015.01.007.dfgdfgfdg>
- [4] Leitzmann M, Powers H, Anderson AS, Berrino F, Boutron-Ruaulte M, Cecchini M, et al. European Code Against Cancer. In: Physical Activity and Cancer; 4th ed. 2015. <http://dx.doi.org/10.1016/j.canep.2015.03.009>
- [5] World Cancer Research Fund/American Institute for Cancer Research (WCRF/AICR). Food, Nutrition, Physical Activity, and the Prevention of Cancer: A Global Perspective. Washington, DC: American Institute for Cancer Research; 2007
- [6] Ning Y, Wang L, Giovannucci EL. A quantitative analysis of body mass index and colorectal cancer: findings from 56 observational studies. *Obes Rev*. 2010; 11:19–30. [PubMed: 19538439]
- [7] World Cancer Research Fund/American Institute for Cancer Research (WCRF/AICR). The Associations between Food, Nutrition and Physical Activity and the Risk of Breast Cancer: WCRF/AICR Systematic Literature Review Continuous Update Report. Washington, DC: American Institute for Cancer Research; 2008.
- [8] Crosbie EJ, Zwahlen M, Kitchener HC, Egger M, Renehan AG. Body mass index, hormone replacement therapy, and endometrial cancer risk: a meta-analysis. *Cancer Epidemiol Biomarkers Prev*. 2010; 19:3119–3130. [PubMed: 21030602]
- [9] Physical Activity Guidelines Advisory Committee. Physical Activity Guidelines Advisory Committee Report (PAGACR), 2008. Washington, DC: US Department of Health and Human Services; 2008.
- [10] Friedenreich CM, Cust AE. Physical activity and breast cancer risk: impact of timing, type and dose of activity and population subgroup effects. *Br J Sports Med*. 2008; 42:636–647. [PubMed: 18487249]
- [11] Wolin KY, Yan Y, Colditz GA, Lee IM. Physical activity and colon cancer prevention: a meta-analysis. *Br J Cancer*. 2009; 100:611–616. [PubMed: 19209175]
- [12] Moore SC, Gierach GL, Schatzkin A, Matthews CE. Physical activity, sedentary behaviours, and the prevention of endometrial cancer. *Br J Cancer*. 2010; 103:933–938. [PubMed: 20877336]
- [13] Anderson, A. S., Key, T. J., Norat, T., Scoccianti, C., Cecchini, M., Berrino, F., ... Romieu, I. (2015). European Code against Cancer 4th Edition: Obesity, body fatness and cancer. *Cancer Epidemiology*, 39 Suppl 1, S34-45. <http://doi.org/10.1016/j.canep.2015.01.017>
- [14] Renehan AG, Soerjomataram I, Tyson M, Egger M, Zwahlen M, Coebergh JW, et al. Incident cancer burden attributable to excess body mass index in 30 European countries. *Int J Cancer*. 2010;126(3):692–702
- [15] Renehan AG, Tyson M, Egger M, Heller RF, Zwahlen M. Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet* 2008;371(9612):569–78
- [16] Chen Y, Liu L, Wang X, Wang J, Yan Z, Cheng J, et al. Body mass index and risk of gastric cancer: a meta-analysis of a population with more than ten million from 24 prospective studies. *CEBP* 2013;22(8):1395–408
- [17] Discacciati A, Orsini N, Wolk A. Body mass index and incidence of localized and advanced prostate cancer-a dose-response meta-analysis of prospective studies. *Ann Oncol* 2012; 23(7):1665–71.
- [18] World Cancer Research Fund International (WCRFI). Diet, nutrition, physical activity and prostate cancer; 2014, <http://www.wcrf.org/sites/default/files/Prostate-Cancer-2014-Report.pdf>.
- [19] Ewertz M, Jensen M-B, Gunnarsdottir KA, et al. Effect of obesity on prognosis after early-stage breast cancer. *J Clin Oncol*. 2011; 29:25–31. [PubMed: 21115856]
- [20] Sinicropo FA, Dannenberg AJ. Obesity and breast cancer prognosis: weight of the evidence. *J Clin Oncol*. 2011; 29:4–7. [PubMed: 21115867]
- [21] Protani M, Coory M, Martin J. Effect of obesity on survival of women with breast cancer: systematic review and meta-analysis. *Breast Cancer Res Treat*. 2010; 123:627–635. [PubMed: 20571870]
- [22] Meyerhardt JA, Ma J, Courmeya KS. Energetics in colorectal and prostate cancer. *J Clin Oncol*. 2010; 28:4066–4073. [PubMed: 20644082]
- [23] Sinicropo FA, Foster NR, Sargent DJ, O'Connell MJ, Rankin C. Obesity is an independent prognostic variable in colon cancer survivors. *Clin Cancer Res*. 2010; 16:1884–1893. [PubMed: 20215553]
- [24] Prizment AE, Flood A, Anderson KE, Folsom AR. Survival of women with colon cancer in relation to precancer anthropometric characteristics: the Iowa Women's Health Study. *Cancer Epidemiol Biomarkers Prev*. 2010; 19:2229–2237. [PubMed: 20826830]
- [25] Cao Y, Ma J. Body mass index, prostate cancer-specific mortality, and biochemical recurrence: a systematic review and meta-analysis. *Cancer Prev Res*. 2011; 4:486–501.
- [26] Courmeya KS. Exercise in cancer survivors: an overview of research. *Med. Sci. Sports Exerc*. 35 (2003) 1846–1852.
- [27] Galvao DA, Newton RU. Review of exercise intervention studies in cancer patients. *J. Clin. Oncol*. 23 (2005) 899–909.
- [28] Knols R, Aaronson NK, Uebelhart D, Fransen J, Aufdemkampe G. Physical exercise in cancer patients during and after medical treatment: a systematic review of randomized and controlled clinical trials. *J. Clin. Oncol*. 23 (2005) 3830–3842.
- [29] Schmitz KH, Holtzman J, Courmeya KS, Masse LC, Duval S, Kane R. Controlled physical activity trials in cancer survivors: a systematic review and meta-analysis. *Cancer Epidemiol. Biomark. Prev*. 14 (2005) 1588–1595.
- [30] Spence RR, Heesch KC, Brown WJ. Exercise and cancer rehabilitation: a systematic review. *Cancer Treat. Rev*. 36 (2010) 185–194

- [31] Klabunde CN, Legler JM, Warren JL, Baldwin LM, Schrag D. A refined comorbidity measurement algorithm for claims-based studies of breast, prostate, colorectal, and lung cancer patients. *Ann Epidemiol* 2007;17:584–90
- [32] Vigneri P, Frasca F, Sciacca L, Pandini G, Vigneri R. Diabetes and cancer. *Endocr Relat Cancer* 2009;16(4):1103–23.
- [33] Deng L, Gui Z, Zhao L, Wang J, Shen L. Diabetes mellitus and the incidence of colorectal cancer: an updated systematic review and meta-analysis. *Dig Dis Sci* 2012;57(6):1576–85.
- [34] Lee JY, Jeon I, Kim JW, Song YS, Yoon JM, Park SM. Diabetes mellitus and ovarian cancer risk: a systematic review and meta-analysis of observational studies. *Int J Gynecol Cancer* 2013;23(3):402–12.
- [35] Massie MJ. Prevalence of depression in patients with cancer. *J Natl Cancer Inst Monogr* 2004;57–71.
- [36] Haisfield-Wolfe ME, McGuire DB, Soeken K, Geiger-Brown J, De Forge BR. Prevalence and correlates of depression among patients with head and neck cancer: a systematic review of implications for research. *Oncol Nurs Forum* 2009;36:E107–25
- [37] Pirl WF. Evidence report on the occurrence, assessment, and treatment of depression in cancer patients. *J Natl Cancer Inst Monogr* 2004;32–9
- [38] Deschields T, Tibbs T, Fan MY, Taylor M. Differences in patterns of depression after treatment for breast cancer. *Psychooncology* 2006; 15:398–406
- [39] Arroyo C, Hu FB, Ryan LM, Kawachi I, Colditz GA, Speizer FE, et al. Depressive symptoms and risk of type 2 diabetes in women. *Diabetes Care* 2004;27:129–33.
- [40] Carnethon MR, Kinder LS, Fair JM, Stafford RS, Fortmann SP. Symptoms of depression as a risk factor for incident diabetes: findings from the National Health and Nutrition Examination Epidemiologic Follow-up Study, 1971–1992. *Am J Epidemiol* 2003;158:416–23.
- [41] Ferketich AK, Frid DJ. Depression and coronary heart disease: a review of the literature. *Clin Geriatrics* 2001;9:1–8
- [42] Fleishman SB. Treatment of symptom clusters: pain, depression, and fatigue. *J Natl Cancer Inst Monogr* 2004;119–23
- [43] Craft, L. L., Vaniterson, E. H., Helenowski, I. B., Rademaker, A. W., & Courneya, K. S. (2012). Exercise effects on depressive symptoms in cancer survivors: a systematic review and meta-analysis. *Cancer Epidemiology, Biomarkers & Prevention : A Publication of the American Association for Cancer Research, Cosponsored by the American Society of Preventive Oncology*, 21(1), 3–19. <http://doi.org/10.1158/1055-9965.EPI-11-0634>
- [44] Stephens J, Allen JK, Dennison Himmelfarb CR. "Smart" coaching to promote physical activity, diet change, and cardiovascular health. *J Cardiovas Nurs* 2011;26(4):282–4
- [45] Patrick K, Raab F, Adams MA, et al. A text message-based intervention for weight loss: randomized controlled trial. *J Med Internet Res* 2009;11(1):e1.
- [46] Riley WT, Rivera DE, Atienza AA, Nilsen W, Allison SM, Mermelstein R. Health behavior models in the age of mobile interventions: are our theories up to the task? *Transl Behav Med* 2011;1(1):53–71.
- [47] Stephens J, Allen J. Mobile phone interventions to increase physical activity and reduce weight: a systematic review. *J Cardiovasc Nurs* 2013;28(4):320–9.
- [48] Hebden L, Cook A, van der Ploeg HP, Allman-Farinelli M. Development of smartphone applications for nutrition and physical activity behavior change. *JMIR Res Protoc* 2012;1(2):e9.
- [49] Backinger CL, Augustson EM. Where there's an app, there's a way? *Am J Prev Med* 2011;40(3):390–1
- [50] Breton E, Fuemmeler B, Abrams L. Weight loss—there is an app for that! But does it adhere to evidence-informed practices? *Transl Behav Med* 2011;1(4):523–9
- [51] Cowan LT, Van Wagenen SA, Brown BA, et al. Apps of steel: are exercise apps providing consumers with realistic expectations?: a content analysis of exercise apps for presence of behavior change theory. *Health Educ Behav* 2013;40(2):133–9.
- [52] Azar, K. M. J., Lesser, L. I., Laing, B. Y., Stephens, J., Aurora, M. S., Burke, L. E., & Palaniappan, L. P. (2013). Mobile Applications for Weight Management. *American Journal of Preventive Medicine*, 45(5), 583–589. <http://doi.org/10.1016/j.amepre.2013.07.005>
- [53] Helf, C., & Hlavacs, H. (2016). Apps for life change: Critical review and solution directions. *Entertainment Computing*, 14, 17–22. <http://doi.org/10.1016/j.entcom.2015.07.001>
- [54] Honda K, Goodwin RD. Cancer and mental disorders in a national community sample: findings from the national comorbidity survey. *Psychother Psychosom* 2004;73:235–42
- [55] Levac, D., Colquhoun, H., & O'Brien, K. K. (2010). Scoping studies: advancing the methodology. *Implementation Science : IS*, 5, 69. <http://doi.org/10.1186/1748-5908-5-69>
- [56] Arksey, H., & O'Malley, L. (2005). Scoping studies: towards a methodological framework. *International Journal of Social Research Methodology*, 8(1), 19–32. <http://doi.org/10.1080/1364557032000119616>
- [57] Daley, A. J., Crank, H., Saxton, J. M., Mutrie, N., Coleman, R., & Roalfe, A. (2007). Randomized trial of exercise therapy in women treated for breast cancer. *Journal of Clinical Oncology*, 25(13), 1713–1721. <http://doi.org/10.1200/JCO.2006.09.5083>
- [58] Jones, L. W., Courneya, K. S., Fairey, A. S., & Mackey, J. R. (2005). Does the theory of planned behavior mediate the effects of an oncologist's recommendation to exercise in newly diagnosed breast cancer survivors? Results from a randomized controlled trial. *Health Psychology*, 24(2), 189–97. <http://doi.org/10.1037/0278-6133.24.2.189>
- [59] Vallance, J. K. H., Courneya, K. S., Plotnikoff, R. C., Yasui, Y., & Mackey, J. R. (2007). Randomized controlled trial of the effects of print materials and step pedometers on physical activity and quality of life in breast cancer survivors. *Journal of Clinical Oncology*, 25(17), 2352–2359. <http://doi.org/10.1200/JCO.2006.07.9988>
- [60] Webb, T. L., Joseph, J., Yardley, L., & Michie, S. (2010). Using the internet to promote health behavior change: a systematic review and meta-analysis of the impact of theoretical basis, use of behavior change techniques, and mode of delivery on efficacy. *Journal of Medical Internet Research*, 12(1), 1–18. <http://doi.org/10.2196/jmir.1376>
- [61] E.I. Konstantinidis, A.S. Billis, C.A. Mouzakidis, V.I. Zilidou, P.E. Antoniou, P.D. Bamidis. Design, Implementation, and Wide Pilot Deployment of FitForAll: An Easy to use Exergaming Platform Improving Physical Fitness and Life Quality of Senior Citizens. *IEEE J Biomed Health Inform.* 2016 Jan; 20(1):189-200. doi: 10.1109/JBHI.2014.2378814. PubMed PMID: 26731797.
- [62] P.D. Bamidis, E.I. Konstantinidis, A.S. Billis, G. Babaropoulos, P.E. Antoniou, C. Pappas, and A. Siountas. Apps, services and new exergaming paradigms for home use by older people in 6th European Conference of the International Federation for Medical and Biological Engineering (MBEC 2014) 7-11 September 2014, Dubrovnik, Croatia.
- [63] E.I. Konstantinidis, P. E. Antoniou and P.D. Bamidis (2015). Exergames for Assessment in Active and Healthy Aging - Emerging Trends and Potentialities. In *Proceedings of the 1st International Conference on Information and Communication Technologies for Ageing Well and e-Health*, ISBN 978-989-758-102-1, pages 325-330. DOI: 10.5220/0005494503250330
- [64] P.E. Antoniou, E. Konstantinidis, A.S. Billis, G. Bamparopoulos, M.S. Tsatali, & A. Sioundas, (2015). Instrumenting the eHome and Preparing Elderly Pilots: The USEFIL Approach. In P. Bamidis, I. Tamasas, L. Hadjileontiadis, & M. Tsolaki (Eds.) *Handbook of Research on Innovations in the Diagnosis and Treatment of Dementia* (pp. 335-361). Hershey, PA: Medical Information Science Reference. doi:10.4018/978-1-4666-8234-4.ch018