

SYSTEMS OF HOLOMORPHIC MULTIVALUED PROJECTIONS ON COMPLEX MANIFOLDS

BY KAMIL DRZYŻGA

Abstract. Let M be a submanifold of a connected Stein manifold X . We construct a global system of holomorphic multivalued projections $X \rightarrow M$. In particular, for every locally bounded family $\mathcal{F} \subset \mathcal{O}(M)$ we get a continuous extension operator $\mathcal{F} \rightarrow \mathcal{O}(X)$.

1. Introduction. Let M be a complex submanifold of a Stein manifold X . Using Bishop's ideas of multivalued projections we proved in [4] that for every domain $U \subset\subset X$ there exists a linear continuous extension operator $\mathcal{O}(M) \rightarrow \mathcal{O}(U)$. Now, we will study the problem of existence of global holomorphic multivalued projections $X \rightarrow M$ (see Definition 5.1 and Theorem 5.5). Note that in the paper [2] the author suggested that a holomorphic multivalued projections could exist. In particular, we prove that there is a continuous extension operator $\mathcal{F} \rightarrow \mathcal{O}(X)$ for each locally bounded family $\mathcal{F} \subset \mathcal{O}(M)$ and moreover as an application we get a **linear** continuous extension operator $L^2(M)^1 \rightarrow \mathcal{O}(X)$.

2. Auxiliary Results. Let M be a d -dimensional analytic subset of a connected Stein manifold X . In the sequel we denote by $\text{Reg}M$ the set of regular points of M . For a compact $K \subset X$, its holomorphic hull (with respect to the space $\mathcal{O}(X)$ of all holomorphic functions on X) will be denoted by $\widehat{K}_{\mathcal{O}(X)}$. Put $\mathbb{D}(r) := \{z \in \mathbb{C} : |z| < r\}$, $\mathbb{D} := \mathbb{D}(1)$.

2010 *Mathematics Subject Classification.* 32D15.

Key words and phrases. Multivalued projections, holomorphic continuation.

${}^1L^2_h(M) := \{f \in \mathcal{O}(M) : \int_M |f|^2 < \infty\}$.

DEFINITION 2.1. Let $f \in \mathcal{O}(X, \mathbb{C}^k)$. We say that a set $P \subset P_0 := M \cap f^{-1}(\mathbb{D}^k)$ is an *analytic polyhedron* in M ($P \in \mathcal{P}(M, k, f)$) if $P \subset\subset M$ and P is the union of a family of connected components of P_0 .

We say that an analytic polyhedron $P \in \mathcal{P}(M, k, f)$ is *special* if $d = k$.

THEOREM 2.2 (cf. [2]). *Assume that $P \in \mathcal{P}(M, k, f)$ and $S \subset P$, $T \subset f^{-1}(\mathbb{D}^k)$ are compact. Then there exists a special analytic polyhedron $Q \in \mathcal{P}(M, d, g)$ such that $S \subset Q \subset P$ and $g(T) \subset \mathbb{D}^d$.*

THEOREM 2.3 (cf. [2]). *Assume that X is Stein, $T \subset X$ is compact, and U is an open neighborhood of T such that $(U \setminus T) \cap \widehat{T}_{\mathcal{O}(X)} = \emptyset$. Let \mathcal{A} stand for the closure of $\mathcal{O}(U)|_T$ in the space $\mathcal{C}(T)$ of all complex valued continuous functions on T . Then for every non-zero homomorphism $\xi : \mathcal{A} \rightarrow \mathbb{C}$ there exists an $x_0 \in T$ such that $\xi(f) = f(x_0)$ for every $f \in \mathcal{A}$. Consequently (cf. [1], Chapter I, Section II, Corollary 10), if $w_1, \dots, w_m \in \mathcal{A}$ have no common zeros on T , then there exist $c_1, \dots, c_m \in \mathcal{A}$ such that $c_1 w_1 + \dots + c_m w_m = 1$.*

DEFINITION 2.4 (cf. [2]). A continuous map $f : X \rightarrow Y$, where X, Y are topological spaces, is called *almost proper* if each connected component of $f^{-1}(S)$ is compact for every compact subset S of Y .

THEOREM 2.5 (cf. [2]). *Let Y be a 0-dimensional analytic subset of $\text{Reg}(M)$. Then there exists an $f \in \mathcal{O}(X, \mathbb{C}^d)$ such that $f|_M$ is almost proper and the mapping f gives local coordinates on M at x for each $x \in Y$.*

THEOREM 2.6 (cf. [2]). *Assume that M is pure d -dimensional and let $f \in \mathcal{O}(X, \mathbb{C}^d)$ be such that $f|_M$ is almost proper. Let $\{S_j\}_{j=1}^\infty$ be an increasing sequence of compact subsets of M , each of which has finitely many connected components and $\bigcup_{j=1}^\infty S_j = M$. Let $\alpha : \mathbb{N} \rightarrow \mathbb{R}_{>0}$ such that*

$$S_j \subset F_j := M \cap f^{-1}(\overline{\mathbb{D}}^d(\alpha(j)))$$

for all $j \in \mathbb{N}$. Let H_j be the union of all those connected components of F_j which intersect S_j . Then H_j is compact. For each $j \in \mathbb{N}$ put

$$G_j := (H_{j+1} \cap F_j) \setminus H_j.$$

Let $\{g_j\}_{j=1}^\infty \subset \mathcal{O}(M)$ and $\{\varepsilon_j\}_{j=1}^\infty \subset \mathbb{R}_{>0}$. Then there exists an $s \in \mathcal{O}(M)$ such that

$$|s(x) - g_j(x)| < \varepsilon_j, \quad x \in G_j, \quad j \in \mathbb{N}.$$

Moreover, given a countable set $A \subset M$, the function s can be chosen to have different values modulo $2\pi i$, i.e. $e^{s(x)} \neq e^{s(y)}$ for $x, y \in M$ and $x \neq y$.

REMARK 2.7. Observe that:

- (a) $H_j \subset H_{j+1}$ for $j \in \mathbb{N}$;
- (b) $\bigcup_{j=1}^\infty H_j = M$.

3. Symmetric products. The aim of this section is to present some properties of the symmetric products. Details can be found in [7], Appendix V.

Let X be a Hausdorff topological space. We define an equivalence relation on X^k by $(x_1, \dots, x_k) \sim (y_1, \dots, y_k) : \iff (y_1, \dots, y_k)$ is a reordering of (x_1, \dots, x_k) . $\overleftarrow{X}^k := X^k / \sim$ is called the k -symmetric product of X . In the case $k = 1$, we get $\overleftarrow{X}^1 = X$. Now, we define the projection $\pi : X^k \longrightarrow \overleftarrow{X}^k$, $\pi(x) := [x]$. We put $[x_1, \dots, x_k] := [(x_1, \dots, x_k)]$, $\{[x_1, \dots, x_k]\} := \{x_1, \dots, x_k\}$. Moreover, we put

$$[x_1 : \mu_1, \dots, x_\ell : \mu_\ell] := \overbrace{[x_1, \dots, x_1]}^{\mu_1\text{-times}}, \dots, \overbrace{[x_\ell, \dots, x_\ell]}^{\mu_\ell\text{-times}},$$

provided that $x_j \neq x_t$ for $j \neq t$, $\mu_1, \dots, \mu_\ell \in \mathbb{N}$, $\mu_1 + \dots + \mu_\ell = k$. We define

$$[A_1, \dots, A_k] := \left\{ [x_1, \dots, x_k] : x_i \in A_i, \quad i = 1, \dots, k \right\}.$$

The topology on \overleftarrow{X}^k is defined by the basis

$$[U_1, \dots, U_m], \quad U_i \text{ is open in } X, \quad i = 1, \dots, k.$$

Observe that π is continuous, open, and \overleftarrow{X}^k is Hausdorff.

DEFINITION 3.1. Let Y be Hausdorff topological space and let $F : X \longrightarrow \overleftarrow{Y}^n$ be continuous. Then we put

$$X_F^{(k)} := \{x \in X : \#\{F(x)\} = k\},$$

$$\chi_F := \max\{k : X_F^{(k)} \neq \emptyset\}, \quad X_F := X_F^{(\chi_F)}.$$

Note that X_F is open.

PROPOSITION 3.2. *Let F be as above. Suppose that*

$$a \in X_F, \quad F(a) = [b_1 : \mu_1, \dots, b_\ell : \mu_\ell], \quad \mu_1 + \dots + \mu_\ell = k \quad \ell := \chi_F.$$

Then there is a neighborhood $U \subset X_F$ of a and there are uniquely defined continuous functions $f_i : U \longrightarrow Y$, $i = 1, \dots, \ell$, such that

$$F(x) = [f_1(x) : \mu_1, \dots, f_\ell(x) : \mu_\ell], \quad x \in U.$$

In the above situation, we will write $F = \mu_1 f_1 \oplus \dots \oplus \mu_\ell f_\ell$ on U .

PROPOSITION 3.3. *Let $F : X^k \longrightarrow Y$ be continuous. Then F is symmetric if and only if there exists a continuous function $\overleftarrow{F} : \overleftarrow{X}^k \longrightarrow Y$ such that $F = \overleftarrow{F} \circ \pi$.*

4. Holomorphic multivalued functions and system of multivalued projections. All propositions below and their proofs are taken from [4]. We recall only those facts which will be used in this paper.

DEFINITION 4.1. Let M, N be complex manifolds with M connected. We say a continuous mapping $F: M \rightarrow \overleftarrow{N}^n$ is *holomorphic on M* ($F \in \mathcal{O}(M, \overleftarrow{N}^n)$) if:

- $M \setminus M_F$ is thin, i.e. every point $x_0 \in M \setminus M_F$ has open connected neighborhood $V \subset M$ and a function $\varphi \in \mathcal{O}(V)$, $\varphi \not\equiv 0$, such that $(M \setminus M_F) \cap V \subset \varphi^{-1}(0)$,
- for every $a \in M_F$, if $F = \mu_1 f_1 \oplus \cdots \oplus \mu_\ell f_\ell$ on V as in Proposition 3.2, then $f_1, \dots, f_\ell \in \mathcal{O}(V)$.

If M is disconnected, then we say that F is *holomorphic on M* if $F|_C \in \mathcal{O}(C, \overleftarrow{N}^n)$ for any connected component $C \subset M$.

PROPOSITION 4.2. Let M, N, K be complex manifolds and let $f \in \mathcal{O}(M, N)$, $g \in \mathcal{O}(N, \overleftarrow{K}^n)$. Assume that $f(M) \cap N_g \neq \emptyset$ and M is connected. Then $g \circ f \in \mathcal{O}(M_{g \circ f}, \overleftarrow{K}^n)$.

PROPOSITION 4.3. Let $f \in \mathcal{O}(M, \overleftarrow{N}^n)$ and $g \in \mathcal{O}(N^n, K)$ be symmetric. Then $\overleftarrow{g} \circ f \in \mathcal{O}(M, K)$.

THEOREM 4.4 (cf. [2]; see also [6], Chapter 7). Assume that $P \in \mathcal{P}(M, d, f)$ is special. Then there exist a $k \in \mathbb{N}$ and a holomorphic mapping $\omega: \mathbb{D}^d \rightarrow \overleftarrow{P}^k$ such that:

- $f^{-1}(z) \cap P = \{\omega(z)\}$, $z \in \mathbb{D}^d$,
- $\#\{\omega(z)\} = k$ for $z \in \mathbb{D}^d \setminus \Sigma'$, where Σ' is a proper analytic set.

The number k in the above theorem is called the *multiplicity of f on P* .

DEFINITION 4.5. Let M be an analytic submanifold of a manifold X . Let $U \subset X$ be a domain such that $U \cap M \neq \emptyset$. We say a holomorphic function

$$\Delta: U \rightarrow \overleftarrow{(M \times \mathbb{C})}^n$$

is a *holomorphic multivalued projection* $U \rightarrow M$ if for any $x \in U \cap M$ such that $\Delta(x) = [(x_1, z_1), \dots, (x_n, z_n)]$ we have $x_{j_0} = x$ for some $j_0 \in \{1, \dots, n\}$ and $z_j = 0$ for any $j \in \{1, 2, \dots, n\} \setminus \{j_0\}$.

Let \mathfrak{P} denote the set of all holomorphic multivalued projections $U \rightarrow M$. Then we define the map

$$\Xi: (U \cap M) \times \mathfrak{P} \rightarrow \mathbb{C}, \quad \Xi(x, \Delta) := z_{j_0}.$$

Observe that Ξ is well defined.

DEFINITION 4.6. We say $\Pi = (\Delta_s)_{s=1}^k$ is a *system of holomorphic multivalued projections* $U \rightarrow M$ if $\Delta_s : U \rightarrow (M \times \mathbb{C})^{\overleftarrow{k_s}}$, $s = 1, \dots, k$, are holomorphic multivalued projections and $\sum_{s=1}^k \Xi(x, \Delta_s) = 1$ for any $x \in U \cap M$.

THEOREM 4.7. *Assume that there exists a system Π of holomorphic multivalued projections on U . Then there exists a linear continuous operator*

$$L_\Pi : \mathcal{O}(M) \rightarrow \mathcal{O}(U)$$

such that $L_\Pi(u)(x) = u(x)$ for $x \in U \cap M$.

THEOREM 4.8. *Let M be an analytic submanifold of a Stein manifold X . Let U be a relatively compact domain of X such that $U \cap M \neq \emptyset$. Then there exists a system of multivalued holomorphic projections $U \rightarrow M$.*

Theorems 4.7 and 4.8 immediately imply the following result.

THEOREM 4.9. *Let M be an analytic submanifold of a Stein manifold X . Let U be a relatively compact domain of X such that $U \cap M \neq \emptyset$. Then there exists a linear continuous extension operator $L : \mathcal{O}(M) \rightarrow \mathcal{O}(U)$.*

PROPOSITION 4.10. *Let ω, f, X, P be as above. Additionally assume that $f(U) \subset \mathbb{D}^d$, where $U \subset X$ is a domain and $U \cap P \neq \emptyset$. Then $\omega \circ f|_U \in \mathcal{O}(U, \overleftarrow{P^k})$.*

PROPOSITION 4.11. *Let ω, f, X, P be as above. Then $\omega \circ f|_P \in \mathcal{O}(P, \overleftarrow{P^k})$.*

5. Global system of holomorphic multivalued projections. Let X be a connected complex manifold and M be a complex submanifold.

DEFINITION 5.1. A sequence $\Pi = (\Delta_{s,j})_{(s,j) \in \{1, \dots, r\} \times \mathbb{N}}$ is called a *global system of holomorphic multivalued projections* $X \rightarrow M$ if for each $j \in \mathbb{N}$ the mapping $\Delta_{s,j} : U_j \rightarrow (M \times \mathbb{C})^{\overleftarrow{k_{s,j}}}$ ($k_{s,j} \in \mathbb{N}$) is a holomorphic multivalued projection (in the sense of Definition 4.5), $s = 1, \dots, r$, having the following properties

- (a) $U_j \subset X$ is a domain with $U_j \cap M \neq \emptyset$, $U_j \subset U_{j+1}$, $\bigcup_{j \in \mathbb{N}} U_j = X$;
- (b) $\lim_{n \rightarrow \infty} \sum_{s=1}^r \Xi(x, \Delta_{s,n}) = 1$, $x \in M$.

REMARK 5.2. Let $\Pi = (\Delta_{s,j})_{(s,j) \in \{1, \dots, r\} \times \mathbb{N}}$ be as above.

- (a) For each $j \in \mathbb{N}$ we get a linear continuous operator (cf. the proof of Theorem 4.7 in [4].)

$$L_{\Pi,j} : \mathcal{O}(M) \rightarrow \mathcal{O}(U_j), \quad L_{\Pi,j} := \sum_{s=1}^r \overleftarrow{u}_{s,j} \circ \Delta_{s,j}, \quad \text{where}$$

$$\overleftarrow{u}_{s,j} : (M \times \mathbb{C})^{\overleftarrow{k_{s,j}}} \rightarrow \mathbb{C}, \quad \overleftarrow{u}_{s,j}([\xi_1, \lambda_1], \dots, [\xi_{k_{s,j}}, \lambda_{k_{s,j}}]) = \sum_{m=1}^{k_{s,j}} u(\xi_m) \lambda_m.$$

(b) Using Definition 5.1(b), for $u \in \mathcal{O}(M)$ and $x \in M$ we get

$$\lim_{j \rightarrow \infty} L_{\Pi,j}(u)(x) = \lim_{j \rightarrow \infty} \sum_{s=1}^r \overleftarrow{u}_{s,j} \circ \Delta_{s,j}(x) = \lim_{j \rightarrow \infty} \sum_{s=1}^r u(x) \Xi(x, \Delta_{s,j}) = u(x).$$

Let $\emptyset \neq \mathcal{F} \subset \mathcal{O}(M)$.

DEFINITION 5.3. We say a global system of holomorphic multivalued projections $\Pi = (\Delta_{s,j})_{(s,j) \in \{1, \dots, r\} \times \mathbb{N}}$ is an \mathcal{F} -extension if for each $u \in \mathcal{F}$ the sequence $(L_{\Pi,j}(u))_{j=1}^{\infty}$ converges locally uniformly in X .

Set $L_{\Pi}(u) := \lim_{j \rightarrow \infty} L_{\Pi,j}(u)$, $u \in \mathcal{F}$.

REMARK 5.4. Let $\Pi = (\Delta_{s,j})_{(s,j) \in \{1, \dots, r\} \times \mathbb{N}}$ be an \mathcal{F} -extension.

- (a) By Remark 5.2(b), $L_{\Pi} : \mathcal{F} \rightarrow \mathcal{O}(X)$ is an extension operator.
- (b) If $u, v \in \mathcal{F}$ and $u + v \in \mathcal{F}$, then $L_{\Pi}(u + v) = L_{\Pi}(u) + L_{\Pi}(v)$.
- (c) If $u \in \mathcal{F}$, $\alpha \in \mathbb{C}$ and $\alpha u \in \mathcal{F}$, then $L_{\Pi}(\alpha u) = \alpha L_{\Pi}(u)$.
- (d) If \mathcal{F} is a vector space, then L_{Π} is linear.
- (e) If $u_1, \dots, u_m \in \mathcal{F}$ are linearly independent (in $\mathcal{O}(M)$), then the formula

$$L_{\Pi}(\alpha_1 u_1 + \dots + \alpha_m u_m) := \alpha_1 L_{\Pi}(u_1) + \dots + \alpha_m L_{\Pi}(u_m), \quad \alpha_1, \dots, \alpha_m \in \mathbb{C},$$

extends the operator L_{Π} to the vector space $\text{span}\{u_1, \dots, u_m\}$.

The main result of the paper is the following theorem.

THEOREM 5.5. *Let X be a Stein manifold and $\mathcal{F} \subset \mathcal{O}(M)$ be locally bounded (i.e. $\sup_{u \in \mathcal{F}} \|u\|_K < +\infty^2$ for every compact set $K \subset M$, e.g. \mathcal{F} is finite). Then there exists an \mathcal{F} -extension $\Pi = (\Delta_{s,j})_{(s,j) \in \{1, \dots, d\} \times \mathbb{N}}$ with $d := \dim M$. Consequently, there exists a continuous extension operator $L_{\Pi} : \mathcal{F} \rightarrow \mathcal{O}(X)$.*

COROLLARY 5.6. *Let X be a Stein manifold and \mathcal{V} be a finitely dimensional vector subspace of $\mathcal{O}(M)$. Then there exists a linear continuous extension operator $L : \mathcal{V} \rightarrow \mathcal{O}(X)$.*

PROPOSITION 5.7. *Assume that $\mathcal{H} \subset \mathcal{O}(M)$ is a Hilbert space such that the unit ball $B := \{f \in \mathcal{H} : \|f\|_{\mathcal{H}} \leq 1\}$ is locally uniformly bounded and the convergence in the sense of \mathcal{H} implies the locally uniform convergence in M . Then there exists a linear continuous extension operator $L : \mathcal{H} \rightarrow \mathcal{O}(X)$. In particular, there exists a linear continuous extension operator $L : L_h^2(M) \rightarrow \mathcal{O}(X)$.*

PROOF. We put $\mathcal{F} := B$. By Theorem 5.5 there exists a continuous extension operator $\tilde{L} : \mathcal{F} \rightarrow \mathcal{O}(X)$. Moreover, since $\text{span}(\mathcal{F}) = \mathcal{H}$, we conclude that there exists a linear continuous extension operator $L : \mathcal{H} \rightarrow \mathcal{O}(X)$.

² $\|f\|_K := \sup_K |f|$.

Indeed, suppose that $(f_j)_{j=1}^\infty \subset \mathcal{F}$ is an orthonormal basis of \mathcal{H} . Set $\tilde{f}_j := \tilde{L}(f_j)$. Let $f \in \mathcal{H}$ be such that $f = \sum_{j=1}^\infty c_j f_j$. Put

- $L(f) := \sum_{j=1}^\infty c_j \tilde{f}_j = C_f \tilde{L}(f/C_f)$,
- $s_N := \sum_{j=1}^N c_j f_j$,

where $C_f := \|f\|_{\mathcal{H}}$. Since $(f_j)_{j=1}^\infty$ is orthonormal, hence

- $f_j, \frac{c_j}{C_f} f_j \in \mathcal{F}$,
- $\frac{c_j}{C_f} f_j + \frac{c_k}{C_f} f_k \in \mathcal{F}$ for $j, k \in \mathbb{N}, j \neq k$.

Therefore, $C_f \tilde{L}(s_N/C_f) = \sum_{j=1}^N c_j \tilde{f}_j$. As $s_N/C_f \rightarrow f/C_f$ locally uniformly and $s_N/C_f \in \mathcal{F}$, we get $L(f) = C_f \tilde{L}(f/C_f)$. By assumption on topologies, L is continuous.

Now, assume that $\mathcal{H} = L_h^2(M)$. It is known that for any compact set $K \subset M$ there are $C_K > 0$ and open neighborhood $K \subset \Omega \subset\subset M$ such that $\|f\|_K \leq C_K \|f\|_{L^2(\Omega, dV)}$. It follows that B is locally uniformly bounded. \square

COROLLARY 5.8. *Let $X \in \{\mathbb{D}^n, \mathbb{B}_n\}$. There exists a linear continuous extension operator $L : L_h^2(\mathbb{D})^3 \rightarrow \mathcal{O}(X)$.*

PROOF OF THEOREM 5.5. Let Y be an arbitrary 0-dimensional analytic subset of M . By Theorem 2.5 there exists a mapping $f \in \mathcal{O}(X, \mathbb{C}^d)$ such that $f|_M$ is almost proper and for each $x \in Y$ the mapping f gives local coordinates on M at x .

Let $S_k, \alpha(k), F_k, H_k$ and G_k be as in Theorem 2.6. Observe that $Q_k := \text{int}H_k = H_k \cap f^{-1}(\mathbb{D}^d(\alpha(k)))$ is a special analytic polyhedron. Let λ_k -denote the multiplicity of f in Q_k , defined via Theorem 4.4 with $\omega_k : \mathbb{D}^d(\alpha(k)) \rightarrow \overleftrightarrow{Q_k^{\lambda_k}}$. Set $\omega_k(f(x)) = [x_1^k, \dots, x_{\lambda_k}^k]$ (counted with multiplicities), $x \in f^{-1}(\mathbb{D}^d(\alpha(k)))$.

Observe that for arbitrary $x \in X$, the set $M \cap f^{-1}(f(x))$ is discrete. Let $(x_\nu)_{\nu=1}^\infty = M \cap f^{-1}(f(x))$ (points are counted with multiplicities). We assume that $x_1 = x$ for $x \in M$. Let

$$\Xi_k(x) := \{j \in \mathbb{N} : x_j \in H_k\}.$$

Observe that for each $k \in \mathbb{N}$ and $x \in f^{-1}(Q_k)$ the set $\Xi_k(x)$ is finite and $\{x_j : j \in \Xi_k(x)\} = \{x_1^k, \dots, x_{\lambda_k}^k\}$.

Put $g_k := \lambda_{k+1} + k^2 + 1$, $k \in \mathbb{N}$. By Theorem 2.6 there exists an $f_{d+1} \in \mathcal{O}(X)$ such that $|f_{d+1} - g_k| < 1$ on G_k , $k \in \mathbb{N}$, and the function $w := e^{-f_{d+1}}$ separates points in $M \cap f^{-1}(f(x))$ for all $x \in Y$.

³ $\mathbb{D} \simeq \{(z, \mathbf{0}) \in \mathbb{C}^n : z \in \mathbb{D}\}$.

LEMMA 5.9. *Let $\mathcal{F} \subset \mathcal{O}(M)$ be locally bounded. Then there exists a function $f_{d+1}^* \in \mathcal{O}(X)$ such that if $h := e^{-f_{d+1}^*}$ and*

$$\tilde{\varphi}_k(x) := \sum_{j \in \Xi_k(x)} \varphi(x_j) h(x_j) \prod_{\substack{\mu \in \Xi_k(x) \\ \mu \neq j}} \left(1 - \frac{w(x_\mu)}{w(x)}\right), \quad \varphi \in \mathcal{F}, x \in X, k \in \mathbb{N},$$

then for every domain $U \subset\subset X$ such that $U \cap M \neq \emptyset$,

- there exists a $k_0 \in \mathbb{N}$ such that $\tilde{\varphi}_k \in \mathcal{O}(U)$ for $k \geq k_0$ and
- the sequence $(\tilde{\varphi}_k)_{k=1}^\infty$ converges uniformly on U .

Suppose for a moment that the lemma is proved. Let $\tilde{\varphi}(x) := \lim_{k \rightarrow \infty} \tilde{\varphi}_k(x)$, $x \in U$. Then $\tilde{\varphi} \in \mathcal{O}(U)$. Since $x_1 = x$ for $x \in M \cap U$, we get

$$\tilde{\varphi}(x) = \varphi(x) h(x) \prod_{\mu=2}^\infty \left(1 - \frac{w(x_\mu)}{w(x)}\right) = \varphi(x) h(x) \tilde{w}_1(x), \quad x \in M \cap U,$$

where

$$\tilde{w}_1(x) := \prod_{\mu=2}^\infty \left(1 - \frac{w(x_\mu)}{w(x)}\right), \quad x \in M.$$

Observe that the condition $|f_{d+1} - (\lambda_{k+1} + k^2 + 1)| < 1$ on G_k , $k \in \mathbb{N}$, implies that the function \tilde{w}_1 is well-defined (cf. the estimate of the function B in the proof of Lemma 5.9). Hence $\tilde{w}_1 \in \mathcal{O}(M)$. Notice that $\tilde{w}_1(x) \neq 0$ for $x \in Y$.

We move to the main part of proof.

First we take $Y = Y_1 \subset M$ having a point in each connected component of M . We get a function $\tilde{w}_1 \in \mathcal{O}(M)$ such that $\tilde{w}_1(x) \neq 0$ for each $x \in Y_1$. In particular $M_1 := \{x \in M : \tilde{w}_1(x) = 0\}$ is $(d-1)$ -dimensional analytic subset of M . Next we take $Y_2 \subset M_1$ having a point in each connected component of $\text{Reg}(M_1)$. We get $\tilde{w}_2 \in \mathcal{O}(M)$ such that $\tilde{w}_2(x) \neq 0$ for each $x \in Y_2$. Thus $M_2 := \{x \in M : \tilde{w}_1(x) = \tilde{w}_2(x) = 0\}$ is a $(d-2)$ -dimensional analytic subset of M . We repeat the procedure and we obtain $\tilde{w}_1, \dots, \tilde{w}_d \in \mathcal{O}(M)$ without common zeros on M . By Theorem 2.3 there exist $c_1, \dots, c_d \in \mathcal{O}(M)$ such that $c_1 \tilde{w}_1 + \dots + c_d \tilde{w}_d = 1$ on M . Assume that h_s is constructed with respect to the family $\mathcal{F}_s := \{uc_s : u \in \mathcal{F}\}$.

We get $f_s, H_{s,k}, Q_{s,k}, \omega_{s,k}, \lambda_{s,k}, (x_{s,j}^k)_{j=1}^{\lambda_{s,k}}, (x_{s,\nu})_{\nu=1}^\infty, \Xi_{s,k}(\cdot), w_s, \tilde{w}_s$ for $s = 1, \dots, d, k \geq 1$.

Now we are going to construct a global system of holomorphic multivalued projections on $X \rightarrow M$ (cf. Definition 5.1). Fix arbitrary domains $U_j \subset U_{j+1} \Subset X$ such that $\bigcup_{j=1}^\infty U_j = X$, $U_j \cap M \neq \emptyset$. Let $(t_j)_{j=1}^\infty \subset \mathbb{N}$ be such that

- $f_s(U_j) \subset \mathbb{D}^d(\alpha_s(t_j))$, where $\alpha_s(t_j) \in (0, +\infty)$;
- $U_j \cap M \subset Q_{s,t_j}$;

- $t_j \leq t_{j+1}$, $s = 1, \dots, d$;
- $t_j \rightarrow +\infty$.

Put $k_{s,j} := \lambda_{s,t_j}$. We define $\Delta_{s,j} : U_j \rightarrow \overleftarrow{(M \times \mathbb{C}^n)^{k_{s,j}}}$ by

$$\Delta_{s,j}(x) := [(F_{s,1}(x), G_{s,1}(x)), \dots, (F_{s,k_{s,j}}(x), G_{s,k_{s,j}}(x))],$$

where $F_{s,m}(x) := x_{s,t_j}^m$,

$$G_{s,m}(x) := \frac{c_s(F_{s,m}(x))h_s(F_{s,m}(x))}{h_s(x)} \prod_{\mu \in \Xi_{t_j}^s(x) \setminus \{p_{j,m,s}\}} \left(1 - \frac{w_s(x_{s,\mu})}{w_s(x)}\right);$$

and $p_{j,m,s} \in \mathbb{N}$ is such that $x_{s,t_j}^m = x_{s,p_{j,m,s}}$.

Then $\Pi := (\Delta_{s,j})_{(s,j) \in \{1, \dots, d\} \times \mathbb{N}}$ is the global system of holomorphic multi-valued projections on $X \rightarrow M$.

Indeed, since $U_j \subset f_s^{-1}(\mathbb{D}^d(\alpha_s(t_j)))$, then similarly as in the proof of Theorem 4.8 we show that $\Delta_{s,j}$ are holomorphic (see [4]). Next, we see that for $x \in M$ we have

$$\lim_{j \rightarrow \infty} \sum_{s=1}^d \Xi(x, \Delta_{s,j}) = \sum_{s=1}^d c_s(x) \tilde{w}_s(x) = 1.$$

The construction of a global system of holomorphic projections has been finished. \square

PROOF OF THE LEMMA 5.9. Fix an arbitrary domain $U \subset \subset X$, $U \cap M \neq \emptyset$ and $k_0 \in \mathbb{N}$ such that $f(\bar{U}) \subset \mathbb{D}^d(\alpha(k_0))$. Let f_{d+1}^* be for a moment arbitrary and let $\varphi \in \mathcal{F}$. Take a $k \geq k_0$.

First, we are going to prove that $\tilde{\varphi}_k \in \mathcal{O}(U)$. Note that if $x \in U$ and $j \in \Xi_k(x)$, then $x_j \in H_k \cap f^{-1}(\mathbb{D}^d(\alpha(k))) = Q_k$. Hence $\{x_j : j \in \Xi_k(x)\} = \{x_1^k, \dots, x_{\lambda_k}^k\} = \{\omega_k(f(x))\}$, $x \in U$. Moreover,

$$\tilde{\varphi}_k(x) = w^{1-\lambda_k}(x) = \sum_{\nu=0}^{\lambda_k-1} \overleftarrow{S}_\nu^{\lambda_k}(\omega_k(f(x))) w^k(x), \quad x \in U,$$

where

$$S_{\lambda_k-1}(t) := \sum_{j=1}^{\lambda_k} \varphi(t_j) h(t_j),$$

$$S_\nu(t) := (-1)^{\lambda_k-1-\nu} \sum_{j=1}^{\lambda_k} \varphi(t_j) h(t_j) \sigma_{k-1-\nu}(w(t_1), \dots, w(t_{j-1}), w(t_{j+1}), \dots, w(t_{\lambda_k})),$$

$$\nu = 0, \dots, \lambda_k - 2, t = (t_1, \dots, t_{\lambda_k}) \in Q_k^{\lambda_k},$$

and $\sigma_1, \dots, \sigma_{\lambda_k-1} : \mathbb{C}^{\lambda_k-1} \rightarrow \mathbb{C}$ are standard symmetric polynomials. Consequently, by Proposition 4.10 we conclude that $\tilde{\varphi}_k \in \mathcal{O}(U)$.

Now we are going to find a function $f_{d+1}^* \in \mathcal{O}(U)$ (independent of U) such that $(\tilde{\varphi}_k)_{k=1}^\infty$ converges uniformly on U .

We construct f_{d+1}^* via Theorem 2.6 in such a way that $|f_{d+1}^* - k^2 \beta_k \lambda_k - 1| < 1$ on G_k , where $\beta_k \geq \sup\{\sup_{G_k} |\varphi| : \varphi \in \mathcal{F}\}$. Our aim is to prove that $\tilde{\varphi}_l(x) - \tilde{\varphi}_k(x) \rightarrow 0$ uniformly on U when $l > k \rightarrow +\infty$. Take $l > k \geq k_0$. For $x \in U$ write

$$\begin{aligned} \tilde{\varphi}_l(x) - \tilde{\varphi}_k(x) &= \sum_{j \in \Xi_l(x) \setminus \Xi_k(x)} \varphi(x_j) h(x_j) \prod_{\substack{\mu \in \Xi_l(x) \\ \mu \neq j}} \left(1 - \frac{w(x_\mu)}{w(x)}\right) \\ &+ \sum_{j \in \Xi_k(x)} \varphi(x_j) h(x_j) \prod_{\substack{\mu \in \Xi_k(x) \\ \mu \neq j}} \left(1 - \frac{w(x_\mu)}{w(x)}\right) \\ &\cdot \left(\left(\prod_{\substack{\mu \in \Xi_l(x) \setminus \Xi_k(x) \\ \mu \neq j}} \left(1 - \frac{w(x_\mu)}{w(x)}\right) \right) - 1 \right) = I_{k,l}(x) + J_{k,l}(x). \end{aligned}$$

We have

$$\begin{aligned} |I_{k,l}(x)| &\leq \left(\sum_{j \notin \Xi_k(x)} |\varphi(x_j) h(x_j)| \right) \cdot \prod_{\mu \in \mathbb{N}} \left(1 + \frac{|w(x_\mu)|}{|w(x)|}\right) =: A_k(x) B(x), \\ |J_{k,l}(x)| &\leq \left(\sum_{j \in \mathbb{N}} |\varphi(x_j) h(x_j)| \right) \cdot \left(\prod_{\mu \in \mathbb{N}} \left(1 + \frac{|w(x_\mu)|}{|w(x)|}\right) \right) \\ &\cdot \left(\left(\prod_{\mu \notin \Xi_k(x)} \left(1 + \frac{|w(x_\mu)|}{|w(x)|}\right) \right) - 1 \right) =: C(x) D(x) (E_k(x) - 1). \end{aligned}$$

Observe that $M \cap f^{-1}(\mathbb{D}^d(\alpha(k_0))) \subset Q_{k_0} \cup \bigcup_{s=k_0}^\infty \text{int} G_s$. Let

$$\gamma := \max_{\bar{U}} \text{Re} f_{d+1}, \quad \delta := \max_{H_{k_0}} (-\text{Re} f_{d+1}).$$

Observe that if $x \in U$ and $x_\mu \in \text{int} G_s$, then we have

$$\log \left(1 + \frac{|w(x_\mu)|}{|w(x)|}\right) \leq \frac{|w(x_\mu)|}{|w(x)|} = e^{-\text{Re} f_{d+1}(x_\mu) + \text{Re} f_{d+1}(x)} \leq e^{-\lambda_{s+1} - s^2 + \gamma} \leq \frac{e^\gamma}{\lambda_{s+1} s^2}.$$

If $x_\mu \in Q_{k_0}$, then

$$\log \left(1 + \frac{|w(x_\mu)|}{|w(x)|}\right) \leq \frac{|w(x_\mu)|}{|w(x)|} = e^{-\text{Re} f_{d+1}(x_\mu) + \text{Re} f_{d+1}(x)} \leq e^{\delta + \gamma}.$$

Thus for all $x \in U$ we have

$$\begin{aligned} \log B(x) &= \sum_{\mu: x_\mu \in Q_{k_0}} \log \left(1 + \frac{|w(x_\mu)|}{|w(x)|} \right) + \sum_{s=k_0}^{\infty} \sum_{\mu: x_\mu \in \text{int}G_s} \log \left(1 + \frac{|w(x_\mu)|}{|w(x)|} \right) \\ &\leq \lambda_{k_0} e^{\delta+\gamma} + \sum_{s=k_0}^{\infty} \frac{e^\gamma}{s^2}, \end{aligned}$$

and therefore the function B is uniformly bounded on U .

Similarly,

$$\begin{aligned} C(x) &= \sum_{\mu: x_\mu \in Q_{k_0}} |\varphi(x_\mu)h(x_\mu)| + \sum_{s=k_0}^{\infty} \sum_{\mu: x_\mu \in \text{int}G_s} |\varphi(x_\mu)h(x_\mu)| \\ &\leq M + \sum_{s=k_0}^{\infty} \sum_{\mu: x_\mu \in \text{int}G_s} \beta_s e^{-\text{Re}f_{d+1}^*(x_\mu)} \leq M + \sum_{s=k_0}^{\infty} \sum_{\mu: x_\mu \in \text{int}G_s} \beta_s e^{-s\beta_s\lambda_s} \\ &\leq M + \sum_{s=k_0}^{\infty} \sum_{\mu: x_\mu \in \text{int}G_s} \beta_s \frac{1}{\beta_s\lambda_s s^2} \leq M + \sum_{s=k_0}^{\infty} \frac{1}{s^2}, \end{aligned}$$

where $M := \lambda_{k_0} \sup_{Q_{k_0}} |\varphi h|$. On the other hand,

$$\begin{aligned} A_k(x) &= \sum_{s=k}^{\infty} \sum_{j: x_j \in \text{int}G_s} \beta_s |h(x_j)| = \sum_{s=k}^{\infty} \sum_{j: x_j \in \text{int}G_s} \beta_s e^{-\text{Re}f_{d+1}^*(x_j)} \\ &\leq \sum_{s=k}^{\infty} \sum_{j: x_j \in \text{int}G_s} \beta_s e^{-\beta_s\lambda_{s+1}-s^2} \leq \sum_{s=k}^{\infty} \sum_{j: x_j \in \text{int}G_s} \beta_s \frac{1}{\beta_s\lambda_{s+1}s^2} \leq \sum_{s=k}^{\infty} \frac{1}{s^2}. \end{aligned}$$

which proves that $A_k(x)B(x) \rightarrow 0$ uniformly on U , when $k \rightarrow +\infty$. We have proved that the product $\prod_{\mu \in \mathbb{N}} \left(1 + \frac{|w(x_\mu)|}{|w(x)|} \right)$ converges uniformly on U . In particular,

$$E_k(x) = \frac{\prod_{\mu \in \mathbb{N}} \left(1 + \frac{|w(x_\mu)|}{|w(x)|} \right)}{\prod_{\mu \in \Xi_k(x)} \left(1 + \frac{|w(x_\mu)|}{|w(x)|} \right)} \rightarrow 1, \quad \text{uniformly for } x \in U. \quad \square$$

Observe that by Remark 5.4, we have the extension operator $L_\Pi : \mathcal{F} \rightarrow \mathcal{O}(X)$. Now we are going to check its continuity. Note that \mathcal{F} and $\mathcal{O}(X)$ are endowed with the locally uniform convergence topologies.

CONTINUITY OF L_Π . Let $\mathcal{F} \ni \varphi_t \rightarrow \varphi \in \mathcal{F}$ locally uniformly. Fix $\varepsilon > 0$ and compact set $K \subset X$. Observe that $\mathcal{F}' := \mathcal{F} \cup (\varphi_t - \varphi)_{t=1}^\infty$ is also locally bounded. Moreover, there is $L'_\Pi : \mathcal{F}' \rightarrow \mathcal{O}(X)$ extension operator such that $L_\Pi = L'_\Pi$ on \mathcal{F} . Indeed, the map f_{d+1}^* is good for the both families \mathcal{F} and \mathcal{F}' if we take $\beta_k = \beta'_k \geq 2 \sup\{\sup_{G_k} |\varphi| : \varphi \in \mathcal{F}\}$, where β'_k is constant in the construction of the operator L'_Π . For $x \in K$ we have

$$\begin{aligned} |L_\Pi(\varphi_t) - L_\Pi(\varphi)|(x) &= |L'_\Pi(\varphi_t - \varphi)|(x) \\ &= \left| \sum_{s=1}^d \frac{1}{h_s(x)} \sum_{j=1}^\infty (\varphi_t - \varphi)(x_{s,j}) c_s(x_{s,j}) h_s(x_{s,j}) \prod_{\substack{\mu \in \mathbb{N} \\ \mu \neq j}} \left(1 - \frac{w_s(x_{s,\mu})}{w_s(x)}\right) \right| \\ &\leq \sum_{s=1}^d \frac{1}{h_s(x)} \prod_{\mu \in \mathbb{N}} \left(1 + \frac{|w_s(x_{s,\mu})|}{|w_s(x)|}\right) \sum_{j=1}^\infty |(\varphi_t - \varphi)(x_{s,j}) c_s(x_{s,j}) h_s(x_{s,j})|. \end{aligned}$$

Let $f(K) \subset \mathbb{D}^d(\alpha(k_0))$ and $k_1 \geq k_0$, where $k_0, k_1 \in \mathbb{N}$. By the proof of the previous lemma we get the following estimate

$$\prod_{\mu \in \mathbb{N}} \left(1 + \frac{|w_s(x_{s,\mu})|}{|w_s(x)|}\right) \leq \lambda_{k_0} e^{\delta+\gamma} + \sum_{s=k_0}^\infty \frac{e^\gamma}{s^2}.$$

Since K is compact, we conclude that the map $x \mapsto \frac{1}{h_s(x)} \prod_{\mu \in \mathbb{N}} \left(1 + \frac{|w_s(x_{s,\mu})|}{|w_s(x)|}\right)$ is bounded on K . On the other hand,

$$\sum_{j=1}^\infty |(\varphi_t - \varphi)(x_{s,j}) c_s(x_{s,j}) h_s(x_{s,j})| \leq M + \sum_{s=k_1}^\infty \frac{1}{s^2},$$

where $M := \lambda_{k_1} \sup_{Q_{k_1}} |(\varphi_t - \varphi) c_s h_s|$. Now we observe that if k_1 and t are sufficiently large, we obtain

$$\|L_\Pi(\varphi_t) - L_\Pi(\varphi)\|_K \leq \varepsilon. \quad \square$$

Acknowledgements. I would like to thank anonymous reviewer for valuable advices on this paper. I would also like to thank my supervisor professor Marek Jarnicki for help in writing the paper.

References

1. Arens R., *Dense inverse limit rings*, Michigan Math. J., **5** (1958), 169–182.
2. Bishop E., *Some global problems in the theory of functions of several complex variables*, Amer. J. Math., **83** (1961), 479–498.
3. Chirka E.M., *Complex analytic sets*, Kluwer Academic Publishers 1989.

4. Drzyzga K., *A remark on Bishop's multivalued projections*, Univ. Iag. Acta Math., **53** (2016), 13–20.
5. Gunning R., Rossi H., *Analytic Functions of Several Complex Variables*, Prentice-Hall, Englewood Cliffs, 1965.
6. Lojasiewicz S., *Introduction to Complex Analytic Geometry*, Birkhäuser, 1991.
7. Whitney H., *Complex Analytic Varieties*, Addison-Wesley Publishing Company, 1972.

Received July 26, 2016

Jagiellonian University
Faculty of Mathematics and Computer Science
Institute of Mathematics
Łojasiewicza 6
30-348 Kraków, Poland
e-mail: kamil.drzyzga@gmail.com