

Sharp, C; Pane, H; Ha, C; Venta, A; Patel, AB; Sturek, J; Fonagy, P; (2011) Theory of Mind and Emotion Regulation Difficulties in Adolescents With Borderline Traits. **J AM ACAD CHILD PSY** , 50 (6) 563 - 573. [10.1016/j.jaac.2011.01.017](https://doi.org/10.1016/j.jaac.2011.01.017).

RESEARCH ARTICLE

Theory of mind and emotion regulation difficulties in adolescents with
borderline traits

*Carla Sharp and Heather Pane**; *Jennifer Sturrek, Carolyn Ha, Ameer Patel and Amanda Venta***; *Peter Fonagy****

*Department of Psychology, University of Houston

**The Adolescent Treatment Program of the Menninger Clinic

***University College London, Research Dept of Clinical, Educational and Health Psychology

Corresponding author:

Carla Sharp, Ph.D.
Department of Psychology
University of Houston
126 Heyne Building
Houston, TX, 77024
Email: csharp2@uh.edu

ABSTRACT

Objective. Dysfunctions in both emotion regulation and social cognition (understanding behavior in mental state terms, or mentalizing) have been proposed as explanations for disturbances of interpersonal behavior in Borderline Personality Disorder (BPD). This study aimed to examine mentalizing in adolescents with emerging BPD from a dimensional and categorical point of view, controlling for sex, age, Axis I and Axis II symptoms, and to explore the mediating role of emotion regulation in the relation between theory of mind and borderline traits.

Method. The newly developed Movie for the Assessment of Social Cognition (MASC) was administered alongside self-report measures of emotion regulation and psychopathology to 111 adolescent inpatients between the ages of 12-17 (mean age = 15.5; $SD = 1.44$). For categorical analyses borderline diagnosis was determined through clinical interview, which showed that 35% of the sample met criteria for BPD.

Results. Findings suggest a unique relationship between borderline traits and 'overmentalizing' (excessive inaccurate mentalizing) independent of age, sex, externalizing, internalizing and psychopathy symptoms. The relation between overmentalizing and BPD traits was mediated by difficulties in emotion regulation, accounting for 43.5% of the overmentalizing to BPD path.

Conclusions. Results suggest that in borderline patients the loss of mentalization is more apparent in the emergence of unusual alternative strategies (overmentalizing)

than in the loss of the capacity *per se* (no mentalizing or undermentalizing). Moreover, for the first time, empirical evidence is provided to support the notion that mentalizing exerts its influence on borderline traits through the mediating role of emotion dysregulation.

Introduction

Disturbances in interpersonal relations are commonly considered one of the three core symptoms of Borderline Personality Disorder (BPD), alongside impulsivity and affective instability^{1, 2, 3, 4}. It has been proposed that dysfunction in mentalizing may lie at the foundation of these disturbances⁵⁻⁷. The concept of mentalizing has been in use in psychoanalytic literature since the 1970s.⁸ It was incorporated into the neurobiological, as well as the developmental literature^{9, 10} in the 1980s and 1990s, where it has been used interchangeably with the more frequently used concept of 'theory of mind' (ToM). Premack and Woodruff¹¹ coined the term 'theory of mind' to refer to the capacity to interpret other people's behavior within a mentalistic framework in order to understand how self and others think, feel, perceive, imagine, react, attribute, infer, and so on. It is through this capacity that we are able to engage in the activities that humans value most, such as family, friendship, love, cooperation, play, and community¹², but perhaps also gaining advantage in intra-species competition for resources¹³.

A wide range of constructs that may be considered aspects of mentalizing have been investigated in relation to BPD in adults and are reviewed elsewhere^{6, 14}. Given the developmental nature of the mentalization theory of BPD¹⁵, and the accumulating evidence of the seriousness of adolescent precursors of BPD¹⁶⁻¹⁸, mentalization could be an important early target for intervention, making it possible to influence the developmental trajectory of BPD^{19, 20}. To our knowledge, ToM (or mentalizing) has not yet been studied in relation to BPD in adolescents. There are two possible reasons for

this paucity of studies. First, the diagnosis of personality disorders in general in adolescents is still associated with controversy²¹⁻²³ and some clinicians appear to be reluctant to consider the diagnosis²⁴. The instability of personality in adolescence²⁵ and the stigma associated with a diagnosis of personality disorder are both reasons for this reluctance²⁶, along with the suggestion that symptoms of BPD are better explained by Axis I symptoms²⁷. However, there has been a steady increase in evidence supporting the diagnosis of juvenile BPD. As summarized in several recent review articles^{18, 28}, this includes evidence for longitudinal continuity^{29, 30}, a genetic basis³¹⁻³³, overlap in the latent variables underlying symptoms^{17, 34, 35} and the risk factors³⁶⁻³⁸ for adolescent BPD and the full-blown adult disorder, and evidence for marked separation of course and outcome of adolescent BPD and other Axis-I and Axis-II disorders^{20, 30, 39}.

A further challenge for studies investigating mentalizing dysfunction in adolescent BPD relates to measurement. Most ToM tasks developed over the last 20 years show ceiling effects in older age groups or lack divergent validity for disorders other than autism spectrum disorders⁴⁰. Developmentally more advanced tests of social cognition have been introduced in recent years⁴¹⁻⁴³ but these tend to measure only singular aspects of mentalizing, and do not resemble the demands of everyday-life social cognition⁴⁴. To address these limitations, Dziobek and colleagues⁴⁴ recently developed a naturalistic, video-based instrument for the assessment of ToM called the Movie for the Assessment of Social Cognition (MASC). The MASC not only allows for the usual dichotomous (right/wrong) response format, which is reflected in its total score, but also opens up the possibility of studying dysfunction in mentalization by including a

qualitative error analysis where wrong choices (distracters) correspond to one of three error categories: (1) undermentalizing involving insufficient mental state reasoning resulting in incorrect, “reduced” mental state attribution; (2) undermentalizing involving a complete lack of ToM; and (3) overmentalizing reflecting over-interpretative mental state reasoning⁴⁵. In addition the test considers different mental state modalities (thoughts, emotions, intentions) with positive, negative and neutral valence⁴⁴.

The first aim of this study was to investigate the relationship between borderline traits and mentalizing as measured by the MASC in a clinical sample of adolescents, in order to assess the specificity of mentalizing dysfunction in psychopathology involving BPD. There is considerable evidence for anomalous social cognition involving over-interpretive mentalizing associated with BPD, including reports of a general hypervigilance and hypersensitivity to social-emotional stimuli⁴⁶⁻⁴⁸, and findings suggesting that these individuals have difficulty with suppressing irrelevant aversive information⁴⁹. We predicted a positive relationship between BPD and overmentalizing or excessive ToM, from both a dimensional (trait) and categorical (diagnosis) perspective.

In examining this relationship, several potential confounding factors had to be controlled for. Studies have shown a correlation between increased ToM understanding and age⁵⁰, and female sex⁵¹. A gender difference has also been reported for BPD traits⁵², although not all studies find predominance of female individuals in adolescent BPD samples³⁰. The most common comorbid disorders with BPD have known social-cognitive deficits, particularly externalizing⁵⁴ and internalizing⁵⁵ problems on Axis I and

psychopathy on Axis II^{56, 57}. Moreover, given the concerns about the borderline construct in adolescence, and the high comorbidity between BPD and Axis I and Axis II conditions⁵³, we wished to control for these confounds in order to establish the specificity of the relationship of borderline personality features and mentalizing dysfunction by statistically controlling for these characteristics. We acknowledge that this is a conservative strategy in so far as these co-occurring demographic and clinical features may not be independent but may be part of this complex disorder. Taken together, we expected borderline traits to associate with overmentalizing, even when controlling for sex, age, symptoms of internalizing and externalizing disorder and psychopathic traits.

The second aim of the study was to investigate whether dysfunctional emotion regulation (ER) was an alternative (separate) or a linked aspect of vulnerability to BPD. The most comprehensive and coherent body of clinical research involving BPD has consistently emphasized the role of ER. Linehan's work⁵⁸ on the role of ER has not only provided a highly efficacious set of clinical interventions focused around this hypothesized dysfunction, but has also provided extensive cross-sectional and some developmental data linking ER to difficulties observed in BPD⁵⁹. ER and mentalizing may be independent predictors of borderline traits in adolescents. However, ER includes the awareness and understanding of emotions, the acceptance of emotions, and the ability to control impulsive behaviors and behave flexibly in accordance with desired goals when experiencing negative emotions⁵⁹, all of which overlap with the mentalizing construct. ER and mentalizing have not been studied in the same

individuals at the same time, either in adolescents or in adults with BPD. We have initially hypothesized that difficulties in emotion regulation may antedate and to some measure underpin mentalizing problems, because ER dysfunction may disrupt the social processes through which mentalizing is normally acquired and thus 'cause' dysfunctions in mentalization (Fonagy, P., Gergely, G., Jurist, E., & Target, M. (2002). *Affect Regulation, Mentalization and the Development of the Self*. New York: Other Press. Subsequently, Sharp and Fonagy^{6, 60} suggested that poor mentalizing capacity in the child may be associated with insecure attachment, which in turn is linked to poor parental mentalizing capacity, and that this may result in the development of psychopathology by bringing about ER difficulties. Mentalizing in our view involves the capacity to understand oneself as well as others in mentalistic terms that extend to states of affect, desire and belief (both epistemic and affective states), and thus it could be argued that mentalizing is an important component of emotion regulation, particularly in the context of social relationships. This study offered the opportunity to test a model in which ER problems play a role in mediating the relationship between mentalizing and BPD.

Methods

Participants

The sample included 111 consecutive admissions (62 girls and 49 boys) to the Adolescent Treatment Program of a private tertiary care inpatient treatment facility specializing in the evaluation and stabilization of adolescents who failed to respond to

previous interventions. Adolescents were between the ages of 12 and 17 (mean age = 15.5; $SD = 1.44$). All patients received a comprehensive psychiatric evaluation at intake. 80% of the sample was diagnosed with a mood disorder (dysthymia, major depressive disorder, bipolar disorder), 52% received an anxiety disorder diagnosis (PTSD, GAD, social phobia, other phobias, OCD) and 24% were diagnosed with an externalizing disorder (ADHD, Conduct Disorder, Oppositional Defiant Disorder). The modal number of diagnoses was two and the average number of diagnoses between two and three. Ten percent of the sample had at least one or more suicide attempts in the last year, while 27% had a lifetime history of one or more suicide attempts. 42% of the sample reported cutting during the last year, and 48% reported ever cutting. 48% of the sample scored above the clinical cut-off (T-score of 65) for internalizing disorders, and 37% for externalizing disorders on the YSR⁶¹, 23% of the sample ($n = 24$) met criteria for BPD on the Child Interview for DSM-IV Borderline Personality Disorder⁶². While the unit was in principle open to all mental disorders, the study adopted the following exclusion criteria: (1) diagnosis of schizophrenia or any psychotic disorder, and/or (2) diagnosis of mental retardation. Inclusion criteria were: (1) age between 12 and 17, and (2) sufficient fluency in English to complete all research. Complete data was absent for 4 individuals who were removed from the final dataset.

Measures

Theory of mind (mentalizing). The MASC⁴⁴ is a computerized test for the assessment of theory of mind or mentalizing abilities that approximates the demands of

everyday life⁶³. Subjects are asked to watch a 15-minute film about four characters getting together for a dinner party. Themes of each segment cover friendship and dating issues. Each character experiences different situations through the course of the film that elicit emotions and mental states such as anger, affection, gratefulness, jealousy, fear, ambition, embarrassment, or disgust. The relationships between the characters vary in the amount of intimacy (friends – strangers) and thus represent different social reference systems on which mental state inferences have to be made.

During administration of the task, the film is stopped at 45 points during the plot and questions referring to the characters' mental states (feelings, thoughts, and intentions) are asked (e.g., “What is Betty feeling?”, “What is Cliff thinking?”).

Participants' correct responses are scored as one point and added to an overall score. In addition to the total score, (i) overmentalizing, (ii) undermentalizing, and (iii) no mentalizing are scored. Similar to a study with young adults⁶³, we used the multiple-choice version of the MASC that offers four options for each query (MASC-MC): the three options described above, in addition to a control question which demands non-social inferences to be made, thereby controlling for verbal understanding of the task stimuli. The MASC is a reliable instrument that has proven sensitive in detecting subtle mindreading difficulties in adults of normal IQ⁴⁴, young adults under stress conditions⁶³, as well as patients with bipolar disorder⁴⁵, and autism⁶⁴.

Borderline Personality Features Scale for Children (BPFS-C). The BPFS-C is a self-report instrument that assesses borderline personality features among children and adolescents aged nine and older⁶⁵. The BPFS-C is based upon the BOR (borderline)

Scale of the Personality Assessment Inventory (PAI⁶⁶), modified for youth. A four-point Likert scale ranging from 1 (*not at all true*) to 5 (*always true*) is used to report on affective instability, identity problems, negative relationships and self-harm. The BPFSC has shown good internal consistency across 12 months as well as construct validity⁶⁵ and criterion validity⁶⁷. In the current study Cronbach's α was .90.

Childhood Interview for DSM-IV Borderline Personality Disorder (CI-BPD). The CI-BPD is a semi-structured interview that assesses DSM-IV BPD in latency-age children and adolescents⁶². It was adapted for use in youth from the Diagnostic Interview for Personality Disorders. After asking a series of corresponding questions, the interviewer rates each DSM-based criterion with a score of 0 ("absent"), 1 ("probably present"), or 2 ("definitely present"). The patient meets criteria for BPD if five or more criteria are met at the 2-level. The CI-BPD has adequate inter-rater reliability and demonstrated a significant relationship to clinician diagnosis at time of discharge in the current sample ($\chi^2 = 20.25, p < .001$). Internal consistency was good with a Cronbach's alpha of .82.

The Youth Self-Report. The Youth Self-Report (YSR)⁶¹ is a self-report measure of psychopathology. The measure contains 112 problem items, each scored on a 3-point scale (0= '*not true*', 1=*somewhat or sometimes true*', or 2=*very or often true*'). The measure yields a Total Problems *t*-score of general psychiatric functioning and two broad subscales of Externalizing behavior problems and Internalizing behavior problems. Externalizing is composed of the subscales Aggressive behavior and Rule-

breaking behavior; and Internalizing is composed of the subscales Anxious/Depressed, Withdrawn/Depressed, and Somatic Complaints.

The Antisocial Process Screening Device (APSD). The APSD⁶⁸ is the most commonly used questionnaire measure of youth psychopathic traits⁶⁹. It is a 20-item self-report measure designed to assess traits associated with the construct of psychopathy similar to those assessed by the PCL-R⁷⁰. Each item on the APSD is scored either 0='not at all true', 1='sometimes true', or 2='definitely true' with the total score indicating overall level of psychopathic traits.

The Difficulties in Emotion Regulation Strategies Scale (DERS). The DERS⁵⁹ provides a comprehensive assessment of difficulties in ER, including awareness and understanding of emotions, acceptance of emotions, the ability to engage in goal-directed behavior and refrain from impulsive behavior when experiencing negative emotions, as well as the flexible use of situationally appropriate strategies to modulate emotional responses. After the original validation study in undergraduate students⁵⁹, the DERS has recently been validated in a community sample of adolescents⁷¹.

Results

Descriptive statistics

Means, standard deviations and ranges for all main study variables are summarized in Table 1.

Table 1 about here

The relationship between mentalizing and borderline traits

Bivariate correlations between study variables are summarized in Table 2.

Table 2 about here

Table 2 shows that borderline traits were positively correlated with both Axis I (internalizing and externalizing problems) and psychopathy. Borderline traits were negatively correlated with the total ToM score (indicating reduced overall ToM capacity associated with increased borderline traits), which was clearly driven by a very strong correlation between ToM errors of the overmentalizing type ($r = .41$; $p < .001$). No other ToM errors correlated with borderline traits. Difficulties in emotion regulation were also strongly correlated with borderline traits ($r = .62$; $p < .001$).

Table 2 furthermore shows that, as expected, overmentalizing also correlated with age, internalizing problems, externalizing problems, and problems in emotion regulation, but not with psychopathy or gender. To determine the specificity of the relationship between overmentalizing and borderline traits, we performed a regression analysis with borderline traits as outcome variable and overmentalizing, internalizing problems, externalizing problems, age and sex as predictor variables. Sex was included in the regression because independent sample t-tests showed a significant difference between boys ($m = 63.90$; $SD = 16.37$) and girls ($m = 73.85$; $SD = 16.31$) on the BPFSC ($t = 3.15$; $df = 107$; $p = .002$).

Results of the hierarchical regression showed a moderately strong overall relationship between predictor variables and borderline traits, which was significantly improved by adding overmentalizing to the equation (R^2 change = .15, $F/t = 4.15$, $p = .04$). Together, predictor variables accounted for 69% of the variation in BPFSC scores (adjusted R^2). Overmentalizing was uniquely associated with borderline traits ($B = .91$;

$p = .002$), alongside sex ($B = -9.99$; $p < .001$), internalizing problems ($B = .39$; $p < .001$) and externalizing problems ($B = .67$; $p < .001$). All variables, however, were independently related to borderline features.

Mentalizing in adolescents meeting criteria for BPD vs. psychiatric controls

Independent sample t-tests showed that adolescents meeting criteria for BPD on the CI-BPD ($n = 28$; $m = 10.13$; $SD = 5.45$) were significantly more likely ($t = -2.27$; $p = .03$) to engage in overmentalizing compared to adolescents not meeting criteria on the CI-BPD ($n = 79$; $m = 7.46$; $SD = 3.36$). Group differences for all other ToM variables were non-significant.

A hierarchical logistic regression analysis with BPD diagnosis as outcome variable, and sex, age, internalizing problems, externalizing problems and overmentalizing as predictor variables confirmed the unique relationship between overmentalizing and BPD. Adding overmentalizing to the equation improved prediction of BPD from $x\%$ to $y\%$ (stats for the omnibus model) with overmentalizing ($\beta = .17$; $SE = .08$; $Wald = 4.04$; $df = 1$, $p = .04$), sex ($\beta = -2.62$; $SE = .77$; $Wald = 11.37$; $df = 1$, $p = .001$) and externalizing ($\beta = .97$; $SE = .35$; $Wald = 7.47$; $df = 1$, $p = .006$) making a significant contribution to the prediction.

Difficulties in emotion regulation as mediator in the relationship between overmentalizing and borderline traits

As we have seen, difficulties in ER (DERS) were strongly associated to BPFSC scores and we wished to examine whether DERS could be seen to serve as a mediator

of the relation between overmentalizing and borderline personality traits. We used standard mediational analyses methods^{72, 73}. Prior to testing for mediation, formal detection-tolerance and the variance inflation factor (VIF) were conducted to test for multicollinearity. It was unnecessary to center the predictor variable^{74, 75} since multicollinearity was not a problem (VIF = 1.082; tolerance = 0.925), with a tolerance of less than 0.20 or 0.10 and a VIF of less than 5 or 10.

Next, overmentalizing was regressed on the dependent variable, borderline personality traits, and then the mediator (DERS). In step one of the hierarchical regression, overmentalizing was significantly related to BPD traits [$t(1, 105) = 4.226, p < .0001$]. When DERS was added in step two, overmentalizing became less significant [$t(2, 105) = 2.934, p < .01$] and DERS was significantly related to BPD traits [$t(2, 105) = .686, p < .0001$]. Thus, DERS appeared to mediate some of the relation between overmentalizing and BPD (see Table 3).

Table 3 about here

Post-hoc probing of the significant mediation model was conducted with Sobel's equation^{72, 73}. The significance of the mediation effect found for DERS in the relation between overmentalizing and BPD traits was tested by regressing: (1) DERS on overmentalizing (B = 2.021, SE = .703); (2) BPD on overmentalizing and DERS (B = .793, SE = .270). To determine whether the mediation effect was statistically reliable, Sobel's test ($z = 2.77$) was performed and was found to be significant at the $p < .01$ level, with approximately 43.5% of the overmentalizing to BPD path accounted for by DERS. The values of path coefficients are visually represented in Figure 1.

Figure 1 about here

Discussion

This study is the first to use a ToM task that resembled the demands of everyday-life social cognition⁴⁴ to examine mentalizing difficulties in relation to borderline traits in adolescents. While other studies have investigated aspects of emotional processing in borderline youth⁷⁶, ours is the first to use a task specifically developed to assess mentalizing impairment in psychiatric disorder by considering potential dysfunctions of mentalizing such as insufficient mental state reasoning resulting in incorrect, “reduced” mental state attribution as opposed to a complete lack of ToM. Neither undermentalizing nor complete absence of mentalizing were linked to borderline traits. By contrast, overmentalizing (over-interpretive mental state reasoning) was strongly associated with BPD features in adolescents. Those with BPD features also showed a tendency to make overly complex inferences based on social cues that resulted in errors. They tended to over-interpret social signs^{45, 63}. Studies using this task have demonstrated general difficulties in ToM for individuals with autism spectrum disorders⁴⁴, and undermentalizing but not overmentalizing in adult euthymic bipolar patients⁴⁵. Although internalizing and externalizing scores were associated with overmentalizing, controlling for these and demographic predictors of mentalizing dysfunction did not eliminate the prediction from overmentalizing to borderline trait scores. Thus the current study adds to the growing body of evidence linking varying types of social cognitive dysfunctions to particular psychiatric disorders and specifically linking overmentalizing to borderline traits in adolescents. Taken together, these results

confirm clinical^{77, 78} and theoretical⁶ evidence that in borderline patients the dysfunction of mentalization is more apparent in the emergence of unusual alternative strategies (overmentalizing) than in the loss of the capacity *per se* (no mentalizing or undermentalizing). This is hardly surprising, since patients with BPD present quite differently from patients with autistic spectrum disorders where undermentalization is most commonly observed.

This is also the first study to examine ToM and difficulties in ER in relation to borderline traits in adolescents. While previous studies have examined these constructs independently from each other in relation to adult BPD, they have not yet been studied together in adolescents. Our results suggest that difficulties in ER at least in part mediate the association between overmentalizing and BPD. Bearing in mind that the cross-sectional nature of the data makes these findings suggestive rather than definitive, the mediational path analyses carried out here are at least consistent with the suggestion that overmentalizing in some adolescents may be indicative of their difficulties in regulating their emotional responses to social situations, either because they misattribute inappropriate epistemic or affective states to others, or because they poorly contextualize and perhaps overinterpret their own emotional reactions. In either case, overmentalizing may cause difficulties in ER, which in turn leads to the emergence of symptoms characteristic of BPD. Results from randomized clinical trials [Bateman, A. W., & Fonagy, P. (2001). Treatment of borderline personality disorder with psychoanalytically oriented partial hospitalization: an 18-month follow-up. *American Journal of Psychiatry*, 158(1), 36-42.; Bateman, A., & Fonagy, P. (2008). 8-year follow-

up of patients treated for borderline personality disorder: mentalization-based treatment versus treatment as usual. *Am J Psychiatry*, 165(5), 631-638. Bateman, A. W., & Fonagy, P. (2009). Randomized controlled trial of outpatient Mentalization-based Treatment versus Structured Clinical Management for borderline personality disorder. *American Journal of Psychiatry*, 166(12), 1355-1364.] testing a psychosocial intervention aimed at improving BPD symptoms by focusing on improving the quality of mentalization in an attachment context [78] are consistent with this model. We have suggested that asking patients to focus on emotional links of thoughts and other mental states specifically in an attachment context can lead to improved emotion regulation [Fonagy, P., & Bateman, A. W. (2006). Mechanisms of change in mentalization-based treatment of BPD. *J Clin Psychol*, 62, 411-430].

An alternative model of the cross-sectional associations we observed might suggest that ER problems may cause mentalization dysfunction. There is extensive neurophysiological evidence from adults and children that emotional arousal disrupts mentalizing in a range of contexts [Bartels, A., & Zeki, S. (2000). The neural basis of romantic love. *Neuroreport*, 11(17), 3829-3834; Bartels, A., & Zeki, S. (2004). The neural correlates of maternal and romantic love. *Neuroimage*, 21(3), 1155-1166; Gobbini, M. I., & Haxby, J. V. (2007). Neural systems for recognition of familiar faces. *Neuropsychologia*, 45(1), 32-41; Gobbini, M. I., Leibenluft, E., Santiago, N., & Haxby, J. V. (2004). Social and emotional attachment in the neural representation of faces. *Neuroimage*, 22(4), 1628-1635; Ortigue, S., Bianchi-Demicheli, F., Hamilton, A. F., & Grafton, S. T. (2007). The neural basis of love as a subliminal prime: an event-related

functional magnetic resonance imaging study. *J Cogn Neurosci*, 19(7), 1218-1230.]. Emotion dysregulation may have causes independent of mentalizing problems. It is well-known that borderline patients commonly have histories of significant trauma^{80, 81}. Recent animal research suggests that early trauma may permanently affect the HPA axis⁸². Research with traumatized children and adult female victims of childhood sexual abuse has also demonstrated persistent changes in the HPA axis⁸³⁻⁸⁶. Indeed, abnormal stress responsivity has been demonstrated in borderline patients^{87, 88}. Increased stress responsivity, in turn, affects mentalizing capacity. A recent study⁶³ used the MASC to show that high cortisol responding women make more mentalizing errors after stress induction – in particular due to a tendency to overmentalize - thereby demonstrating that stress responsivity modulates social cognition.

A less parsimonious but ultimately more plausible model would assume that mentalizing and emotion dysregulation represent separate but interacting difficulties in individuals with a vulnerability to BPD. In a dynamic developmental model, we may consider early affect dysregulation to undermine an individual's capacity to utilize social environments that are likely to enhance the development of mentalizing, particularly family environments [Dunn, J., & Brown, J. (2001). Relationships, talk about feelings, and the development of affect regulation in early childhood. In J. Garber & K. Dodge (Eds.), *Affect regulation and dysregulation in childhood* (pp. 89–108). Cambridge: Cambridge University Press. Dunn, J., Deater-Deckard, K., Pickering, K., & Golding, J. (1999). Siblings, parents, and partners: family relationships within a longitudinal community study. ALSPAC study team. *Avon Longitudinal Study of Pregnancy and*

Childhood. *J Child Psychol Psychiatry*, 40(7), 1025-1037], leading to dysfunctional mentalization. Overmentalizing, which involves over-interpreting social cues in others, in turn, derails the emotion regulation system spinning the adolescent into a vicious cycle of over-interpreting what others are thinking and being unable to regulate the anxious rumination caused by this over-interpretation.

There are several limitations to this study, most notably the cross-sectional nature of the mediational model. Further, we are just beginning to appreciate the complexities of the normal development of mentalizing in adolescence [Blakemore, S. J. (2008). The social brain in adolescence. *Nat Rev Neurosci*, 9(4), 267-277. Blakemore, S. J., den Ouden, H., Choudhury, S., & Frith, C. (2007). Adolescent development of the neural circuitry for thinking about intentions. *Soc Cogn Affect Neurosci*, 2(2), 130-139], which must provide the background for the anomalies observed in this group. Longitudinal studies will be needed to elaborate our understanding of the dynamic interplay of emotion regulation and mentalization across development. Notwithstanding this limitation, the current study is important as the first to examine mentalizing and emotion dysregulation in adolescent BPD. It has been suggested that disturbed relationships may be a phenotype for BPD in the same way that impulsivity and affective instability have been conceptualized¹. The psychological endophenotype of mentalizing offers an important bridge from the neurobiology of relationships to the more specific interpersonal impairments of BPD. It also provides a valuable target for treatment in adolescents with emerging BPD. Given that the MASC has recently been

adapted for fMRI⁸⁹, a natural next step would be to examine the neural correlates of overmentalizing in adults or adolescents with BPD.

Table 1. Means, standard deviations and ranges for all main study variables

	Mean	SD	Range
Age	15.49	1.44	12-17
Total BPFSC	69.47	17.00	30-112
YSR Internalizing	62.45	13.11	32-89
YSR Externalizing	60.96	10.81	34-91
Total APSD	15.32	5.74	0-33
Total ToM	31.84	5.48	10-39
Excessive ToM	8.11	4.08	2-26
No ToM	1.93	1.65	0-7
Less ToM	3.12	2.45	0-18
Control ToM	4.51	1.24	1-6
DERS total	102.18	31.08	38-173

Table 2. Bivariate correlations between main study variables (n = 107)

	Age	BPF	Int	Ext	YPI	TotTo	ExTo	8	9	10	1
	e	SC				M	M				1
1. Age	1	-									
2. BPFSC	-.03	1									
3. Int	.11	.53*	1								
		*									
4. Ext	.07	.60*	.35	1							
		*	**								
5. APSD	.13	.36*	.26	.61	1						
		*	*	**							
6. Tot	.27	-	-.03	-	.06	1					
ToM	**	.22*		.12							
7. Ex ToM	-	.41*	.25	.27	.16	-.78**	1				
	.25	*	**	**							
	**										
8. No ToM	.02	-.08	-.13	-	-.04	-.38**	-.02	1			
			.03								
9. Less	-.14	-.13	-	-	-	-.49	.04	.17	1		
ToM			.29	.16	.33*						
			**		*						
10. Cont	.12	.14	.11	-	-	.36**	-.24*	-	-	1	


ToM				.02	.001			.23	.25		
								*	**		
11.DERS	-.02	.75*	.62	.48	-	-.11	.25**	-.09	-.09	.14	1
		*	**	**			.32*				
						*					

Table 3. Summary of hierarchical regression analysis for mediation of overmentalizing to Borderline Personality Traits (n = 107)

Variable	B	SE B	β
Step 1			
	1.56	.370	.383**
Overmentalizing			
Step 2			
	.793	.270	.194*
Overmentalizing			
DERS	.375	.036	.686**

Note. $R^2 = .15$ for Step 1 ($p < .0001$); $\Delta R^2 = .58$ for Step 2 ($p < .0001$). MASC = Movie for the Assessment of Social Cognition; DERS = Difficulties in Emotion Regulation Scale. * $p < .01$, ** $p < .0001$.

Figure 1.



Note: Values on each path are standardized β 's (path coefficients). Those coefficients inside of parentheses are standardized partial regression coefficients from equations that include both variables with direct effects on the criterion or dependent variable.

References

1. Gunderson JG. Disturbed relationships as a phenotype for borderline personality disorder. *Am J Psychiatry*. Nov 2007;164(11):1637-1640.
2. Skodol AE, Siever LJ, Livesley WJ, Gunderson JG, Pfohl B, Widiger TA. The borderline diagnosis II: biology, genetics, and clinical course. *Biol Psychiatry*. Jun 15 2002;51(12):951-963.
3. Gunderson JG. Borderline personality disorder: The ontogeny of a diagnosis. *American Journal of Psychiatry*. 2009;166:530-539.
4. Lieb K, Zanarini MC, Schmahl C, Linehan MM, Bohus M. Borderline personality disorder. *Lancet*. Jul 31 2004;364(9432):453-461.
5. Fonagy P, Steele H, Moran G, Steele M, Higgitt A. The capacity for understanding mental states: The reflective self in parent and child and its significance for security of attachment. *Infant Mental Health Journal*. 1991;12(3):201-218.
6. Sharp C, Fonagy P. Social cognition and attachment-related disorders. In: Sharp C, Fonagy P, Goodyer I, eds. *Social cognition and developmental psychopathology*. Oxford: Oxford University Press; 2008:269-302.
7. Fonagy P, Target M. Playing with reality III: The persistence of dual psychic reality in borderline patients. *Int J Psychoanal*. 2000;81(5):853-874.
8. Allen JG. Mentalizing. *Bulletin of the Menninger Clinic*. 2003;67(2):91-112.
9. Morton J. The origins of autism. *New Scientist*. 9 December 1989 1989.

10. Frith CD. *The Cognitive Neuropsychology of Schizophrenia*. Hillsdale, NJ: Erlbaum; 1992.
11. Premack D, Woodruff G. Does the chimpanzee have a 'theory of mind'? *Behavior and Brain Sciences*. 1978;4:515-526.
12. Sharp C, Fonagy P, Goodyer IM. Introduction: Social cognition and developmental psychopathology. In: Sharp C, Fonagy P, Goodyer IM, eds. *Social cognition and Developmental Psychopathology*. Oxford: Oxford University Press; 2008.
13. Alexander RD. *The biology of moral systems*: De Gruyter; 1987.
14. Fonagy P, Bateman A. The development of borderline personality disorder-a mentalizing model. *J Personal Disord*. Feb 2008;22(1):4-21.
15. Fonagy P, Luyten P. A developmental, mentalization-based approach to the understanding and treatment of borderline personality disorder. *Dev Psychopathol*. Fall 2009;21(4):1355-1381.
16. Paris J. The development of impulsivity and suicidality in borderline personality disorder. *Development and Psychopathology*. 2005;17:1091-1104.
17. Gratz KL, Tull MT, Reynolds EK, et al. Extending extant models of the pathogenesis of borderline personality disorder to childhood borderline personality symptoms: the roles of affective dysfunction, disinhibition, and self- and emotion-regulation deficits. *Dev Psychopathol*. Fall 2009;21(4):1263-1291.

18. Bondurant H, Greenfield B, Tse SM. Construct validity of the adolescent borderline personality disorder: A review. *The Canadian Child and Adolescent Psychiatry Review*. 2004;13:53-57.
19. Winograd G, Cohen P, Chen H. Adolescent borderline symptoms in the community: prognosis for functioning over 20 years. *J Child Psychol Psychiatry*. Sep 2008;49(9):933-941.
20. Crawford TN, Cohen PR, Chen H, Anglin DM, Ehrensaft M. Early maternal separation and the trajectory of borderline personality disorder symptoms. *Dev Psychopathol*. Summer 2009;21(3):1013-1030.
21. Sharp C, Bleiberg E. Borderline Personality Disorder in children and adolescents. In: Martin A, Volkmar F, eds. *Lewis Child and Adolescent Psychiatry: Comprehensive Textbook*. Baltimore: Lippincott Williams and Wilkins 2007:680-691.
22. Paris J. Personality disorders over time: Precursors, course and outcome. *Journal of Personality Disorders*. 2003;17(6):479-488.
23. Vito E, Ladame F, Orlandini A. Adolescence and personality disorders: Current perspectives on a controversial problem. In: Derksen J, Maffei C, Groen H, eds. *Treatment of personality disorders*. New York: Kluwer Academic/Plenum; 1999:77-95.
24. Chanen AM, Jackson HJ, McCutcheon LK, et al. Early intervention for adolescents with borderline personality disorder using cognitive analytic therapy: randomised controlled trial. *Br J Psychiatry*. Dec 2008;193(6):477-484.

25. Meijer M, Goedhart AW, Treffers PDA. The persistence of borderline personality disorder in adolescence. *J Personal Disord.* 1998;12:13-22.
26. Chanen AM, Jackson HJ, McGorry PD, Allot KA, Clarkson V, Yuen HP. Two-year stability of personality disorder in older adolescent outpatients. *J Pers Disord.* Dec 2004;18(6):526-541.
27. Chanen AM, Jovev M, Jackson HJ. Adaptive functioning and psychiatric symptoms in adolescents with borderline personality disorder. *J Clin Psychiatry.* Feb 2007;68(2):297-306.
28. Sharp C, Romero C. Borderline personality disorder: a comparison between children and adults. *Bull Menninger Clin.* Spring 2007;71(2):85-114.
29. Cohen P, Chen H, Gordon K, Johnson J, Brook J, Kasen S. Socioeconomic background and the developmental course of schizotypal and borderline personality disorder symptoms. *Dev Psychopathol.* Spring 2008;20(2):633-650.
30. Bornovalova MA, Hicks BM, Iacono WG, McGue M. Stability, change, and heritability of borderline personality disorder traits from adolescence to adulthood: a longitudinal twin study. *Dev Psychopathol.* Fall 2009;21(4):1335-1353.
31. Torgersen S, Czajkowski N, Jacobson K, et al. Dimensional representations of DSM-IV cluster B personality disorders in a population-based sample of Norwegian twins: A multivariate study. *Psychological Medicine.* 2008;38:1617-1625.

32. Kendler KS, Aggen SH, Czajkowski N, et al. The structure of genetic and environmental risk factors for DSM-IV personality disorders: a multivariate twin study. *Arch Gen Psychiatry*. Dec 2008;65(12):1438-1446.
33. Distel MA, Trull TJ, Derom CA, et al. Heritability of borderline personality disorder features is similar across three countries. *Psychol Med*. Sep 2008;38(9):1219-1229.
34. Leung S-W, Leung F. Construct validity and prevalence rate of borderline personality disorder among Chinese adolescents. *Journal of Personality Disorder*. 2009;23(5):494-513.
35. Bradley R, Zittel Conklin C, Westen D. The borderline personality diagnosis in adolescents: gender differences and subtypes. *J Child Psychol Psychiatry*. Sep 2005;46(9):1006-1019.
36. Rogosch FA, Cicchetti D. Child maltreatment, attention networks, and potential precursors to borderline personality disorder. *Dev Psychopathol*. Fall 2005;17(4):1071-1089.
37. Carlson EA, Sroufe LA, Egeland B. The construction of experience: a longitudinal study of representation and behavior. *Child Dev*. Jan-Feb 2004;75(1):66-83.
38. Carlson EA, Egeland B, Sroufe LA. A prospective investigation of the development of borderline personality symptoms. *Dev Psychopathol*. Fall 2009;21(4):1311-1334.

39. Crick NR, Murray-Close D, Woods K. Borderline personality features in childhood: a short-term longitudinal study. *Dev Psychopathol.* Fall 2005;17(4):1051-1070.
40. Sharp C. Mentalizing problems in childhood disorders. In: Allen JG, Fonagy P, eds. *Handbook of mentalization-based treatments.* Chichester: John Wiley & Sons; 2006:101-121.
41. Baron Cohen S, Wheelwright S, Hill J, Raste Y, Plumb I. The "Reading the mind in the eyes" Test revised version: A study with normal adults, and adults with Asperger syndrome or high-functioning autism. *Journal of Child Psychology and Psychiatry and Allied Disciplines.* 2001;42(2):241-251.
42. Sharp C, Croudace TJ, Goodyer IM. Biased mentalising in children aged 7-11: Latent class confirmation of response styles to social scenarios and associations with psychopathology. *Social Development.* 2007;16(1):81-202.
43. Happé FGE. An advanced test of theory of mind: Understanding of story characters' thoughts and feelings by able autistic, mentally handicapped, and normal children and adults. *Journal of Autism and Developmental Disorders.* 1994;24(2):129-154.
44. Dziobek I, Fleck S, Kalbe E, et al. Introducing MASC: a movie for the assessment of social cognition. *J Autism Dev Disord.* Jul 2006;36(5):623-636.
45. Montag C, Ehrlich A, Neuhaus K, et al. Theory of mind impairments in euthymic bipolar patients. *J Affect Disord.* Sep 12 2009.

46. Donegan NH, Sanislow CA, Blumberg HP, et al. Amygdala hyperreactivity in borderline personality disorder: implications for emotional dysregulation. *Biol Psychiatry*. Dec 1 2003;54(11):1284-1293.
47. Lynch TR, Rosenthal MZ, Kosson DS, Cheavens JS, Lejuez CW, Blair RJ. Heightened sensitivity to facial expressions of emotion in borderline personality disorder. *Emotion*. Nov 2006;6(4):647-655.
48. Bland AR, Williams CA, Scharer K, Manning S. Emotion processing in borderline personality disorders. *Issues Ment Health Nurs*. Oct-Nov 2004;25(7):655-672.
49. Domes G, Winter B, Schnell K, Vohs K, Fast K, Herpertz SC. The influence of emotions on inhibitory functioning in borderline personality disorder. *Psychol Med*. Aug 2006;36(8):1163-1172.
50. Astington JW, Jenkins JM. Theory of mind development and social understanding. *Cognition and Emotion*. 1995;9(2-3):151-165 URLJ: <http://www.tandf.co.uk/journals/02699931.html>.
51. Baron-Cohen S. *The essential difference: The truth about the male and female brain*. New York: Basic Books; 2003.
52. Paris J. Gender differences in personality traits and disorders. *Curr Psychiatry Rep*. Feb 2004;6(1):71-74.
53. Grant BF, Chou SP, Goldstein RB, et al. Prevalence, correlates, disability, and comorbidity of DSM-IV borderline personality disorder: results from the Wave 2 National Epidemiologic Survey on Alcohol and Related Conditions. *J Clin Psychiatry*. Apr 2008;69(4):533-545.

54. Sharp C. Theory of mind and conduct problems in children: Deficits in reading the 'emotions of the eyes'. *Cognition and Emotion*. 2008;22(6):1149-1158.
55. Kyte Z, Goodyer I. Social cognition in depressed children and adolescents. In: Sharp C, Fonagy P, Goodyer I, eds. *Social cognition and developmental psychopathology*. Oxford: Oxford University Press; 2008:201-237.
56. Dadds MR, Perry Y, Hawes DJ, et al. Attention to the eyes and fear-recognition deficits in child psychopathy. *British Journal of Psychiatry* 2006;189:280-281.
57. Richell RA, Mitchell DG, Newman C, Leonard A, Baron-Cohen S, Blair RJ. Theory of mind and psychopathy: can psychopathic individuals read the 'language of the eyes'? *Neuropsychologia*. 2003;41(5):523-526.
58. Linehan MM. *Cognitive-behavioral treatment of borderline personality disorder*. New York: The Guildford Press; 1993.
59. Gratz KL, Roemer L. Multidimensional assessment of emotion regulation and dysregulation: Development, factor structure and initial validation of the Difficulties in Emotion Regulation Scale. *Journal of Psychopathology and Behavioral Assessment*. 2004;26(1):41-54.
60. Sharp C, Fonagy P. The parent's capacity to treat the child as a psychological agent: Constructs, measures and implications for developmental psychopathology. *Social Development*. 2008;17(3):737-754.
61. Achenbach TM, Rescorla LA. *Manual for ASEBA school-age forms and profiles*. Burlington: University of Vermont, Research Center for Children, Youth and Families; 2001.

62. Zanarini MC. *The Child Interview for DSM-IV Borderline Personality Disorder*. Belmont, MA: McLean Hospital; 2003.
63. Smeets T, Dziobek I, Wolf OT. Social cognition under stress: Differential effects of stress-induced cortisol elevations in healthy young men and women. *Horm Behav*. Feb 6 2009.
64. Dziobek I, Fleck S, Rogers K, Wolf OT, Convit A. The 'amygdala theory of autism' revisited: Linking structure to behavior. *Neuropsychologia*. 2006;44:1891-1899.
65. Crick NR, Murray-Close D, Woods K. Borderline personality features in childhood: a short-term longitudinal study. *Development and Psychopathology*. Fall 2005;17(4):1051-1070.
66. Morey L. *Personality Assessment Inventory*. Odessa, FL: Psychological Assessment Resources 1991.
67. Chang B, Sharp C, Ha C. The criterion validity of the Borderline Personality Feature Scale for Children in an adolescent inpatient setting. *Journal of Personality Disorders*. in press.
68. Frick PJ, Hare RD. *The Antisocial Process Screening Device*. Toronto: Multi-Health Systems; 2001.
69. Sharp C, Kine S. The assessment of juvenile psychopathy: Strengths and weaknesses of currently used questionnaire measures. *Child and Adolescent Mental Health*. 2008;13(2):85-95.

70. Hare RD. *The Hare Psychopathy Checklist-Revised Manual*. Toronto: Multi-Health Systems; 1991.
71. Neumann A, van Lier PAC, Gratz KL, Koot HM. Multidimensional assessment of emotion regulation difficulties in adolescents Using the Difficulties in Emotion Regulation Scale
Assessment. 2010;17(1):138-149
72. Hombeck GN. Toward terminological, conceptual and statistical clarity in the study of mediators and moderators: Examples from the child-clinical and pediatric psychology literatures. *Journal of Consulting and Clinical Psychology*. 1997;65(599-610).
73. Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: Conceptual, strategic and statistical considerations. . *Journal of Personality and Social Psychology*. 1986;51:1173-1182.
74. Hombeck GN. Post-hoc probing of significant moderational and mediational effects in studies of pediatric populations. *Journal of Pediatric Psychology* 2002;27(87-96).
75. Aiken LS, West SG. *Multiple regression: Testing and interpreting interactions*. Newbury Park, CA: Sage; 1991.
76. von Ceumern-Lindenstjerna IA, Brunner R, Parzer P, Mundt C, Fiedler P, Resch F. Initial orienting to emotional faces in female adolescents with borderline personality disorder. *Psychopathology*.43(2):79-87.

77. Allen J. *Traumatic attachments*. New York: John Wiley; 2002.
78. Bateman A, Fonagy P. *Psychotherapy for borderline personality disorder: Mentalization-based treatment*. Oxford: Oxford University Press; 2004.
79. Baron Cohen S, Leslie AM, Frith U. Does the autistic child have a "theory of mind"? *Cognition*. 1985;21(1):37-46 URLJ: <http://www.elsevier.com/inca/publications/store/35/30/35/36/32/36/>.
80. Zanarini MC. Childhood experiences associated with the development of borderline personality disorder. *Psychiatr Clin North Am*. Mar 2000;23(1):89-101.
81. Zanarini MC, Gunderson JG, Marino MF, Schwartz EO, Frankenburg FR. Childhood experiences of borderline patients. *Comprehensive Psychiatry*. Jan-Feb 1989;30(1):18-25.
82. Oitzl MS, Workel JO, Flutters M, Frosch F, De Kloet ER. Maternal deprivation affects behaviour from youth to senescence: Amplification of individual differences in spatial learning and memory in senescent Brown Norway rats. *European Journal of Neuroscience*. 2000;12:3771-3780.
83. De Bellis MD, Chrousos GP, Dorn LD, Burke L, Helmers K, Kling MA. Hypothalamic-pituitary-adrenal axis dysregulation in sexually abused girls. *Journal of Clinical Endocrinology and Metabolism*. 1994;78:249-255.
84. Kaufman J, Birmaher B, Perel J, et al. The corticotropin-releasing hormone challenge in depressed abused, depressed nonabused, and normal control children. *Biological Psychiatry*. 1997;42:669-679.

85. Heim C, Newport DJ, Heit S, et al. Pituitary-adrenal and autonomic responses to stress in women after sexual and physical abuse in childhood. *JAMA*. 2000;284(592-597).
86. Heim C, Newport DJ, Bonsall R, Miller AH, Nemeroff CB. Altered pituitary-adrenal axis responses to provocative challenge tests in adult survivors of childhood abuse. *American Journal of Psychiatry*. 2001;158(575-581).
87. Rinne T, de Kloet ER, Wouters L, Goekoop JG, DeRijk RH, van den Brink W. Hyperresponsiveness of hypothalamic-pituitary-adrenal axis to combined dexamethasone/corticotropin-releasing hormone challenge in female borderline personality disorder subjects with a history of sustained childhood abuse. *Biol Psychiatry*. Dec 1 2002;52(11):1102-1112.
88. Rinne T, Westenberg HG, den Boer JA, van den Brink W. Serotonergic blunting to meta-chlorophenylpiperazine (m-CPP) highly correlates with sustained childhood abuse in impulsive and autoaggressive female borderline patients. *Biol Psychiatry*. Mar 15 2000;47(6):548-556.
89. Wolf I, Dziobek I, Heekeren HR. Neural correlates of social cognition in naturalistic settings: a model-free analysis approach. *Neuroimage*. Jan 1 2009;49(1):894-904.