

1
2
3
4 **The impact of chemotherapy-related nausea on patients'**
5 **nutritional status, psychological distress and quality of life**
6
7
8
9

10
11
12
13 Carole Farrell¹, Sarah G Brearley², Mark Pilling³, Alex Molassiotis^{3*},
14
15

16
17 ¹Christie NHS Foundation Trust, Manchester, UK
18

19 ²Faculty of Health and Medicine, Lancaster University, Lancaster, UK
20

21 ³School of Nursing, Midwifery & Social Work, University of Manchester, Manchester,
22
23 UK
24
25
26
27
28
29

30 ***Corresponding author contact details:**
31

32 Alexander Molassiotis, RN, PhD
33

34 Professor of Cancer & Supportive Care
35

36 University of Manchester
37

38 School of Nursing
39

40 University Place
41

42 MANCHESTER M13 9PL,
43

44 UK
45

46 email: alex.molassiotis@manchester.ac.uk
47

48 tel: +44 (0)161-3067830; fax: +44 (0)161-3067894
49

50 **Acknowledgements:**

51 This study was funded by the European Oncology Nursing Society through its Major
52 Grant Award.
53

54 **Conflict of Interest**

55 Authors declare no conflict of interest.
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

Abstract:

Purpose: Nausea is a troublesome and distressing symptom for patients receiving chemotherapy. While vomiting is well-controlled with current antiemetics, nausea is a more difficult symptom to manage. The aim of this study was to assess the impact of nausea on nutritional status, quality of life and psychological distress.

Methods: This was a prospective observational study over two cycles of chemotherapy. Patients completed the MASCC Antiemesis Tool, a measure of nutritional status (PG-SGA), the FACT-G quality of Life scale and the Hospital Anxiety & Depression Scale at the end of each chemotherapy cycle (around day 10 post-chemotherapy).

Results: The sample consisted of 104 patients, primarily female, receiving anthracycline-based chemotherapy. While vomiting was minimal (5.2-14.6% of patients), high levels of nausea were observed (55.2%-72.9%), and severe nausea (>6 on a 0-10 scale) was reported by 20.5%-29.2% of the participants. Severe nausea had a borderline significant impact in relation to physical functioning (p=0.025) and a significant impact on nutritional status (severe acute nausea, p=0.003; severe delayed nausea, p=0.017). Clinically meaningful changes were observed in relation to the FACT-G total score.

Conclusion: Chemotherapy induced nausea does have an impact on nutritional status and physical functioning and can impair anxiety and quality of life. As a key symptom associated with other symptoms it is imperative that greater attention is given to managing treatment-related nausea through innovative non-pharmacological and nutritional interventions.

49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Key words: nausea, chemotherapy, nutrition, anxiety, depression, quality of life, physical functioning

The impact of chemotherapy-related nausea on patients' nutritional status, psychological distress and quality of life

Background

Chemotherapy-induced nausea is a significant problem in clinical practice, with 42-52% of patients experiencing nausea post-chemotherapy in routine practice [15,25]. Despite the availability of effective anti-emetics, chemotherapy-induced nausea and vomiting (CINV) is one of the most feared adverse events associated with chemotherapy [8,12,37]. The subjective and unobservable nature of CINV creates challenges in assessment [17,19] and means that clinicians tend to underestimate patients' experiences [17]. Generally, the clinical assessment of nausea is quite poor with clinical attention focused primarily on managing chemotherapy-induced vomiting rather than on the potential impact of chemotherapy-induced nausea. Unlike vomiting, nausea is more subjective and difficult for clinicians to evaluate and treat.

CINV can have a profoundly negative impact on social, physical and emotional functioning and on quality of life [9,15,22]. Moreover, despite recognition that nausea and vomiting are two related but separate entities, little attention has been directed to the concept of chemotherapy-induced nausea [2,23]. The impact of combined nausea and vomiting on quality of life (QOL) has been highlighted in the literature. Osoba et al [30] in a study of 832 chemotherapy-naïve patients showed that those patients who had both nausea and vomiting had worse physical, cognitive and social functioning, global quality of life, fatigue, anorexia and dyspnoea compared to those who did not experience nausea and vomiting. Similarly, in a study of 119 patients receiving chemotherapy it was reported that those experiencing either nausea or vomiting had decreased QOL in several functioning and symptom subscales of the EORTC-QLQ-C30 scale [36]. A more recent study focusing on delayed nausea and vomiting also showed QOL impairments in patients experiencing these symptoms, highlighting that nausea had a stronger negative impact than vomiting on patients' daily lives [5]. Furthermore, there are indications that the presence of 'nutrition impact symptoms' that include nausea and impede oral intake are linked with poorer QOL [40].

1
2 Initial evidence from a study involving 220 lung cancer patients suggests that
3 nausea forms a cluster of symptoms together with appetite loss, fatigue, weight
4 loss, taste changes and vomiting [16]. Another study also highlighted that nausea,
5 vomiting, feeling bloated, appetite loss, difficulty swallowing and taste changes are
6 part of a gastrointestinal symptom cluster [27]. No work has focused to date on the
7 nausea experience and nutritional impairment during chemotherapy, and
8 establishing a link between nutritional symptoms and nausea seems imperative.
9
10
11
12
13
14
15

16 Hence, the aim of this study is to assess the impact of nausea on patients'
17 nutritional status, quality of life and psychological distress.
18
19
20

21 **Methods**

22 A prospective observational study of patients over two cycles of chemotherapy
23 treatment was undertaken using a quantitative descriptive exploratory design.
24
25
26

27 **Population and setting**

28 The study was carried out in a large cancer centre in the UK after approval from the
29 local NHS Ethics and Research Committee. Consecutive patients were recruited if
30 they were about to receive moderate or highly emetogenic adjuvant chemotherapy
31 every three weeks as an outpatient and were willing to participate in the study.
32 Patients were excluded if chemotherapy was palliative/disease stage was IV, if they
33 had head and neck or upper gastrointestinal cancer, or if they were experiencing
34 nausea from other causes.
35
36
37
38
39
40
41
42

43 Patients were recruited by a research assistant who provided detailed information
44 about the study, and if patients agreed to participate they signed a consent form.
45 Participants completed the study instruments prior to the first cycle of
46 chemotherapy, at the end of cycle 1 and the end of cycle 2 and returned to the
47 investigators either directly in clinic or using a pre-paid envelope.
48
49
50
51
52
53

54 **Study assessments**

55 Information about socio-demographic characteristics, type of cancer and
56 chemotherapy regimens were obtained from the patients' medical records or the
57 patients themselves.
58
59
60
61
62
63
64
65

1 Nausea and vomiting was assessed using the Multinational Association of
2 Supportive Care in Cancer (MASCC) Antiemesis Tool (MAT) [26]. This 8-item
3 scale assesses presence (yes/no) and severity (0-10) of acute and delayed
4 nausea, and presence (yes/no) and severity (number of times) of vomiting during
5 chemotherapy. Severe nausea was calculated based on the MAT visual analogue
6 scale (VAS) score of 6-10 in the respective items. The MAT also defines nausea
7 for the patients as '*the feeling that you might vomit*'.

16 Nutritional status was assessed using the Patient Generated-Subjective Global
17 Assessment (PG-SGA) [12,31]. This includes weight changes, alterations in food
18 intake, gastrointestinal symptoms, changes in functional capacity and physical
19 signs of malnutrition. Higher scores on the PG-SGA suggest a greater risk of
20 malnutrition and scores of ≥ 9 are indicative of malnutrition, in critical need of
21 symptom improvement and/or nutritional intervention. In addition, objective
22 measurements of weight, height, body mass index (BMI) and albumin levels were
23 also recorded.

32 The Hospital Anxiety and Depression Scale (HADS) [42], a 14-item scale, was
33 used to assess the presence and severity of anxiety and depression (7 items
34 respectively). Scores < 8 are within the normal range, scores of 8-10 indicate
35 borderline cases needing further assessment before establishing
36 psychopathology, and scores 11+ indicate cases of clinical anxiety or depression
37 respectively.

45 The Functional Assessment of Cancer Therapy-General (FACT-G) scale [7] was
46 used to measure quality of life. This covers four dimensions of QOL: physical,
47 social and family, emotional and functional well-being. The FACT-G questionnaire
48 is scored using a 5-point scale from 0 = 'not at all' to 4 = 'very much'.

55 **Data analysis**

57 Data was coded and entered into SPSS (v.15) and descriptive statistics used to
58 summarize the data with missing values omitted from the calculation of

1 percentages. One-way analysis of covariance (ANCOVA) was used to assess the
2 effect of nausea on four sets of binary groups (defined by MAT subscales: acute
3 nausea (yes/no), severe (≥ 6) acute nausea, delayed nausea (yes/no), severe (≥ 6)
4 delayed nausea), in relation to nutritional status, psychological distress and quality
5 of life, using the baseline scores as covariates. The Bonferroni adjustment was
6 used to set the level of significance at $5\%/4=0.0125$. Friedman's Chi-squared (χ^2)
7 non-parametric test was used to compare differences between groups at multiple
8 time-points (i.e. repeated measures).
9
10
11
12
13
14
15
16
17

18 Results

19
20
21 106 cancer patients participated in the study, however the analysis was undertaken
22 on data from 104 patients, as 2 sets of questionnaires were incomplete and
23 unusable. 92% completed all questionnaires at end of cycle 1 (n=96, 8% attrition)
24 and 85% at end of cycle 2 (n=88, 15% attrition). Attrition was related primarily to the
25 changing health status of the patients and being overwhelmed with the
26 chemotherapy. Based on a power calculation for a bivariate test, in order to achieve
27 an alpha of 0.05 and power of 0.80, when the lowest correlation is 0.24, an
28 estimated number of 126 patients, as seen in past correlational studies, was
29 required. In practise, only 104 patients were available for analysis which led to a
30 small reduction in power ($=0.72$).
31
32
33
34
35
36
37
38
39
40

41 The participants were mainly women (90.3%, 93/103). 73% (73/100) were married
42 or with a partner, 58% were working and 25% had retired. The majority had
43 completed secondary (41.1%, 37/90) or college (40%) education, while the
44 remaining 18.9% had university or higher education. The mean age was 53.2 (SD
45 11.6, range=29-79). 83 (80.6%, 83/103) patients had breast cancer, 10 (9.7%) had
46 bladder cancer and 8 (7.8%) ovarian cancer. Chemotherapy regimens were mainly
47 combinations of cytotoxic drugs: 81 (78.7%) contained anthracyclines, 3 (2.9%)
48 contained taxanes, and 19 (18.5%) were platinum-based. All patients received as
49 primary antiemetics IV 8mg ondansetron with IV 8mg dexamethasone prior to
50 chemotherapy administration and twice daily oral ondansetron (8mg) and
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

dexamethasone (4mg) with oral metoclopramide 10mg (as required) for 2-3 days after chemotherapy.

Nausea and Vomiting

MAT scores highlighted a decrease in acute vomiting between cycle 1 (14.6%, 14/96) and cycle 2 (9.1%, 8/88) of chemotherapy, although the incidence of delayed vomiting increased slightly from 5.2 to 8.0% (5/96 to 7/87). However, the incidence of nausea was much greater with 55.2-64.8% of participants reporting acute nausea and 72.9-67.8% reporting delayed nausea after their first and second cycles of chemotherapy respectively [Figure 1].

INSERT FIGURE 1 HERE

The severity of acute and delayed vomiting was measured by the number of times participants vomited in the first 24 hours after chemotherapy. In terms of acute vomiting this ranged from one episode (n=6/96, 6.2% in cycle 1; n=6/88, 6.8% in cycle 2), two episodes (5% in cycle 1, 2.3% in cycle 2), and three episodes (3% in cycle 1, 0% in cycle 2). In terms of delayed vomiting after cycle 1 this ranged from 0-8 episodes (mean 0.24, SD= 1.17). The duration was less after cycle 2 (range 0-5 episodes, mean 0.17, SD= 0.70). In contrast, almost a third (20.5-29.2%) of patients experienced severe nausea (either acute or delayed) [Figure 2].

INSERT FIGURE 2 HERE

Nutritional status

Baseline weight ranged from 50-125kg (mean 74.05, SD 14.96) with little change over time and the BMI ranged from 19-42 kg/m² (mean 27.63, SD 5.06). Clinically, 27/85 (32%) were of normal weight, 37 (44%) overweight and 21 (25%) were obese. Serum albumin was used to assess for possible malnutrition (<35g/dL) based on a normal reference range of 35 – 50 g/L. All participants but one had a serum albumin within the normal range at each time point.

Participants' scores on the PG-SGA indicated deteriorating nutritional status; 25.3% had symptoms of malnutrition after one cycle of chemotherapy and this was

1 maintained across a further chemotherapy cycle [Table 1]. There was a statistically
2 significant deterioration in PG-SGA scores from baseline (median=1) to the end of
3 cycles 1 and 2 (median=5 and 4 respectively, $p<0.0001$ Friedman χ^2). There is no
4 trend in relation to presence of nausea and serum albumin levels. However, weight
5 and BMI at cycle 1 has a trend to be lower for patients with poor nutritional status as
6 can be seen in Table 1, although this did not reach statistical significance (ANOVA
7 for weight and BMI respectively at cycle 2; $p=0.2$, $p=0.1$). On their own, however,
8 albumin was lower after baseline (Friedman test, $p<0.001$), whereas weight was
9 lower at the start of cycle 2 ($p=0.013$) as it was for BMI ($p=0.006$).
10
11
12
13
14
15
16
17

18 INSERT TABLE 1 HERE
19
20

21 **Psychological distress**

22 HADS scores showed variability in the range and mean scores at baseline, end of
23 cycle 1 and end of cycle 2 chemotherapy . Levels of depression were quite low at
24 baseline (score for clinical case of $>10=1.9\%$; $n=2/103$), although it showed a
25 significant deterioration after baseline (7.3% after cycle 1 [$n=7/96$] and 14.1% after
26 cycle 2 [$n=12/85$]) ($p=0.0005$, Friedman χ^2 over time). In contrast, the incidence of
27 anxiety was much higher at baseline ($n=19/102$, 18.6%), with a slight increase at
28 the end of Cycle 1 ($n=19/96$, 19.8%), although this change was not statistically
29 significant.
30
31
32
33
34
35
36
37

38 **Quality of Life**

39 FACT-G total scores can range from 0-108, with higher scores indicating better QOL.
40 Participants' total scores indicated deterioration in QOL after the start of
41 chemotherapy. Mean scores were 83.68 (SD=14.5) at baseline, 77.8 (SD=17.7) at
42 the end of cycle 1 and 78.4 (SD=16.4) at end of cycle 2; this change was statistically
43 significant ($p=0.002$, Friedman χ^2).
44
45
46
47
48
49

50 **Impact of nausea on quality of life and nutritional status**

51 The proportion of patients at the end of cycle 1 who had a score ≥ 9 in the PG-SGA,
52 indicative of malnutrition, was higher in those experiencing acute nausea and delayed
53 nausea (58.3% and 79.2% respectively) than those who did not experience these
54 symptoms (41.7% and 20.8% respectively). This indicates a difference of 16.6% and
55 58.4% respectively in patients experiencing the two symptoms also having a PG-SGA
56
57
58
59
60
61
62
63
64
65

1 score ≥ 9 . Similar data is also evident in relation to cycle 2 (72.7% vs. 27.3% for both
2 acute and delayed nausea) . This trend is also present in relation to psychological
3 distress at cycle 1 (62.5% and 75% for acute and delayed nausea respectively vs.
4 37.5 and 25%). Fisher's tests of difference for the above were all non-significant.
5
6
7

8
9 ANCOVA was used to identify significant associations between MAT nausea scores
10 and QOL, nutritional status and psychological distress after cycle 1 with the level of
11 significance set at 0.0125 due to testing four different binary groupings of the same
12 patients. In general, baseline covariates were significant in the ANCOVA. Severe
13 acute nausea had a borderline significant impact on physical QOL ($p=0.025$), and a
14 significant impact on nutritional status ($p=0.003$ for severe acute and $p=0.017$ for
15 severe delayed nausea) [Table 3]. At cycle 2, ANCOVA calculations were not
16 statistically significant for either acute or delayed nausea / severe nausea.
17
18
19
20
21
22
23
24

25 INSERT TABLE 2 HERE
26
27

28 **Discussion**

29
30

31 The findings of the present study clearly indicate that chemotherapy-induced
32 nausea is associated with poor quality of life, nutritional status and psychological
33 distress, with clinically meaningful differences being observed in those
34 experiencing nausea compared to those who did not. This is the first study looking
35 exclusively at the impact of chemotherapy-related nausea (rather than combined
36 with vomiting as in past literature) on quality of life outcomes and one of the few
37 focusing on nutritional aspects of nausea as a single entity. There was low
38 incidence of acute vomiting after cycle 1 (14.6%) and cycle 2 (9.1%) of
39 chemotherapy, with 5-8% of patients reporting delayed vomiting, reflecting that
40 management of chemotherapy-induced vomiting has greatly improved over the
41 last decade. In contrast, the incidence of chemotherapy-induced nausea was very
42 high and severe nausea affected 20-30% of the patients over the two cycles of
43 chemotherapy. However, these results should be viewed with some caution, as
44 international antiemetic recommendations for anthracycline-based chemotherapy
45 propose a three-drug combination, including aprepitant [35] and our sample
46 received only two of them Plus metoclopramide); hence the incidence of nausea
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 reported here may be over-estimated, although it reflects widely used routine
2 clinical practices [25].
3
4

5 Although 20-30% of patients with cancer are known to experience psychological
6 distress [21], this sample showed a low incidence of clinical depression (1.9-7.3%)
7 but higher scores of clinical anxiety (14.1-19.8%). The psychological impact of
8 chemotherapy is well recognised and associated with considerable uncertainty for
9 patients [14]. However, clinical experience suggests patients' levels of anxiety
10 decreases after the start of chemotherapy, which is reflected in these results. The
11 borderline significant link with nausea highlights the distressing nature of this
12 symptom.
13
14
15
16
17
18
19
20

21 Generally, while serum albumin levels were not significantly linked with nausea,
22 patients' weight and BMI had a trend of decreasing from baseline to cycles 1 and 2
23 reflecting nutritional changes. Furthermore, the PG-SGA scores indicated
24 deteriorating nutritional status after the start of chemotherapy, with 25% showing
25 scores indicative of malnutrition after the first cycle of chemotherapy. This
26 suggests that current standard measurements of nutritional status (weight and
27 serum albumin) are inadequate indicators of malnutrition, and that clinicians
28 should look to adopt alternative strategies to assess patients, such as the PG-
29 SGA. Body composition might change without reducing BMI and reflects the
30 influence of nausea in nutritional status. Hence, BMI decreases and low serum
31 albumin may not be sensitive indicators of malnutrition in this population which
32 often experiences fluid retention during chemotherapy. Indeed, weight gain is
33 common in women with breast cancer receiving chemotherapy [10,28,39]. Fat
34 mass and lean body mass, components of body composition, are also difficult to
35 measure in a clinical setting. The options are anthropometry using skin callipers
36 measured at 4 sites and placed into an equation to calculate fat and fat free mass
37 or bioelectrical impedance monitoring. The former is subject to a high degree of
38 inter-rater reliability and requires some experience and training in the technique to
39 produce valid results. The latter is subject to alterations in fluid balance, which
40 would occur in chemotherapy patients. Therefore its usefulness would be of limited
41 value and subject to strong opposition in relation to validity in oncology patients
42 undergoing treatment. The Patient-generated SGA derived from the professional
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

tool produced by Desksy [11] does incorporate a subjective assessment of body composition. This evaluates fat stores and muscle from pre-determined criteria that outline the visual assessment in some detail. The Patient-generated SGA is also validated against numerous nutritional status parameters, it has been used widely in oncology patients [3,32] and has been shown to be significantly associated with change in quality of life and change in lean body mass [4].

Generally, the nutritional impact of chemotherapy is poorly addressed by clinicians [37,40]; assessments are usually based on CTC toxicity scores that do not take into account important details of nutrition. If patients' weight generally remains stable and patients recover in between cycles of chemotherapy, current assessment approaches may fail to detect nutritional problems. This may be compounded by the lack of dieticians available for patients attending outpatient appointments. It is imperative that proactive nutritional assessment is integrated in clinical practice. Good nutrition practices may contribute to the relief of symptoms associated with nutrition (nausea, vomiting, diarrhoea, dysphagia, change in taste and smell, etc) and potentially improve quality of life, while poor nutrition may increase the incidence and severity of treatment side effects [1].

We have found that severe nausea has significant associations with nutritional status in this study, as judged by the proportions presented in table 3 and the statistically significant scores of the PG-SGA presented in table 4. As nausea is a key symptom whose presence (particularly severe acute and delayed nausea) has been demonstrated to be associated with decreased physical functioning and nutritional status, it is important to develop further research introducing nutritional interventions for the management of chemotherapy-related nausea. These may include education about what foods to eat/not to eat while nauseous, frequency, quantity, etc. This approach may be particularly appropriate, as the management of nausea is not satisfactory with current antiemetics.

Findings from this study showed that patients' QOL was sensitive to change within a short period of time, with some deterioration evident soon after the start of chemotherapy. Areas of impairments in quality of life such as physical functioning in those experiencing severe acute and delayed nausea confirm findings from past studies [30,36], suggesting that little has changed in this area over the past 10-15

1 years. While we have followed a conservative approach to statistical significance to
2 decrease error in the results due to using multiple tests, there are indications that
3 nausea, particularly severe nausea, has an association with functional, emotional
4 and overall quality of life. This would need to be further ascertained in future
5 research with the use of a larger and more heterogeneous sample. However,
6 studies have established minimally important clinical change in relation to the
7 FACT-G scale and in a sample with breast cancer patients this was around 5-6
8 points [13,18]. According to this estimate of clinically meaningful and important
9 change, our results of the impact of nausea on patients' quality of life are clinically
10 important. However, such estimates are not available for the nutritional and
11 psychological distress outcome measures we have used.
12
13
14
15
16
17
18
19
20
21

22 Considering these data of unacceptably high levels of nausea and its clear
23 association with quality of life and nutritional status, more interventions should be
24 directed to the management of nausea. Evidence exists about the effects of
25 acupuncture/acupressure, relaxation techniques, hypnosis, guided imagery,
26 exercise, cognitive distraction, systematic desensitisation or behavioural
27 approaches [20,24,33,34] in managing nausea. Careful clinical management is
28 also required to improve patients' concordance with treatment and prevent
29 anticipatory nausea/vomiting developing in subsequent chemotherapy cycles.
30
31
32
33
34
35
36
37

38 Appropriate symptom management begins with symptom assessment and an
39 accurate understanding of the prevalence and severity of symptoms experienced
40 by patients. A number of instruments are available for the clinician to use, and
41 critical reviews have provided evidence of their validity and usefulness, which
42 clinicians could consult in order to select the most appropriate assessment tool for
43 their practice [6, 41]. Furthermore, nausea can also be assessed as part of routine
44 clinical practice when patients complete instruments that rapidly assess multiple
45 symptoms.
46
47
48
49
50
51
52
53

54 This sample was mainly patients with breast cancer receiving anthracycline-based
55 chemotherapy; therefore generalisability would be increased by using a more
56 heterogeneous sample. As the sample is gender and breast cancer skewed, results
57 may not be generalisable to males and other cancers. A larger sample may provide
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

stronger evidence of the links between nausea and domains of quality of life, although our results are consistent with past literature using larger sample sizes but examining nausea and vomiting combined. The results are somewhat tentative, as the design of the study does not allow for a cause-and-effect outcome to be established, and the consequences of chemotherapy itself, stressors and other symptoms and comorbidities may also impact on the outcomes assessed in this study. Future research could focus on testing non-pharmacological interventions as adjunct therapies to antiemetics, improve the utility of available antiemetics by following evidence-based clinical guidelines and understand more fully the symptom of nausea from a pathophysiological perspective [29].

Conclusions

Our study confirms that nausea still remains a key quality of life problem for patients, it is distressing and leads to significant nutritional changes. This is clearly an important clinical problem, which this study assessed on its own rather than combined with the symptom of vomiting, recognising that nausea is related to vomiting but a different entity. The impact of nausea on nutritional status and quality of life is evident within days of chemotherapy administration. Given that most patients are treated in the outpatient setting, this requires careful assessment and clinical management. Specific nutritional interventions and use of a combination of pharmacological and non-pharmacological therapies for the management of this distressing symptom are necessary.

References

1. American Cancer Society: Nutrition for the Person with Cancer: A Guide for Patients and Families. Atlanta, Ga: American Cancer Society, Inc., 2000.
2. ASHP (1999) Therapeutic guidelines on the pharmacologic management of nausea and vomiting in adult and pediatric patients receiving chemotherapy or radiation therapy or undergoing surgery. *Am J Health Syst Pharm* 56:729-764.
3. Bauer J CS, and Ferguson M: . Use of the scored Patient-Generated Subjective Global Assessment (PG-SGA) as a nutrition assessment *Eur J Clin Nutr* 2002;56:779–85.
4. Bauer JD CS. Nutrition intervention improves outcomes in patients with cancer cachexia receiving chemotherapy—a pilot study. *Support Care Cancer* 2005; 13:270-74.
5. Bloechl-Daum B, Deuson RR, Mavros P, Hansen M, Herrstedt J. Delayed nausea and vomiting continue to reduce patients' quality of life after highly and moderately emetogenic chemotherapy despite antiemetic treatment. *J Clin Oncol* 2006; 24: 4472-4478.
6. Brearley SG, Clements CV, Molassiotis A. A review of patient self-report tools for chemotherapy-induced nausea and vomiting. *Support Care Cancer*. 2008; 16: 1213-1229.
7. Cella D, Tulsky D, Gray G, Sarafian B, Linn E, Bonomi A et al (1993) The Functional Assessment of Cancer Therapy Scale : development and validation of the general measure. *J Clin Oncol* 11 :570-579.
8. de Boer-Dennert M, de Wit R, Schmitz PI, Djontono J, v Beurden V, Stoter G & Verweij J (1997) Patient perceptions of the side-effects of chemotherapy: the influence of 5HT3 antagonists. *Br J Cancer* 76:1055-1061.
9. Decker GM, DeMeyer ES, & Kisko DL (2006) Measuring the maintenance of daily life activities using the functional living index-emesis (FLIE) in patients receiving moderately emetogenic chemotherapy. *J Support Oncol* 4:35-41
10. Demark-Wahnefried W, Peterson BL, Winer EP, Marks L, Aziz N, Marcom PK, Blackwell K, Rimer BK. Changes in weight, body composition, and factors influencing energy balance among premenopausal breast cancer patients receiving adjuvant chemotherapy. *J Clin Oncol*. 2001;19(9):2381-9.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
11. Detsky AS, Baker JP, Mendelson RA (1984). Evaluating the accuracy of nutritional assessment techniques applied to hospitalized patients: Methodology and comparisons. *JPEN J Parenter Enteral Nutr.* ;8(2):153-9.
12. Detsky AS, McLaughlin JR, Johnstone N, Whittaker S, Mendelson RA, Jeejeebhoy KN (1987) What is subjective global assessment of nutritional status. *JPEN J Parenter Enteral Nutr* 11:8-13.
13. Eton DT, Cella D, Yost KJ, et al. (2004). A combination of distribution- and anchor-based approaches determined minimally important differences (MIDs) for four endpoints in a breast cancer scale. *J Clin Epidemiol.*; 57(9):898-910.
14. Farrell C, Heaven C, Beaver K, Maguire P (2005). Identifying the concerns of women undergoing chemotherapy. *Patient Education and Counseling*, 56: 72-7.
15. Glaus A, Knipping C, Morant R, Bohme C, Lebert B, Beldermann F, Glawogger B, Ortega PF, Husler A & Deuson R (2004) Chemotherapy-induced nausea and vomiting in routine practice: a European perspective. *Support Care Cancer* 12:708-715.
17. 16. Gift AG, Jablonski A, Stommel M, Given CW (2004) Symptom clusters on elderly patients with lung cancer. *Oncol Nurs Forum* 31 :203-212. Grunberg SM, Deuson RR, Mavros P, Geling O, Hansen M, Cruciani G, Daniele B, De Pouvourville G, Rubenstein EB & Daugaard G (2004) Incidence of chemotherapy-induced nausea and emesis after modern antiemetics. *Cancer* 100:2261–2268.
18. King MT, Stockler MR, Cella DF, et al. (2010). Meta-analysis provides evidence-based effect sizes for a cancer-specific quality-of-life questionnaire, the FACT-G. *J Clin Epidemiol.*; 63(3):270-281.
19. Liau CT, Chu NM, Liu HE, Deuson R, Chen JS (2005) Incidence of chemotherapy-induced nausea and vomiting in Taiwan: physicians' and nurses' estimation vs patient's reported outcomes. *Supp Care Cancer* 13:277-286.
20. Lotfi-Jam K, Carey M, Jefford M, Schofield P, Charleson C, Aranda S. Nonpharmacologic strategies for managing common chemotherapy adverse effects: a systematic review. *J Clin Oncol.* 2008; 26: 5618-5629
21. Maguire P (2000). Managing psychological morbidity in cancer patients. *European Journal of Cancer* 36: 556-58.


- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
22. Martin CG, Rubenstein EB, Elting LS, Kim YJ & Osoba D (2003) Measuring chemotherapy-induced nausea and emesis. Psychometric properties of a quality of life questionnaire. *Cancer* 98:645–655.
 23. Miller M, Kearney N (2004) chemotherapy-related nausea and vomiting- past reflections, present practice and future management. *Eur J Cancer Care* 13:71-81.
 24. Molassiotis A, Yung HP, Yam BM, Chan FY, Mok TS. The effectiveness of progressive muscle relaxation training in managing chemotherapy-induced nausea and vomiting in Chinese breast cancer patients: a randomised controlled trial. *Support Care Cancer*. 2002; 10: 237-246.
 25. Molassiotis A, Saunders M, Valle J, Wilson G, Lorigan P, Wardley A, Levine E, Cowan R, Loncaster J & Rittenberg C (2008) A Prospective observational study of chemotherapy-induced nausea and vomiting in routine practice in a UK cancer centre. *Support Care Cancer* 16: 201-208.
 26. Molassiotis A, Coventry PA, Stricker CT, Clements C, Eaby B, Velders L, Rittenberg C & Gralla RJ (2007) Validation and psychometric assessment of a short clinical scale to measure chemotherapy-induced nausea and vomiting: the MASCC Antiemesis Tool (MAT). *J Pain Symptom Manage* 34:148-159.
 27. Molassiotis A, Wengström Y, Kearney N. Symptom cluster patterns during the first year after diagnosis with cancer. *J Pain Symptom Manage*. 2010; 39: 847-858.
 28. Nissen MJ, Shapiro A, Swenson KK. Changes in weight and body composition in women receiving chemotherapy for breast cancer. *Clin Breast Cancer*. 2011; 11(1):52-60.
 29. Olver I, Molassiotis A, Aapro M, Herrstedt J, Grunberg S, Morrow G. Antiemetic research: future directions. *Support Care Cancer*. 2011; 19 Suppl 1: S49-55.
 30. Osoba D, Zee B, Warr D, Latreille J, Kaizer L, Pater J. Effect of postchemotherapy nausea and vomiting on health-related quality of life. The Quality of Life and Symptom Control Committees of the National Cancer Institute of Canada Clinical Trials Group. *Support Care Cancer* 1997; 5: 307-313.
 31. Ottery F (2000) Patient-generated subjective global assessment. In Polisena C (ed) *The clinical guide to oncology nutrition*, American Dietetic Association: Chicago, pp. 1-23.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
32. Read JA, Crockett N, Volker DH, MacLennan P, Choy STB, Beale P, et al. Nutritional Assessment in Cancer: Comparing the Mini-Nutritional Assessment (MNA) With the Scored Patient-Generated Subjective Global Assessment (PGSGA). *Nutrition and Cancer* 2005;53(1):51-56.
 33. Richardson J, Smith JE, McCall G, Richardson A, Pilkington K, Kirsch I. Hypnosis for nausea and vomiting in cancer chemotherapy: a systematic review of the research evidence. *Eur J Cancer Care* 2007; 16: 402-412.
 34. Roscoe JA, Morrow GR, Hickok JT, Bushunow P, Pierce HI, Flynn PJ, Kirshner JJ, Moore DF, Atkins JN. The efficacy of acupressure and acustimulation wrist bands for the relief of chemotherapy-induced nausea and vomiting. A University of Rochester Cancer Center Community Clinical Oncology Program multicenter study. *J Pain Symptom Manage*. 2003; 26: 731-742.
 35. Roila F, Herrstedt J, Aapro M, Gralla RJ, Einhorn LH, Ballatori E, Bria E, Clark-Snow RA, Espersen BT, Feyer P, Grunberg SM, Hesketh PJ, Jordan K, Kris MG, Maranzano E, Molassiotis A, Morrow G, Olver I, Rapoport BL, Rittenberg C, Saito M, Tonato M, Warr D; ESMO/MASCC Guidelines Working Group. Guideline update for MASCC and ESMO in the prevention of chemotherapy- and radiotherapy-induced nausea and vomiting: results of the Perugia consensus conference. *Ann Oncol*. 2010; 21 Suppl 5:v232-43.
 36. Rusthoven JJ, Osoba D, Butts CA, Yelle L, Findlay H, Grenville A. The impact of postchemotherapy nausea and vomiting on quality of life after moderately emetogenic chemotherapy. *Support Care Cancer* 1998; 6: 389-395.
 37. Shaw C. *Nutrition and cancer*. Blackwell Publishers, Oxford, UK, 2011.
 38. Smith HS (2005) A receptor-based paradigm of nausea and vomiting. *J Cancer Pain Symptom Pall* 1:11–23.
 39. Trédan O, Bajard A, Meunier A, Roux P, Fiorletta I, Gargi T, Bachelot T, Guastalla JP, Lallemand Y, Faure C, Pérol D, Bachmann P. Body weight change in women receiving adjuvant chemotherapy for breast cancer: a French prospective study. *Clin Nutr*. 2010;29(2):187-91.
 40. Tong H, Isenring E, Yates P. The prevalence of nutrition impact symptoms and their relationship to quality of life and clinical outcomes in medical oncology patients. *Support Care Cancer* 2009; 17: 83-90.
 41. Wood JM, Chapman K, Eilers J. Tools for assessing nausea, vomiting, and retching. *Cancer Nurs*. 2011; 34: E14-24.

42. Zigmund A S & Snaith R P. (1983) Hospital Anxiety and Depression Scale.
Acta Psych Scand **67**: 361-370.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2 **Figure 1:** MAT Acute and delayed nausea and vomiting (% of 'Yes' answers)
3 with Exact upper 95%CI bars. Cycle1=96 responses. Cycle2=88 acute, 87 delayed
4 responses.
5
6
7
8
9



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 2: Incidence of severe nausea (6-10 on the MAT item of nausea) (% of patients), with Exact upper 95%CI bars. Cycle1=96 responses. Cycle2=88 acute, 87 delayed responses.




Table 1: Nutritional status indicators: changes over time

	Median	Total	PG-SGA Score 0- 8 n (%)	PG-SGA Score \geq 9 n (%)
PG-SGA Baseline	1	101	91 (90.1)	10 (9.9)
PG-SGA Cycle 1	5	95	71 (74.7)	24 (25.3)
PG-SGA Cycle 2	4	87	65 (74.7)	22 (25.3)
	Median	Total	Median	Median
Weight at Baseline	72.0	96	72	72.1
Weight pre Cycle1	71.7	87	71.7	72.7
Weight pre Cycle2	72.7	42	72.8	64.5
BMI at Baseline	27.4	85	27.5	28.5
BMI pre Cycle1	27.1	74	26.7	27.5
BMI pre Cycle2	26.1	37	26.4	24.4
Albumin at Baseline	43	95	43	45
Albumin pre Cycle1	42	93	42.5	42
Albumin pre Cycle2	42	87	42	43

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 2: Mean scores of Quality of Life, nutritional status and psychological distress in relation to acute and delayed nausea (ANCOVA analysis) for Cycle 1 (n=96). Patients are analysed in four different binary groupings.

	Acute Nausea		Severe Acute Nausea		Delayed Nausea		Severe Delayed Nausea	
	No	yes	<6	>=6	No	yes	<6	>=6
<i>FACT-G: Physical</i>	20.5	17.8	20.2	16.1	20.7	18.3	20.1	15.5
<i>FACT-G: Social</i>	24.0	24.3	24.2	24.0	23.6	24.4	24.1	24.3
<i>FACT-G: Emotional</i>	19.0	17.0	18.4	16.6	18.8	17.6	18.1	17.4
<i>FACT-G: Functional</i>	17.4	15.8	17.2	15.0	17.1	16.3	17.2	14.2
<i>FACT-G: Total</i>	80.9	75.0	80.1	71.8	80.2	76.8	79.5	71.8
<i>PG-SGA</i>	5.0	6.5	5.0	8.0	4.3	6.4	5.1	8.0
<i>HADS: Anxiety</i>	4.8	7.2	5.7	7.1	5.5	6.3	5.8	7.0
<i>HADS: Depression</i>	4.0	5.2	4.5	5.2	4.7	4.6	4.6	5.0
<i>HADS: Total</i>	8.8	12.3	10.1	12.3	10.2	11.0	10.4	12.0

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

	Difference in mean	P value	Difference in mean	P value	Difference in mean	P value	Difference in mean	P value
FACT-G: Physical	2.7	0.052	4.1	0.025*	2.4	0.397	4.6	0.217
FACT-G: Social	-0.3	0.518	0.2	0.248	-0.8	0.185	-0.2	0.038
FACT-G: Emotional	2	0.907	1.8	0.193	1.2	0.754	0.7	0.595
FACT-G: Functional	1.6	0.675	2.2	0.412	0.8	0.682	3	0.409
FACT-G: Total	5.9†	0.603	8.3†	0.566	3.4	0.907	7.7†	0.628
PG-SGA	-1.5	0.05	-3	0.003**	-2.1	0.051	-2.9	0.017*
HADS: Anxiety	-2.4	0.494	-1.4	0.978	-0.8	0.961	-1.2	0.638
HADS: Depression	-1.2	0.579	-0.7	0.794	0.1	0.405	-0.4	0.554
HADS: Total	-3.5	0.592	-2.2	0.74	-0.8	0.547	-1.6	0.429

** Statistically significant (<5%/4=0.0125)

* Borderline statistically significant (<5%)

† FACT-G scale: Clinically meaningful change, although not statistically significant