

American Journal of  
Transplantation

**Targeting PI3K $\delta$  Function For Amelioration of Murine  
Chronic Graft-Versus-Host Disease**

Journal:	<i>American Journal of Transplantation</i>
Manuscript ID	AJT-B-18-01422.R2
Wiley - Manuscript type:	B - Brief Communication
Date Submitted by the Author:	24-Jan-2019
Complete List of Authors:	<p>Paz, Katelyn; Masonic Cancer Center, Department of Paediatrics          Flynn, Ryan; University of Minnesota, Pediatric Blood and Marrow          Transplantation          Du, Jing; University of Minnesota Twin Cities,          TANNHEIMER, Stacey; Gilead Sciences, Inc.          Johnson, Amy; Gilead Sciences, Inc.          Dong, Shuai; Ohio State University, Division of Pharmeceutics and          Pharmaceutical Chemistry          Start, Anne-Katrien; University of Cambridge, Department of Pathology          Okkenhaug, Klaus; University of Cambridge, Department of Pathology          Panoskaltzis-Mortari, Angela; University of Minnesota, Pediatrics          Sage, Peter; Harvard Medical School, Department of Microbiology and          Immunobiology          Sharpe, Arlene; Harvard Medical School, Microbiology and          Immunobiology          Luznik, Leo; Johns Hopkins, Oncology          Ritz, Jerome; Harvard Medical School          Soiffer, Robert J.; Dana-Farber Cancer Institute Department of Medical          Oncology          cutler, Corey; Dana-Farber Cancer Institute Department of Medical          Oncology          Koreth, John; Harvard Medical School          Antin, Joseph; Dana-Farber Cancer Institute Department of Medical          Oncology          Miklos, David; Stanford University, School of Medicine          MacDonald, Kelli; University of Queensland, Department of          Immunology          Hill, Geoffrey; QIMR Berghofer Medical Research Institute, Immunology          Maillard, Ivan; University of Pennsylvania, Perelman School of Medicine          Serody, Jonathan; University of North Carolina School of Medicine,          Lineberger Cancer Center          Murphy, William; University of California Davis Health System,          Munn, David; Augusta University, Medical College of Georgia          Feser, Colby; University of Minnesota, Pediatric Blood and Marrow          Transplantation          Zaiken, Michael; Masonic Cancer Center, Department of Paediatrics          Vanhaesebroeck, Bart; University College London Medical School          Turka, Laurence; Massachusetts General Hospital, Transplantation          Biology Research Center          Byrd, John; Ohio State University, Division of Pharmeceutics and</p>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

	Pharmaceutical Chemistry Blazar, Bruce; University of Minnesota, Cancer Center and Department of Pediatrics, Division of BMT
Keywords - Scope:	basic (laboratory) research / science
Keywords - Discipline:	immunobiology
Keywords - Focus:	graft-versus-host disease (GVHD), immunosuppressant - other



1  
2  
3 **Targeting PI3K $\delta$  Function For Amelioration of**  
4  
5 **Murine Chronic Graft-Versus-Host Disease**  
6

7 Katelyn Paz<sup>1</sup>, Ryan Flynn<sup>1</sup>, Jing Du<sup>1</sup>, Stacey Tannheimer<sup>2</sup>, Amy J. Johnson<sup>3</sup>, Shuai Dong<sup>4</sup>, Anne-  
8 Katrien Stark<sup>5</sup>, Klaus Okkenhaug<sup>5</sup>, Angela Panoskaltzis-Mortari<sup>1</sup>, Peter T. Sage<sup>6</sup>, Arlene H. Sharpe<sup>7,8,9</sup>,  
9 Leo Luznik<sup>10</sup>, Jerome Ritz<sup>11</sup>, Robert J. Soiffer<sup>11</sup>, Corey S. Cutler<sup>11</sup>, John Koreth<sup>11</sup>, Joseph H. Antin<sup>11</sup>,  
10 David B. Miklos<sup>12</sup>, Kelli P. MacDonald<sup>13</sup>, Geoffrey R. Hill<sup>13</sup>, Ivan Maillard<sup>14</sup>, Jonathan S. Serody<sup>15</sup>,  
11 William J. Murphy<sup>16</sup>, David H. Munn<sup>17</sup>, Colby Feser<sup>1</sup>, Michael Zaiken<sup>1</sup>, Bart Vanhaesebroeck<sup>18</sup>,  
12 Laurence A. Turka<sup>19</sup>, John C. Byrd<sup>3</sup>, Bruce R. Blazar<sup>1</sup>  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

- 24 1. Division of Blood and Marrow Transplantation, Department of Pediatrics, University of  
25 Minnesota, Minneapolis, Minnesota, USA  
26  
27 2. Gilead Sciences, Inc., Foster City, California, USA  
28  
29 3. Division of Hematology, Department of Internal Medicine and Comprehensive Cancer Center,  
30 and Division of Medicinal Chemistry, College of Pharmacy, The Ohio State University, Columbus,  
31 Ohio, USA  
32  
33 4. Division of Pharmaceutics and Pharmaceutical Chemistry, College of Pharmacy. The Ohio  
34 State University, Columbus, Ohio, USA  
35  
36 5. Department of Pathology, University of Cambridge, Cambridge, UK  
37  
38 6. Transplantation Research Center, Renal Division, Brigham and Women's Hospital, Harvard  
39 Medical School, Boston, Massachusetts, USA  
40  
41 7. Department of Microbiology and Immunobiology, Harvard Medical School, Boston,  
42 Massachusetts, USA  
43  
44 8. Evergrande Center for Immunologic Diseases, Harvard Medical School and Brigham and  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 Women's Hospital, Boston, Massachusetts, USA  
4  
5 9. Department of Pathology, Brigham and Women's Hospital, Boston, Massachusetts, USA  
6  
7 10. Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins, The Johns Hopkins  
8  
9 University School of Medicine, Baltimore, Maryland, USA  
10  
11 11. Stem Cell/Bone Marrow Transplantation Program, Division of Hematologic Malignancy, Dana-  
12  
13 Farber Cancer Institute, Harvard Medical School, Boston, Massachusetts, USA  
14  
15 12. Stanford Cancer Center, Stanford University School of Medicine, Stanford, CA;  
16  
17 13. Department of Immunology, QIMR Berghofer Medical Research Institute and School of  
18  
19 Medicine, University of Queensland, Brisbane, Australia  
20  
21 14. Division of Hematology-Oncology, Department of Medicine, University of Pennsylvania,  
22  
23 Perelman School of Medicine, Philadelphia, Pennsylvania, USA.  
24  
25 15. Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC, USA  
26  
27 16. Departments of Dermatology and Internal Medicine, Division of Hematology and Oncology,  
28  
29 University of California Davis School of Medicine, Sacramento, CA, USA  
30  
31 17. Georgia Cancer Center and Department of Pediatrics, Medical College of Georgia, Augusta  
32  
33 University, Augusta, Georgia  
34  
35 18. UCL Cancer Institute, University College London, London, UK  
36  
37 19. Center for Transplantation Sciences, Department of Surgery, Massachusetts General Hospital,  
38  
39 Boston, Massachusetts, USA  
40  
41  
42  
43  
44  
45  
46  
47  
48

49 Correspondence: Bruce Blazar, [blaza001@umn.edu](mailto:blaza001@umn.edu)  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Abbreviations list**

BID	Twice daily
BO	Bronchiolitis obliterans
cGVHD	chronic graft versus host disease
EAE	Experimental autoimmune encephalitis
GC	Germinal center
GVL	Graft versus Leukemia
Ig	Immunoglobulin
KO	Knock out
IL	Interleukin
OCT	Optimum cutting temperature
PC	Plasma cell
PFT	Pulmonary function test
PI3K	Phosphoinositide-3-kinase
SD	Standard deviation
Teff	Teffector cells
TCD	T cell depleted
TGF- $\beta$	Transforming growth factor- beta
Th	T helper
Tfh	T follicular helper
Treg	T regulatory
Tfr	T follicular regulatory
WT	Wild type

**Abstract**

Chronic graft-versus-host disease is a leading cause of morbidity and mortality following allotransplant. Activated donor effector T-cells can differentiate into pathogenic T helper (Th)-17 cells and germinal center -promoting Tfollicular helper cells, resulting in cGVHD. Phosphoinositide-3-kinase- $\delta$ , a lipid kinase, is critical for activated T-cell survival, proliferation, differentiation, and metabolism. We demonstrate PI3K $\delta$  activity in donor T-cells that become Tfh is required for cGVHD in a non-sclerodermatous multi-organ system disease model that includes bronchiolitis obliterans, dependent upon GC B-cells, Tfh, and counterbalanced by Tfollicular regulatory cells, each requiring PI3K $\delta$  signaling for function and survival. Although B-cells rely on PI3K $\delta$  pathway signaling and GC formation is disrupted resulting in a substantial decrease in Ig production, PI3K $\delta$  kinase-dead mutant donor bone marrow derived GC B-cells still supported BO cGVHD generation. A PI3K $\delta$ -specific inhibitor, compound GS-649443 that has superior potency to idelalisib while maintaining selectivity, reduced cGVHD in mice with active disease. In a Th1-dependent and Th17-associated scleroderma model, GS-649443 effectively treated mice with active cGVHD. These data provide a foundation for clinical trials of FDA-approved PI3K $\delta$  inhibitors for cGVHD therapy in patients.

## Introduction

Graft-versus-host disease (GVHD) is a major obstacle for allogeneic hematopoietic stem cell transplant patients, greatly impacting their quality of life. GVHD is a primary cause of mortality, second only to primary disease relapse. Chronic GVHD (cGVHD) is a leading cause of morbidity, occurring in 20-70% of aHSCT patients<sup>1,2</sup>. CGVHD clinical presentations are varied and virtually every organ in the body can be affected; amongst the more severe outcomes are cGVHD of the lung, manifesting as bronchiolitis obliterans (BO) and skin as scleroderma<sup>3</sup>. Due to this broad and varied pathogenesis, multiple murine models have been developed to recapitulate a larger portion of the disease spectrum<sup>4-6</sup>. A common feature among models and in patients is the driving role of chronically stimulated alloreactive T cells in disease pathogenesis<sup>3,7</sup>. Activated alloreactive donor CD4<sup>+</sup> T-cells differentiate into T follicular helper (Tfh) and IL-17-producing helper T-cells (Th17s) that have known pathogenic roles in cGVHD<sup>4,8-10</sup>.

Tfh cells are a specialized CD4<sup>+</sup> Th cell subset that provide essential signals to support germinal center (GC) B-cell, memory B-cell or antibody-producing plasma cell (PC) development<sup>11-13</sup>. A subpopulation of T regulatory (Treg), T follicular regulatory (Tfr) cells, suppress Tfh and GC B-cells to regulate the GC reaction<sup>14</sup>. Immunoglobulin (Ig) produced by PCs and deposited in target tissues, such as the lung, liver, and colon contributes to organ damage in BO cGVHD and skin in the scleroderma model<sup>15</sup>. We previously reported that Tfh and GC B-cells are required for the development of murine BO cGVHD, a model that recapitulates many aspects of human cGVHD pathology, with the predominant exception of scleroderma<sup>15-19</sup>. In this BO cGVHD model, weight loss and mortality are low (around or less than 20%). Th17 cells, a source of the pro-inflammatory cytokine IL-17 that contributes to autoimmunity<sup>20</sup>, are also involved in BO as well as our sclerodermatous model of cGVHD<sup>21,22</sup>.

1  
2  
3  
4  
5 Phosphoinositide-3-kinases (PI3Ks) are a family of lipid kinases that that regulate numerous signaling  
6 cascades via the phosphorylation of 3-hydroxyl group of phosphatidylinositol lipid substrates<sup>23</sup>.  
7  
8 Structural and substrate preferences divide the PI3Ks into three classes (I, II, III)<sup>24</sup>. Within the class I  
9  
10 PI3Ks, present in all cell types, there are several isoforms, each comprised of regulatory and catalytic  
11  
12 subunit heterodimers<sup>23</sup>. The p110 $\delta$  catalytic subunit, referred to as PI3K $\delta$ , is an isoform preferentially  
13  
14 expressed in leukocytes, regulating immune cell signalling<sup>25,26</sup>. PI3K $\delta$  is activated upon T-cell receptor  
15  
16 engagement, CD28 costimulation, and cytokine receptor signaling to sustain an activated Teff  
17  
18 phenotype and promote the function of these cells, including regulation of survival, cell cycle  
19  
20 progression, differentiation and metabolism<sup>27-30</sup>. Loss of PI3K $\delta$  diminishes Teffector (Teff) activity<sup>31,32</sup>.  
21  
22 Relevant to our models of cGVHD, PI3K $\delta$  signaling has been found to be necessary for both murine  
23  
24 and human IL-17 production<sup>32-34</sup>. Recent work has demonstrated that PI3K $\delta$  mutant T-cells have  
25  
26 impaired alloimmune activity and that PI3K $\delta$  inhibition was able to effectively suppress alloreactive  
27  
28 Teffs to prevent solid organ heart transplant rejection<sup>35</sup>. In non-chronic models of GVHD, PI3K $\delta$   
29  
30 inhibition ameliorated lethality and reduced severity of clinical signs and organ damage<sup>36,37</sup>.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40 Similar to its role in immune cells, PI3K signaling controls proliferation, survival and metabolism of  
41  
42 cancer cells. Certain hematological malignancies have been found to have upregulated PI3K $\delta$   
43  
44 activity<sup>38,39</sup>. Idelalisib is a PI3K $\delta$  specific inhibitor that has been approved to treat hematological  
45  
46 malignancies, such as chronic lymphocytic leukemia, follicular lymphoma (that can be of GC B- or T-  
47  
48 cell origin) and small lymphocytic lymphoma<sup>40,41</sup>. While demonstrating therapeutic benefit, there are  
49  
50 also concerning toxicities associated with Idelalisib, including hepatotoxicity, diarrhea/colitis,  
51  
52 pneumonitis and intestinal perforation. Due to these off target effects, efforts are being made to develop  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 new drugs. One such compound utilized here is GS-649443, a PI3K $\delta$  isoform-specific inhibitor that has  
4  
5 demonstrated superior potency to idelalisib while maintaining selectivity<sup>42,43</sup>. In vitro and in vivo  
6  
7 studies demonstrated that this inhibitor reduces inflammatory cytokines, including IFN $\gamma$  and IL-17<sup>43,44</sup>.  
8  
9

10  
11  
12 The role of PI3K $\delta$  in the pathophysiology of cGVHD is unknown and deserves investigation in order to  
13  
14 develop new therapeutics to treat steroid-resistant or refractory cGVHD. In this study, we sought to  
15  
16 determine the requirement of PI3K $\delta$  function in cGVHD pathogenesis. We show that donor T-cells  
17  
18 deficient for PI3K $\delta$  activity are unable to induce cGVHD. Further, we demonstrate that the PI3K $\delta$   
19  
20 specific inhibitor, GS-649443, used for treatment of ongoing cGVHD, diminished the GC reaction and  
21  
22 antibody production in BO cGVHD. GS-649443 was also efficacious in sclerodermatous cGVHD  
23  
24 model, reducing pro-inflammatory IL-17 production. Together, these results provide basic mechanistic  
25  
26 insights regarding cGVHD pathophysiology and pre-clinical support for testing of PI3K $\delta$  inhibitors as a  
27  
28 therapeutic strategy for steroid-refractory or resistant cGVHD.  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Materials and Methods

### Mice

C57Bl/6 (B6, H2<sup>b</sup>) and Balb/c (H2<sup>d</sup>) mice were purchased from the National Cancer Institute. B10.BR (H2<sup>k</sup>) and B10.D2 (H2<sup>d</sup>) mice were purchased from Jackson Laboratory. Mice were housed in a specific-pathogen-free facility used with the approval of the University of Minnesota's animal care committee. To explore the effects of PI3K $\delta$  loss in donor cells in cGVHD, we used bone marrow (BM) and/or splenocytes from catalytically inactive p110 $\delta$ <sup>D910A/D910A</sup> (further referred to as p110 $\delta$ <sup>D910</sup>) homozygous mutant<sup>45</sup> and p110 $\delta$ <sup>D910A/WT</sup> (wildtype) heterozygous mutant mice, shipped overnight from Drs. Amy Johnson, Klaus Okkenhaug, Anne-Katrien Stark, and Bart Vanhaesebroeck.

### Bone Marrow Transplantation

For the BO cGVHD, B10.BR recipients were conditioned with cyclophosphamide (Sigma St. Louis, MO) 120mg/kg/day intraperitoneally, on days -3 and -2, and TBI 8.3 Gy, day -1. Recipients then received 10 x 10<sup>6</sup> B6 T-cell-depleted (TCD) BM only or with 7.5 x 10<sup>4</sup> purified splenic T-cells (cGVHD). For the B10.D2→Balb/c scleroderma model, Balb/c recipients were conditioned with TBI, 7 Gy, day -1 and then received 10 x 10<sup>6</sup> B10.D2 TCD BM only or with 1.8 x 10<sup>6</sup> CD4 and 0.9 x 10<sup>6</sup> CD8 T-cells on day 0<sup>22,46,47</sup>. Mice were monitored daily for survival and weighed twice weekly. In the scleroderma model, mice were assessed twice weekly for clinical and cutaneous GVHD, as previously described<sup>48</sup>.

### Pulmonary Function Tests

Pulmonary function tests (PFTs) were performed as previously described<sup>49</sup>. Briefly, mice were anesthetized with Nembutal, intubated and ventilated using the Flexivent system (Scireq Montreal,

1  
2  
3 QC). Pulmonary resistance, elastance and compliance were reported using Flexivent software version  
4  
5 7. We observe that cGVHD controls have increased pulmonary resistance and elastance along with  
6  
7 decreased compliance as compared to BM only controls in our BO cGVHD model<sup>15</sup>.  
8  
9

### 12 **PI3K $\delta$ Inhibition**

14 GS-649443<sup>42</sup>, provided by Gilead, was delivered in a vehicle consisting of 10% Ethanol, 20%  
15  
16 cremophor EL and 70% normal saline. Mice were given GS-649443 (10mg/kg) twice daily (BID) by  
17  
18 oral gavage from days 28-56 (BO model) or days 21-50 (scleroderma model). Mice in the vehicle  
19  
20 control group were treated with the same volume of vehicle.  
21  
22  
23  
24  
25

### 26 **Histopathology and Immunostaining**

28 Tissue sections were embedded in Optimal Cutting Temperature (OCT) compound, snap-frozen in  
29  
30 liquid nitrogen and stored at -80°C. Lungs were inflated by 75% OCT before harvest and freezing. For  
31  
32 Trichrome staining, 6- $\mu$ m cryosections were fixed overnight in Bouin's solution and stained with  
33  
34 Masson's Trichrome staining kit (Sigma HT15). Collagen deposition was quantified as a ratio of blue  
35  
36 area to total area using ImageJ. For Histopathology, acetone-fixed 6- $\mu$ m cryosections were hemotoxylin  
37  
38 and eosin stained and evaluated<sup>50</sup> without knowledge of treatment by APM. For immunoglobulin  
39  
40 deposition immunostaining, acetone-fixed 6 $\mu$ m cryosections were stained with goat anti-mouse IgG  
41  
42 (BD55401). Confocal images were acquired on Olympus Confocal Laser Scanning Microscope at 20X  
43  
44 and quantified by ImageJ.  
45  
46  
47  
48  
49  
50  
51

### 52 **Statistical Analysis**

GraphPad Prism 7 was used to conduct statistical analysis. One-way ANOVA with Bonferroni correction and Student's t-test were used for statistical analysis as indicated. Error bars indicate mean  $\pm$  standard deviation (SD). Significance: \* $P < .05$ ; \*\* $P < .01$ ; \*\*\* $P < .001$ ; \*\*\*\* $P < .0001$ .

## Results

### Fully intact donor T-cell PI3K $\delta$ activity is essential for BO cGVHD generation

The prominent contribution of PI3K $\delta$  activity to T-cell survival and function prompted us to determine whether donor T-cells with decreased or absent PI3K $\delta$  kinase activity would fail to cause cGVHD in the BO model. T cells from p110 $\delta^{D910A/wt}$  mice that have a knock-in mutation in one allele leading to heterozygote levels of catalytically inactive, mutant PI3K $\delta$  were given to a cohort of mice and compared to BM only and cGVHD controls. Mice receiving WT BM and either heterozygous p110 $\delta^{D910A/wt}$  or WT T-cells had  $\geq 90\%$  survival and  $\leq 5\%$  weight loss compared to day 0 body weights (not shown). BO cGVHD pulmonary dysfunction was comparable to WT T cell controls (Figure S1).

Next, we asked if PI3K $\delta$  activity in donor BM was required for cGVHD. Homozygous p110 $\delta^{D910A}$  BM with WT T-cells still resulted in pulmonary dysfunction consistent with cGVHD (Figure 1A). As compared to cGVHD only controls, mice receiving p110 $\delta^{D910A}$  BM with WT T-cells had significantly lower Treg and Tfr frequencies (Figure 1B-C). Tfh frequencies in mice that received p110 $\delta^{D910A}$  BM with WT T-cells were reduced from that of the cGVHD but still increased from their BM only control. An unfavorable Tfr:Tfh ratio, similar to that of the cGVHD control (Figure 1D-E), was observed. Since the magnitude of antibody responses, that originate in the GC, can be functionally predicted by the Tfr/Tfh ratio in a wide range of diseases in both mice and humans<sup>14</sup>, the low Tfr:Tfh ratio associated with an increased GC B-cell frequency (Figure 1F) was anticipated. Lung pathology scores correlated

1  
2  
3 with pulmonary function tests, with WT BM only compared to WT BM plus supplemental WT T-cells  
4 (0.1 ± 0.1 vs 2 ± 0.1581, p= <0.001) and p110δ<sup>D910A</sup> BM compared to p110δ<sup>D910A</sup> BM WT T-cells (0.2  
5  
6 ± 0.1225 vs 0.8 ± 0.255, p= 0.067)(data not shown). Whereas the statistical difference between the first  
7  
8 two groups was significant, statistical comparison in the histopathology scores between the recipients  
9  
10 receiving p110δ<sup>D910A</sup> BM only reached a statistical trend. These latter data suggest either a modest  
11  
12 effect of the KO BM on altering cGVHD severity or sample size limitations. Infusion of p110δ<sup>D910A/wt</sup>  
13  
14 T-cells with p110δ<sup>D910A/wt</sup> BM cells did not avert cGVHD pulmonary dysfunction (Figure S1).  
15  
16  
17  
18  
19

20  
21 Since haploinsufficient T-cells and BM cells did not have evidence of reduced cGVHD, we proceeded  
22  
23 to studies using homozygous p110δ<sup>D910A</sup> T-cells. We hypothesized that donor T-cells lacking all PI3Kδ  
24  
25 kinase activity would be inferior in inducing and sustaining cGVHD as compared to their WT  
26  
27 counterparts. We observed no significant changes in weight or survival between cGVHD controls and  
28  
29 mice that received p110δ<sup>D910A</sup> donor T-cells (Figure S2A-B). Mice that received p110δ<sup>D910A</sup> donor T-  
30  
31 cells did not develop pulmonary dysfunction associated with BO cGVHD (Figure 2A). Loss of PI3Kδ  
32  
33 activity resulted in a significant decrease in the frequency of splenic Tfh cells (Figure 2B) with  
34  
35 unaltered Treg (Figure S2C) and Tfr frequencies (Figure 2C). We observed an increased Tfr:Tfh ratio  
36  
37 (Figure 2D) and decreased GC B cell frequencies (Figure 2E) in mice that received p110δ<sup>D910A</sup> versus  
38  
39 WT donor T cells, consistent with studies demonstrating that the ratio of Tfr:Tfh controls the GC  
40  
41 reaction<sup>51</sup>. As expected by the significant improvement in pulmonary function parameters, recipients of  
42  
43 p110δ<sup>D910A</sup> donor T cells had significantly reduced histopathology scores (Figure 2F). T cells and BM  
44  
45 cells that had haplosufficient PI3Kδ expression did not provide adequate protection from cGVHD,  
46  
47 suggesting that high level PI3Kδ inhibition will be required to treat cGVHD in the clinic.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### **Therapeutic administration of GS-649443 ameliorates cGVHD in a non-sclerodermatous, BO model**

To validate if PI3K $\delta$  can be targeted as a novel therapeutic strategy, we tested the novel PI3K $\delta$  inhibitor, GS-649443, in our BO model of cGVHD. GS-649443 given at 10mg/kg, PO, BID beginning on day 28, the time of established cGVHD<sup>15</sup>, was well-tolerated as shown by weight and survival curves (Figure S3A, B). Treatment at a lower dose of 5mg/kg, PO, BID did not improve pulmonary function (Figure S4). Vehicle treatment alone had no significant effect on cGVHD outcome for any parameters tested. GS-649443 improved PFTs (Figure 3A), reduced the lung pathology associated with cGVHD (Figure 3B) and decreased Tfh (Figure 3C) frequencies. Both the Treg (Figure S3C) as well as Tfr (Figure 3D) frequencies were decreased by GS-649443 treatment. Although the Tfr:Tfh ratio was similar to that of the vehicle controls (Figure 3E), the GC B-cell frequency in GS-649443 treated mice was significantly decreased (Figure 3F). Together, these data point to either to a direct effect of GS-649443 on GC B-cells and/or reduction of Tfh frequency below threshold limits to cause a GC response.

### **Reduced Ig and collagen lung deposition in GS-649443-treated mice phenocopies findings in recipients given p110 $\delta$ <sup>D910A</sup> donor T-cells**

cGVHD has several autoimmune-like features, including but not limited to the deposition of antibodies and fibrosis of target organs, including the lung<sup>52</sup>. In accordance with improved PFTs and immune analysis, we demonstrated that lung IgG (Figure 4A) and collagen deposition (Figure 4B) was decreased in mice that received WT BM plus p110 $\delta$ <sup>D910A</sup> donor T-cells. Mice that received GS-649443 treatment also had reduced lung IgG and collagen deposition (Figure 4).

1  
2  
3 **Therapeutic administration of the PI3K $\delta$ -specific inhibitor GS-649443 ameliorates**  
4  
5 **sclerodermatous cGVHD**  
6

7  
8 A major clinical and histopathological manifestation absent from the multi-organ system BO cGVHD  
9  
10 model is scleroderma<sup>53</sup>. We utilized a multiple minor histocompatibility mismatch model  
11  
12 (B10.D2 $\rightarrow$ BALB/c) that presents with a cutaneous cGVHD and associated increased Th17 T cells and  
13  
14 systemic inflammatory response<sup>22</sup>. GS-649443 treatment significantly improved skin and clinical  
15  
16 scores of mice (Figure 5A-B). GS-649443 treatment decreased IL-17<sup>+</sup> T-cell frequency (Figure 5C),  
17  
18 characteristic of cGVHD in this model and IL-17<sup>+</sup>IFN $\gamma$ <sup>+</sup> double positive cells (Figure S5A), which can  
19  
20 contribute to autoimmunity<sup>22,54</sup>. IFN $\gamma$ <sup>+</sup> T-cells remained increased in mice treated with GS-649443  
21  
22 (Figure S5B) indicating potentially only a partial amelioration of disease. Nonetheless, decreased IL-  
23  
24 17-producing T-cells resulted in correspondingly lower, although not quite significant, IgG deposition  
25  
26 in the skin of scleroderma mice (Figure S5C).  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Discussion

PI3K $\delta$  is a key regulator of T<sub>H</sub>17 function, found here to be required for cGVHD development. Here, we have demonstrated that cGVHD generated in distinct murine models that simulate several, but not all, cGVHD manifestations, are dependent upon PI3K $\delta$  activity. We demonstrated that PI3K $\delta$  activity in donor T-cells but not B-cells is necessary to initiate and/or sustain the GC response critical for cGVHD in the BO model. We utilized the PI3K $\delta$  isoform-specific inhibitor GS-649443 to show that PI3K $\delta$  inhibition is effective in treating ongoing, established cGVHD in both the BO and sclerodermatous models. Overall, our data show that the PI3K $\delta$  signaling pathway is required to generate and maintain murine cGVHD in two, independent models with distinct pathophysiology and few overlapping cGVHD manifestations.

PI3K $\delta$  has roles in T<sub>H</sub>17s and other immune cell types, notably B-cells, Tregs and macrophages. Mice lacking functional PI3K $\delta$  exhibit B-cell defects. Such mice have fewer mature B-cells, reduced B-cell receptor-induced proliferation, decreased B-cell differentiation into antibody-producing cells, substantially reduced Ig production and disrupted GCs in response to antigen challenge<sup>45,55,56</sup>. Interestingly, p110 $\delta$ <sup>D910A</sup> BM with WT T-cells still induced pulmonary dysfunction that was significantly worse than their p110 $\delta$ <sup>D910A</sup> BM only counterpart. The magnitude of the GC B-cells was sufficient to induce pulmonary dysfunction. Because Tregs also reside in the BM, p110 $\delta$ <sup>D910A</sup> BM would produce Tregs or T<sub>H</sub>17s defective in suppressing T<sub>H</sub>17s that may have contributed to GC B-cell driven pulmonary dysfunction. Related to this possibility, PI3K $\delta$  signaling supports Treg development and function. We previously showed Tregs and T<sub>H</sub>17s are critical in controlling GC reactions and cGVHD<sup>57</sup> and that PI3K $\delta$  inhibition results in diminished *in vitro* and *in vivo* suppressor function and Treg survival<sup>31,35</sup>. Indeed, both the Treg and T<sub>H</sub>17 populations were decreased in mice that received



1  
2  
3 p110 $\delta$ <sup>D910A</sup> BM alone or with T-cells. The resulting overall unfavorable Tfr:Tfh ratio creates an  
4  
5 environment in the B-cell follicle permissive for an increased GC B-cell frequency<sup>14</sup>. In this study, we  
6  
7 observed decreased Tfr and GC B-cell frequencies associated with the therapeutic benefit of GS-  
8  
9 649443 treatment.  
10  
11  
12  
13

14 Macrophages are known to be key mediators of several types of inflammatory immune responses,  
15  
16 including those culminating in fibrosis. Indeed, macrophages were proven to be a source of  
17  
18 Transforming Growth Factor-beta (TFG- $\beta$ ), a mediator of tissue fibrosis<sup>21</sup>. Macrophage depletion<sup>21</sup>  
19  
20 or inhibition of macrophage migratory capacity<sup>58</sup> precluded the generation of cGVHD in both the BO  
21  
22 and scleroderma models. Optimal macrophage function has been associated with various PI3K  
23  
24 isoforms, including PI3K $\beta$ , PI3K $\delta$  and PI3K $\gamma$ <sup>40</sup> and in particular PI3K $\delta$  has been shown to inhibit  
25  
26 macrophage migration<sup>59</sup>. Although the improvement in cGVHD outcome with GS-649443 correlated  
27  
28 with a reduction in GC reaction, decreased macrophage migration may have contributed to disease  
29  
30 amelioration. Such may occur by a direct effect by PI3K $\delta$  inhibition on donor macrophage function or  
31  
32 indirectly inhibit macrophage migration as a consequence of low GCs, Ig deposition in cGVHD organs  
33  
34 and subsequently lower levels of macrophage chemoattractants. Further studies will be required to  
35  
36 determine how PI3K $\delta$  affects macrophage migration and function in the context of cGVHD. Additional  
37  
38 studies are needed to determine whether altered Tfr/Tfh, reduced Th17 cell as seen in the scleroderma  
39  
40 model, or impaired macrophage migration are the dominant or critical mechanism(s) of by which  
41  
42 PI3K $\delta$  inhibition ameliorates cGVHD BO.  
43  
44  
45  
46  
47  
48  
49  
50

51 Increased PI3K $\delta$  signaling has been found in autoimmune diseases<sup>60</sup> and has been of interest for  
52  
53 therapeutics in autoimmune and inflammatory disease mouse models. In models of experimental  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 autoimmune encephalitis (EAE), PI3K $\delta$  mutant mice were noted to have a defective Th17 response and  
4 reduced disease severity<sup>34</sup>. PI3K $\delta$  inhibition slowed disease progression and organ damage in a murine  
5 model of systemic lupus erythematosus, an autoimmune disease with T- and B- cell involvement similar  
6 to several immunological abnormalities associated with cGVHD<sup>61</sup>. Loss of PI3K $\delta$  activity improved  
7 outcomes in multiple sclerosis, rheumatoid arthritis, psoriasis and autoimmune (type 1) diabetes  
8 models<sup>40</sup>. We observed similar results with PI3K $\delta$  inhibition in cGVHD models studied here, including  
9 decreased damage to the lung, Ig deposition and IL-17. Of note, prior *in vitro* assays have shown that  
10 pharmacologic pan-PI3K inhibition was more effective than more selective inhibition of p110 $\delta$  alone  
11 for preventing differentiation of Th1 cells, as determined by IFN $\gamma$  production; in contrast, IL-17 was  
12 completely blocked by both inhibitor types<sup>34</sup>. Moreover, p110 $\delta$ <sup>D910A</sup> mice had greater reduction in Th17  
13 compared to Th1 responses in an EAE model<sup>34</sup>. While cytokines were not directly measured in our BO  
14 cGVHD, previously we have reported that IL-17 contributes to cGVHD in the BO model, as  
15 demonstrated by the lack of cGVHD using RORC deficient T cells and reversal of established disease  
16 using small molecule ROR $\gamma$ t inhibitors or neutralizing anti-IL-17 mAb treatment<sup>10</sup>.

17  
18  
19 In addition to regulation of IL-17 production, sustained PI3K $\delta$  activation has been found to be  
20 necessary for optimal IFN $\gamma$  production<sup>32</sup>. In the scleroderma model, inhibition of the  $\delta$  isoform with  
21 GS-649443 did not impact the frequencies of IFN $\gamma$  expressing donor T-cells. These data are however  
22 consistent with the reduced efficacy in IFN $\gamma$  suppression seen in CD8+ T cell later after TCR  
23 stimulation. Importantly, our data indicate that IFN $\gamma$  inhibition alone is not essential for reducing  
24 disease severity. In the cGVHD BO model, the role of IFN $\gamma$  in mediating disease has not been  
25 elucidated. However, in acute GVHD models, the lack of donor IFN $\gamma$  production increased pulmonary  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 GVHD and GVL responses, while reducing GI GVHD<sup>62</sup>. Thus, we do not favor the explanation that  
4 reduced IFN $\gamma$  production by PI3K $\delta$  inhibition is fundamentally important for cGVHD with BO.  
5

6  
7 A sizable population of allo-BMT patients have a hematological malignancy, many of whom will  
8 develop cGVHD and hence are potential candidates for PI3K $\delta$  treatment for post-BMT relapse and/or  
9 cGVHD. Because donor T-cells are principal protectors against relapse providing the beneficial graft-  
10 versus-leukemia (GVL) response<sup>63</sup>, the GVL response could be diminished by PI3K $\delta$  inhibition in  
11 cGVHD patients in whom PI3K $\delta$  activity is not a driving force in malignancy. However, for many  
12 cGVHD patients, especially those with long-standing disease, the GVL effect already may have  
13 eliminated residual malignant cells by the time that therapy would begin and for patients with steroid-  
14 resistant or refractory cGVHD, profound immune suppression may subvert existing GVL responses.  
15  
16 Future studies will need to be conducted to determine how inhibition of PI3K $\delta$  will impact on GVL and  
17 other immune function in the context of cGVHD treatment.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

33 Several important issues remain to be addressed. For example, PI3K signaling is involved in many  
34 different aspects of immunity and therefore inhibition could impact immune reconstitution. The impact  
35 of this therapy on cells of the immune system will be an important consideration going forward. GS-  
36 649443 ameliorated cGVHD in both the BO and scleroderma models, treatment was initiated at early  
37 times after disease establishment. The efficacy of PI3K $\delta$  inhibition in patients with steroid-refractory or  
38 advanced cGVHD remains to be determined. While the toxicities associated with PI3K $\delta$  inhibitors are  
39 of concern for future therapeutic applications, structural modifications, such as the one utilized in this  
40 study, offer the promise to decrease off target effects related with treatment and improve the  
41 historically poor outcome of cGVHD patients failing to respond to steroids. Nonetheless, careful  
42 pharmacological toxicology studies must be performed given the potential broader implications of  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 PI3K $\delta$  inhibition on systems beyond immunity and inflammation. Lastly, the potential broader off target  
4 effects of GS-649443 on other PI3K isoforms and other kinases for those drugs destined for clinical  
5 applications deserve thorough exploration.  
6  
7

8  
9  
10 In conclusion, these results demonstrate that PI3K $\delta$  activity is necessary for the development of  
11 cGVHD in murine models. We have demonstrated that targeting PI3K $\delta$  can result in a decreased GC  
12 reaction. Inhibiting PI3K $\delta$  improved cGVHD disease outcome by reducing pathogenic Tfh/GC B-cells  
13 resulting in decreased antibody and collagen deposition in the lungs. PI3K $\delta$  inhibition is also able to  
14 decrease inflammatory cytokines associated with cGVHD. These studies add to current knowledge of  
15 application of PI3K $\delta$  inhibition for disease treatment and present support for targeting PI3K $\delta$  for  
16 cGVHD therapy.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

### 29 **Acknowledgments**

30  
31 This work was supported by the National Institutes of Health grants P01 AI 056299, P01 CA 142106  
32 and T32 CA009138. Thank you to Dr. Christophe Queva for providing comments and edits.  
33  
34  
35  
36  
37

### 38 **Disclosure**

39  
40 The authors of this manuscript have conflicts of interest to disclose as described by the *American*  
41 *Journal of Transplantation*. S.T. is an employee of Gilead Sciences, Inc. B.V. is a consultant to Karus  
42 Therapeutics (Oxford, UK). The other authors have no conflicts of interest to disclose.  
43  
44  
45  
46  
47  
48  
49

### 50 **Data Availability Statement**

51  
52 The data that support the findings of this study are available from the corresponding author upon  
53 reasonable request.  
54  
55  
56  
57  
58  
59  
60

## Figure Legend

### Figure 1. Mice receiving p110 $\delta$ <sup>D910A</sup> BM develop cGVHD

B10.BR mice were conditioned with Cytoxan and TBI and infused with BM alone or with WT purified splenic T-cells (cGVHD) along with mice receiving p110 $\delta$ <sup>D910A</sup> BM alone or with WT T-cells. (A) Day 56 PFTs show that mice that received p110 $\delta$ <sup>D910A</sup> BM with WT T cells still developed BO comparable to cGVHD controls. (B-C) The frequency of splenic Tregs and Tfr demonstrate that these populations are reduced in Tfh both groups that received p110 $\delta$ <sup>D910A</sup> BM. (D) The splenic Tfh frequency was decreased in p110 $\delta$ <sup>D910A</sup> BM supplemented with T-cell group compared to the cGVHD control. (E) The Tfh frequency was still increased from the p110 $\delta$ <sup>D910A</sup> BM resulting in a Tfr:Tfh ratio similar to that of the cGVHD control. (F) The frequency of splenic GC B-cells were decreased in mice that received the p110 $\delta$ <sup>D910A</sup> T cells compared to cGVHD control but still increased from p110 $\delta$ <sup>D910A</sup> BM. A-E Data are from 2 pooled, independent experiments, with 5-7 mice per group per experiment. In F data are representative from 1 experiment. Data shown with mean  $\pm$  SD. One-way ANOVA with Bonferroni correction for multiple comparisons used with significance: \* $P > .05$ ; \*\* $P > .01$ ; \*\*\* $P > .001$ .

### Figure 2. PI3K $\delta$ is necessary in donor T-cells for cGVHD development

B10.BR mice were conditioned with Cytoxan and TBI and infused with BM alone or with WT purified splenic T-cells (cGVHD) or catalytically inactive T-cells. (A) Pulmonary function tests performed on day 56 show that the p110 $\delta$ <sup>D910A</sup> T cells did not induce BO cGVHD. (B) The frequency of splenic Tfh was decreased in mice that received the p110 $\delta$ <sup>D910A</sup> T cells. The Tfr frequency was not changed among any of the groups (C), however the Tfr:Tfh ratio was significantly improved (D). (E) The frequency of splenic GC B-cells was also decreased in mice that received p110 $\delta$ <sup>D910A</sup> T cells (F) Hematoxylin and eosin staining of lungs show that mice receiving p110 $\delta$ <sup>D910A</sup> T-cells had had improved histopathology.

1  
2  
3 Data are representative of 2 independent experiments with similar result with 4-5 mice per group,  
4  
5 shown with mean  $\pm$  SD. Student's t-test was used when comparing two groups with significance:  $*P >$   
6  
7  $.05$ ;  $**P >$   $.01$ ;  $***P >$   $.001$ .  
8  
9

10  
11  
12 **Figure 3. Therapeutic administration of PI3K $\delta$  specific inhibitor GS-649443 ameliorates disease**  
13  
14 **in a non-sclerodermatous, BO model of cGVHD**

15  
16  
17 B10.BR mice were conditioned with Cytoxan and TBI received BM alone or with B6 purified splenic  
18  
19 T-cells (cGVHD) treated mice received vehicle or PI3K $\delta$  specific inhibitor GS-649443 (10mg/kg/BID)  
20  
21 beginning on day 28 after transplant. (A) Day 56 PFTs show that GS-649443 improved lung function  
22  
23 of cGVHD mice. (B) Hemotoxylin and eosin staining of lungs show that mice treated with the inhibitor  
24  
25 had improved histopathology. (C) The frequency of splenic Tfh was significantly decreased in mice  
26  
27 treated with GS-649443. (D) These mice still had reduced frequency of Tfr cells and the ratio of  
28  
29 Tfr:Tfh was not improved (E). (F) The frequency of splenic GC B cells was significantly reduced in  
30  
31 mice treated with GS-649443. A and F are pooled from 3 independent experiments. B-E are pooled  
32  
33 from 2 independent experiments, with 4-6 mice per group per experiment. Data are shown with mean  $\pm$   
34  
35 SD. One-way ANOVA with Bonferroni correction for multiple comparisons used with significance:  
36  
37  $*P >$   $.05$ ;  $**P >$   $.01$ ;  $***P >$   $.001$ .  
38  
39  
40  
41  
42  
43

44  
45 **Figure 4. Histopathology and immunoglobulin (Ig) deposition of GS-649443 treated mice**  
46  
47 **phenocopies mice that received p110 $\delta$ <sup>D910A</sup> donor T-cells**

48  
49 Transplant set up was the same as figures 2 and 3. (A) Representative images of Ig deposition staining.  
50  
51 Ig deposition was quantified in ImageJ. (B) Representative images of Masson's Trichrome staining.  
52  
53 Collagen was identified as area stained blue and quantified using ImageJ indicating decreased collagen  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 deposited in the lungs of mice that received p110 $\delta$ <sup>D910A</sup> T-cells and mice treated with GS-649443. Data  
4  
5 are from one experiment with 3-5 mice per group, shown with mean  $\pm$  SD. One-way ANOVA with  
6  
7 Bonferroni correction for multiple comparisons used with significance: \* $P > .05$ ; \*\* $P > .01$ ; \*\*\* $P > .001$ .  
8  
9

10  
11  
12 **Figure 5. Therapeutic administration of the PI3K $\delta$  specific inhibitor GS-649443 ameliorates**  
13  
14 **sclerodermatous cGVHD**  
15

16  
17 Balb/c mice received TBI and received WT B10.D2 BM alone (BM only) or with  $1.8 \times 10^6$  CD4+ and  
18  
19  $0.9 \times 10^6$  CD8+ T-cells. Treatment groups received PI3K $\delta$  specific inhibitor GS-649443  
20  
21 (10mg/kg/BID) starting at day 21. (A) Mice treated with GS-649443 had improved skin scores. (B) GS-  
22  
23 649443 improved clinical scores in treated mice. Analysis of lymph nodes taken at day 50 post-  
24  
25 transplant, each sample is pooled from 2 mice, with 8-12 mice per group (C) Mice treated with GS-  
26  
27 649443 had reduced IL-17 frequency. (D) IL-17 and IFN $\gamma$  double positive population frequency were  
28  
29 also decreased. (E) IFN $\gamma$  positive population frequency was not decreased with treatment. (F)  
30  
31 Representative images of Ig deposition in the skin of mice treated with GS-649443 quantified using  
32  
33 ImageJ (G). Data in (A) is pooled data from two independent experiments, (B-C) are representative  
34  
35 from 2 independent experiments. Data are shown with mean  $\pm$  SD. Student's t-test was used with  
36  
37 significance: \* $P > .05$ ; \*\* $P > .01$ ; \*\*\* $P > .001$ .  
38  
39  
40  
41  
42  
43  
44

45 **Supporting Information**

46  
47 Additional supporting information may be found online in the Supporting Information section at the  
48  
49 end of this article.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

1. Arai S, Arora M, Wang T, et al. Increasing incidence of chronic graft-versus-host disease in allogeneic transplantation: a report from the Center for International Blood and Marrow Transplant Research. *Biol Blood Marrow Transplant*. 2015;21(2):266-274.
2. Zeiser R, Blazar BR. Pathophysiology of Chronic Graft-versus-Host Disease and Therapeutic Targets. *N Engl J Med*. 2017;377(26):2565-2579.
3. Socié G, Ritz J. Current issues in chronic graft-versus-host disease. *Blood*. 2014;124(3):374-384.
4. MacDonald KP, Hill GR, Blazar BR. Chronic graft-versus-host disease: biological insights from preclinical and clinical studies. *Blood*. 2017;129(1):13-21.
5. Zeiser R, Blazar BR. Preclinical models of acute and chronic graft-versus-host disease: how predictive are they for a successful clinical translation? *Blood*. 2016;127(25):3117-3126.
6. Schroeder MA, DiPersio JF. Mouse models of graft-versus-host disease: advances and limitations. *Dis Model Mech*. 2011;4(3):318-333.
7. Coghill JM, Sarantopoulos S, Moran TP, Murphy WJ, Blazar BR, Serody JS. Effector CD4+ T cells, the cytokines they generate, and GVHD: something old and something new. *Blood*. 2011;117(12):3268-3276.
8. Cooke KR, Luznik L, Sarantopoulos S, et al. The Biology of Chronic Graft-versus-Host Disease: A Task Force Report from the National Institutes of Health Consensus Development Project on Criteria for Clinical Trials in Chronic Graft-versus-Host Disease. *Biol Blood Marrow Transplant*. 2017;23(2):211-234.
9. MacDonald KP, Blazar BR, Hill GR. Cytokine mediators of chronic graft-versus-host disease. *J Clin Invest*. 2017;127(7):2452-2463.
10. Forcade E, Paz K, Flynn R, et al. An activated Th17-prone T cell subset involved in chronic graft-versus-host disease sensitive to pharmacological inhibition. *JCI Insight*. 2017;2(12).
11. Craft JE. Follicular helper T cells in immunity and systemic autoimmunity. *Nat Rev Rheumatol*. 2012;8(6):337-347.
12. Weinstein JS, Herman EI, Lainez B, et al. TFH cells progressively differentiate to regulate the germinal center response. *Nat Immunol*. 2016;17(10):1197-1205.
13. Crotty S. T follicular helper cell differentiation, function, and roles in disease. *Immunity*. 2014;41(4):529-542.
14. Sage PT, Sharpe AH. T follicular regulatory cells in the regulation of B cell responses. *Trends Immunol*. 2015;36(7):410-418.
15. Srinivasan M, Flynn R, Price A, et al. Donor B-cell alloantibody deposition and germinal center formation are required for the development of murine chronic GVHD and bronchiolitis obliterans. *Blood*. 2012;119(6):1570-1580.
16. Flynn R, Du J, Veenstra RG, et al. Increased T follicular helper cells and germinal center B cells are required for cGVHD and bronchiolitis obliterans. *Blood*. 2014;123(25):3988-3998.
17. Dubovsky JA, Flynn R, Du J, et al. Ibrutinib treatment ameliorates murine chronic graft-versus-host disease. *J Clin Invest*. 2014;124(11):4867-4876.
18. Flynn R, Allen JL, Luznik L, et al. Targeting Syk-activated B cells in murine and human chronic graft-versus-host disease. *Blood*. 2015;125(26):4085-4094.
19. Flynn R, Paz K, Du J, et al. Targeted Rho-associated kinase 2 inhibition suppresses murine and human chronic GVHD through a Stat3-dependent mechanism. *Blood*. 2016;127(17):2144-2154.
20. Aggarwal S, Ghilardi N, Xie MH, de Sauvage FJ, Gurney AL. Interleukin-23 promotes a distinct CD4 T cell activation state characterized by the production of interleukin-17. *J Biol Chem*. 2003;278(3):1910-1914.
21. Alexander KA, Flynn R, Lineburg KE, et al. CSF-1-dependant donor-derived macrophages mediate chronic graft-versus-host disease. *J Clin Invest*. 2014;124(10):4266-4280.
22. Radojic V, Pletneva MA, Yen HR, et al. STAT3 signaling in CD4+ T cells is critical for the pathogenesis of chronic sclerodermatous graft-versus-host disease in a murine model. *J Immunol*. 2010;184(2):764-774.




23. Engelman JA, Luo J, Cantley LC. The evolution of phosphatidylinositol 3-kinases as regulators of growth and metabolism. *Nat Rev Genet*. 2006;7(8):606-619.
24. Fruman DA, Meyers RE, Cantley LC. Phosphoinositide kinases. *Annu Rev Biochem*. 1998;67:481-507.
25. Huang YH, Sauer K. Lipid signaling in T-cell development and function. *Cold Spring Harb Perspect Biol*. 2010;2(11):a002428.
26. Okkenhaug K. Signaling by the phosphoinositide 3-kinase family in immune cells. *Annu Rev Immunol*. 2013;31:675-704.
27. Wu LX, La Rose J, Chen L, et al. CD28 regulates the translation of Bcl-xL via the phosphatidylinositol 3-kinase/mammalian target of rapamycin pathway. *J Immunol*. 2005;174(1):180-194.
28. Appleman LJ, van Puijenbroek AA, Shu KM, Nadler LM, Boussiotis VA. CD28 costimulation mediates down-regulation of p27kip1 and cell cycle progression by activation of the PI3K/PKB signaling pathway in primary human T cells. *J Immunol*. 2002;168(6):2729-2736.
29. Delgoffe GM, Pollizzi KN, Waickman AT, et al. The kinase mTOR regulates the differentiation of helper T cells through the selective activation of signaling by mTORC1 and mTORC2. *Nat Immunol*. 2011;12(4):295-303.
30. Jacobs SR, Herman CE, Maciver NJ, et al. Glucose uptake is limiting in T cell activation and requires CD28-mediated Akt-dependent and independent pathways. *J Immunol*. 2008;180(7):4476-4486.
31. Patton DT, Garden OA, Pearce WP, et al. Cutting edge: the phosphoinositide 3-kinase p110 delta is critical for the function of CD4+CD25+Foxp3+ regulatory T cells. *J Immunol*. 2006;177(10):6598-6602.
32. Soond DR, Bjørgo E, Moltu K, et al. PI3K p110delta regulates T-cell cytokine production during primary and secondary immune responses in mice and humans. *Blood*. 2010;115(11):2203-2213.
33. Park SJ, Lee KS, Kim SR, et al. Phosphoinositide 3-kinase  $\delta$  inhibitor suppresses interleukin-17 expression in a murine asthma model. *Eur Respir J*. 2010;36(6):1448-1459.
34. Haylock-Jacobs S, Comerford I, Bunting M, et al. PI3K $\delta$  drives the pathogenesis of experimental autoimmune encephalomyelitis by inhibiting effector T cell apoptosis and promoting Th17 differentiation. *J Autoimmun*. 2011;36(3-4):278-287.
35. Uehara M, McGrath MM, Otori S, et al. Regulation of T cell alloimmunity by PI3K $\gamma$  and PI3K $\delta$ . *Nat Commun*. 2017;8(1):951.
36. Herrero-Sánchez MC, Rodríguez-Serrano C, Almeida J, et al. Targeting of PI3K/AKT/mTOR pathway to inhibit T cell activation and prevent graft-versus-host disease development. *J Hematol Oncol*. 2016;9(1):113.
37. Doisne JM, Hüber CM, Okkenhaug K, Colucci F. Immunomodulation of Selective Naive T Cell Functions by p110 $\delta$  Inactivation Improves the Outcome of Mismatched Cell Transplantation. *Cell Rep*. 2015.
38. Borlado LR, Redondo C, Alvarez B, et al. Increased phosphoinositide 3-kinase activity induces a lymphoproliferative disorder and contributes to tumor generation in vivo. *FASEB J*. 2000;14(7):895-903.
39. Uddin S, Hussain AR, Siraj AK, et al. Role of phosphatidylinositol 3'-kinase/AKT pathway in diffuse large B-cell lymphoma survival. *Blood*. 2006;108(13):4178-4186.
40. Stark AK, Srisantharajah S, Hessel EM, Okkenhaug K. PI3K inhibitors in inflammation, autoimmunity and cancer. *Curr Opin Pharmacol*. 2015;23:82-91.
41. Yang Q, Modi P, Newcomb T, Quéva C, Gandhi V. Idelalisib: First-in-Class PI3K Delta Inhibitor for the Treatment of Chronic Lymphocytic Leukemia, Small Lymphocytic Leukemia, and Follicular Lymphoma. *Clin Cancer Res*. 2015;21(7):1537-1542.
42. Patel L, Chandrasekhar J, Everts J, et al. Discovery of Orally Efficacious Phosphoinositide 3-Kinase  $\delta$  Inhibitors with Improved Metabolic Stability. *J Med Chem*. 2016.
43. Yahiaoui A, Meadows SA, Sorensen RA, et al. PI3K $\delta$  inhibitor idelalisib in combination with BTK inhibitor ONO/GS-4059 in diffuse large B cell lymphoma with acquired resistance to PI3K $\delta$  and BTK inhibitors. *PLoS One*. 2017;12(2):e0171221.
44. Way EE, Trevejo-Nunez G, Kane LP, et al. Dose-Dependent Suppression of Cytokine production from T cells by a Novel Phosphoinositide 3-Kinase Delta Inhibitor. *Sci Rep*. 2016;6:30384.


- 1  
2  
3 45. Okkenhaug K, Bilancio A, Farjot G, et al. Impaired B and T cell antigen receptor signaling in p110delta PI  
4 3-kinase mutant mice. *Science*. 2002;297(5583):1031-1034.
- 5 46. Zhang Y, McCormick LL, Desai SR, Wu C, Gilliam AC. Murine sclerodermatous graft-versus-host disease,  
6 a model for human scleroderma: cutaneous cytokines, chemokines, and immune cell activation. *J Immunol*.  
7 2002;168(6):3088-3098.
- 8 47. Hill GR, Olver SD, Kuns RD, et al. Stem cell mobilization with G-CSF induces type 17 differentiation and  
9 promotes scleroderma. *Blood*. 2010;116(5):819-828.
- 10 48. Le Huu D, Matsushita T, Jin G, et al. IL-6 blockade attenuates the development of murine  
11 sclerodermatous chronic graft-versus-host disease. *J Invest Dermatol*. 2012;132(12):2752-2761.
- 12 49. Panoskaltis-Mortari A, Tram KV, Price AP, Wendt CH, Blazar BR. A new murine model for bronchiolitis  
13 obliterans post-bone marrow transplant. *Am J Respir Crit Care Med*. 2007;176(7):713-723.
- 14 50. Blazar BR, Taylor PA, McElmurry R, et al. Engraftment of severe combined immune deficient mice  
15 receiving allogeneic bone marrow via In utero or postnatal transfer. *Blood*. 1998;92(10):3949-3959.
- 16 51. Sage PT, Tan CL, Freeman GJ, Haigis M, Sharpe AH. Defective TFH Cell Function and Increased TFR Cells  
17 Contribute to Defective Antibody Production in Aging. *Cell Rep*. 2015;12(2):163-171.
- 18 52. McDonald-Hyman C, Turka LA, Blazar BR. Advances and challenges in immunotherapy for solid organ  
19 and hematopoietic stem cell transplantation. *Sci Transl Med*. 2015;7(280):280rv282.
- 20 53. Martires KJ, Baird K, Steinberg SM, et al. Sclerotic-type chronic GVHD of the skin: clinical risk factors,  
21 laboratory markers, and burden of disease. *Blood*. 2011;118(15):4250-4257.
- 22 54. Lee YK, Turner H, Maynard CL, et al. Late developmental plasticity in the T helper 17 lineage. *Immunity*.  
23 2009;30(1):92-107.
- 24 55. Clayton E, Bardi G, Bell SE, et al. A crucial role for the p110delta subunit of phosphatidylinositol 3-  
25 kinase in B cell development and activation. *J Exp Med*. 2002;196(6):753-763.
- 26 56. Jou ST, Carpino N, Takahashi Y, et al. Essential, nonredundant role for the phosphoinositide 3-kinase  
27 p110delta in signaling by the B-cell receptor complex. *Mol Cell Biol*. 2002;22(24):8580-8591.
- 28 57. McDonald-Hyman C, Flynn R, Panoskaltis-Mortari A, et al. Therapeutic regulatory T-cell adoptive  
29 transfer ameliorates established murine chronic GVHD in a CXCR5-dependent manner. *Blood*.  
30 2016;128(7):1013-1017.
- 31 58. Du J, Paz K, Flynn R, et al. Pirfenidone ameliorates murine chronic GVHD through inhibition of  
32 macrophage infiltration and TGF- $\beta$  production. *Blood*. 2017;129(18):2570-2580.
- 33 59. Mouchemore KA, Sampaio NG, Murrey MW, Stanley ER, Lannutti BJ, Pixley FJ. Specific inhibition of PI3K  
34 p110 $\delta$  inhibits CSF-1-induced macrophage spreading and invasive capacity. *FEBS J*. 2013;280(21):5228-5236.
- 35 60. Suárez-Fueyo A, Barber DF, Martínez-Ara J, Zea-Mendoza AC, Carrera AC. Enhanced phosphoinositide  
36 3-kinase  $\delta$  activity is a frequent event in systemic lupus erythematosus that confers resistance to activation-  
37 induced T cell death. *J Immunol*. 2011;187(5):2376-2385.
- 38 61. Wang Y, Zhang L, Wei P, Zhang H, Liu C. Inhibition of PI3K $\delta$  improves systemic lupus in mice.  
39 *Inflammation*. 2014;37(3):978-983.
- 40 62. Wang H, Asavaroengchai W, Yeap BY, et al. Paradoxical effects of IFN-gamma in graft-versus-host  
41 disease reflect promotion of lymphohematopoietic graft-versus-host reactions and inhibition of epithelial tissue  
42 injury. *Blood*. 2009;113(15):3612-3619.
- 43 63. Johnson BD, Hanke CA, Truitt RL. The graft-versus-leukemia effect of post-transplant donor leukocyte  
44 infusion. *Leuk Lymphoma*. 1996;23(1-2):1-9.
- 45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Figure 1.**


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60





**Figure 3.**







# Figure 4.

A.




B.





- BM only
- BM + T (Vehicle)
- ▲ BM + T (GS-649443)



## Supporting Information

### Figure S1. Heterozygous p110 $\delta$ <sup>D910A/WT</sup> cells still induce pulmonary dysfunction associated with cGVHD

Transplant set up the same as figure 2, with additional groups of mice receiving WT BM with p110 $\delta$ <sup>D910A/WT</sup> T-cells, p110 $\delta$ <sup>D910A</sup> PI3K $\delta$  BM alone or with WT T cells, and p110 $\delta$ <sup>D910A/WT</sup> BM and T-cells. Day 56 PFTs show that mice that one functional copy of PI3K $\delta$  in donor BM or T-cells paired with WT BM or T-cells is sufficient for causing disease however, one copy in donor BM and donor T-cells had a modest effect on pulmonary outcome. Data are representative from one experiment, with 4-6 mice per group. Data are shown with mean  $\pm$  SD. Significance: \* $P > .05$ ; \*\* $P > .01$ ; \*\*\* $P > .001$ .

### Figure S2. Weight, survival and Treg frequency of mice that received p110 $\delta$ <sup>D910A</sup> donor T-cells was not significantly changed from cGVHD mice

(A) Weights and (B) survival from mice that received p110 $\delta$ <sup>D910A</sup> donor T-cells was not significantly changed from mice that received WT BM supplemented with WT donor T-cells. (C) Treg population is not changed in mice that received p110 $\delta$ <sup>D910A</sup> donor T-cells. Data are representative from two independent experiments, with 8-10 mice per group.

### Figure S3. Therapeutic administration of PI3K $\delta$ specific inhibitor GS-649443 did not significantly impact weight and survival, however Treg frequency was decreased

B10.BR mice were conditioned with Cytoxan and TBI received BM alone or with B6 purified splenic T-cells (cGVHD) treated mice received vehicle or PI3K $\delta$  specific inhibitor GS-649443 (10mg/kg/BID) beginning on day 28 after transplant. (A-B) Weight and survival curves of transplanted mice




1  
2  
3 demonstrate that GS-649443 did not have toxic effect. Data are pooled from three independent  
4  
5 experiments, with 8-10 mice per group. (C) Treg population is significantly decreased in mice that  
6  
7 received GS-649443 compared to vehicle control.  
8  
9

10  
11  
12 **Figure S4. Therapeutic administration of PI3K $\delta$  specific inhibitor GS-649443 at 5mg/kg does not**  
13  
14 **reduce pulmonary dysfunction associated with BO model of cGVHD**

15  
16 Conditioned mice received BM alone (BM only) or supplemented with purified splenic T cells. Mice  
17  
18 were treated with vehicle or the PI3K $\delta$  specific inhibitor GS-649443 (5mg/kg/BID) beginning on day  
19  
20 28 after transplant. (A) Day 56 PFTs show that GS-649443 improved lung function of cGVHD mice.  
21  
22 Data are representative of 2 independent experiments, with 7-9 mice per group, shown with mean  $\pm$  SD.  
23  
24 Significance: \* $P > .05$ ; \*\* $P > .01$ ; \*\*\* $P > .001$   
25  
26  
27  
28  
29  
30


31 **Figure S5. Therapeutic administration of the PI3K $\delta$  specific inhibitor GS-649443 reduced**  
32  
33 **inflammatory cell subsets as well as Ig deposition in skin**

34  
35 Balb/c mice received TBI and received WT B10.D2 BM alone (BM only) or with  $1.8 \times 10^6$  CD4+ and  
36  
37  $0.9 \times 10^6$  CD8+ T-cells. Treatment groups received PI3K $\delta$  specific inhibitor GS-649443  
38  
39 (10mg/kg/BID) starting at day 21. A-B Analysis of lymph nodes taken at day 50 post-transplant, each  
40  
41 sample is pooled from 2 mice, with 8-12 mice per group. (A) IL-17 and IFN $\gamma$  double positive  
42  
43 population frequency was decreased with treatment however IFN $\gamma$  positive population frequency was  
44  
45 not reduced (B). (C) Representative images of Ig deposition in the skin of mice treated with GS-649443  
46  
47 quantified using ImageJ. Data is representative from 2 independent experiments. Data are shown with  
48  
49 mean  $\pm$  SD. Student's t-test was used with significance: \* $P > .05$ ; \*\* $P > .01$ ; \*\*\* $P > .001$ .  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


A.



B.




C.




# Figure S3.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**A.**




**B.**




- BM only
- cGVHD
- ▲ cGVHD (Vehicle)
- ▼ cGVHD (10mg/kg GS-649443)

**C.**



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

