

1 **IL-17 and IL-22 Elicited by a DNA vaccine encoding ROP13 associate with protection**
2 **against *Toxoplasma gondii* in BALB/c mice**

3
4 Paria Alizadeh^{1,2}, Ehsan Ahmadpour^{*1}, Ahmad Daryani³, Tohid Kazemi¹, Adel Spotin⁴,
5 Mahmoud Mahami Oskouei⁵, Robin J Flynn⁶, Yaghob Azadi², Saba Rajabi², Siamak
6 Sandoghchian¹

7
8 ¹ Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

9 ² Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

10 ³ Toxoplasmosis Research Center, Mazandaran University of Medical Sciences, Sari, Iran

11 ⁴ Infectious and Tropical Disease Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

12 ⁵ Department of Parasitology and Mycology, Tabriz University of Medical Sciences, Tabriz, Iran.

13 ⁶ Department of Infection Biology, Institute of Infection and Global Health, University of Liverpool,
14 L3 5RF, UK

15

16

17

18 ***Corresponding author:** Ehsan Ahmadpour, Ph.D

19 Address: Department of Parasitology and Mycology, Tabriz university of Medical Sciences,
20 Tabriz, Iran.

21 Tel/ Fax: +98 41 33373745

22 Email address: ehsanahmadpour@gmail.com, ahmadpoure@tbzmed.ac.ir

23 **Conflict of Interest:** Authors declare there is no any conflict of interest.

24 **Abstract:**

25 *Toxoplasma gondii*, an intracellular parasitic protozoan, is capable of infecting man and all
26 warm blooded animals. Cell-mediated immunity is vital in mounting protective responses
27 against *T. gondii* infection. Recent studies have shown that T-helper (Th) 17 responses may
28 play a key role in parasite control. In this current study, we constructed a DNA vaccine
29 encoding *T. gondii* ROP13 in a pcDNA vector. Groups of BALB/c mice were immunized
30 intramuscularly with pcROP13 or controls and challenged with the RH strain of *T. gondii*.
31 The results showed that immunization with pcROP13 could elicit an antibody response
32 against *T. gondii*. The expression of the canonical Th17 cytokines, IL-17 and IL-22, were
33 significantly increased after immunization with pcROP13 compared to control groups
34 ($P<0.05$). Furthermore, vaccination resulted in a significant decrease in parasite load
35 ($P<0.05$). The induction of Th17 related cytokines, using a ROP13 DNA vaccine, against *T.*
36 *gondii* should be considered as a potential vaccine approach for the control of toxoplasmosis.

37

38

39

40

41 **Keywords:** ROP13, *Toxoplasma gondii*, Th17, IL-22, IL-17, DNA vaccine, Gene Expression

42

43

44 **1. Introduction**

45 Toxoplasmosis is globally widespread parasitic infection caused by the intracellular
46 protozoan pathogen, *T. gondii*, infecting humans and the other warm-blooded animals [1]. In
47 immunocompetent people *Toxoplasma* infection is benign and mostly presents with no
48 clinical manifestations. However, dependent on the status of host immune system it can cause
49 serious and irreversible effects [1-2]. Toxoplasmosis in immunocompromised individuals is
50 an opportunistic infection that may cause severe ocular and life-threatening neurological
51 disorders [2]. Due to the high prevalence of *T. gondii* and the resulting pathogenesis of
52 infection it is considered as a public health hazard. Despite extensive research, effective anti-
53 *Toxoplasma* therapeutics without side effects remains a barren area [3]. Hence, the most
54 effective strategy to reduce disease burden and clinical outcomes is the development of
55 vaccine formulations against *T. gondii* [4].

56 Current approaches to immunization against *Toxoplasma* infection takes several forms
57 including attenuated live vaccines, killed vaccines and subunit vaccines [4-5]. Owing to the
58 safety issues with the use of attenuated or killed forms of the pathogen, subunit vaccines have
59 attracted considerable attention [4, 6]. In particular, DNA vaccines have been developed in
60 recent years [5]. Results from several studies have raised the possibility of developing a DNA
61 vectored vaccine to protect against *T. gondii* infection. The most investigated compounds as
62 vaccine candidates include excreted-secreted antigens (ESA) and surface antigens of
63 tachyzoites [7-9]. Previous findings indicate that ESA play a significant role in disease
64 pathogenesis, and escape of the parasite from host immunity [10]. In particular Rhoptries
65 (ROP) are unique secretory organelles that involved in host cell penetration by *T. gondii* and
66 parasitophorous vacuole formation allowing survival and multiplication [11-12].

67 The protective mechanisms against *T. gondii* involve both CD4⁺ and CD8⁺ T-cell responses
68 [13]. IFN- γ is known to be the major effector as a result of T-helper 1 (Th1) cell and NK cell
69 activation. Th17 cells are a subset of CD4⁺ T-cells conditioned to produce the cytokines IL-
70 17, IL-21, and IL-22 which trigger responses causing the elimination of infection [13-16].
71 However, T cell-dependent production of IL-17 has been implicated in both protective and
72 pathogenic responses during infection with *T. gondii* [17]. Subsequent studies identified NK
73 cells as the innate IL-17 secreting cells in mice challenged with *Toxoplasma* [17-18].
74 Moreover, IL-17 mediated signaling was reported to play an important role during the initial
75 stages of *T. gondii* infection through neutrophil recruitment and activation [19].

76 Given the essential roles of ROP proteins in the pathogenesis of the *Toxoplasma* infection,
77 these critical antigens are appropriate vaccine candidates [20]. ROP13 is a relatively recently
78 recognized antigen of *T. gondii* and in tandem few studies have evaluated the Th17 response
79 in *T. gondii* DNA vaccine [21]. Hence, the two major objectives of the present study was to
80 construct a DNA vaccine vector expressing *T. gondii* ROP13 for use in immunization and to
81 thereafter analyze the protective immune responses induced by vaccination and challenge
82 with *T. gondii* RH strain.

83 **2. Materials and methods**

84 **2.1 Mice and parasite**

85 The highly virulent RH strain of *T. gondii* (type I) was used in all experiments. The RH
86 tachyzoites were provided by Toxoplasmosis Research Center in Mazandaran University of
87 Medical Sciences, Sari, Iran. The parasite was maintained by serial passage and
88 intraperitoneal inoculation and female 6 to 8 week-old BALB/c mice. The animals were
89 obtained from the Pasteur Institute of Iran and maintained under standard conventional

90 conditions. The animal experiments were approved by the local Ethics Committee of Tabriz
91 University of Medical Sciences, Tabriz, Iran (No. IR.TBZMED.REC.1395.578).

92 **2.2 Cloning of ROP13 and construction of plasmids**

93 DNA was extracted from tachyzoites by using an AccuPrep genomic DNA extraction kit
94 (Bioneer, Korea) according to the manufacturer's instructions. The NCBI GenBank database
95 was used to determine the complete sequence of ROP13 gene of RH strain and to design
96 specific primers (GeneBank accession number: JN051278.1). The ROP13 gene was
97 amplified using the primer pair Forward: 5' -GGATCCATGAAGAGAACAGAGCTTTG- 3',
98 and Reverse: 5' -TCTAGATCACAATAGCCTCAAGGAATTC- 3' with six base pair,
99 underlined, recognition sites for *Bam HI* and *Xba I* respectively in the primers. The coding
100 sequence of ROP13 gene was 1203bp in length which was inspected by using 1% agarose gel
101 electrophoresis to ensure the fidelity of the PCR product. The ROP13 PCR product was then
102 inserted into the linearized pTG19-T vector (Vivantis) between the *Bam HI* and *Xba I* sites.

103 The pTROP13 plasmid was then transferred into competent Top10 *E. coli* cells. Transformed
104 bacteria were plated on LB-agar plates containing ampicillin, X-gal, and IPTG and incubated
105 overnight. Blue/white screening was used to select transformed colonies harboring pTROP13
106 were isolated and subjected to PCR to confirm the correct insertion was present [21-22].

107 To generate the vaccine plasmid pTROP13 was recovered from *E. coli* and subject to
108 miniprep plasmid extraction (Gene All). The ROP13 coding sequence removed from the
109 vector by double digestion cleavage using *BamHI* / *XbaI* (Jena Bioscience). The coding
110 sequence was subject to gel purification and extraction (Bioneer, AccuPrep® Gel Purification
111 Kit) before confirmation by DNA sequencing. To construct the vaccine vector, the ROP13
112 gene sequence was ligated into the pcDNA3, yielded the plasmid pcROP13.

113 **2.5 Transfection of CHO cells**

114 Chinese Hamster Ovary (CHO) cells were transfected with the pcROP13 plasmid. Cells, 1-2
115 $\times 10^4$ per well were plated into a 96-well tissue culture plate and used when the cells were 50-
116 80% confluent. Transfection was performed using jetPrime (Polyplus, France) according to
117 the manufacturer's instructions. Uptake of pcROP13 and expression from pcROP13 was
118 detected 24-48 hours after transfection by immunofluorescence [21].

119 **2.6 In vitro expression of pcROP13**

120 PcROP13 plasmid expression was detected by indirect immunofluorescence assay. Serial
121 dilutions, beginning at 1/10, of the human anti-*T. gondii* antiserum were applied to
122 transfected cells. Anti-sera were coated on a slide where the Transfected cells, on slides, were
123 fixed, followed by incubation with anti-sera in a humidified chamber for 30 minutes; slides
124 were then washed with PBS and dried at room temperature. The slides were subsequently
125 incubated with secondary antibody of goat anti-human IgG conjugated with fluorescein
126 isothiocyanate (FITC) for 30 minutes in the dark. After washing 3 times with PBS, the cells
127 were mounted using buffered glycerol and examined for fluorescence detection under
128 CYTATION5 imaging reader [21].

129 **2.7 Mice immunization and challenge**

130 Forty female 6-8-week-old BALB/c mice were divided into four groups; group A was
131 vaccinated with 100 μ g of pcROP13 DNA plasmid suspended in PBS, by intramuscular
132 injection. Group B received PBS, Group C received empty pcDNA3 vector in PBS, and
133 Group D received 20 μ g of TLA (*T. gondii* lysate antigen). All mice were immunized three

134 times, two weeks apart prior to parasite challenge. Animals were infected with *T. gondii* RH
135 strain by intraperitoneal injection with 1×10^4 parasites.

136 **2.8 Immune responses and determination of parasite load**

137 Serum IgG antibody levels were determined by ELISA as previously described (REF).
138 Samples were obtained from mice at two individual time points including the pre-vaccination
139 period (day 0) and on day 42 after immunization but prior to infection.

140 To evaluate parasite load 3 days after challenge, DNA was extracted from the blood using the
141 Blood Genomic DNA Extraction kit (YTA, Iran, Cat No: YT9040) according to the
142 manufacturer's instructions. Parasite load was determined by quantification of tachyzoites
143 using real time PCR amplification of the highly conserved RE gene of *T. gondii* as previously
144 described [23-24]. Briefly, forward primer: 5`AGGGACAGAAGTCGAAGGGG-3` and
145 reverse primer: 5`GCAGCCAAGCCGGAAACATC-3` specified to amplify a 164-bp
146 fragment of the RE gene using SYBR green chemistry, with all amplifications in triplicate.
147 Q-PCR was performed using the following thermal cycling protocol: 10 minutes at 95°C, 40
148 cycles at 94°C for 30 seconds (denaturation), 55°C for 30 seconds (annealing) and 72°C for
149 30 seconds (amplification). Melting curve analysis was performed to verify the correct gene
150 product ensuring the absence of side products. The threshold cycle (CT) value at which the
151 fluorescence passes the fixed threshold was used to calculate the number of parasites in the
152 samples according to a standard curve obtained with tachyzoites prepared for DNA samples
153 over a range of 5×10^6 to 5×10^1 /ml. The results were reported as *T. gondii* tachyzoite-
154 equivalents per ml of blood.

155 **Th17 cytokine gene expression was monitored using blood samples collected from**
156 **animals**

157 Subsequently, the blood RNA was extracted (YTA, Iran, Cat No: YT9075) and cDNA was
158 synthesized (YTA, Iran, Cat No: YT4500) Real-time PCR for IL-17, IL-22 and GAPDH (as
159 internal control) was performed using SYBR Green chemistry (YTA, Iran) on a Roche Real-
160 time PCR system (Applied Biosystems). The primers for IL-17 and IL-22 based on real-time
161 PCR were as follow: IL-17 Forward primer: TCTCTGATGCTGTTGCTGCT, IL-17 Reverse
162 primer: CGTGGAACGGTTGAGGTAGT, IL-22 Forward primer:
163 TTGAGGTGTCCAACCTCCAGCA, IL-22 Reverse primer:
164 AGCCGGACGTCTGTGTTGTTA. The PCR cycling was carried out in a final volume of 20
165 μ l reaction by an initial denaturation step at 95°C for 3 min followed by 45 cycles at 95°C for
166 10 seconds, 58°C for 30 seconds, and 72°C for 20 seconds. Relative mRNA expression was
167 measured by the $2^{-(\Delta\Delta CT)}$ method, using GAPDH as a reference gene.

168 **2.11 Statistical analysis**

169 Statistical analysis was performed using GraphPad Prism 5.0 (San Diego, CA). Multiple
170 comparisons between groups were conducted by 1-way ANOVA with post-hoc testing. $P <$
171 0.05 was reported to be statistically significant.


172 **3. Results**

173 **3.1 Vaccine Construct**

174 The total DNA extracted from *Toxoplasma* tachyzoites and the coding sequence of ROP13
175 gene was amplified using PCR, a 1203-bp PCR product corresponding to the ROP13 coding
176 sequence was obtained (Figure 1A). This was inserted into the expression vector pcDNA3
177 between the *Bam* *HI* and *Xba* *I* cloning sites. The pcROP13 was transferred into CHO cells
178 and the protein expression was confirmed using IFAT (Figure. 1).

179

180




181

182 **Figure 1.** (Left) Gel electrophoresis analysis on PCR product, Lane 1: DNA size marker, Lane 2:
183 ROP13 gene with the expected band size. Lane 3: negative control. (Right) Indirect
184 immunofluorescence (IFA) detection of *Toxoplasma gondii* ROP13 on CHO cells (A: cells were
185 transfected with pcROP13, B: empty vector).

186 **3.2 Immunization with pcROP13**

187 Groups of BALB/c mice were immunized with pcROP13 or appropriate controls.
188 Immunization resulted in the seroconversion of animals as determined by ELISA. A specific
189 antibody response in both TLA and pcROP13 immunized groups was detected after the third
190 immunization (Figure 2). The total IgG levels for both groups was significantly different
191 ($P<0.05$) when compared to the negative control groups (PBS and pcDNA3).




192

193 **Figure 2.** Measurement of the specific anti-*Toxoplasma* IgG antibody in the sera of
 194 BALB/c mice before (left) and after (right) immunization at a 1:100 dilution. The results
 195 are shown as mean of the OD₄₅₀ ± SD of three independent experiments. **P* < 0.05
 196 Statistically significant differences compared to control group were determined by a 1-
 197 way anova. There were no detectable antibodies against *T. gondii* in the sera of control
 198 groups.

199

200 3.3 Blood Cytokine mRNA expression

201 48hr after the third immunization peripheral blood was collected from tail vein to evaluate the
 202 expression level of Th17 cytokines. The expression level of IL-22 mRNA in pcROP13 and
 203 TLA groups was found to be respectively 4 and nearly 2.5 folds higher than that observed in
 204 PBS and pcDNA3 groups (Figure 3). The expression of IL-17 was also significantly elevated
 205 among pcROP13 and TLA groups (2.8 and nearly 2 folds, respectively) compared with
 206 negative controls (*P*<0.05).




207

208 **Figure 3.** Relative mRNA expression of IL-17 (A) and IL-22 (B) in immunized mice with pcROP13,
 209 TLA, pcDNA3 and PBS. Results are expressed as mean and SD. *P*-value was determined using 1-way
 210 anova analysis (**P*<0.05, ** *P*<0.01, *** *P*<0.001).

211 3.4 Determination of parasite load in immunized mice

212 To determine the protective efficacy of the pcROP13 vaccine to induce protection against *T.*
 213 *gondii*, immunized mice were challenged via the intraperitoneal route with 1×10^4
 214 tachyzoites 2 weeks after the third immunization. Blood parasite load was then determined by
 215 qPCR three days after challenge. Figure 4 clearly shows that immunization with either TLA
 216 or pcROP13 induced protection in mice as measured by the significantly different parasite
 217 burden (*P*<0.05). The PBS and pcDNA groups harbored on average 22201 and 18436
 218 parasite/mL, respectively. The pcROP13 and TLA immunized groups harbored 1694 and 812
 219 parasites, respectively. However no significant difference was observed between pcROP13
 220 and TLA groups (Figure 4).



221

222 **Figure 4:** The parasite load, tachyzoites/mL of blood in immunized mice following a 3-day i.p.
 223 infection with RH strain of *T. gondii*. Significant differences in parasite load between groups was
 224 detected by 1-way anova (* $P < 0.05$).

225 4. Discussion

226 Immunization with *T. gondii* ROP13 gene has been previously shown to induce a strong
 227 protective humoral and cellular response against infection in the recent study when
 228 adjuvanted with IL-18 [21]. In the current study, we constructed a vaccine plasmid pcROP13
 229 expressing protein ROP13 of *T. gondii* and evaluated the immune response induced in
 230 BALB/c mice. Our findings demonstrated that in addition to the induction of a humoral
 231 response, there is also an increased gene expression of Th17 cytokines (IL-17 and IL-22). In
 232 agreement with previous studies we found that immunization with pcROP13, as a DNA
 233 vaccine, successfully decreased the parasite load of blood in immunized mice.

234 In the past decade, DNA vaccines have been widely studied and have been shown to elicit an
 235 efficient immune response against target antigens in various animal models [5, 25]. Various
 236 antigens of *T. gondii* have been assessed as potential candidates for vaccine development [8,

237 22, 26-27]. Rhoptries are found in apical secretory organelle and function in the
238 establishment of infection through formation of specific compartments known as
239 parasitophorous vacuoles in which parasite evades intracellular killing [11]. ROP13 is a
240 unique soluble effector protein known to implicate in host cell invasion that can be detected
241 in the cytoplasm of host cells [21]. A previous study evaluated the immunogenicity of a DNA
242 vaccine expressing ROP13 of *T. gondii*, pVAX-ROP13, in Kunming mice. The pVAX-
243 ROP13 could induce humoral and cellular immunity against *T. gondii* [21]. The mice were
244 assessed for production of cytokines specific for Th1 (IFN- γ and IL-2) and Th2 (IL-4 and IL-
245 10) after immunization and the results showed that the protective efficacy of the DNA
246 vaccine expressing *T. gondii* ROP13 were related to Th1-driven immune response in
247 Kunming mice, confirming the importance of the cellular immune response.

248 Elimination of intracellular parasites is mainly conferred by Th1 immunity leading to the
249 secretion of cytokines interferon-gamma (IFN- γ), interleukin-12 (IL-12) and tumor necrosis
250 factor-alpha (TNF- α) [16, 18]. IL17-producing T cells termed Th17 cells secrete a set of anti-
251 microbial cytokines including IL-22 and IL-21 and also mediate host protection against
252 parasites and other pathogens [28-29]. They are involved in immunity to intracellular
253 infections including *Cryptosporidium*, *Plasmodium* spp. and *Trypanosoma cruzi* [30-31].
254 However, pathogenic roles of Th17 responses have also been reported in the context of some
255 parasitic infections [30, 32]. Among the complex network of cytokines that have been
256 described in the immune responses to *T. gondii*, the pro-inflammatory cytokine, IFN- γ was
257 shown to block intracellular development of the parasite and is considered as the main
258 mediator of resistance to *T. gondii* [33]. There is evidence indicating that IL-22 has anti-
259 parasite effects during infection with intracellular parasite, *Eimeria falciformis* that belongs to
260 the same phylum with *T. gondii* [34]. In contrast with these findings, IL-22 but not IL-17 is

261 shown to drive inflammation and tissue injury following mice infection with *T. gondii* [35].
262 Furthermore, recently published data from a mouse model of rickettsial infection, an obligate
263 intracellular bacterium, demonstrates that either Th1 or Th17 responses can have protective
264 effects. Surprisingly cells producing IL-17A or IL-22 are as protective as IFN- γ producing
265 Th1 cells, if the immunopathologic effects of TNF- α are controlled [36]. This compliments
266 recent novel findings that Th17 cells provide stronger protection, compared with Th1
267 responses, against the intracellular microorganism *T. cruzi* [32]. These findings open the
268 possibility that Th17 mediated protection during *T. gondii* is a prospect for vaccination.

269 In the present study, we found significantly raised levels of both IL-17 and IL-22 mRNA in
270 mice immunized with pcROP13 compared with control mice immediately prior to infection.
271 These elevated levels of IL-17 and IL-22 in pcROP13 immunized mice associated with lower
272 parasite burdens ($P < 0.05$) compared with PBS and pcDNA3 treated mice. This set of
273 responses also provoked secretion of specific IgG antibodies detected in the sera of mice
274 immunized with pcROP13 after the last immunization compared to control groups ($P < 0.05$).
275 Early reports supporting our data indicating a key role for IL-17 in the recruitment of
276 neutrophils which is required for resistance to *T. gondii* [37]. Neutrophils are critical for
277 successful host protection during early *T. gondii* infection [38] and experimental models have
278 demonstrated that IL-17R^{-/-} mice show significantly decreased migration of neutrophils into
279 the peritoneal cavity after *T. gondii* infection [19], indicating that neutrophil response is
280 dependent on IL-17-induced signaling. Studies on the NK response demonstrated that the
281 need for Il-6 in driving IL-17 responses against *T. gondii* was conserved between across the
282 T-cell populations [17].

283 In contrast to the perception that Th17 cells only function against extracellular pathogens, we
284 have demonstrated that Th17 effectors, IL-17 and IL-22, may be important in the defense

285 against *T. gondii* infection as conferred by a ROP13 DNA-based vaccine. Multiple subtypes
286 of innate and adaptive immune cells such as NK cells, $\gamma\delta$ and CD4 T cells have been found as
287 a source of IL-17; what subset of IL-17 producers is specifically implicated in pathogenic or
288 protective immunity to *T. gondii* remained unclear. Further research is required to achieve a
289 more detailed understanding of the exact correlates of protection against *T. gondii* infection in
290 our system. This may enable us to revise the previously described harmful effects of IL-17
291 and IL-22 producing T cells during infection with intracellular pathogens in particular *T.*
292 *gondii* infection.

293 **Acknowledgment**

294 This study was prepared from Paria Alizadeh's MSc thesis and supported by Immunology
295 research center, Tabriz University of Medical Sciences, Tabriz, Iran (Grant No. 94/2-5/7).
296 The sponsor or funding organization had no role in the design or conduct of this research.

297

298

299

300

301

302

303

304

305

306 **References**

- 307 [1] J.G. Montoya, O. Liesenfeld, Toxoplasmosis, *The Lancet* 363(9425) (2004) 1965-1976.
- 308 [2] E. Ahmadpour, A. Daryani, M. Sharif, S. Sarvi, M. Aarabi, A. Mizani, M.T. Rahimi, A. Shokri,
309 Toxoplasmosis in immunocompromised patients in Iran: a systematic review and meta-analysis, *The Journal of*
310 *Infection in Developing Countries* 8(12) (2014) 1503-1510.
- 311 [3] M. Montazeri, M. Sharif, S. Sarvi, S. Mehrzadi, E. Ahmadpour, A. Daryani, A systematic review of in vitro
312 and in vivo activities of anti-Toxoplasma drugs and compounds (2006–2016), *Frontiers in microbiology* 8
313 (2017) 25.
- 314 [4] N.-Z. Zhang, J. Chen, M. Wang, E. Petersen, X.-Q. Zhu, Vaccines against *Toxoplasma gondii*: new
315 developments and perspectives, *Expert review of vaccines* 12(11) (2013) 1287-1299.
- 316 [5] E. Hiszczyńska-Sawicka, L. Holec-Gasior, J. Kur, DNA vaccines and recombinant antigens in prevention of
317 *Toxoplasma gondii* infections--current status of the studies, *Wiadomosci parazytologiczne* 55(2) (2009) 125-
318 139.
- 319 [6] F. Azmi, A.A.H. Ahmad Fuaad, M. Skwarczynski, I. Toth, Recent progress in adjuvant discovery for
320 peptide-based subunit vaccines, *Human vaccines & immunotherapeutics* 10(3) (2014) 778-796.
- 321 [7] M. Meng, S. He, G. Zhao, Y. Bai, H. Zhou, H. Cong, G. Lu, Q. Zhao, X.-Q. Zhu, Evaluation of protective
322 immune responses induced by DNA vaccines encoding *Toxoplasma gondii* surface antigen 1 (SAG1) and 14-3-
323 3 protein in BALB/c mice, *Parasites & vectors* 5(1) (2012) 273.
- 324 [8] M.T. Rahimi, S. Sarvi, M. Sharif, S. Abediankenari, E. Ahmadpour, R. Valadan, M. Fasihi-Ramandie, S.-A.
325 Hosseini, A. Daryani, Immunological evaluation of a DNA cocktail vaccine with co-delivery of calcium
326 phosphate nanoparticles (CaPNs) against the *Toxoplasma gondii* RH strain in BALB/c mice, *Parasitology*
327 *research* 116(2) (2017) 609-616.
- 328 [9] H. Waldeland, J. Frenkel, Live and killed vaccines against toxoplasmosis in mice, *The Journal of*
329 *parasitology* (1983) 60-65.
- 330 [10] K.N. Deilami, A. Daryani, E. Ahmadpour, M. Sharif, Y. Dadimoghaddam, S. Sarvi, A. Alizadeh,
331 Excretory–secretory antigens: A suitable candidate for immunization against ocular toxoplasmosis in a murine
332 model, *Comparative immunology, microbiology and infectious diseases* 37(5-6) (2014) 369-374.
- 333 [11] H. Dlugonska, *Toxoplasma* rhoptries: unique secretory organelles and source of promising vaccine proteins
334 for immunoprevention of toxoplasmosis, *BioMed Research International* 2008 (2008).
- 335 [12] P.-Y. Wang, Z.-G. Yuan, E. Petersen, J. Li, X.-X. Zhang, X.-Z. Li, H.-X. Li, Z.-C. Lv, T. Cheng, D. Ren,
336 Protective efficacy of a *Toxoplasma gondii* rhoptry protein 13 plasmid DNA vaccine in mice, *Clinical and*
337 *Vaccine Immunology* 19(12) (2012) 1916-1920.
- 338 [13] R. Gazzinelli, Y. Xu, S. Hieny, A. Cheever, A. Sher, Simultaneous depletion of CD4+ and CD8+ T
339 lymphocytes is required to reactivate chronic infection with *Toxoplasma gondii*, *The Journal of Immunology*
340 149(1) (1992) 175-180.
- 341 [14] E.Y. Denkers, R.T. Gazzinelli, Regulation and function of T-cell-mediated immunity during *Toxoplasma*
342 *gondii* infection, *Clinical microbiology reviews* 11(4) (1998) 569-588.
- 343 [15] T. Korn, E. Bettelli, M. Oukka, V.K. Kuchroo, IL-17 and Th17 Cells, *Annual review of immunology* 27
344 (2009) 485-517.
- 345 [16] M. Munoz, O. Liesenfeld, M.M. Heimesaat, Immunology of *Toxoplasma gondii*, *Immunological reviews*
346 240(1) (2011) 269-285.
- 347 [17] S.T. Passos, J.S. Silver, A.C. O’Hara, D. Sehy, J.S. Stumhofer, C.A. Hunter, IL-6 promotes NK cell
348 production of IL-17 during toxoplasmosis, *The Journal of Immunology* 184(4) (2010) 1776-1783.
- 349 [18] J. Alves Silva, K. Rezende-Oliveira, M.V. da Silva, mez-Hern, C. ndez, sar, B. Crema Peghini, nea, N.M.
350 Silva, J. Mineo, Roberto, R. J. V. nior, IL-17-Expressing CD4+ and CD8+ T Lymphocytes in Human
351 Toxoplasmosis, *Mediators of Inflammation* 2014 (2014) 7.
- 352 [19] M.N. Kelly, J.K. Kolls, K. Happel, J.D. Schwartzman, P. Schwarzenberger, C. Combe, M. Moretto, I.A.
353 Khan, Interleukin-17/interleukin-17 receptor-mediated signaling is important for generation of an optimal

- 354 polymorphonuclear response against *Toxoplasma gondii* infection, *Infection and immunity* 73(1) (2005) 617-
355 621.
- 356 [20] J.C. Boothroyd, J.-F. Dubremetz, Kiss and spit: the dual roles of *Toxoplasma* rhoptries, *Nature reviews.*
357 *Microbiology* 6(1) (2008) 79.
- 358 [21] J.M. Turetzky, D.K. Chu, B.E. Hajagos, P.J. Bradley, Processing and secretion of ROP13: A unique
359 *Toxoplasma* effector protein, *International journal for parasitology* 40(9) (2010) 1037-1044.
- 360 [22] E. Ahmadpour, S. Sarvi, M.B.H. Soteh, M. Sharif, M.T. Rahimi, R. Valadan, M. Tehrani, A. Khalilian, M.
361 Montazeri, M. Fasihi-Ramandi, Enhancing immune responses to a DNA vaccine encoding *Toxoplasma gondii*
362 GRA14 by calcium phosphate nanoparticles as an adjuvant, *Immunology letters* 185 (2017) 40-47.
- 363 [23] Y. Dadimoghaddam, A. Daryani, M. Sharif, E. Ahmadpour, Z. Hossienikhah, Tissue tropism and parasite
364 burden of *Toxoplasma gondii* RH strain in experimentally infected mice, *Asian Pacific Journal of Tropical*
365 *Medicine* 7(7) (2014) 521-524.
- 366 [24] A. Daryani, M. Sharif, Y. Dadimoghaddam, M.B.H. Souteh, E. Ahmadpour, A. Khalilian, S. Sarvi, T.
367 Farazmand, H. Kalani, M. Rasouli, Determination of parasitic load in different tissues of murine toxoplasmosis
368 after immunization by excretory–secretory antigens using Real time QPCR, *Experimental parasitology* 143
369 (2014) 55-59.
- 370 [25] A. Xenopoulos, P. Pattnaik, Production and purification of plasmid DNA vaccines: is there scope for
371 further innovation?, *Expert review of vaccines* 13(12) (2014) 1537-1551.
- 372 [26] P. Sonaimuthu, X.T. Ching, M.Y. Fong, R. Kalyanasundaram, Y.L. Lau, Induction of protective immunity
373 against toxoplasmosis in BALB/c mice vaccinated with *Toxoplasma gondii* rhoptry-1, *Frontiers in microbiology*
374 7 (2016) 808.
- 375 [27] H.-K. Yan, Z.-G. Yuan, H.-Q. Song, E. Petersen, Y. Zhou, D. Ren, D.-H. Zhou, H.-X. Li, R.-Q. Lin, G.-L.
376 Yang, Vaccination with a DNA vaccine coding Perforin-like Protein 1 (TgPLP1) and MIC6 induces significant
377 protective immunity against *Toxoplasma gondii*, *Clinical and Vaccine Immunology* (2012) CVI. 05578-11.
- 378 [28] G.F. Sonnenberg, L.A. Monticelli, M.M. Elloso, L.A. Fouser, D. Artis, CD4+ lymphoid tissue-inducer cells
379 promote innate immunity in the gut, *Immunity* 34(1) (2011) 122-134.
- 380 [29] G. Zhao, Y. Fang, U. Ryan, Y. Guo, F. Wu, S. Du, D. Chen, Q. Lin, Dynamics of Th17 associating
381 cytokines in *Cryptosporidium parvum*-infected mice, *Parasitology research* 115(2) (2016) 879-887.
- 382 [30] T. Keswani, S. Sarkar, A. Sengupta, A. Bhattacharyya, Role of TGF- β and IL-6 in dendritic cells, Treg and
383 Th17 mediated immune response during experimental cerebral malaria, *Cytokine* 88 (2016) 154-166.
- 384 [31] G.-H. Zhao, W.-Y. Cheng, W. Wang, Y.-Q. Jia, Y.-Q. Fang, S.-Z. Du, S.-K. Yu, The expression dynamics
385 of IL-17 and Th17 response relative cytokines in the trachea and spleen of chickens after infection with
386 *Cryptosporidium baileyi*, *Parasites & vectors* 7(1) (2014) 212.
- 387 [32] C.W. Cai, J.R. Blase, X. Zhang, C.S. Eickhoff, D.F. Hoft, Th17 cells are more protective than Th1 cells
388 against the intracellular parasite *Trypanosoma cruzi*, *PLoS pathogens* 12(10) (2016) e1005902.
- 389 [33] C.D. Dupont, D.A. Christian, C.A. Hunter, Immune response and immunopathology during toxoplasmosis,
390 *Seminars in immunopathology*, Springer, 2012, pp. 793-813.
- 391 [34] J. Stange, M.R. Hepworth, S. Rausch, L. Zajic, A.A. Köhl, C. Uyttenhove, J.-C. Renauld, S. Hartmann, R.
392 Lucius, IL-22 mediates host defense against an intestinal intracellular parasite in the absence of IFN- γ at the cost
393 of Th17-driven immunopathology, *The Journal of Immunology* 188(5) (2012) 2410-2418.
- 394 [35] M. Muñoz, M.M. Heimesaat, K. Danker, D. Struck, U. Lohmann, R. Plickert, S. Bereswill, A. Fischer, I.R.
395 Dunay, K. Wolk, Interleukin (IL)-23 mediates *Toxoplasma gondii*-induced immunopathology in the gut via
396 matrixmetalloproteinase-2 and IL-22 but independent of IL-17, *Journal of Experimental Medicine* 206(13)
397 (2009) 3047-3059.
- 398 [36] K. Moderzynski, L. Heine, J. Rauch, S. Papp, S. Kuehl, U. Richardt, B. Fleischer, A. Osterloh, Cytotoxic
399 effector functions of T cells are not required for protective immunity against fatal *Rickettsia typhi* infection in a
400 murine model of infection: Role of TH1 and TH17 cytokines in protection and pathology, *PLoS neglected*
401 *tropical diseases* 11(2) (2017) e0005404.
- 402 [37] E.Y. Denkers, B.A. Butcher, L. Del Rio, S. Bennouna, Neutrophils, dendritic cells and *Toxoplasma*,
403 *International journal for parasitology* 34(3) (2004) 411-421.

404 [38] S.K. Bliss, L.C. Gavrilescu, A. Alcaraz, E.Y. Denkers, Neutrophil depletion during *Toxoplasma gondii*
405 infection leads to impaired immunity and lethal systemic pathology, *Infection and immunity* 69(8) (2001) 4898-
406 4905.
407
408
409