



Genomic Analysis of Multiresistant *Staphylococcus capitis* Associated with Neonatal Sepsis

Glen P. Carter,^a James E. Ussher,^{b,c} Anders Gonçalves Da Silva,^{a,d} Sarah L. Baines,^a Helen Heffernan,^e Thomas V. Riley,^f Roland Broadbent,^g Antje van der Linden,^c Jean Lee,^a Ian R. Monk,^a Timothy P. Stinear,^a Benjamin P. Howden,^{a,d} Deborah A. Williamson^{a,d}

^aDepartment of Microbiology and Immunology, Doherty Institute for Infection and Immunity, The University of Melbourne, Melbourne, Australia

^bDepartment of Microbiology and Immunology, University of Otago, Dunedin, New Zealand

^cSouthern Community Laboratories, Dunedin, New Zealand

^dMicrobiological Diagnostic Unit Public Health Laboratory, Department of Microbiology & Immunology, Doherty Institute, The University of Melbourne, Melbourne, Australia

^eInstitute of Environmental Science and Research, Wellington, New Zealand

^fDepartment of Microbiology, PathWest Laboratory Medicine, Perth, Western Australia, Australia

^gSchool of Medicine, University of Otago, Dunedin, New Zealand

ABSTRACT Coagulase-negative staphylococci (CoNS), such as *Staphylococcus capitis*, are major causes of bloodstream infections in neonatal intensive care units (NICUs). Recently, a distinct clone of *S. capitis* (designated *S. capitis* NRCS-A) has emerged as an important pathogen in NICUs internationally. Here, 122 *S. capitis* isolates from New Zealand (NZ) underwent whole-genome sequencing (WGS), and these data were supplemented with publicly available *S. capitis* sequence reads. Phylogenetic and comparative genomic analyses were performed, as were phenotypic assessments of antimicrobial resistance, biofilm formation, and plasmid segregational stability on representative isolates. A distinct lineage of *S. capitis* was identified in NZ associated with neonates and the NICU environment. Isolates from this lineage produced increased levels of biofilm, displayed higher levels of tolerance to chlorhexidine, and were multidrug resistant. Although similar to globally circulating NICU-associated *S. capitis* strains at a core-genome level, NZ NICU *S. capitis* isolates carried a novel stably maintained multidrug-resistant plasmid that was not present in non-NICU isolates. Neonatal blood culture isolates were indistinguishable from environmental *S. capitis* isolates found on fomites, such as stethoscopes and neonatal incubators, but were generally distinct from those isolates carried by NICU staff. This work implicates the NICU environment as a potential reservoir for neonatal sepsis caused by *S. capitis* and highlights the capacity of genomics-based tracking and surveillance to inform future hospital infection control practices aimed at containing the spread of this important neonatal pathogen.

KEYWORDS coagulase-negative staphylococci, genomics, multidrug resistance, neonates, plasmids

Coagulase-negative staphylococci (CoNS) are the most common pathogens involved in nosocomial sepsis in neonatal intensive care units (NICUs) and are often associated with the presence of intravascular devices (1). Although *Staphylococcus epidermidis* is the most frequently isolated pathogen in CoNS bacteremia in NICUs, studies have highlighted the emergence of methicillin-resistant *Staphylococcus capitis* as an important opportunistic pathogen in the NICU setting (2–5). A genetically distinct clone of *S. capitis*, designated NRCS-A by pulse-field gel electrophoresis (PFGE), has been identified as a common cause of NICU-associated bacteremia in France and more recently in the United Kingdom, Belgium, and Australia (3, 5).

Received 3 May 2018 Returned for modification 18 July 2018 Accepted 18 August 2018

Accepted manuscript posted online 27 August 2018

Citation Carter GP, Ussher JE, Da Silva AG, Baines SL, Heffernan H, Riley TV, Broadbent R, van der Linden A, Lee J, Monk IR, Stinear TP, Howden BP, Williamson DA. 2018. Genomic analysis of multiresistant *Staphylococcus capitis* associated with neonatal sepsis. *Antimicrob Agents Chemother* 62:e00898-18. <https://doi.org/10.1128/AAC.00898-18>.

Copyright © 2018 Carter et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Deborah A. Williamson, deborah.williamson@unimelb.edu.au.

G.P.C. and J.E.U. contributed equally to the work.

Like other regions, *S. capitis* has been associated with neonatal bacteremia in the NICU setting in New Zealand (NZ) and has been responsible for a sustained series of neonatal bacteremia episodes in the NICU located in Dunedin Hospital in the South Island of NZ. PFGE analysis has confirmed that the Dunedin NICU isolate is part of the NRCS-A clone of *S. capitis* (5). To date, however, the global population structure of *S. capitis* has not been comprehensively assessed using whole-genome sequencing (WGS), and the genomic relatedness of *S. capitis* isolates in NZ relative to those circulating in other NICUs around the world is unknown. Here, we describe the emergence and microevolution of *S. capitis* in NZ in the context of a globally diverse data set. Further, we identify genotypic and phenotypic factors that may contribute to the emergence and spread of *S. capitis* in the NICU setting and facilitate the success of *S. capitis* as a neonatal pathogen.

RESULTS

Population structure of NICU-associated *S. capitis* and emergence in NZ. WGS was performed on 122 *S. capitis* isolates that were collected between January 2007 and July 2016 (additional information relating to isolate collection is detailed in Supplemental Methods and Data Set S1 in the supplemental material). These included *S. capitis* isolates obtained from neonates ($n = 56$), staff ($n = 13$), and the environment ($n = 9$) in the Dunedin NICU (see below and Data Set S1). For context, the Dunedin NICU isolates were supplemented with a collection of 12 *S. capitis* isolates from adult patients in Dunedin (representing 12 adult blood culture contaminants), 21 *S. capitis* isolates from adults in two other cities in NZ (Auckland and Christchurch), and 11 *S. capitis* isolates from adults in one city in Australia (Perth) (Data Set S1). In addition, 13 publicly available *S. capitis* genomes (available at the time of this study) were included in the genomic analysis. In total, 135 *S. capitis* isolates were included in phylogenetic analyses.

A complete *S. capitis* reference genome (NZ-SC16875) was generated using PacBio single-molecule real-time (SMRT) technology. NZ-SC16875 was one of the earliest isolates collected in this study, being obtained in 2011 from a neonate in Dunedin with bacteremia. Sequence reads were mapped to NZ-SC16875, and 75,177 single nucleotide polymorphisms (SNPs) were present within the core genome and were used to construct a maximum likelihood (ML) phylogeny following filtering for recombination. There was extensive recombination among the study isolates (see Fig. S1 in the supplemental material), with recombination most evident within the ancestral branches leading to the main Bayesian Analysis of Population Structure (BAPS) groups, rather than within any of the specific groups. With respect to the NICU-associated BAPS 3 (BP3) group, recombination was inferred across large regions of the chromosome, predominantly within the first ~600 kb and the last ~1 Mb of the genome. These events generally mapped to the ancestral branch leading up to the BP3 group. This suggests, therefore, that the major lineages of *S. capitis* may have arisen following extensive recombination events.

The recombination-filtered population structure was assessed using a Bayesian hierarchical clustering approach, which identified four distinct groups (BAPS groups). NICU-associated isolates (clinical, screening, and environmental) were found predominantly within BAPS group 3 (BP3), while BAPS group 1, 2, and 4 (BP1, BP2, and BP4, respectively) isolates were most often associated with non-NICU samples (Fig. 1A). Importantly, with one exception (a staff member identified in 2014 as part of screening), isolates obtained from Dunedin NICU staff clustered predominantly within BP1 rather than BP3 (Fig. 1A), suggesting that the Dunedin NICU staff who were screened did not form a major reservoir for *S. capitis* NICU sepsis.

The NICU environment is a potential reservoir for neonatal sepsis caused by multidrug-resistant BP3 lineage *S. capitis*. To further understand the population structure and clonal dissemination of the BP3 group, a second phylogenetic analysis was performed on BP3 isolates only using the local NZ-SC16875 reference genome (Fig. 1B). Within BP3, NICU-associated isolates from NZ clustered together in a distinct clade, designated BP3.1 (Fig. 1B). Importantly, many of the NZ BP3.1 lineage isolates obtained





FIG 1 Maximum likelihood (ML) trees demonstrating the population structure of *Staphylococcus capitis*. ML trees were inferred from core-genome single nucleotide polymorphisms (SNPs) and rooted using the minimal ancestor deviation method (29). (A) ML tree of all 135 included *S. capitis* isolates, with metadata indicating source, highest level BAPS grouping, and presence/absence of antimicrobial resistance (AMR) genes (*mecA* and plasmid-carried *fusB*, *aadD*, and *qacA*) and *embp*. (B) ML tree of isolates classified as part of BAPS group 3, with metadata indicating source and BAPS grouping of isolates within BAPS group 3 (indicated in panel A).

from neonates (both blood culture isolates and screening specimens) were indistinguishable at a core-genome level from those isolates collected from the NICU environment over a 12-month period. This included isolates collected from stethoscopes and the internal surface of neonatal incubators, suggesting that the NICU environment and associated fomites might act as a possible reservoir of *S. capitis* neonatal infections.

In addition to NICU-associated isolates from Dunedin, the BP3.1 lineage contained isolates that were temporally and geographically diverse. These included isolates from adults from other cities in NZ and five international isolates from Australia, Belgium, France, the United States, and the UK which were previously associated with neonatal sepsis (Fig. 1B). However, these isolates showed marked variation in their accessory genome compared to NZ NICU isolates and did not harbor the NZ NICU-associated pSC16875 plasmid (see below).

Complete genome sequence of multidrug-resistant NICU-associated *S. capitis* NZ-SC16875. The genome of NZ-SC16875 comprised a circular chromosome of 2,516,968 bp and a single circular plasmid (termed pSC16875) of 28,248 bp. A nucleotide comparison of the chromosomes of representative draft and complete *S. capitis* genome sequences in GenBank (<http://www.ncbi.nlm.nih.gov/GenBank/>) relative to NZ-SC16875 is illustrated in Fig. 2A. Visual assessment of the alignment revealed high conservation of the genome between members of the species, with variable regions associated predominantly with mobile genetic elements (MGEs) (Fig. 2A). These MGEs

A.



B.




FIG 2 (A) Comparison of representative *S. capitis* chromosome sequences relative to *S. capitis* NZ-SC16875, illustrating a high level of conservation among the international NICU-associated *S. capitis* BP3 isolates (inner six rings). Variable regions are annotated in the outer ring. These regions (Continued on next page)

included two presumptive prophages and a 60.9-kb composite staphylococcal cassette chromosome (SCC) that displayed 99.9% nucleotide identity to the composite SCC*mec*-SCC*cad*-SCC*Cars*-SCC*cop* element previously described in NCRS-A CR01, isolated from the blood culture of an infant diagnosed with sepsis in 2007 in France (GenBank accession no. [KF049201](https://www.ncbi.nlm.nih.gov/nuccore/KF049201)) (6). This SCC*mec*-SCC*cad*-SCC*Cars*-SCC*cop* element was also present in five other publicly available NICU-associated *S. capitis* genomes from different countries (Fig. 2A). Previous work has suggested that this element is a hybrid between a 38.8-kb SCC*mec* V region of a sequence type 398 (ST398) methicillin-resistant *Staphylococcus aureus* (MRSA) isolate and a 22.2-kb SCC*cad*-SCC*Cars*-SCC*cop* region from staphylococcal species, such as *S. epidermidis* and *S. haemolyticus* (6). However, a BLAST search (<http://blast.ncbi.nlm.nih.gov>) of both the SCC*mec* region and the SCC*cad*-SCC*Cars*-SCC*cop* region against publicly available genomic data revealed a high level of identity to SCC elements in numerous other staphylococcal species, including *Staphylococcus schleiferi* and *Staphylococcus pseudintermedius* (Fig. 2B). The conservation of the large SCC*mec*-SCC*cad*-SCC*Cars*-SCC*cop* element across NICU-associated *S. capitis* isolates from separate geographic origins is suggestive of widespread clonal dissemination of a distinct *S. capitis* lineage rather than multiple independent introductions of this element into locally circulating *S. capitis* isolates. This hypothesis is in keeping with the findings of Butin et al., who recently used PFGE analysis on multiple international *S. capitis* isolates to highlight the global dissemination of the NCRS-A *S. capitis* clone (3).

NZ-SC16875 also harbored genes associated with biofilm formation, including the *icaADBC* operon, which produces polysaccharide intercellular adhesion (PIA), essential for biofilm formation in *S. epidermidis* (7). In all included genomes, the *icaADBC* region displayed >99% nucleotide identity to a previously sequenced *S. capitis* *icaADBC* operon (GenBank accession no. [JF930147](https://www.ncbi.nlm.nih.gov/nuccore/JF930147)) (8). In addition, NZ-SC16875 contained a 25.5-kb gene (*embp*) encoding the large extracellular matrix-binding protein (Embp) (9). The *embp* gene was also present in 10 publicly available genomes (Fig. 2A), five of which were obtained from NICU-associated *S. capitis* isolates. Notably, the *icaADBC* operon was present in *S. capitis* isolates from all BAPS groups, whereas *embp* was almost exclusively present in the BP3 group (Fig. 1A). This observation is in keeping with the findings of Simões et al., who recently identified the *ebh* gene (an alternative name for *embp*) as a potential virulence factor in a small number of other recently sequenced *S. capitis* isolates (10).

Characterization of the NZ-associated pSC16875 multidrug resistance plasmid.

A total of 42 NZ NICU isolates (clinical and environmental) harbored a plasmid (pSC16875) which carried antimicrobial resistance determinants, namely, *bla_Z*, *fusB*, and *aadD* (Fig. 1B). In keeping with these genotypic findings, isolates containing pSC16875 were phenotypically resistant to amoxicillin, fusidic acid, and aminoglycosides. In addition, pSC16875 also harbored the *qacA* gene, associated with reduced susceptibility to chlorhexidine (see below). Comparison with the other publicly available closed NICU-associated *S. capitis* plasmid (CRO1; GenBank accession no. [LN866850](https://www.ncbi.nlm.nih.gov/nuccore/LN866850)) revealed minimal nucleotide identity with pSC16875 (Fig. S1), suggesting independent, possibly local acquisition of this novel plasmid, or the expansion of a successful clone following its introduction into NZ. All isolates within the BP3 NICU-associated clade (in addition to other global NICU-associated *S. capitis* isolates) harbored the chromosomal *aac(6')*-*aph(2')* gene and were phenotypically resistant to gentamicin.

FIG 2 Legend (Continued)

include two putative bacteriophages (phage 1 and phage 2), the SCC*cad*-SCC*Cars*-SCC*cop* region, and the *embp* gene (B) Schematic diagram illustrating the genetic organization of the composite SCC*cad*-SCC*Cars*-SCC*cop* element in BP3 lineage *S. capitis* isolates NZ-SC16875 and CR01. Also included are the SCC*mec* V elements of *S. schleiferi* TSCC54, *S. pseudintermedius* 06-3228, *S. aureus* JCSC6944, and MRSA P126, and *S. haemolyticus* NW19A, as well as the SCC*cad*-SCC*Cars*-SCC*cop* regions of *S. haemolyticus* S167 and 5H621 and *S. simulans* FDAARGOS_124 for comparison. Green and yellow arrows indicate sequences present within SCC*mec* V regions; orange arrows are CRISPR-associated genes; and blue, purple, and red arrows indicate genes within SCC*cad*-SCC*Cars*-SCC*cop* regions. The direction of the arrows indicates the direction of transcription for open reading frames. Only coding sequences of >200 bp are shown. Shaded areas represent regions that share >99% nucleotide sequence identity (pink denotes the same orientation, and blue denotes the reverse orientation). Note that the *S. capitis* NZ-SC16875 region is shown at the top and bottom of the figure for ease of comparison.

Phenotypic properties of NZ NICU-associated *S. capitis*. The results from our genomic analyses were used to inform the assessment of phenotypic traits that may have contributed to the success of the BP3 *S. capitis* NICU lineage in NZ. First, given the finding of the *embp* gene predominantly in BP3 isolates (Fig. 1A), we compared biofilm formation in representative *S. capitis* BP3 isolates with representative isolates from the *S. capitis* BP1, BP2, and BP4 groups. Unlike BP3 and BP4 isolates, BP1 and BP2 isolates did not produce detectable biofilm (Fig. S2A). BP3 isolates produced significantly more biofilm than isolates from BP4 (mean \pm standard error of the mean [SEM] optical density at 590 nm [OD₅₉₀], 2.775 \pm 0.01 versus 0.795 \pm 0.16, respectively; $P < 0.001$) (Fig. S2A), suggesting a possible advantage for these isolates in the NICU setting, as well as more broadly.

Second, to determine the stability of pSC16875 with and without the selective pressure of antimicrobials, plasmid segregational stability was assessed. pSC16875 was stably maintained under the conditions tested, with no plasmid loss observed after 1 week of serial passage, regardless of exposure to antimicrobials (Fig. S2B). This demonstrates that the multiresistant phenotype conferred by pSC16875 carriage can be maintained in this clone even in the absence of direct antimicrobial selection (Fig. S2B).

Third, to investigate the association between the carriage of *qacA* of pSC16875 and chlorhexidine tolerance, chlorhexidine susceptibility testing was performed. Isolates that harbored *qacA* (on pSC16875) had significantly higher chlorhexidine MICs and minimum bactericidal concentrations (MBCs) than those of isolates without *qacA* (median MIC, 4 mg/liter versus 1 mg/liter, respectively, $P < 0.001$; median MBC, 4 mg/liter versus 1.5 mg/liter, respectively, $P < 0.0001$) (Fig. S2C). In addition, isolates that harbored *fusB* on pSC16875 also had significantly higher fusidic acid MICs than those without *fusB* (Fig. S2D).

Finally, none of the tested isolates displayed heterogeneous intermediate vancomycin resistance, based on criteria defined for *S. aureus*, since there are no criteria for *S. capitis* or CoNS generally.

DISCUSSION

Taken together, our analyses support the worldwide dissemination of a distinct *S. capitis* NICU-associated clone, NRCS-A, previously reported by Butin et al., which we believe has then undergone successful local adaptation in the NZ setting, predominantly through the acquisition of a multidrug-resistant plasmid (5). The global dissemination of this health care-associated clone has parallels with the emergence and local adaptation of other staphylococcal clones that are ecologically associated with the hospital environment. For example, epidemic MRSA-15 (EMRSA-15) belonging to sequence type 22 (ST22), emerged as an important cause of nosocomial bacteremia in the UK in the 1990s. It has previously been suggested that subsequent global spread of EMRSA-15 may have been facilitated by the movement of health care workers across the globe (11). In our study, the mode of transmission of *S. capitis* could not be determined, although our finding that NICU-associated isolates from multiple continents were highly related at a core-genome level (in contrast to the diverse population structure of *S. capitis* from adults) suggests global dissemination, possibly via the intercontinental movement of health care workers, in keeping with the findings of others (5). However, staff screening undertaken as part of infection control measures in Dunedin demonstrated that apart from one isolate, staff isolates were a different lineage of *S. capitis* (BP1) rather than the BP3 neonatal lineage, although it is possible that staff colonization is transient or is present in only a small number of individuals. However, given our finding of highly related *S. capitis* isolates from neonates and the NICU environment, it is also possible that the environment is an important reservoir for this multidrug-resistant clone. As such, an alternative possibility for the global dissemination of this pathogen is through point-source contamination of equipment or products used in the NICU setting. This theory has parallels with the international clonal dissemination of another pathogen, *Mycobacterium chimaera*, which has spread across the globe through contaminated heater-cooler devices used in cardiac surgery (12).

However, seven of eight positive environmental samples were from stethoscopes, while the other positive site was an incubator. These items were in contact with the skin of colonized babies; therefore, it is not possible to determine if the environment is the reservoir of *S. capitis* or if the colonized neonates have contaminated their environment. Future studies should investigate the potential reservoirs and vehicles that have permitted the global dissemination of this strain.

Another key finding from our study was the presence of the multidrug-resistant plasmid pSC16875 in NZ *S. capitis* NICU-associated isolates. The observation that pSC16875 was only found in NZ NICU isolates suggests possible acquisition in NZ, perhaps under selection pressure from local antimicrobial and antiseptic use, although we acknowledge that sample bias might be a confounding factor in this observation. Of note, NZ NICU isolates were commonly resistant to fusidic acid, mediated by plasmid-borne *fusB*. Fusidic acid is used extensively in NZ (13), and consequently, NZ has one of the highest rates of fusidic acid resistance among *S. aureus* bacteria in the developed world (14). Under selection pressure from fusidic acid, dominant clones of ST1 methicillin-susceptible *S. aureus* and ST5 MRSA have emerged in the NZ setting (14, 15). Like *S. aureus*, it is plausible that widespread use of fusidic acid in NZ has contributed to the local selection of fusidic acid-resistant *S. capitis*. pSC16875 also harbored *qacA*, which encodes a multidrug resistance efflux pump that can extrude chlorhexidine (16). The presence of *qacA* has been variably correlated with increased chlorhexidine MICs and MBCs in *S. aureus* (16), but chlorhexidine susceptibility has not previously been systematically compared in *qacA*-positive and *qacA*-negative *S. capitis* isolates. Our finding that *S. capitis* isolates with *qacA* had significantly higher chlorhexidine tolerance is of concern in the NICU setting, where chlorhexidine is often used extensively for staff and environmental decontamination. Given the plasmid colocation of *qacA* and other resistance determinants, future work should assess the impact of chlorhexidine usage on coselection for multiresistant *S. capitis* isolates. Unlike previous studies assessing antimicrobial resistance in *S. capitis* (2, 3, 17), no resistance or heterogeneous vancomycin-intermediate resistance was observed in NICU-associated *S. capitis* isolates from NZ. A previous study suggested that heterogeneous vancomycin-intermediate resistance in *S. capitis* NRCS-A isolates may be driven by local use of vancomycin (17), and it is possible that low use of vancomycin in the NICU setting in NZ, which occurs only occasionally after a CoNS isolate is cultured, has mitigated the clonal spread of heteroresistant *S. capitis*.

In addition to antimicrobial resistance, biofilm formation has also been suggested to be an important virulence characteristic of *S. capitis* in the NICU environment (8). In keeping with this suggestion, we observed significantly more biofilm formation in NICU-associated BP3 *S. capitis* isolates than in other *S. capitis* isolates. Interestingly, all study isolates had identical *icaADBC* operons, but *S. capitis* BP3 isolates contained the *embp* gene, which was absent in most non-BP3 strains. The product of *embp* (Embp) was recently demonstrated to be a multifunctional cell surface fibronectin binding protein involved in attachment to host extracellular matrix, biofilm accumulation, and evasion of host phagocytic responses in *S. epidermidis* (9). In *S. aureus*, the giant protein Ebh (encoded by *ebh*) is a homologue of Embp, and mutations in *ebh* are associated with increased susceptibility to complement-mediated killing, in addition to attenuated virulence in mice (18). It is therefore plausible that Embp production confers a selective advantage to *S. capitis* in the NICU setting. Future work, including mutagenesis studies, should be conducted to test this hypothesis.

In summary, our genotypic and phenotypic analyses of multidrug-resistant *S. capitis* strains provide valuable insights into factors contributing to the unprecedented global emergence of *S. capitis* in the NICU setting and furthermore suggest that the NICU environment might act as a reservoir for neonatal sepsis caused by this important pathogen. Importantly, compared to previous studies using PFGE alone, the higher resolution afforded by WGS in our study has enabled important differences in the accessory genomes between NICU-associated *S. capitis* strains across different geographic regions to be uncovered, and it has allowed us to explore phenotypic traits that may provide an advantage,

particularly in the local NZ NICU environment. Our phylogenomic framework will provide a valuable foundation for future studies investigating the dissemination, pathogenicity, and adaptation of this important neonatal pathogen.

MATERIALS AND METHODS

Epidemiological context of NICU-associated *S. capitis* in NZ. New Zealand is an island nation in the South West Pacific, with a population of approximately 4.4 million. Located in Dunedin (the second largest city in the South Island of NZ), Dunedin Hospital serves a catchment population of approximately 320,000 residents. Within Dunedin Hospital, there is a 16-bed NICU providing tertiary-level neonatal care. Between January 2007 and July 2016, *S. capitis* was identified in 40 of 127 (31.5%) CoNS blood culture isolates in the Dunedin Hospital NICU.

From 2010, infection control measures within the NICU were intensified to include staff undertaking a 2-min scrub with chlorhexidine on entry to the NICU, staff cleaning their arms to the elbows with an alcohol-based hand sanitizer (especially when placing arms in an incubator), and daily cleaning of ports on incubators and plastic exterior catches with benzalkonium chloride (Viraclean) wipes. In September 2013, a neonatal screening program was initiated, with axillary swabs collected from all new admissions and thereafter weekly from neonates who had previously tested negative. During the period of December 2013 to March 2015, 214 neonates were screened, and 61 (28.5%) neonates became colonized. In December 2013, the NICU moved to a new facility, with extensive cleaning of all equipment conducted prior to the move. Despite these measures, ongoing blood culture isolation of *S. capitis* and neonatal colonization episodes continued to occur. Therefore, in July and August 2014, voluntary anonymous screening of NICU staff was undertaken to investigate potential staff reservoirs. Briefly, staff were asked to swab their external ear (both inside the pinna and behind the ear) and, following hand washing with soap to remove transient skin microbiota, to make a plate impression of their thumb on a blood agar plate. As part of this staff screening program, a total of 41 isolates of *S. capitis* were identified.

In addition, between September and December 2013, a total of 40 environmental samples were collected. Environmental surfaces were swabbed; the swab was premoistened for dry surfaces, and swabs were cultured on blood agar at 36°C in 5% CO₂ for 48 h. Samples were collected from stethoscopes ($n = 13$), incubators ($n = 9$), milk warming bottles ($n = 5$), breast pumps ($n = 3$), sinks ($n = 3$), ultrasound gel ($n = 2$), weighing scales ($n = 2$), and other sources ($n = 3$). Of the 40 samples, 8 (20%) samples grew the NICU-endemic *S. capitis* strain, with 7 from 9 stethoscope samples (78%) and 1 from 9 incubator samples (11%). Two samples from the ear pieces of stethoscopes grew unrelated *S. capitis* strains.

From the above-described epidemiological investigations, a total of 21 blood culture isolates (from 2007 to 2014), 34 isolates from neonatal axillary swabs or other nonsterile site neonatal clinical specimens, 9 environmental isolates from the NICU, and 13 isolates from the staff screening (3 from the thumb imprint and 10 from the pinna) were selected for further genotypic and phenotypic analyses (Table S1); where more isolates were available, isolates were selected for sequencing to ensure coverage over time and minor differences in PFGE patterns.

Bacterial isolates and susceptibility testing. Initial identification of *S. capitis* isolates was performed using matrix-assisted laser desorption ionization–time of flight (MALDI-TOF) mass spectrometry (Bruker Daltonics, USA), and urease and maltose fermentation tests were used to differentiate between *S. capitis* subsp. *urealyticus* and *S. capitis* subsp. *capitis*. Antimicrobial susceptibility testing was performed using the Vitek 2 system (bioMérieux, Marcy l'Etoile, France), or broth microdilution for fusidic acid, and was interpreted according to CLSI (19) or EUCAST guidelines (www.eucast.org). To identify heterogeneous vancomycin-intermediate resistance, all strains underwent testing using the Etest macromethod, using previously described methods and cutoffs for screening *S. aureus* (20). The vancomycin-susceptible *S. aureus* strain ATCC 29213 was used as a negative control, and the heterogeneous vancomycin-intermediate-resistant *S. aureus* strain Mu3 was used as a positive control (21). Chlorhexidine phenotypic susceptibility testing was performed in triplicate using standard CLSI broth microdilution methods to determine chlorhexidine MICs and minimum bactericidal concentrations (MBCs) (see Supplemental Methods in the supplemental material) (22, 23).

Genomic and phylogenetic analysis. Genomic DNA extraction, library preparation, and whole-genome sequencing on a NextSeq 500 platform (Illumina, Inc.), using 2 × 150-bp chemistry, was performed as previously described (24). One isolate (designated NZ-SC16875) underwent genomic DNA extraction with the GenElute bacterial genomic DNA kit (Sigma-Aldrich), followed by sequencing on the RSII (Pacific Biosciences) with P6-C4 chemistry according to the 20-kb template preparation using the BluePippin size selection system protocol (Pacific Biosciences). Sequences were initially analyzed using the Nullarbor pipeline (T. Seeman; <https://github.com/tseemann/nullarbor>). Phage regions were detected using Phaster (25). Maximum likelihood trees were inferred using IQ-TREE (26). Recombination was detected using Gubbins (version 2.2.0-1) (27), and population structure was investigated using hierarchical Bayesian analysis with hierBAPS (28). Additional information on bioinformatic analyses is provided in the Supplemental Methods.

Biofilm assessment and assessment of plasmid segregational stability. A quantitative determination of biofilm production was performed using a 96-well microtiter plate assay (see Supplemental Methods) on a representative sampling of isolates from each *S. capitis* BP group. Where possible, every isolate within BP groups was included; however, for the larger BP groups, representative isolates were selected that provided the greatest coverage of strain variation within the BP group. To assess plasmid

stability, triplicate cultures of each test strain were grown overnight in brain heart infusion (BHI) broth, with appropriate antimicrobial selection for plasmid maintenance, followed by five daily passages under either selective or nonselective conditions. The proportion of plasmid-free cells was determined after each passage (see Supplemental Methods).

Ethical approval. All samples were collected as part of normal clinical care or as part of an infection prevention and control investigation. Access to clinical data for this study was approved by the University of Otago Human Ethics Committee (Health) (project HD16/050).

Data availability. All bacterial isolates used in this study are available upon request, and all sequence data generated have been deposited in the European Nucleotide Archive (ENA) under study accession no. [PRJNA398301](https://doi.org/10.1128/AAC.00898-18).

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/AAC.00898-18>.

SUPPLEMENTAL FILE 1, PDF file, 1.2 MB.

SUPPLEMENTAL FILE 2, XLS file, 0.1 MB.

ACKNOWLEDGMENTS

We thank the microbiology staff at Southern Community Laboratories, Dunedin, New Zealand, and the Microbiological Diagnostic Unit Public Health Laboratory, Melbourne, Australia, as well as staff and patients in the NICU at Dunedin Hospital. We thank Susan Taylor and David Murdoch for contributing isolates. We also thank the staff at the Nosocomial Infections Laboratory at the Institute of Environmental Science and Research, New Zealand.

We declare no conflicts of interest.

This study was supported by a grant from the William Sherriff Charitable Trust (to J.E.U.) and internal funding from the Department of Microbiology and Immunology at the University of Melbourne and The Institute of Environmental Science and Research, New Zealand. The National Health and Medical Research Council, Australia, funded Practitioner Fellowship GNT1105905 to B.P.H. and an Early Career Fellowship GNT1123854 to D.A.W.

REFERENCES

- Vergnano S, Menson E, Kennea N, Embleton N, Russell AB, Watts T, Robinson MJ, Collinson A, Heath PT. 2011. Neonatal infections in England: the NeonIN surveillance network. *Arch Dis Child Fetal Neonatal Ed* 96:F9–F14. <https://doi.org/10.1136/adc.2009.178798>.
- Rasigade JP, Raulin O, Picaud JC, Tellini C, Bes M, Grandi J, Ben Said M, Claris O, Etienne J, Tigaud S, Laurent F. 2012. Methicillin-resistant *Staphylococcus capitis* with reduced vancomycin susceptibility causes late-onset sepsis in intensive care neonates. *PLoS One* 7:e31548. <https://doi.org/10.1371/journal.pone.0031548>.
- Butin M, Rasigade JP, Martins-Simoes P, Meugnier H, Lemriss H, Goering RV, Kearns A, Deighton MA, Denis O, Ibrahim A, Claris O, Vandenesch F, Picaud JC, Laurent F. 2016. Wide geographical dissemination of the multiresistant *Staphylococcus capitis* NRCS-A clone in neonatal intensive-care units. *Clin Microbiol Infect* 22:46–52. <https://doi.org/10.1016/j.cmi.2015.09.008>.
- Ben Said M, Hays S, Bonfils M, Jourdes E, Rasigade JP, Laurent F, Picaud JC. 2016. Late-onset sepsis due to *Staphylococcus capitis* 'neonatalis' in low-birthweight infants: a new entity? *J Hosp Infect* 94:95–98. <https://doi.org/10.1016/j.jhin.2016.06.008>.
- Butin M, Martins-Simoes P, Rasigade JP, Picaud JC, Laurent F. 2017. Worldwide endemicity of a multidrug-resistant *Staphylococcus capitis* clone involved in neonatal sepsis. *Emerg Infect Dis* 23:538–539. <https://doi.org/10.3201/eid2303.160833>.
- Martins Simões P, Rasigade JP, Lemriss H, Butin M, Ginevra C, Lemriss S, Goering RV, Ibrahim A, Picaud JC, El Kabbaj S, Vandenesch F, Laurent F. 2013. Characterization of a novel composite staphylococcal cassette chromosome *mec* (SCC*mec*-SCC*cad/ars/cop*) in the neonatal sepsis-associated *Staphylococcus capitis* pulsotype NRCS-A. *Antimicrob Agents Chemother* 57:6354–6357. <https://doi.org/10.1128/AAC.01576-13>.
- Cue D, Lei MG, Lee CY. 2012. Genetic regulation of the intercellular adhesion locus in staphylococci. *Front Cell Infect Microbiol* 2:38. <https://doi.org/10.3389/fcimb.2012.00038>.
- Cui B, Smooker PM, Rouch DA, Daley AJ, Deighton MA. 2013. Differences between two clinical *Staphylococcus capitis* subspecies as revealed by biofilm, antibiotic resistance, and pulsed-field gel electrophoresis profiling. *J Clin Microbiol* 51:9–14. <https://doi.org/10.1128/JCM.05124-11>.
- Christner M, Franke GC, Schommer NN, Wendt U, Wegert K, Pehle P, Kroll G, Schulze C, Buck F, Mack D, Aepfelbacher M, Rohde H. 2010. The giant extracellular matrix-binding protein of *Staphylococcus epidermidis* mediates biofilm accumulation and attachment to fibronectin. *Mol Microbiol* 75:187–207. <https://doi.org/10.1111/j.1365-2958.2009.06981.x>.
- Simões PM, Lemriss H, Dumont Y, Lemriss S, Rasigade J-P, Assant-Trouillet S, Ibrahim A, El Kabbaj S, Butin M, Laurent F. Single-molecule sequencing (PacBio) of the *Staphylococcus capitis* NRCS-A clone reveals the basis of multidrug resistance and adaptation to the neonatal intensive care unit environment. *Front Microbiol* 7:1991. <https://doi.org/10.3389/fmicb.2016.01991>.
- Hart J, Christiansen KJ, Lee R, Heath CH, Coombs GW, Robinson JO. 2014. Increased EMRSA-15 health-care worker colonization demonstrated in retrospective review of EMRSA hospital outbreaks. *Antimicrob Resist Infect Control* 3:7. <https://doi.org/10.1186/2047-2994-3-7>.
- Williamson D, Howden B, Stinear T. 2017. *Mycobacterium chimaera* spread from heating and cooling units in heart surgery. *N Engl J Med* 376:600–602. <https://doi.org/10.1056/NEJMc1612023>.
- Williamson DA, Ritchie SR, Best E, Upton A, Leversha A, Smith A, Thomas MG. 2015. A bug in the ointment: topical antimicrobial usage and resistance in New Zealand. *N Z Med J* 128:103–109.
- Williamson DA, Monecke S, Heffernan H, Ritchie SR, Roberts SA, Upton A, Thomas MG, Fraser JD. 2014. High usage of topical fusidic acid and rapid clonal expansion of fusidic acid-resistant *Staphylococcus aureus*: a cautionary tale. *Clin Infect Dis* 59:1451–1454. <https://doi.org/10.1093/cid/ciu658>.
- Carter GP, Schultz MB, Baines SL, Goncalves da Silva A, Heffernan H, Tiong A, Pham PH, Monk IR, Stinear TP, Howden BP, Williamson DA. 2017. Topical antibiotic use co-selects for the carriage of mobile genetic elements conferring resistance to unrelated antimicrobials in *Staphylo-*

- coccus aureus*. Antimicrobial Agents Chemother 62:e02000-17. <https://doi.org/10.1128/AAC.02000-17>.
16. Horner C, Mawer D, Wilcox M. 2012. Reduced susceptibility to chlorhexidine in staphylococci: is it increasing and does it matter? J Antimicrob Chemother 67:2547–2559. <https://doi.org/10.1093/jac/dks284>.
 17. Butin M, Martins-Simoes P, Picaud JC, Kearns A, Claris O, Vandenesch F, Laurent F, Rasigade JP. 2015. Adaptation to vancomycin pressure of multi-resistant *Staphylococcus capitis* NRCS-A involved in neonatal sepsis. J Antimicrob Chemother 70:3027–3031. <https://doi.org/10.1093/jac/dkv217>.
 18. Cheng AG, Missiakas D, Schneewind O. 2014. The giant protein Ehb is a determinant of *Staphylococcus aureus* cell size and complement resistance. J Bacteriol 196:971–981. <https://doi.org/10.1128/JB.01366-13>.
 19. Clinical and Laboratory Standards Institute (CLSI). 2015. Performance standards for antimicrobial susceptibility testing, 26th ed. CLSI document M100S. Clinical and Laboratory Standards Institute, Wayne, PA.
 20. Walsh TR, Bolmstrom A, Qvarnstrom A, Ho P, Wootton M, Howe RA, MacGowan AP, Diekema D. 2001. Evaluation of current methods for detection of staphylococci with reduced susceptibility to glycopeptides. J Clin Microbiol 39:2439–2444. <https://doi.org/10.1128/JCM.39.7.2439-2444.2001>.
 21. Hiramatsu K, Aritaka N, Hanaki H, Kawasaki S, Hosoda Y, Hori S, Fukuchi Y, Kobayashi I. 1997. Dissemination in Japanese hospitals of strains of *Staphylococcus aureus* heterogeneously resistant to vancomycin. Lancet 350:1670–1673. [https://doi.org/10.1016/S0140-6736\(97\)07324-8](https://doi.org/10.1016/S0140-6736(97)07324-8).
 22. Clinical and Laboratory Standards Institute (CLSI). 2015. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically, 10th ed. CLSI document M07-A10. Clinical and Laboratory Standards Institute, Wayne, PA.
 23. Clinical and Laboratory Standards Institute (CLSI). 1999. Methods for determining bactericidal activity of antimicrobial agents; approved guideline. CLSI document M26-A. Clinical and Laboratory Standards Institute, Wayne, PA.
 24. Baines SL, Howden BP, Heffernan H, Stinear TP, Carter GP, Seemann T, Kwong JC, Ritchie SR, Williamson DA. 2016. Rapid emergence and evolution of *Staphylococcus aureus* clones harboring *fusC*-containing staphylococcal cassette chromosome elements. Antimicrob Agents Chemother 60:2359–2365. <https://doi.org/10.1128/AAC.03020-15>.
 25. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. Nucleic Acids Res 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.
 26. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol 32:268–274. <https://doi.org/10.1093/molbev/msu300>.
 27. Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, Parkhill J, Harris SR. 2015. Rapid phylogenetic analysis of large samples of recombinant bacterial whole genome sequences using Gubbins. Nucleic Acids Res 43:e15. <https://doi.org/10.1093/nar/gku1196>.
 28. Cheng L, Connor TR, Siren J, Aanensen DM, Corander J. 2013. Hierarchical and spatially explicit clustering of DNA sequences with BAPS software. Mol Biol Evol 30:1224–1228. <https://doi.org/10.1093/molbev/mst028>.
 29. Tria FDK, Landan G, Dagan T. 2017. Phylogenetic rooting using minimal ancestor deviation. Nat Ecol Evol 1:0193. <https://doi.org/10.1038/s41559-017-0193>.



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Carter, GP; Ussher, JE; Da Silva, AG; Baines, SL; Heffernan, H; Riley, T; Broadbent, R; van der Linden, A; Lee, J; Monk, IR; Stinear, TP; Howden, BP; Williamson, DA

Title:

Genomic Analysis of Multiresistant Staphylococcus capitis Associated with Neonatal Sepsis

Date:

2018-11-01

Citation:

Carter, GP; Ussher, JE; Da Silva, AG; Baines, SL; Heffernan, H; Riley, T; Broadbent, R; van der Linden, A; Lee, J; Monk, IR; Stinear, TP; Howden, BP; Williamson, DA, Genomic Analysis of Multiresistant Staphylococcus capitis Associated with Neonatal Sepsis, ANTIMICROBIAL AGENTS AND CHEMOTHERAPY, 2018, 62 (11)

Persistent Link:

<http://hdl.handle.net/11343/220639>

File Description:

Published version