

Vet Pathol 44:521–524 (2007)

## Characterization of a Population of Unique Granular Lymphocytes in a Bitch Deciduoma, Using a Panel of Histo- and Immunohistochemical Markers

S. ARRIGHI, F. CREMONESI, G. BOSI, D. GROPPETTI, AND A. PECILE

Department of Veterinary Sciences and Technologies for Food Safety, Laboratory of Anatomy, Faculty of Veterinary Medicine, Milan, Italy (SA,GB); Department of Veterinary Clinical Science, Section of Obstetrics and Gynecology, Faculty of Veterinary Medicine, Milan, Italy (FC,DG,AP)

**Abstract.** The ovaries and uterus were collected after ovariohysterectomy from a 16-month-old Labrador bitch in diestrus that never mated. Discrete swellings were found in the uterine horns, with the macroscopic appearance of normal early pregnancy. At histologic examination, the endometrium, devoid of any conceptus and chorion, showed a marked proliferation, on the basis of which a diagnosis of deciduoma was made. A remarkable population of stromal eosinophilic granular lymphocytes was present, especially in the axis of the endometrial folds. Periodic acid–Schiff and *Dolichos biflorus*–lectin histochemical reaction and a panel of 10 immunohistochemical markers were used to characterize eosinophilic granular cells. Our findings allowed us to compare these granular cells with the granulated decidual cells, whose presence was until now described only in primates, rodents, or a few other epitheliochorial species. On the basis of our results, the importance of eosinophilic granular cells in a decidualization process is hypothesized to occur also in the bitch.

**Key words:** Bitch; decidual cells; deciduoma; eosinophilic granular cells; histology; immunohistochemistry; uterus.

The histopathologic features of decidua-like proliferation in nonpregnant animals of many species has been referred to as deciduoma.<sup>7</sup> In human medicine, “deciduoid mesothelioma” is a term described by Nascimento et al.<sup>6</sup> for indicating a pathologic feature that bears a remarkable cytomorphologic resemblance to decidua or decidualized tissue that is localized elsewhere.


The stromal cell infiltration of this pathology has not yet been characterized in the bitch. In carnivores, indeed, the endotheliochorial or semidecidualized placentation, in the absence of a marked decidual reaction, is characterized by segmented neutrophil migration into the glandular zone of the endometrial stroma and by active proliferation of the uterine epithelium, causing an “inflammation-like” aspect of the endometrium in response to embryo implantation.<sup>12</sup> The aim of this work was to histo- and immunohistochemically characterize a peculiar stromal eosinophilic granule cell population, which was present in the axis of the endometrial folds of a naturally occurring deciduoma in a diestrus bitch.

The uterus and ovaries were collected from a 16-month-old Labrador bitch undergoing a routine ovariohysterectomy. The owner reported that the bitch was confined, and no matings had occurred during the last estrus. Clinical hematology and biochemistry collected 1 day before surgery were within normal limits. At surgery, the uterus revealed 3 thumb-sized segmental swellings in one of the horns, simulating an early pregnancy. The ovaries were typically in the luteal phase, with many corpora lutea covering the surface.

Formalin-fixed, paraffin-embedded sections (5  $\mu$ m thick) from each uterine enlargement were stained with HE for routine histopathology. At histologic examination, the uterine wall between the swellings showed the normal endometrial spiral appearance of the luteal phase, with a tall columnar epithelial lining (Fig. 1a). At the areas of gross

swelling, endometrial proliferations with long folds protruding into the lumen and intermingled with each other were evident with a lamina propria extending into their axis, indicating that they were permanent nondistensible structures. Surface epithelium lining the folds was cuboidal to columnar, with a cytoplasmic foamy appearance and bleb-like apical protrusions (Fig. 1b). The extensive branching of folds and subfolds was partially occluded and appeared to dissolve into the lumen, devoid of conceptus but containing masses of cellular debris (Fig. 1c). This structural pattern of the mucosa was shared by all the 3 uterine swellings. In some section of the swellings, hyperemic aspects were noticed, with a large quantity of blood cells in the endometrial stroma (Fig. 1d) and in the lumen of the swelling, where blood cells were mixed with debris of the mucosal folds (Fig. 1e). The macroscopic and microanatomic aspects allowed us to refer to this highly organized maternal placenta-like endometrial hyperplasia as a naturally occurring deciduoma.<sup>5</sup> For histochemical selective staining of neutral glycoconjugates, paraffin sections were processed for periodic acid–Schiff (PAS) reaction. The *Dolichos biflorus* (DBA)–lectin reactivity was tested as well, by the histochemical procedure according to Domeneghini et al.<sup>1</sup>

For immunohistochemistry, deparaffinized sections were treated with hydrogen peroxide followed by incubation in 1:20 normal goat serum (DakoCytomation, Glostrup, Denmark) in Tris-buffered saline (TBS: 0.05 M Tris/HCl, 0.15 M NaCl) for 30 minutes to prevent background prior to incubation with primary antisera. Sections were then incubated in a humidity chamber at 4°C for 18–22 hours, using a panel of primary antisera followed by peroxidase-conjugated antibodies (DakoCytomation) (Table 1). After exposure to an appropriate chromogen, the slides were counterstained with Mayer’s hematoxylin and mounted using Eukitt (Sigma, Italy).



**Table 1.** Primary antisera tested, their sources, and working dilution conditions of antigen retrieval and immunocomplex formation.

Primary Antisera Tested	Source	Code	Working Dilutions	Antigen Retrieval
Monoclonal mouse anti-human*				
CD31, endothelial cell	Dako, Denmark	N1596	Ready-to-use	90–95°C for 10'†
CD34	Dako, Denmark	M7165	1 : 50	—
CD68 (clone PG-M1)	Dako, Denmark	H-7122	Ready-to-use	90–95°C for 10'†
CD74 (clone LN-2)	Sigma, USA	C-2955	1 : 200	—
Macrophages (clone LN-5)	Sigma, USA	M-1919	1 : 400	—
Policlonal rabbit anti-‡				
Human T cell CD3	Sigma, USA	C-7930	1 : 400	Pronase 0.05%
Human immunoglobulin A	Dako, Denmark	A0262	1 : 200	Pronase 0.05%
Neuronal nitric oxide synthase (NOS1) (R-20)	Santa Cruz, USA	Sc-648	1 : 200	—
Inducible nitric oxide synthase (NOS2) (N-20)	Santa Cruz, USA	Sc-651	1 : 200	—
Endothelial nitric oxide synthase (NOS3) (N-20)	Santa Cruz, USA	Sc-653	1 : 200	—

\* Secondary antisera were goat anti-mouse immunoglobulins 1 : 25; immunocomplex was mouse Peroxidase-anti-peroxidase (PAP) 1 : 50.

† Two microwave cycles of 5 minutes each at 600W in 0.01-M citrate buffer, pH 6.0 with an intermediate step at room temperature to cool the sections.

‡ Secondary antisera were goat anti-rabbit immunoglobulins 1 : 100; immunocomplex was rabbit PAP 1 : 200.

A peculiar stromal quantity of leucocytes, especially lymphocytes and cells with cytoplasmic acidophilic granules and polymorphic, mostly bilobed nuclei, likely an unusual type of eosinophilic granular cells, appeared evident at histologic examination (Fig. 1f, g). Cytoplasmic granules stained positively either by PAS reaction (Fig. 1h), demonstrating neutral glycoconjugates, or DBA lectin (Fig. 1i), which has high selectivity for glycoconjugates

containing *N*-acetyl *D*-galactosamine in the terminal position. DBA-lectin binding was indicated as strongly characterizing the transient lymphocyte population of uterine natural killer cells (uNK) in mice.<sup>9</sup> Due to its high affinity for a specific sugar, DBA lectin is able to discriminate uNK from circulating natural killer cells (cNK), which makes it suitable as antibody reagents for phenotypic identification of uNK cells.<sup>9</sup>

←

**Fig. 1.** Canine deciduoma: histologic, histochemical, and immunohistochemical results obtained on serial sections. **Fig. 1a.** Histologic aspect of the uterine wall. Arrow indicates the point where the swelling (on the right) started, adjacent to a part of uterine horn with a normal aspect (on the left). Uterine glands (ug) have a spiral shape. HE. Bar = 500 µm. **Fig. 1b.** The marked endometrial proliferation is composed of long folds protruding into the lumen (L). Surface epithelium of the folds is composed of columnar cells, having a foamy cytoplasm and bleblike apical protrusions (*arrows*). The connective axis of the folds is filled with a large quantity of cells. HE. Bar = 50 µm. **Fig. 1c.** The mucosal folds (f) seem to dissolve into the lumen, devoid of conceptus but containing masses of cellular debris (*asterisks*). HE. Bar = 100 µm. **Fig. 1d.** Hyperemic aspects of the swellings. A large quantity of blood cells (bc) is present in the mucosal lamina propria, surrounding the sections of uterine glands (ug). HE. Bar = 100 µm. **Fig. 1e.** Higher magnification of the lumen content shown in Fig. 1d. Blood cells are present, mixed with debris of the mucosal folds (md). HE. Bar = 20 µm. **Fig. 1f, g.** Connective tissue sustaining the mucosal folds of the uterine swellings. Mixed with blood cells, a cell population is present, with cytoplasmic acidophilic granules and polymorphic, mostly bilobed nuclei (*arrows*). HE. Bar = 20 µm. **Fig. 1h.** By periodic acid–Schiff (PAS)-stain reaction, cytoplasmic granules of eosinophilic granular cells stain positively (*arrows*). Glycocalix of surface epithelial cells is also PAS-positive (*asterisks*). Bar = 20 µm. **Fig. 1i.** Cytoplasmic granules of eosinophilic granular cells show a *Dolichos biflorus* (DBA)-lectin reactivity (*arrows*). The endometrial surface epithelium (ep) is DAB-positive, too. Bar = 20 µm. (j–o) Panel of immunohistochemical reactions. Scale bar = 20 µm. **Fig. 1j.** Anti-CD 3 immunoreactivity can be seen in some eosinophilic granular cells (*arrows*) and T cells (*arrowhead*). **Fig. 1k.** A strong anti-CD 74 immunoreactivity can be seen in all the eosinophilic granular cells. **Fig. 1l.** An immunoglobulin A-immunoreactive plasma cell is indicated by the arrow. Polymorphonuclear cell populations are unreactive. **Fig. 1m.** Anti-macrophage immunoreactivity can be seen in cytoplasmic granules of eosinophilic granular cells (*arrows*). **Fig. 1n.** Anti-nitric oxide synthase II (NOSII) immunoreactivity is strongly present in the cytoplasm of polymorphonuclear cells, including eosinophilic granular cells (*arrows*). **Fig. 1o.** Anti-NOSIII immunoreactivity is weakly present in the cytoplasm of eosinophilic granular cells (*arrows*), whereas neutrophils are negative (*arrowhead*).

By immunohistochemical examination, eosinophilic granular cell appearance was that of granulated lymphocytes, characterized by a high content of lytic molecules. Anti-CD3 immunoreactivity was present in a small number of eosinophilic granular cells and in few small cells with roundish nuclei, most likely T cells (Fig. 1j). Anti-CD74 immunoreactivity was present in all of the eosinophilic granular cells, showing a strong immunoreactivity, while blood granulocytes were negative (Fig. 1k). Anti-immunoglobulin A immunoreactivity was present only in plasma cells (Fig. 1l), while polymorphonuclear cell populations were unreactive. Anti-macrophage serum immunoreacted with the granules of a large part of the eosinophilic granular cells (Fig. 1m), as well as in connective cells identifiable as macrophages. Nitric oxide synthase I (NOS I) stained some nerve fiber bundles located among the myometrial smooth musculature, especially evident in between swollen parts of the uterine wall. This nitrergic innervation, which was principally restricted to the vessel wall, seemed to be lost at the level of the lesion. Anti-NOS II immunoreactivity was strongly present in the cytoplasm of the totality of polymorphonuclear cells (Fig. 1n), without the possibility of distinguishing among eosinophilic granular cells and neutrophils. Anti-NOS III immunoreactivity was present in cytoplasmic granules of eosinophilic granular cells, whereas neutrophil granulocytes were unstained (Fig. 1o).

Our data indicate that the peculiar lymphocyte population observed in the bitch deciduoma shows similarities with the granulated decidual cells, until now described only in primates and rodents. In rodents, it is reported that the decidualization process is closely associated with a dramatic increase of macrophages and T cells, as well as with an unusual leukocyte population.<sup>4</sup> The latter is described as large lymphocytes with a reniform nucleus and prominent cytoplasmic granules. They have been immunohistochemically characterized and called large granular lymphocytes or uNK.<sup>10</sup> Yet, few reports exist about the presence of uNK in species with epitheliochorial placentation,<sup>3,8,11</sup> showing that the sow, only, shares similarities with humans and rodents, despite the noninvasive nature of the pig placenta.<sup>2</sup>

The etiopathogenesis of naturally occurring bitch deciduoma is unclear. It is possible that the reaction was induced by early pregnancy and resorption, but the case history strongly suggests that this was not the case. Suggestions have been put forward by Nuroma and Funahashi<sup>7</sup> and Nomura and Nishida<sup>8</sup> in search of the possible cause-effect relationships between pyometra and deciduoma. According to these authors, a possibility exists that canine pyometra is a kind of deciduoma induced by a naturally occurring infection, early during the luteal phase. In this article on the stromal cell population, we have identified the presence of an unusual type of granulated leukocytes, highly likely to be the granulated decidual cells described in humans and rodents.

On this basis, the presence and contribution of uterine-associated lymphocytes provided novel information about a process of decidualization that occurs as the mechanisms triggering the naturally occurring deciduoma in the bitch.

#### Acknowledgements

We thank Dr. John R. Whatts for critically revising the manuscript. This work was supported by grants from the University of Milan.

#### References

- 1 Domeneghini C, Arrighi S, Radaelli G, Bosi G, Veggetti A: Histochemical study of glycoconjugate secretions in the alimentary canal of *Anguilla anguilla* L. *Acta Histochem* **106**:477–487, 2005
- 2 Engelhardt H, Croy BA, King GJ: Evaluation of natural killer cell recruitment to embryonic attachment sites during early porcine pregnancy. *Biol Reprod* **66**:1185–1192, 2002
- 3 Engelhardt H, King GJ: Uterine natural killer cells in species with epitheliochorial placentation. *Nat Immun* **15**:53–69, 1996–97
- 4 King A: Uterine leukocytes and decidualization. *Hum Reprod Update* **6**:28–36, 2000
- 5 Koguchi A, Nomura K, Fujiwara T, Kawai Y, Okaniwa A: Maternal placenta-like endometrial hyperplasia in a beagle dog (Canine deciduoma). *Exp Anim* **44**:251–253, 1995
- 6 Nascimento AG, Keeney GL, Fletcher CD: Deciduioid peritoneal mesothelioma: an unusual phenotype affecting young females. *Am J Surg Pathol* **18**:439–445, 1994
- 7 Nomura K, Funahashi H: Histological characteristics of canine deciduoma induced by intrauterine inoculation of *E. coli* suspension. *J Vet Med Sci* **61**:433–438, 1999
- 8 Nomura K, Nishida A: Histological variations of canine deciduoma induced in non pregnant horn at different stages of unilateral pregnancy. *J Vet Med Sci* **60**:623–626, 1998
- 9 Paffaro VA Jr, Bizinotto MC, Joazeiro PP, Yamada AT: Subset classification of mouse uterine natural killer cells by DBA lectin reactivity. *Placenta* **24**:479–488, 2003
- 10 Slukvin II, Breburda EE, Golos TG: Dynamic changes in primate endometrial leukocyte populations: differential distribution of macrophages and natural killer cells at the rhesus monkey implantation site and in early pregnancy. *Placenta* **25**:297–307, 2004
- 11 Tekin S, Hansen PJ: Natural killer-like cells in the sheep: functional characterization and regulation by pregnancy-associated proteins. *Exp Biol Med* **227**:803–811, 2002
- 12 Zybina TG, Zybina EV, Kiknadze II, Zhelezova AI: Polyploidization in the trophoblast and uterine glandular epithelium of the endotheliochorial placenta of Silver fox (*Vulpes fulvus Desm.*), as revealed by DNA content. *Placenta* **22**:490–498, 2001

Request reprints from S. Arrighi, DVM, Associate Professor of Domestic Animals Anatomy, Department of Veterinary Sciences and Technologies for Food Safety-Laboratory of Anatomy, Faculty of Veterinary Medicine, Via Trentacoste, 2I-20134 Milan (Italy). E-mail: silvana.arrighi@unimi.it.