

## ACTIVATION OF THE ORPHAN NUCLEAR RECEPTOR ROR $\alpha$ COUNTERACTS THE PROLIFERATIVE EFFECT OF FATTY ACIDS ON PROSTATE CANCER CELLS: CRUCIAL ROLE OF 5-LIPOXYGENASE

Roberta M. MORETTI<sup>1</sup>, Marina MONTAGNANI MARELLI<sup>1</sup>, Angelo SALA<sup>2</sup>, Marcella MOTTA<sup>1</sup> and Patrizia LIMONTA<sup>1\*</sup>

<sup>1</sup>Center for Endocrinological Oncology, Department of Endocrinology, University of Milano, Milano, Italy

<sup>2</sup>Department of Pharmacological Sciences, University of Milano, Milano, Italy

**The incidence of prostate carcinoma is very low in Eastern countries, such as Japan, suggesting that life style conditions may play a crucial role in the development of this pathology. Dietary  $\omega$ -6 polyunsaturated fatty acids, such as linoleic (LA) and arachidonic (AA) acids, have been shown to stimulate the proliferation of prostate cancer cells after being converted into 5-HETE by means of the 5-lipoxygenase (5-LOX) pathway. Blockade of 5-LOX activity has been proposed as an attractive target for the prevention of the mitogenic action of dietary fats on prostate cancer. The 5-LOX gene has been shown to carry a response element for the orphan nuclear receptor ROR $\alpha$  (for its ROR $\alpha$ 1 isoform in particular) in its promoter region. We attempt to clarify whether activation of ROR $\alpha$  might modulate the expression of 5-LOX, thus interfering with the mitogenic activity of fatty acids in prostate cancer cells. We show that in androgen-independent DU 145 prostate cancer cells, LA, AA and their metabolite 5-HETE exert a strong stimulatory action on cell proliferation. This effect is completely counteracted by the simultaneous treatment of the cells with a non redox inhibitor of 5-LOX activity. We then demonstrate that: i) ROR $\alpha$ , and specifically its ROR $\alpha$ 1 isoform, is expressed in DU 145 cells; ii) activation of ROR $\alpha$ , by means of the thiazolidinedione derivative CGP 52608 (the synthetic ROR $\alpha$  activator), significantly reduces 5-LOX expression, both at mRNA (as evaluated by comparative RT-PCR) and at protein (as investigated by Western blot analysis) level (this was confirmed by the reduced activity of 5-LOX in CGP 52608 treated cells); and iii) the treatment of DU 145 cells with CGP 52608 completely abrogated the proliferative action of both LA and AA. These results have been confirmed in another androgen-independent prostate cancer cell line (PC3). Our data indicate that, by decreasing the expression of 5-LOX, activation of ROR $\alpha$  might interfere with the mitogenic activity of fatty acids on prostate cancer. We have shown previously that CGP 52608 reduces the proliferation and the metastatic behavior of DU 145 cells. These observations indicate that the orphan nuclear receptor ROR $\alpha$  might be considered as a molecular target for the development of new chemopreventive or chemotherapeutic strategies for prostate carcinoma.**

© 2004 Wiley-Liss, Inc.

**Key words:** prostate cancer; ROR $\alpha$ ; 5-lipoxygenase; linoleic acid; arachidonic acid

Prostate cancer incidence is particularly high in the United States and Europe and much lower in Japan and in other Asian countries.<sup>1</sup> In addition to circulating levels of androgens, environmental factors, such as dietary fats, have been suggested to be responsible for this difference.<sup>2–4</sup> Among polyunsaturated fatty acids, the  $\omega$ -6 acids that are present in animal fat and in vegetable oils, have been shown to enhance the risk of prostate cancer occurrence. On the contrary,  $\omega$ -3 acids, that can be found in high amounts in fish oils, seem to exert a protective action against the growth of the tumor.<sup>5–7</sup>

Linoleic acid (LA), a member of the  $\omega$ -6 polyunsaturated fatty acid family, is converted to arachidonic acid (AA), which is then incorporated into the phospholipids of the cell membrane. Arachidonic acid is in turn metabolized to the series of eicosanoids after having been released from phospholipids by PLA2.<sup>8,9</sup> Eicosanoids have been reported to mediate the mitogenic activity of both LA and AA on different tumor cell lines.<sup>5,10–14</sup>

In prostate cancer, AA has been shown to stimulate cell proliferation mainly through its conversion to 5-HETE (5-hydroxyeicosatetraenoic acid), via the 5-lipoxygenase (5-LOX) pathway.<sup>15–17</sup> Moreover, 5-LOX has been reported recently to be overexpressed in prostate carcinoma.<sup>18</sup> Taken together, these observations led to the suggestion that 5-LOX might be considered as an attractive target for the development of new chemopreventive or chemotherapeutic strategies.

ROR $\alpha$  is a transcription factor belonging to the family of orphan nuclear receptors, for which currently endogenous ligands are unknown.<sup>19,20</sup> The gene for the human ROR $\alpha$  receptor encodes at least 4 distinct isoforms: ROR $\alpha$  1, 2, 3, and 4 (also named RZR $\alpha$ ).<sup>21,22</sup> The 4 splicing variants share common DNA- and putative ligand-binding domains, but possess distinct amino-terminal domains.<sup>22</sup> ROR $\alpha$  receptors bind preferentially to response elements (ROREs) composed of a typical hexameric core-binding site RGGTCA (R = A or G), and an A/T rich 5'-flanking sequence.<sup>21,22</sup> By binding to ROREs in the promoter region of target genes, ROR $\alpha$  has been shown to be involved in the control of several physiological processes, such as cerebellar development,<sup>23,24</sup> muscle cell differentiation,<sup>25</sup> bone metabolism<sup>26</sup> and immune functions.<sup>27</sup>

A RORE sequence has also been identified in the promoter region of the human 5-LOX gene. In particular, this RORE specifically binds ROR $\alpha$ 1 and RZR $\alpha$ , but not ROR $\alpha$ 2 and 3.<sup>28</sup> Our experiments attempt to verify whether in prostate cancer cells, the activation of ROR $\alpha$  might interfere with the mitogenic action of fatty acids through the modulation of 5-LOX expression. We used the thiazolidinedione derivative CGP 52608, which has been shown previously to activate the nuclear receptor.<sup>29,30</sup>

### MATERIAL AND METHODS

#### Materials

The thiazolidinedione derivative CGP 52608 (1-[3-allyl]-4-oxo-thiazolidine-2-ylidene]-4-methyl-thisemcarbazon) was kindly donated by Dr. D. Evans (Novartis, Basel, Switzerland). This compound has been demonstrated previously to activate the orphan nuclear receptor ROR $\alpha$ .<sup>29,30</sup> Linoleic acid (LA), arachidonic acid (AA), 5-hydroxyeicosatetraenoic acid (5[S]-HETE and 5[R]-HETE) and delipidized bovine serum albumin (BSA) were purchased from Sigma Chemical (St. Louis, MO). MK886, the inhibitor of 5-lipoxygenase activating protein (FLAP), was obtained from Merck Frosst (Quebec, Canada).

Grant sponsor: COFIN; Grant sponsor: EC.

\*Correspondence to: Center for Endocrinological Oncology, Department of Endocrinology, Via Balzaretti 9, 20133 Milano, Italy.  
Fax: +39-02-50318204. E-mail: patrizia.limonta@unimi.it

Received 1 April 2003; Accepted 7 April 2004

DOI 10.1002/ijc.20387  
Published online 2 June 2004 in Wiley InterScience (www.interscience.wiley.com).

### Cell culture

The human androgen-independent DU 145 and PC3 prostate cancer cell lines were obtained from American Type Culture Collection (Rockville, MD). Cells were routinely grown in RPMI-1640 medium (Seromed Biochrom, Berlin, Germany) supplemented with 5% fetal bovine serum (FBS) (Gibco, Paisley, Scotland, UK), glutamine (1 mM) and antibiotics (100 U/ml penicillin G sodium, 100 µg/ml streptomycin sulphate) in a humidified atmosphere of 5% CO<sub>2</sub>/95% air.

### Cell proliferation studies

**Effects of fatty acids and 5-HETEs on cell proliferation.** To study the effects of fatty acids and 5-HETEs on cell proliferation, DU 145 cells were plated at a density of  $2 \times 10^4$  cells/plate in RPMI-1640 supplemented with 5% FBS. Forty-eight hours after plating, the medium was changed to RPMI-1640 supplemented with 0.5% FBS and delipidized BSA (5 µM). Cells were then treated with LA ( $10^{-8}$ – $10^{-5}$  M), AA ( $10^{-8}$ – $10^{-6}$  M), 5(S)-HETE or 5(R)-HETE ( $5 \times 10^{-8}$ – $10^{-6}$  M) for 7 days. Fatty acids and 5-HETEs were pre-complexed with lipid-free BSA before addition to the medium. Final concentration of BSA was maintained at 5 µM throughout the experiments. Control cells were treated with the 0.5% FBS added medium containing BSA only. At the end of the treatment, cells were harvested and counted by hemocytometer.

To confirm the role of 5-LOX in the growth modulatory activity of fatty acids, DU 145 cells were treated, for 7 days, with AA ( $10^{-7}$  M), either in the absence or in the presence of the 5-LOX inhibitor MK886 ( $10^{-7}$ – $10^{-5}$  M). Among the biochemical inhibitors of 5-LOX, this compound has been shown to act by interacting with FLAP and interfering with the presentation of AA to 5-LOX enzyme at the nuclear envelope membrane.<sup>31</sup>

**Effects of RORα activation on the mitogenic activity of fatty acids.** DU 145 cells were treated as described above, either with LA ( $10^{-6}$  M) or AA ( $10^{-7}$  M) in the absence or in the presence of CGP 52608 ( $10^{-7}$  M). Cells were harvested and counted after 7 days of treatment. All the cell proliferation experiments were repeated at least 3 times. Five plates/dose were included in each experiment.

### RT-PCR for the RORα isoforms

The specific isoform of the RORα receptor that might be expressed in DU 145 cells was investigated by RT-PCR. After phenol-chloroform extraction, 1 µg of total RNA from cancer cells was used in a reverse transcription reaction. The same RNA has been used in all the RT-PCR reactions. cDNA synthesis was carried out using the Gene AMP kit (Perkin Elmer Cetus, Norwalk, CT), with an oligo(dT)<sub>16</sub> as a primer for the reverse transcriptase. Samples containing cDNAs were then amplified in a 100 µl solution containing PCR buffer (50 mM KCl, 10 mM Tris-HCl), 2 mM MgCl<sub>2</sub>, 15 pmol of a pair of specific primers and 2.5 U Taq Polymerase. Thirty-five cycles of amplification were carried out in a programmable heat block (Perkin Elmer Cetus) (1-min denaturation at 94°C, 45-sec primer annealing at 60°C and 2-min primer extension at 72°C). The sense primers were: 5'-AAACATG-GAGTCAGCTCCG-3' for RORα1; 5'-CTCCAAATACTCCAT-CAGTGTATCC-3' for RORα2; 5'-CAACTTGAGCACATA-AACTGG-3' for RORα3; 5'-TGTATTTTGTGATCGCAGAG-3' for RORα4. For all the isoforms, the antisense primer used was: 5'-CATAAAGCTGTCTCTCTGC-3'.<sup>28</sup> The same RT-PCR conditions were used to amplify β-actin cDNA; in this case the primers used were 5'-TGACGGGGTCACCCACACTGTGC-CCATCTA-3' (sense) and 5'-CTAGAAGCATTTGCGGTGGAC-GATGGAGGG-3' (antisense).<sup>32</sup> After RT-PCR, the amplified DNA products were separated on a 1.5% agarose gel and stained with ethidium bromide.

### Immunoprecipitation and western blot analysis for RORα1

To confirm the expression of the RORα1 isoform at the protein level, DU 145 cells were harvested and solubilized in RIPA buffer

(50 mM Tris-HCl, pH 7.7), 150 mM Na<sub>3</sub>VO<sub>4</sub>, 50 mM NaF, 0.3 mM phenylmethylsulfonyl fluoride and 5 mM iodoacetic acid. RORα has been immunoprecipitated with 10 µg/10 µl of a polyclonal antibody against the RORα1 receptor (sc-6062, Santa Cruz Biotechnology, Santa Cruz, CA) and electrophoresed on 10% polyacrylamide gel under reducing conditions. Precipitated proteins were transferred onto a nitrocellulose filter, which was then probed with the sc-6062 antibody for 2 hr at room temperature. Filters were then incubated for 1 hr at room temperature with an anti-goat IgG (1:10,000). Antibody bound to RORα1 was detected using the Supersignal detection system (Celbio, Pierce, Rockford, IL) after a 5–10-min exposure to a Hyperfilm X-ray film (Amersham Pharmacia Biotech, Milano, Italy) at room temperature.

### RT-PCR for 5-LOX expression

In preliminary experiments, RT-PCR was carried out to choose the adequate number of cycles for amplification of 5-LOX and β-actin (as an internal control) cDNAs. To this purpose, RNA (1 µg) from DU 145 cells was reverse transcribed and cDNAs were amplified by PCR as described above. The primers used for 5-LOX were: 5'-CACTGACGACTACATCTACC-3' (sense) and 5'-GT-TCCACTCCATCCATCGAT-3' (antisense);<sup>28</sup> the primers for β-actin were the same as described above.<sup>32</sup> PCR reactions consisted of 1-min denaturation at 94°C, 30-sec annealing at 60°C and 2-min primer extension at 72°C. The number of cycles carried out were 30–45 for 5-LOX and 15–30 for β-actin cDNAs, respectively. After RT-PCR, the amplified cDNAs were separated on an agarose gel, stained with ethidium bromide and photographed under UV light.

To study the effects of RORα activation on the expression of 5-LOX, DU 145 cells were plated in 100 mm-dishes at a density of  $8 \times 10^5$  cells/dish. Twenty-four hours after plating, the media were changed to experimental media (RPMI-1640 supplemented with 5% FBS) either in the absence or in the presence of CGP 52608 ( $10^{-7}$  M) for 6, 12, 24, 36, 48 and 60 hr. At the end of the treatment, cells were harvested and RNAs were extracted and submitted to RT-PCR at adequate cycles. Equal RT reaction solutions were utilized in PCR reaction in the presence of the specific primers for 5-LOX and for β-actin.

### Western blot analysis of 5-LOX

To study the effects of RORα activation on the level of 5-LOX protein, DU 145 cells were plated at a density of  $8 \times 10^4$  cells/plate. Twenty-four hours later, the media were changed to experimental media and cells were treated daily with CGP 52608 ( $10^{-7}$  M) for 4 days (96 hr). At the end of the treatment, cells were harvested and protein fractions were prepared as described for the RORα protein. Proteins were electrophoresed and transferred onto a nitrocellulose filter. Filters were probed with the rabbit polyclonal 5-LOX antibody (LO32) (kindly provided by Dr. R.N. Young, Merk Frosst) for 2 hr at room temperature. Filters were then incubated for 1 hr at room temperature with an anti-rabbit IgG (1:10,000; Santa Cruz Biotechnology, Santa Cruz, CA). Antibody bound to 5-LOX was detected using the Supersignal detection system, as described for the RORα protein.

### Evaluation of 5-LOX enzymatic activity

5-LOX activity was evaluated after the formation of 5(S)-HETE upon incubation of DU 145 cells with AA and the calcium ionophore A23187, after a daily treatment, for 4 days (96 hr), with CGP 52608 ( $10^{-7}$  M) or with its vehicle. Briefly, cells were washed and incubated with 1 ml of PBS containing 1 µM of AA and the calcium ionophore A23187 (2 µM) for 5 min at 37°C. After centrifugation at 12,000g, 4°C for 5 min, supernatants were loaded onto a solid phase extraction cartridge (1 ml OASIS, Waters, Milford, MA), preactivated with 1 ml methanol and 1 ml of water. Arachidonic acid metabolites were eluted using 1 ml of methanol/acetonitrile (50/50, v:v) and taken to dryness using a SpeedVac Rotary Evaporator (Savant, Farmingdale, NY). After reconstitution in 200 µl of HPLC mobile phase (see below) samples were

injected into a Beckman Liquid Chromatograph mod 110B (Beckman Analytical, Palo Alto, CA). Separation of mono-HETEs was carried out on Ultrasphere ODS column ( $4.6 \times 250$  mm,  $5 \mu\text{m}$ , Beckman Analytical) eluted isocratically at 1 ml/min using a mobile phase of acetonitrile:water:acetic acid 60:40:0.01, pH 5 with ammonium hydroxide. UV absorbance was monitored at 236 nm with a Beckman diode-array UV Detector and full scans (220–300 nm) were acquired every 2 sec. Retention time of mono-HETEs was determined using synthetic 5(S)-HETE and 15(S)-HETE.

#### Experiments in PC3 cells

Experiments were carried out in PC3 cells, to confirm the effects of ROR $\alpha$  activation on the proliferative action of fatty acids in a second experimental model of prostate cancer cells. In preliminary studies, PC3 cells were treated with CGP 52608 ( $10^{-11}$ – $10^{-6}$  M), to confirm the antiproliferative effects of the thiazolidinedione derivative on androgen-independent prostate cancer cells, as previously reported for DU 145 cells.<sup>33</sup> PC3 cells were then treated with AA ( $10^{-8}$  M– $5 \times 10^{-6}$  M), harvested and counted after 7 days of treatment. Finally, PC3 cells were treated with AA ( $5 \times 10^{-6}$  M), either in the absence or in the presence of CGP52608 ( $10^{-7}$  M) (as described for DU 145 cells) and counted after 7 days of treatment.

#### Statistical analysis

Data from cell proliferation studies were analyzed according to the Dunnett's test after one-way analysis of variance.<sup>34</sup>

## RESULTS

#### Effects of fatty acids and 5-HETE on DU 145 cell proliferation


Figure 1a shows that LA significantly stimulates cell proliferation in the range of doses  $5 \times 10^{-7}$ – $5 \times 10^{-6}$  M. AA exerted its mitogenic action on DU 145 cells at the concentrations of  $5 \times 10^{-7}$  M and  $10^{-7}$  M (Fig. 1b). Finally, 5(S)-HETE induced a significant increase of cell growth when used at the dose of  $5 \times 10^{-7}$  M (Fig 1c). 5(R)-HETE did not modify DU 145 cell proliferation at any dose used (data not shown). DU 145 cells were then treated with AA ( $10^{-7}$  M) either in the absence or in the presence of the 5-LOX inhibitor MK886 ( $10^{-7}$ – $10^{-5}$  M). Figure 2 shows that MK886 counteracts the stimulatory action of AA on prostate cancer cells in a dose-dependent way, being significantly effective at  $5 \times 10^{-6}$  and  $10^{-6}$  M. These results confirm the proliferative effect of fatty acids on prostate cancer cells and the crucial role played by 5-LOX in this activity.

#### Expression of the ROR $\alpha$ 1 isoform in DU 145 cells

The 5-LOX gene carries a ROR $\alpha$ 1 response element on its promoter region.<sup>28</sup> We have shown previously that the ROR $\alpha$  family is expressed in DU 145 prostate cancer cells.<sup>33</sup> Experiments have now been carried out to identify the specific isoform of the orphan nuclear receptor that might be present in these cells. By RT-PCR, we have observed that only one cDNA band, corresponding to the ROR $\alpha$ 1 cDNA, was present in DU 145 cells (Fig. 3a, upper panel, lane 1). No band corresponding to ROR $\alpha$ 2, 3, or 4 cDNAs could be obtained (Fig. 3a, upper panel, lanes 2–4).  $\beta$ -Actin cDNAs were amplified in all the 4 samples, indicating that RNA was not degraded (Fig. 3a, lower panel, lanes 1–4). To confirm the expression of ROR $\alpha$ 1 in prostate cancer cells, the receptor was first immunoprecipitated and then submitted to gel electrophoresis and to Western blot analysis. As shown in Figure 3b, a major protein band with molecular weight of about 80 kDa, recognizing the sc-6062 antibody (Santa Cruz Biotechnology), was present in DU 145 cells. These results indicate that ROR $\alpha$ 1 is the major isoform of the receptor to be expressed in these cells.

#### Effects of ROR $\alpha$ activation on 5-LOX expression

Because the 5-LOX gene carries a RORE in its promoter region, we reasoned that the activation of the nuclear receptor might affect the expression of the enzyme. We verified this hypothesis both at




**FIGURE 1** – Effects of linoleic acid (LA) (a), arachidonic acid (AA) (b), and 5(S)-hydroxyeicosatetraenoic acid, 5(S)-HETE (c) on DU 145 cell proliferation. Each experiment was repeated 3 times with identical results. Data are expressed as the mean cell number/plate  $\pm$  SEM. \* $p < 0.05$  vs. controls (C).


the mRNA (comparative RT-PCR analysis) and at the protein (Western blot analysis) level.

**RT-PCR analysis.** In preliminary experiments, we carried out RT-PCR analysis at various cycles to choose the adequate number of cycles for the detection of the mRNA expression of 5-LOX and of  $\beta$ -actin (as an internal standard) mRNA expression. The intensity of each band increased in parallel with the increasing number of PCR cycles (data not shown). We concluded that the adequate number of cycles was 35 for 5-LOX and 20 for  $\beta$ -actin.

To study the effects of ROR $\alpha$  activation on the expression of 5-LOX, DU 145 cells were treated with the thiazolidinedione derivative CGP 52608 ( $10^{-7}$  M), the exogenous ROR $\alpha$  ligand and activator,<sup>29,30</sup> and RNA was extracted at different time intervals (from 6–60 hr) for comparative RT-PCR. As shown in Figure 4, in control DU 145 cells, changing of the medium at the beginning of the experiments induces an increase of the expression of 5-LOX mRNA at 36, 48 and 60 hr. It is well known that FBS contains fatty acids and these have been shown to increase the expression of the




**FIGURE 2** – Effects of the 5-LOX inhibitor MK 886 on the stimulatory activity of AA ( $10^{-7}$  M) on DU 145 cell proliferation. Each experiment was repeated 3 times with identical results. Data are expressed as the mean cell number/plate  $\pm$  SEM. \* $p < 0.05$  vs. controls (C); \*\* $p < 0.05$  vs. AA.




**FIGURE 3** – Expression of the different ROR $\alpha$  isoforms in DU 145 cells. (a) RT-PCR analysis of the different isoforms of ROR $\alpha$  and of  $\beta$ -actin. (Upper panel) Ethidium bromide-stained gel of the amplified cDNAs obtained in the presence of oligonucleotide primers specific for ROR $\alpha$ 1 (lane 1), ROR $\alpha$ 2 (lane 2), ROR $\alpha$ 3 (lane 3), and ROR $\alpha$ 4 (lane 4). (Lane 5) RT-PCR control. (Lower panel) Ethidium bromide-stained gel of the amplified cDNAs obtained in the presence of oligonucleotide primers specific for  $\beta$ -actin. (b) Western blot analysis of ROR $\alpha$  after immunoprecipitation using the sc-6062 antibody.

enzyme.<sup>15–17</sup> Treatment of the cells with CGP 52608 completely counteracted the increase at all the time intervals (Fig. 4). As expected, the level of expression of  $\beta$ -actin was not modified either in controls or in treated cells at any time interval considered (Fig. 4).

**Western blot analysis.** The effects of ROR $\alpha$  activation on the level of the 5-LOX protein were evaluated at 96 hr after the treatment of DU 145 cells with the thiazolidinedione derivative ( $10^{-7}$  M), by using a specific antibody against the enzyme. As shown in Figure 5, the treatment resulted in a decreased level of expression of the protein, when compared to that found in untreated controls. Treatment with CGP 52608 for shorter time intervals did not induce any significant variation in 5-LOX levels (data not shown). From these results, it seems that the decrease of 5-LOX levels at the protein level is more pronounced than that observed at the mRNA level. It must be underlined, however, that for the RT-PCR experiments the cells received only one treatment at the beginning of the observation period. Cells were treated daily for 4 days for the Western blot analysis.



**FIGURE 4** – Effects of CGP 52608 on 5-LOX mRNA expression in DU 145 cells. Cells were treated with CGP 52608 ( $10^{-7}$  M); at the indicated time intervals, total RNA was extracted and analyzed by RT-PCR at the appropriate cycles, as described in Material and Methods, for 5-LOX and  $\beta$ -actin. One representative of 3 experiments is reported.




**FIGURE 5** – Effects of CGP 52608 on 5-LOX protein expression in DU 145 cells. Cells were treated daily, for 4 days (96 hr), with CGP 52608 ( $10^{-7}$  M); protein extracts were then prepared and submitted to Western blot analysis, as described in Material and Methods. One representative of 3 experiments is reported. Lane 1, controls; lane 2, CGP 52608.

#### Effects of ROR $\alpha$ activation on 5-LOX enzymatic activity

Incubation of DU 145 cells with AA in the presence of the calcium ionophore A23187 resulted in the formation of 5(S)-HETE, as assessed by retention time using isocratic RP-HPLC separation (Fig. 6a,b) and UV spectral analysis (Fig. 6b, inset). Treatment of cells with CGP 52608 ( $10^{-7}$  M), the exogenous ROR $\alpha$  ligand and activator, showed a marked decrease ( $68 \pm 6\%$ , mean  $\pm$  SD) of the peak corresponding to 5(S)-HETE (Fig. 6c), supporting the hypothesis that 5-LOX mRNA, protein and activity are all downregulated by ROR $\alpha$  activation.

#### Effects of ROR $\alpha$ activation on the mitogenic activity of fatty acids

Because fatty acids are mitogenic on prostate cancer cells through their conversion into 5-HETE by 5-LOX activity, and because ROR $\alpha$  activation significantly reduces 5-LOX expression in DU 145 cells, we reasoned that CGP 52608 might counteract the proliferative action of fatty acids on these cells. DU 145 cells were




**FIGURE 6**—Effects of CGP 52608 on 5-LOX enzymatic activity in DU 145 cells. Cells were treated daily, for 4 days (96 hr) with CGP 52608 ( $10^{-7}$  M); then, they were washed and incubated with AA (1  $\mu$ M) and A23187 (2  $\mu$ M) for 5 min at 37°C. Supernatants were analyzed by RP-HPLC as described in Material and Methods. (a) Chromatographic tracing of UV absorbance at 236 nm from standard 15(S)-HETE and 5(S)-HETE. (b) Chromatographic tracing of UV absorbance at 236 nm from control DU 145 cells; inset shows the full UV spectrum of chromatographic peak eluting at the retention time of standard 5(S)-HETE. (c) Chromatographic tracing of UV absorbance at 236 nm from DU 145 cells treated with CGP 52608 ( $10^{-7}$  M) for 96 hr.

treated with LA ( $10^{-6}$  M) or AA ( $10^{-7}$  M), either in the absence or in the presence of CGP 52608 ( $10^{-7}$  M). As expected, we found that CGP 52608 completely counteracts the stimulatory effect of both LA (Fig. 7a) and AA (Fig. 7b).

#### Experiments in PC3 cells

In preliminary experiments we have shown that the ROR $\alpha$ 1 isoform of the nuclear receptor is expressed in PC3 cells, both at mRNA and at protein levels (data not shown). The treatment of PC3 cells with CGP 52608 dose-dependently inhibited PC3 cell




**FIGURE 7**—Effects of CGP 52608 ( $10^{-7}$  M) on the stimulatory activity of LA ( $10^{-6}$  M) (a) and of AA ( $10^{-7}$  M) (b) on DU 145 cell proliferation. Each experiment was repeated 3 times with identical results. Data are expressed as the mean cell number/plate  $\pm$  SEM. \* $p < 0.05$  vs. controls (c); \*\* $p < 0.05$  vs. LA (a) or AA (b).

proliferation, as previously described by our group for DU 145 cells (Fig. 8a).<sup>33</sup> AA stimulated PC3 cell growth, although at a higher dose ( $5 \times 10^{-6}$  M) than that required to increase DU 145 cell proliferation (Fig. 8b). This dose corresponds to that previously reported by other authors to be required for AA to maximally stimulate PC3 cell growth.<sup>15</sup>

Finally, PC3 cells were treated with AA ( $5 \times 10^{-6}$  M), either in the absence or in the presence of CGP 52608 ( $10^{-7}$  M). The thiazolidinedione derivative completely counteracted the stimulatory effect of AA (Fig. 8c), confirming the results obtained in DU 145 cells.

#### DISCUSSION

Dietary fatty acids, and in particular  $\omega$ -6 polyunsaturated fatty acids, are associated with an increased risk of the tumor of the prostate.<sup>7</sup> Our data supports this observation by showing that both LA and AA exert a significant proliferative effect on prostate cancer cells. This is in line with those data reported previously on PC3 cells, as well as on LNCaP androgen-dependent prostate cancer cells.<sup>5,15</sup> At variance with our data, Rose and Connolly<sup>5</sup> did not observe any effect of LA on DU 145 cell growth. At the



**FIGURE 8** – Effects of CGP 52608 ( $10^{-11}$ – $10^{-6}$  M) alone (a), of AA ( $10^{-8}$ – $5 \times 10^{-6}$  M) alone (b), and of CGP 52608 ( $10^{-7}$  M) either alone or in the presence of AA ( $5 \times 10^{-6}$  M) (c) on PC3 cell proliferation. Each experiment was repeated 3 times with identical results. Data are expressed as the mean cell number/plate  $\pm$  SEM. \* $p < 0.05$  vs. controls (c); \*\* $p < 0.05$  vs. AA.

moment, the reason for this discrepancy is not clear; however it might be related to methodological differences (e.g., cell culture conditions, length of the treatment, etc.) present in the 2 studies.

Our results, together with previous reports<sup>15–17</sup> also confirm that both LA and AA, to exert their mitogenic effect, need to be metabolized through the 5-lipoxygenase pathway. Therefore, modulation of 5-lipoxygenase expression or activity has been suggested as a possible target for the development of new chemopreventive or chemotherapeutic strategies for prostate cancer.<sup>17</sup>

We have stated that the promoter region of 5-LOX contains a response element for the orphan nuclear receptor ROR $\alpha$ 1,<sup>28,35</sup> suggesting that the expression of the enzyme might also be regulated through the activation of the nuclear receptor. We

have shown that this specific isoform of the receptor is expressed in DU 145 cells, both at the mRNA and at the protein level. We have demonstrated that 5-LOX expression is induced by FBS added medium; it is actually well known that FBS contains fatty acids and these have been shown to increase the expression of the enzyme.<sup>15–17</sup> The activation of ROR $\alpha$  significantly counteracted this overexpression, both at the mRNA and at protein level. In line with these results, the treatment with the thiazolidinedione derivative markedly decreased 5-LOX enzymatic activity, as shown by the decreased production of 5(S)-HETE observed in DU 145 cells exposed to exogenous AA in the presence of the calcium ionophore A23187. These data indicate that in the presence of serum, and therefore in the case of high supply of fatty acids, the activation of the orphan nuclear receptor ROR $\alpha$  significantly decreases the overexpression of 5-LOX and, consequently, its activity.

On the basis of these observations, we reasoned that the thiazolidinedione derivative CGP 52608, through the reduction of the expression and of the activity of 5-LOX, might affect the proliferative effects of  $\omega$ -6 polyunsaturated fatty acids on prostate cancer cells. We verified this hypothesis by showing that, in DU 145 cells, the mitogenic activity of both LA and AA can be completely counteracted by the simultaneous treatment of the cells with the thiazolidinedione derivative. These results have been further confirmed in another androgen-independent prostate cancer cell line (PC3).

So far, only inhibitors of 5-LOX activity or compounds that reduce the association of the enzyme with arachidonic acid and FLAP<sup>31,36</sup> have been proposed as effective agents to inhibit fatty acid-induced cancer cell proliferation.<sup>5,11,15–17,37</sup> Our data indicate that ROR $\alpha$ 1 activating compounds might be considered as possible antagonists of the mitogenic activity of fatty acids on prostate cancer because of their ability to inhibit the 5-lipoxygenase expression.

In previous studies, we have shown that the activation of the orphan nuclear receptor ROR $\alpha$  reduces the proliferation of DU 145 cells, both *in vitro* and *in vivo*, through the modulation of the expression of cell cycle-related genes (p21<sup>WAF1/CIP1</sup>, cyclin A).<sup>34</sup> Moreover, CGP 52608 also decreases the metastatic behavior of DU 145 cells by affecting the expression of cell adhesion molecules.<sup>38</sup> These observations indicate that the activation of this receptor might counteract cancer growth through complementary and additive mechanisms: antiproliferative, antimetastatic and antagonistic toward the mitogenic effect of dietary fatty acids. Therefore, ROR $\alpha$  might be considered as a possible molecular target for the development of new, and hopefully successful, strategies for prostate cancer chemoprevention or chemotherapy. It is interesting to underline that thiazolidinedione derivatives, such as CGP 52608, by modulating ROR $\alpha$  activity, have also been shown to possess strong antiinflammatory and immunostimulatory activities.<sup>27,29,30</sup>

In 1995, Wiesenberg *et al.*<sup>35</sup> indicated melatonin as the possible endogenous ligand for ROR $\alpha$ ; however, this observation remained controversial. More recently, while analyzing the crystal structure of the ligand binding domain of ROR $\alpha$ , Kallen *et al.*<sup>39</sup> have identified cholesterol as the possible endogenous ligand for the nuclear receptor. If confirmed, this finding will certainly improve the understanding of the multiple biological functions of this transcription factor.

The molecular mechanisms through which ROR $\alpha$  and specifically the ROR $\alpha$ 1 isoform, might regulate the expression of 5-LOX are still unknown. It has been shown that the nuclear receptor can interact with either coactivators or corepressors. It has been then proposed that ROR $\alpha$ 1 can assume 2 different conformations: an active state that promotes binding to coactivator complexes, and a repressive state that allows the interaction with corepressor complexes.<sup>40</sup> Very recently, it has been shown that ROR $\alpha$ 1 induces *Rev-erba* gene expression.<sup>41</sup> *Rev-erba*, in turn, is known to inter-

act with the nuclear receptor corepressor N-CoR.<sup>42,43</sup> It is then possible that in prostate cancer cells the activated ROR $\alpha$ 1 might also reclute specific corepressors to inhibit the expression of the 5-LOX gene.

Our data indicate that activation of the orphan nuclear receptor ROR $\alpha$ 1, which is expressed in prostate cancer cells, decreases the

expression of 5-LOX enzyme thus counteracting the mitogenic action of  $\omega$ -6 polyunsaturated fatty acids on these cells.

#### ACKNOWLEDGEMENTS

This work was supported by COFIN (R.M.M., M.M.M., M.M., P.L.), and by EC grant QLG1 CT 2001-01521 (A.S.).

#### REFERENCES

- Greenlee RT, Murray T, Bolden S, Wingo PA. Cancer statistics, 2000. *CA Cancer J Clin* 2000;50:7–33.
- Meikle AW, Smith JA. Epidemiology of prostate cancer. *Urol Clin North Am* 1990;17:709–18.
- Nomura AMY, Kolonel LN. Prostate cancer: a current perspective. *Epidemiol Rev* 1991;13:200–27.
- Giovannucci E, Rimm EB, Colditz GA, Stampfer MJ, Ascherio A, Chute, CC, Willett WC. A prospective study of dietary fat and risk of prostate cancer. *J Natl Cancer Inst* 1993;85:1571–9.
- Rose DP, Connolly JM. Effects of fatty acids and eicosanoid synthesis inhibitors on the growth of two human prostate cancer cell lines. *Prostate* 1991;18:243–54.
- Hughes-Fulford M, Chen Y, Tjandrawinata RR. Fatty acid regulates gene expression and growth of human prostate cancer PC-3 cells. *Carcinogenesis* 2001;22:701–7.
- Newcomer LM, King IB, Wicklund KG, Stanford JL. The association of fatty acids with prostate cancer risk. *Prostate* 2001;47:262–8.
- Needleman P, Turk J, Jakschik BA, Morrison AR, Lefkowitz JB. Arachidonic acid metabolism. *Ann Rev Biochem* 1986;55:69–102.
- Samuelsson B, Dahlen SE, Lindgren JA, Rouzer CA, Serhan CN. Leukotrienes and lipoxins: structures, biosynthesis, and biological effects. *Science* 1987;237:1171–76.
- Connolly JM, Liu X-H, Rose DP. Dietary linoleic acid-stimulated human breast cancer cell growth and metastasis in nude mice and their suppression by indomethacin, a cyclooxygenase inhibitor. *Nutr Cancer* 1996;25:23–40.
- Earashi M, Noguchi M, Tanaka M. *In vitro* effects of eicosanoid synthesis inhibitors in the presence of linoleic acid on MDA-MB-231 human breast cancer cells. *Breast Cancer Res Treat* 1996;37:29–37.
- Nie D, Hillman G, Geddes T, Tang K, Pierson C, Grignon DJ, Honn KV. Platelet-type 12-lipoxygenase in a human prostate carcinoma stimulates angiogenesis and tumor growth. *Cancer Res* 1998;58:4047–51.
- Cuendet M, Pezzuto, JM. The role of cyclooxygenase and lipoxygenase in cancer chemoprevention. *Drug Metabol Drug Interact* 2000;17:109–57.
- Dannenberg AJ, Altorki NK, Boyle JO, Dang C, Howe LR, Weksler BB, Subbaramaiah K. Cyclo-oxygenase 2: a pharmacological target for the prevention of cancer. *Lancet Oncol* 2001;2:544–51.
- Ghosh J, Myers CE. Arachidonic acid stimulates prostate cancer cell growth: critical role of 5-lipoxygenase. *Biochem Biophys Res Commun* 1997;235:418–23.
- Ghosh J, Myers CE. Inhibition of arachidonate 5-lipoxygenase triggers massive apoptosis in human prostate cancer cells. *Proc Natl Acad Sci USA* 1998;95:13182–7.
- Anderson KM, Seed T, Vos M, Mulshine J, Meng J, Alrefai W, Ou D, Harris JE. 5-Lipoxygenase inhibitors reduce PC-3 cell proliferation and initiate non necrotic cell death. *Prostate* 1998;37:161–73.
- Gupta S, Srivastava M, Ahmad N, Sakamoto K, Bostwick DG, Mukhtar H. Lipoxygenase-5 is overexpressed in prostate adenocarcinoma. *Cancer* 2001;91:737–43.
- Giguere V. Orphan nuclear receptor: from gene to function. *Endocr Rev* 1999;20:689–725.
- Sladek R, Giguere V. Orphan nuclear receptors: an emerging family of metabolic regulators. *Adv Pharmacol* 2000;47:23–87.
- Carlberg C, Hooft van Huijsduijnen R, Staple J, De Lamarter JF, Becker-André M. RZR $\alpha$ s, a novel class of retinoid related orphan receptors that function as both monomers and homodimers. *Mol Endocrinol* 1994;8:757–70.
- Giguere V, Tini M, Flock G, Ong E, Evans RM, Otulakowski G. Isoform-specific amino-terminal domains dictate DNA-binding properties of ROR $\alpha$ , a novel family of orphan hormone nuclear receptors. *Genes Dev* 1994;8:538–53.
- Matsui T. Transcriptional regulation of a Purkinje cell-specific gene through a functional interaction between ROR $\alpha$  and RAR. *Genes Cells* 1997;2:263–72.
- Steinmayr M, André E, Conquet F, Rondi-Reig L, Delhaye-Bouchaud N, Auclair N, Daniel H, Crepel F, Mariani J, Sotelo C, Becker-André M. Staggerer phenotype in retinoid-related orphan receptor  $\alpha$ -deficient mice. *Proc Natl Acad Sci USA* 1998;95:3960–5.
- Lau P, Bailey P, Dowhan DH, Muscat GE. Exogenous expression of a dominant negative ROR $\alpha$ 1 vector in muscle cells impairs differentiation: ROR $\alpha$ 1 directly interacts with p300 and myoD. *Nucleic Acid Res* 1999;27:411–20.
- Meyer T, Kneissel M, Mariani J, Fournier B. *In vitro* and *in vivo* evidence for orphan nuclear receptor ROR $\alpha$  function in bone metabolism. *Proc Natl Acad Sci USA* 2000;97:9107–202.
- Garcia-Maurino S, Pozo D, Calvo JR, Guerrero JM. Correlation between nuclear melatonin receptor expression and enhanced cytokine production in human lymphocytic and monocytic cell lines. *J Pineal Res* 2000;29:129–37.
- Steinhilber D, Brungs M, Werz O, Wiesenberg I, Danielsson C, Kahlen J-P, Nayeri S, Schrader M, Carlberg C. The nuclear receptor for melatonin represses 5-lipoxygenase gene expression in human B lymphocytes. *J Biol Chem* 1995;270:7037–40.
- Missbach M, Jagher B, Sigg I, Nayeri S, Carlberg C, Wiesernberg I. Thiazolidine diones, specific ligands of the nuclear receptor retinoid Z receptor/retinoid acid receptor-related orphan receptor  $\alpha$  with potent antiarthritic activity. *J Biol Chem* 1996;271:13515–22.
- Wiesenberg I, Chiesi M, Missbach M, Spanka C, Pignat W, Carlberg C. Specific activation of the nuclear receptors PPAR $\gamma$  and RORA by the antidiabetic thiazolidinedione BRL 49653 and the antiarthritic thiazolidinedione derivative CGP 52608. *Mol Pharmacol* 1998;53:1131–8.
- Ford-Hutchinson AW, Cresser M, Young RN. 5-Lipoxygenase. *Annu Rev Biochem* 1994;61:383–417.
- Ponte P, Ng S-Y, Engel J, Gunning P, Kedes L. Evolutionary conservation of the untranslated regions of actin mRNAs: DNA sequence of a human beta-actin cDNA. *Nucleic Acid Res* 1984;12:1687–96.
- Moretti RM, Montagnani Marelli M, Motta M, Polizzi D, Monestiroli S, Pratesi G, Limonta P. Activation of the orphan nuclear receptor ROR $\alpha$  induces growth arrest in androgen-independent DU 145 prostate cancer cells. *Prostate* 2001;46:327–35.
- Dunnett CW. A multiple comparison procedure for comparing several treatments with a control. *J Am Stat Assoc* 1955;50:1096–121.
- Wiesenberg I, Missbach M, Kahlen J-P, Schrader M, Carlberg C. Transcription activation of the nuclear receptor RZR $\alpha$  by the pineal gland hormone melatonin and identification of CGP 52608 as a synthetic ligand. *Nucleic Acid Res* 1995;23:327–33.
- Mancini JA, Abramovitz M., Cox ME, Wong E, Charleson S, Perrier H, Wang Z, Prasit P, Vickers PJ. 5-Lipoxygenase activating protein is an arachidonate binding protein. *FEBS Lett* 1993;318:277–81.
- Avis I, Hong SH, Martinez A, Moody T, Choi YH, Trepel J, Das R, Jett M, Mulshine JL. Five-lipoxygenase inhibitors can mediate apoptosis in human breast cancer cell lines through complex eicosanoid interactions. *FASEB J* 2001;15:2007–9.
- Moretti RM, Montagnani Marelli M, Motta M, Limonta P. Role of the orphan nuclear receptor ROR $\alpha$  in the control of the metastatic behavior of androgen-independent prostate cancer cells. *Oncol Rep* 2002;9:1139–43.
- Kallen JA, Schlaeppli J-M, Bitsch F, Geisse S, Geiser M, Delhon I, Fournier B. X-ray structure of the hROR $\alpha$  LBD at 1.63 Å: structural and functional data that cholesterol or a cholesterol derivative is the natural ligand for ROR $\alpha$ . *Structure* 2002;10:1697–707.
- Jetten AM, Kurebayashi S, Ueda E. The ROR nuclear orphan receptor subfamily: critical regulators of multiple biological processes. *Prog Nucleic Acid Res Mol Biol* 2001;69:205–47.
- Delerive P, Chin WW, Suen CS. Identification of Rev-erb( $\alpha$ ) as a novel ROR( $\alpha$ ) target gene. *J Biol Chem* 2002;277:35013–8.
- Harding HP, Lazar MA. The monomer-binding orphan receptor Rev-erb represses transcription as a dimer on a novel direct repeat. *Mol Cell Biol* 1995;15:4791–802.
- Downes M, Burke LJ, Bailey PJ, Muscat GE. Two receptor interaction domains in the corepressor, N-CoR/RIP13, are required for an efficient interaction with Rev-erb $\alpha$  and RVR: physical association is dependent on the E region of the orphan receptors. *Nucleic Acid Res* 1996;24:4379–86.