

Biextensions and 1-motives

CARLO MAZZA

(joint work with Cristiana Bertolin)

The notion of biextension of two abelian groups by another abelian group is a classical one coming from [SGA7, VII], later extended by Deligne in [D, 10.2] to the case where K_0 and K_1 are two complexes of abelian sheaves and H is another abelian sheaf. His result $\text{Biext}^1(K_1, K_2; H) \cong \text{Ext}^1(K_1 \otimes^L K_2, H)$ suggests that the biextension of two 1-motives is related to their tensor product, which is outside the category of 1-motives for trivial reasons.

Cristiana Bertolin further extended Deligne's work by defining the biextensions of two 1-motives (M_1 and M_2) by another 1-motive (M_3) and proved in [B] that

$$\text{Biext}^1(M_1, M_2; M_3) \cong \text{Ext}^1(M_1 \otimes^L M_2, M_3).$$

It was stated by Voevodsky, later proved by Orgogozo in [O], and generalized by Barbieri Viale and Kahn in [BVK], that there is a fully faithful embedding Tot of the category of 1-motives into the category DM . Cristiana Bertolin and I proved in [BM] that, after tensoring with \mathbb{Q} , that the embedding realizes the connection between biextension and the tensor product of two 1-motives, i.e.,

$$\text{Biext}^1(M_1, M_2; M_3) \otimes \mathbb{Q} \cong \text{Hom}_{DM(\mathbb{Q})}(\text{Tot}(M_1) \otimes \text{Tot}(M_2), M_3).$$

This also answers a question in [BVK] because applying $L\text{Alb}$ on the right gives

$$\text{Biext}^1(M_1, M_2; M_3) \otimes \mathbb{Q} \cong \text{Hom}_{1\text{-isoMot}}(M_1 \otimes^1 M_2, M_3).$$

Here is a sketch of our proof: taking into account the different degree conventions by Deligne and Voevodsky, by [B] we get

$$\text{Biext}^1(M_1, M_2; M_3) \otimes \mathbb{Q} \cong \text{Ext}^1(M_1[1] \otimes^L M_2[1], M_3[1]) \otimes \mathbb{Q},$$

and by simple homological algebra we have

$$\text{Ext}^1(M_1[1] \otimes^L M_2[1], M_3[1]) \otimes \mathbb{Q} \cong \text{Hom}_{D(\mathcal{S}h) \otimes \mathbb{Q}}(M_1 \otimes^L M_2, M_3),$$

where $D(\mathcal{S}h)$ is the derived category of abelian sheaves. By [BVK, 4.4.1]:

$$\text{Hom}_{D(\mathcal{S}h) \otimes \mathbb{Q}}(M_1 \otimes^L M_2, M_3) \cong \text{Hom}_{DM(\mathbb{Q})}(\text{Tot}(M_1) \otimes \text{Tot}(M_2), M_3).$$

REFERENCES

- [B] C. Bertolin, *Homological interpretation of extensions and biextensions of complexes*, submitted.
- [BM] C. Bertolin and C. Mazza, *Biextensions of 1-motives in Voevodsky's category of motives*, Int. Math. Res. Not. IMRN, 2009, **19**, pp. 3747–3757.
- [BVK] L. Barbieri-Viale and B. Kahn, *On the derived category of 1-motives*, in preparation <http://arxiv.org/abs/1009.1900v1>.
- [D] P. Deligne, *Théorie de Hodge III*, pp. 5–77, Inst. Hautes Études Sci. Publ. Math. No. 44, 1974.
- [O] F. Orgogozo, *Isomotifs de dimension inférieure ou égale à 1*, pp. 339–360, Manuscripta Math. 115, no. 3, 2004.
- [SGA7] A. Grothendieck and others, *Groupes de Monodromie en Géométrie Algébrique*, SGA 7 I, Lecture Notes in Mathematics, Vol. 288. Springer-Verlag, Berlin-New York, 1972.