



Bilateral sudden sensorineural hearing loss and chronic venous cerebrospinal insufficiency : a case report.

Journal:	<i>Phlebology</i>
Manuscript ID:	Draft
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Alpini, Dario; Don C. Gnocchi Foundation, Scientific Institute S.Maria Nascente Bavera, Pietro; Don C. Gnocchi Foundation, Scientific Institute S.Maria Nascente Di Berardino, Federica; Università degli Studi di Milano, Dipartimento di Scienze Chirurgiche Specialistiche Barozzi, Stefania; Università degli Studi di Milano , Dipartimento di Scienze Chirurgiche Specialistiche Cesarani, Antonio; Università degli Studi di Milano , Dipartimento di Scienze Chirurgiche Specialistiche; Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Audiology Unit
Revised Keywords:	CCSVI
Keyword:	Deep venous insufficiency, Patients- views

1
2
3 **Bilateral sudden sensorineural hearing loss and chronic venous cerebrospinal insufficiency : a**
4
5 **case report**
6
7

8
9 Alpini D. ^a, Bavera P.M. ^a, Di Bernardino F. ^b, Barozzi S. ^b, Cesarani A. ^{b,c}
10
11

12
13
14 ^a Scientific Institute S.Maria Nascente, Don C. Gnocchi Foundation, via Capecelatro 66, Milan,
15
16 Italy
17

18 ^b Dipartimento di Scienze Chirurgiche Specialistiche, Università degli Studi di Milano, Milan, Italy
19

20 ^c Audiology Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy
21
22

23
24
25 **Short title:** Sudden hearing loss and CCSVI
26
27

28
29
30 Key-words: Sudden hearing loss, CCSVI,
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 **Corresponding author:**

46 Federica Di Bernardino

47
48 Audiology Unit - Ospedale Maggiore Policlinico

49
50 Viale Piceno 14 - 20129 Milan – Italy

51
52 Phone- Fax: +39 02 70126502

53
54 e-mail: federica.diberardino@unimi.it
55

56 **Abstract**
57
58
59
60

1
2
3 **Objectives.** We report a case of bilateral sudden sensorineural hearing loss in a patient suffering
4 from chronic venous cerebrospinal insufficiency. **Methods.** Audiometric testing confirmed bilateral
5 sensorineural hearing loss with hypoexcitability to caloric stimulation on the left side and echo-
6 colour Doppler examination showed abnormal cerebral venous deficiency. **Results.** The patient's
7 condition improved after fifteen days following medical treatment. **Conclusions.** Chronic venous
8 cerebrospinal insufficiency may explain the anatomical background which provides a predisposing
9 factor for sudden sensorineural hearing loss although further studies are needed to verify whether
10 this observation is casual or coincidental.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Under Review

Introduction

Sudden sensorineural hearing loss (SSHL) is defined as a sensorineural hearing loss of 30dB or greater over at least three contiguous audiometric frequencies occurring within a 72 hour period.¹ Various etiologies have been proposed to explain SSHL including vascular mechanisms (microembolism, hemorrhage, thrombosis), inflammation, meningitis, syphilis, HIV infection, autoimmune inner ear disease, ototoxic drugs, tumors, traumatic and idiopathic mechanisms.² SSHL has also been associated with vertebro basilar insufficiency (VBI).³⁻¹⁰ However, for the majority of patients an etiologic factor is not identified.

Chronic venous cerebrospinal insufficiency (CCSVI) was initially described as being strongly associated with multiple sclerosis (MS).¹¹ It is a syndrome characterized by stenosis of the internal jugular veins (IJVs) and/or azygous vein (AZ) with opening of collaterals and insufficient drainage.¹²⁻¹⁴

We report a case of bilateral SSHL with vertigo, showing evidence of the CCSVI pattern but not associated with MS. To the best of the authors' knowledge, this kind of association has never been reported.

Case Report

In 2010, a 56-year-old female came to our observation for a sudden sensorineural hearing loss (SSHL) in the right ear associated with aspecific dizziness and lightheadedness. In 2003, without any previous history of hearing difficulties, the patient had experienced an episode of SSHL in the left ear. In both episodes, there was no prior history of temporal bone fracture, meningitis, autoimmune disease or exposure to ototoxic drugs. General and neurological evaluations were normal; extensive blood tests including liver function tests, complete blood count, urea nitrogen, creatinine, erythrocyte sedimentation rate, circulating immune complex, and serum electrolytes all

1
2
3 gave normal results. Antinuclear antibody, rheumatoid factor, C reactive protein, HIV and VDRL
4
5 tests were unremarkable as were clotting tests. A transthoracic echocardiogram and an
6
7 electrocardiogram showed no abnormalities.
8

9
10 Audiometric testing, performed on admission, showed sensorineural hearing loss with a pure-tone
11
12 threshold (PTA) at 500- 1000-2000 Hz at 55 dB with 90% speech discrimination in the left ear and
13
14 PTA 45 dB with 95% speech discrimination in the right ear. Stapedial reflex thresholds at multiple
15
16 frequencies were normal on both sides.
17

18
19 Normal waveform responses were evoked bilaterally on brain stem auditory evoked potentials.
20
21 Electronystagmography showed hypoexcitability (more than 22% asymmetry in our laboratory) to
22
23 caloric stimulation on the left side. There were no abnormalities in saccades, pursuit, or optokinetic
24
25 nystagmus. Magnetic resonance imaging of the brain, including diffusion images, was normal for a
26
27 subject of this age.
28

29
30 The echo-colour Doppler for assessing cerebral venous return, according to the protocol of
31
32 Zamboni P et al., (15) showed an abnormal cerebral venous deficiency. A type B pattern
33
34 characterized by significant stenoses of both IJVs and of the proximal azygous was observed.
35

36
37 Abnormal reflux of the vertebral veins and bilateral stenosis of the brachiocefalic junction were also
38
39 detected.
40

41
42 The problem we observed concerning the IJVs appeared to consist mainly of a malfunction of the
43
44 valve leaflets that usually allow the correct outflow of the venous bloodstream and impede back
45
46 stream. In both cases the leaflets appeared to have a reduced motility and, de facto, caused a
47
48 stenosis of the vessel lumen. Secondly, the reflux of the vertebral vein system was quite evident,
49
50 especially when the examination was performed in the upright position, since there was a poor
51
52 outflow into the azygos venous system.
53
54
55
56
57
58
59
60

1
2
3 The patient was treated with systemic steroid, hyperbaric oxygen and platelet aggregation inhibitor
4 (16) and an improvement of 35 dB of PTA was obtained in the right ear after fifteen days. No
5
6
7 modifications were observed in the left ear.
8
9

10 11 **Discussion**

12
13 In 2007, Shim et al.¹⁷ presented a case of unilateral profound sensorineural hearing loss due to a
14
15 venous malformation of the internal auditory canal. As expected, no malformations of the internal
16
17 auditory canals were detected in our patient and therefore we had to focus on the possible effects of
18
19 CCSVI on the outflow of the inner ear, that is completely drained by the vein of the cochlear
20
21 aqueduct.¹⁸ In fact, the common modiolar vein enters the bony channel immediately adjacent to the
22
23 aqueduct to become the vein of the cochlear aqueduct which in turn drains via the inferior petrous
24
25 sinus into the IJVs. Injury or occlusion of this vessel would be particularly significant since it is
26
27 widely believed to provide virtually the entire venous drainage of the cochlea.^{13, 19} The cochlear
28
29 aqueduct and the internal auditory canal communicate with the subarachnoidal space; in the guinea
30
31 pig model, an occlusion of the veins of the cochlear aqueduct results in an increase of perilympatic
32
33 endolymphatic pressure, a decrease of cochlear blood flow and endolymphatic potential.²⁰
34
35

36
37 Furthermore, since many of the venous vessels in the scala tympani have little or no bony covering
38
39 and are essentially exposed to the perilympatic space, the venous system is a route of entry for the
40
41 cells participating in the inner ear inflammatory process.²¹
42
43

44
45 Another interesting point is that the blood leaves the brain by using the back propulsion of the
46
47 residual arterial pressure (vis a tergo), complemented by anterograde respiratory mechanisms (vis a
48
49 fronte).^{1, 2, 4-7} The latter consist of the thoracic pump increased venous outflow during inspiration:
50
51 the increase of negative thoracic pressure improves the aspiration of blood toward the right atrium.
52
53

54
55 In addition to vis a tergo and vis a fronte, postural mechanisms play a fundamental role in ensuring
56
57 a correct cerebral venous return. Several ultrasound studies of healthy volunteers have demonstrated
58
59 that the pattern of cerebral venous drainage changes, even under physiological conditions,
60

1
2
3 depending on the body position. ^{1, 2, 4 - 7} In the prone or supine position, the outflow through the
4
5 IJVs is favoured, whereas passing to the upright position transfers most of the encephalic drainage
6
7 to the vertebral veins. ^{1, 4-7}
8

9
10 On the basis of these considerations, it would be interesting to study the CCSVI in certain cases of
11
12 bilateral SSHL following general anesthesia for non-otologic surgery involving the use of
13
14 cardiopulmonary bypass of unknown origin ²² since the CCSVI might explain the anatomical
15
16 background which provides a predisposing factor for SSHL. Further studies are needed to verify
17
18 whether this observation is causal or coincidental.
19

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Under Review

References

1. Zivadinov R, Poloni GU, Marr K, Schirda CV, Magnano CR, Carl E, Bergsland N, Hojnacki D, Kennedy C, Beggs CB, Dwyer MG, Weinstock-Guttman B. Decreased brain venous vasculature visibility on susceptibility-weighted imaging venography in patients with multiple sclerosis is related to chronic cerebrospinal venous insufficiency. *BMC Neurol*. 2011 Oct 19;11:128.
2. Zamboni P, Consorti G, Galeotti R, et al. Venous collateral circulation of the extracranial cerebrospinal outflow routes. *Curr Neurovasc Res* 2009;3:204–12
3. Zamboni P, Galeotti R, Menegatti E, et al. Chronic cerebrospinal venous insufficiency in patients with multiple sclerosis. *J Neurol Neurosurg Psychiatry* 2009;80:392–9
4. Zamboni P, Menegatti E, Galeotti R, et al. The value of cerebral Doppler venous haemodynamics in the assessment of multiple sclerosis. *J Neurol Sci* 2009; 282:21–7
5. Koç A, Sanisoğlu O. Sudden sensorineural hearing loss: literature survey on recent studies. *J Otolaryngol*. 2003 Oct;32(5):308-13. Review.
6. Kuhn M, Heman-Ackah SE, Shaikh JA, Roehm PC. Sudden Sensorineural Hearing Loss: A Review of Diagnosis, Treatment, and Prognosis. *Trends Amplif*. 2011 May [Epub ahead of print] PubMed PMID: 21606048.
7. Stephan H, Wappenschmidt J, Kiefer H. Occlusion of the basilar artery. *Neurosurg Rev* 1981;4:41–52.)
8. Huang MH, Huang CC, Ryu SJ, et al. Sudden bilateral hearing impairment in vertebrobasilar occlusive disease. *Stroke* 1993;24:132–7.
9. Deplanque D, Godefroy D, Guerouaou D, et al. Sudden bilateral deafness: lateral inferior pontine infarction. *J Neurol Neurosurg Psychiatry* 1998;64:817–8.
10. Sunose H, Toshima M, Mitani S, et al. Sudden bilateral hearing loss and dizziness occurred with cerebellar infarction. *Otolaryngol Head Neck Surg* 2000;122:146–7.

- 1
2
3 11. Lee H, Whitman GT, Lim JG, et al. Bilateral sudden deafness as a prodrome of anterior inferior
4 cerebellar artery infarction. Arch Neurol 2001;58:1287–9.
5
- 6
7 12. Toyoda K, Hirano T, Kumai Y, et al. Bilateral deafness as a prodromal symptom of basilar
8 artery occlusion. J Neurol Sci 2002;193:147–50.
9
- 10
11 13. Buttner U, Ott M, Helmchen CH, et al. Bilateral loss of eighth nerve function as the only
12 clinical sign of vertebrobasilar dolichoectasia. J Vestib Res 1995;5:47–51.
13
- 14
15 14. Fetterman BL, Luxford WM, Saunders JE. Sudden bilateral sensorineural hearing loss.
16 Laryngoscope 1996;106:1347–50.
17
- 18
19 15. Zamboni P., Galeotti R. The chronic cerebrospinal venous insufficiency syndrome. Phlebology
20 2010;25:269–279.
21
- 22
23 16. Conlin AE, Parnes LS. Treatment of sudden sensorineural hearing loss: I. A systematic review.
24 Arch Otolaryngol Head Neck Surg. 2007 Jun;133(6):573-81. Review.
25
- 26
27 17. Shim HJ, Song DK, Lee SW, Lee DY, Park JH, Shin JH, Kim S. A case of unilateral
28 sensorineural hearing loss caused by a venous malformation of the internal auditory canal. Int J
29 Pediatr Otorhinolaryngol. 2007 Sep;71(9):1479-83. Epub 2007 Jun 22.
30
- 31
32 18. Baloh RW. Dizziness, hearing loss, and tinnitus. New York: Oxford, 1988:89–105.
33
- 34
35 19. Schuknecht HF. Pathology Of The Ear. 2nd ed. Philadelphia, PA: Lea and Febiger; 1993.
36
- 37
38 20. Watanabe Y, Nakashima T, Yanagita N. Venous communications of the cochlea after acute
39 occlusion of the vein of the cochlear aqueduct. Arch Otorhinolaryngology 1988;245:340-343
40
- 41
42 21. Harris JP, Fukuda S, Keithley EM. Spiral modiolar vein: its importance in inner ear
43 inflammation. Acta Otolaryngol. 1990 Nov-Dec;110(5-6):357-65.
44
- 45
46 22. Son HJ, Joh JH, Kim WJ, Chin JH, Choi DK, Lee EH, Sim JY, Choi IC. Temporary bilateral
47 sensorineural hearing loss following cardiopulmonary bypass -A case report-. Korean J Anesthesiol.
48 2011 Aug;61(2):162-5.
49
50
51
52
53
54
55
56
57
58
59
60