

An aerial photograph of a coastal fortification. A concrete perimeter wall surrounds a central area containing a tall, reddish-brown rock tower. The fortification is situated on a rocky coastline overlooking the North Sea. The sky is blue with scattered white clouds.

**ZOOPLANKTON PERFORMANCE IN A  
CHANGING OCEAN:  
ADAPTIVE CAPACITIES TO A SHIFTING FOOD  
REGIME IN THE NORTH SEA**

**Rita Melo Franco Santos**

**Universität Bremen**

**November 2018**



**ZOOPLANKTON PERFORMANCE IN A CHANGING OCEAN:  
ADAPTIVE CAPACITIES TO A SHIFTING  
FOOD REGIME IN THE NORTH SEA**

**Rita Melo Franco Santos**

Dissertation zur Erlangung des akademischen Grades eines  
Doktors der Naturwissenschaften (Dr. rer. nat.)

am Fachbereich 2  
(Biologie/Chemie)  
der Universität Bremen



November 2018

**Datum des Kolloquiums (Defense date): 29/11/2018**

**Gutachter 1:** PD Dr. Barbara Niehoff

(Expert 1) Alfred-Wegener-Institut, Helmholtz-Zentrum für Polar- und  
Meeresforschung, Deutschland

**Gutachter 2:** Prof Dr. Tom Moens

(Expert 2) Universiteit Gent, Marine Biology Section, Belgium

**Prüfer 1:** Prof. Dr. Wilhelm Hagen

(Examiner 1) Universität Bremen, Fachbereich 2, Deutschland

**Prüfer 2:** Prof. Dr. Marleen De Troch

(Examiner 2) Universiteit Gent, Marine Biology Section, Belgium

## MAR PORTUGUÊS

Ó mar salgado, quanto do teu sal  
São lágrimas de Portugal!  
Por te cruzarmos, quantas mães choraram,  
Quantos filhos em vão rezaram!  
Quantas noivas ficaram por casar  
Para que fosses nosso, ó mar!

Valeu a pena? Tudo vale a pena  
Se a alma não é pequena.  
Quem quer passar além do Bojador  
Tem que passar além da dor.  
Deus ao mar o perigo e o abismo deu,  
Mas nele é que espelhou o céu.

Fernando Pessoa



**CONTENTS**

<b>LIST OF ABBREVIATIONS</b> .....	i
<b>ABSTRACT</b> .....	v
<b>SAMENVATTING</b> .....	vii
<b>ACKNOWLEDGEMENTS</b> .....	xv
<b>1 SCIENTIFIC BACKGROUND AND OBJECTIVES</b> .....	1
1.1 ZOOPLANKTON COMMUNITIES .....	1
1.1.1 Zooplankton species investigated .....	1
1.1.2 Common prey items for the studied zooplankton species .....	4
1.2 NUTRITIONAL ECOLOGY.....	6
1.2.1 Effects of nutrient limitation on prey quality.....	7
1.2.2 Effects of prey quality on predator performance .....	7
1.2.3 Measuring predator performance.....	9
1.3 FOOD REGIME SHIFTS IN THE NORTH SEA AND AROUND HELGOLAND....	11
1.4 OBJECTIVES AND THESIS OUTLINE.....	13
Chapter 1: copepods feeding on a diet with a molar C:N ratio close to their own body composition will have the highest possible growth and reproduction rates and the lowest possible egestion, respiration and excretion rates. ....	15
Chapter 2: how does prey nutritional quality affect the biochemical composition, the dynamics of lipid utilization and assimilation, and the reproductive output of copepods?...	15
Chapter 3: how is lipid C assimilated in the larvae of polychaetes? Can the concept of FATM be applied to the study of lipid dynamics in these lipid-poor meroplanktonic larvae? .....	16

## **CONTENTS**

---

<b>2</b>	<b>MATERIALS AND METHODS</b> .....	17
2.1	SAMPLING AREA – THE NORTH SEA AND THE ISLAND OF HELGOLAND...	17
2.2	ZOOPLANKTON SAMPLING.....	19
2.3	PREY CULTURE .....	20
2.4	EXPERIMENTAL DESIGNS .....	21
2.4.1	Feeding experiments with <i>T. longicornis</i> females .....	21
2.4.2	Feeding experiment with <i>L. conchilega</i> larvae .....	24
2.5	ANALYTICAL WORK.....	25
2.5.1	Carbon, nitrogen, and FA content analyses .....	25
2.5.2	Compound-specific stable isotope analysis (CSIA) .....	25
2.5.3	Statistical analyses .....	26
<b>3</b>	<b>PUBLICATIONS</b> .....	27
	CHAPTER I.....	31
	Bioenergetics of the copepod <i>Temora longicornis</i> under different nutrient regimes.....	31
	CHAPTER II.....	49
	Life history strategies of the copepod <i>Temora longicornis</i> as inferred from lipid carbon assimilation experiments .....	49
	CHAPTER III.....	97
	Fatty acid bioconversion and homeostasis in the larvae of the sand mason worm <i>Lanice conchilega</i> .....	97
	CHAPTER IV .....	139



Temperature driven changes in the diet preference of omnivorous copepods: no more meat when it's hot?.....	139
<b>4 SYNOPSIS DISCUSSION</b> .....	151
4.1 ADVANTAGES AND CAVEATS OF USING COMPOUND-SPECIFIC STABLE ISOTOPE ANALYSIS.....	152
<i>Methodological constrains of CSIA in this PhD</i> .....	154
4.2 ECOLOGICAL STOICHIOMETRY AND HOMEOSTASIS AS TOOLS TO STUDY ZOOPLANKTON PHYSIOLOGY .....	158
<i>ES framework for the study of performance in adult females of T. longicornis</i> .....	160
<i>Biochemical (lipid) homeostasis patterns in zooplankton</i> .....	164
4.3 CLIMATE CHANGE AND MARINE REGIME SHIFTS .....	168
<i>Regime shifts</i> .....	168
<i>The regime shifts in the southern North Sea</i> .....	170
<i>Investigating zooplankton performance in relation to diet nutritional quality</i> .....	171
<b>5 PERSPECTIVES</b> .....	177
<b>REFERENCES</b> .....	179
<b>ERKLÄRUNG</b> .....	199



**LIST OF ABBREVIATIONS**

AA	Amino acid
AE <sub>C</sub>	Absorption efficiency for carbon
AE <sub>N</sub>	Absorption efficiency for nitrogen
ANOSIM	Analysis of similarity
ANOVA	Analysis of variance
ASW	Artificial sea water
atom%	Atom percent
APE	Atom percent excess
AWI	Alfred Wegener Institute
BAH	Biologische Anstalt Helgoland
BSIA	Bulk stable isotope analysis
C	Carbon
CPR	Continuous Plankton Recorder
CSIA	Compound-specific stable isotope analysis
DOC	Dissolved organic carbon
DON	Dissolved organic nitrogen
EPR	Egg production rate
ES	Ecological stoichiometry
FA	Fatty acid

## ***LIST OF ABBREVIATIONS***

---

FAME	Fatty acid methyl ester
FATM	Fatty acid trophic markers
FP	Faecal pellets
FPR	Faecal pellet production rate
GC	Gas chromatograph
GC-c-IRMS	Gas chromatograph/combustion/isotope ratio mass spectrometer
GGE	Gross growth efficiency
LC-PUFA	Long-chain polyunsaturated fatty acids
MUFA	Monounsaturated fatty acids
N	Nitrogen
NEPE <sub>C</sub>	Net egg production efficiency for carbon
NEPE <sub>N</sub>	Net egg production efficiency for nitrogen
nMDS	Non-metric multidimensional scaling
NPOC	Non-purgeable organic carbon
NSGE <sub>C</sub>	Net somatic growth efficiency for carbon
NSGE <sub>N</sub>	Net somatic growth efficiency for nitrogen
P	Phosphorus
PERMANOVA	Permutational multivariate analysis of variance
PUFA	Polyunsaturated fatty acids
RQ	Respiratory quotient
SFA	Saturated fatty acids

---

***LIST OF ABBREVIATIONS***

SIA	Stable isotope analysis
SIMPER	Similarity percentages
SST	Sea surface temperature
TDN	Total dissolved nitrogen
TAG	Triacylglycerol
UGhent	University of Ghent



**ABSTRACT**

The zooplankton is a key group in marine ecosystems – it has high biomass and is the link between primary producers and higher trophic levels. Zooplankton organisms are often vulnerable to fluctuations in food supply, and their population dynamics is directly influenced by changes in phytoplankton availability and nutritional quality. Changes in parameters such as water temperature and nutrient loading can affect the nutritional quality of algae as food for herbivores, as well as phytoplankton community composition and dominance. Alterations in the base of the food web will, consequently, affect all trophic levels. The adaptive capacity of the zooplankton to shifting food regimes will likely be determined by the effects of food quality (a.o.) on their physiology. Understanding such effects is a pressing issue given the gap in the literature regarding how species cope with inadequate (low quality) food supply. In the southern North Sea the zooplankton community is dominated by calanoid copepods, with important contributions from polychaete larvae. This PhD project investigated how food quality influenced the performance, the elemental and biochemical compositions, and the assimilation and turnover of carbon (C) in females of the calanoid copepod *Temora longicornis* and in the meroplanktonic larvae of the polychaete *Lanice conchilega*.

Zooplankton samples were collected between May and June 2016 off the German island of Helgoland, in the southern North Sea, and copepods and polychaetes were sorted for five-day laboratory feeding experiments. Two experiments were conducted with *T. longicornis* and one with *L. conchilega*, during which the individuals were fed with diets of different quality - diatoms (*Conticribra weissflogii*) and dinoflagellates (*Oxyrrhis marina*) cultured in nutrient-replete and in nitrogen (N)-depleted conditions. In order to be able to follow dietary lipid C assimilation, prey items were enriched with  $^{13}\text{C}$  before being offered as food to copepods and polychaetes. Grazing, respiration, excretion, and egg and faecal pellet production rates were also recorded in incubations conducted during the experiments with copepods. These rates were used to construct C and N budgets in order to evaluate their intake and usage of energy. Performance was also discussed in terms of the efficiency with which these processes occurred.

Copepods feeding on prey with a molar C:N ratio that resembled their own showed the highest investment in somatic and reproductive growth. However, they also showed the highest expenditures with egestion and catabolism, which were expected for copepods feeding on prey

## ***ABSTRACT***

---

with the highest molar C:N ratio. The major result regarding copepod performance in relation to food quality is that dinoflagellates are a food source of superior or similar quality to diatoms under nutrient-replete or N-depleted conditions, respectively. This result is, however, limited to the temperature tested during the laboratory experiments, as different temperatures will influence copepod metabolism and C requirements. Furthermore, the experiments with *T. longicornis* revealed that egestion is a major pathway for eliminating excess C, and that low food quality can influence respiration and the intensity and speed of DOC leakage from faecal pellets.

The experiments investigating C assimilation and turnover in *T. longicornis* revealed a higher absolute lipid C assimilation in copepods fed with prey whose molar C:N ratio was the most different from their own. The relative lipid C assimilation (as % food ingested), however, was similar between copepods fed with different diets. This result suggests that prey biochemical composition does not have an effect over lipid C assimilation by *T. longicornis*, only over ingestion rates. The high C turnover rates measured indicate an opportunistic feeding strategy for *T. longicornis*, which would accumulate dietary C when it is readily available with a speed that decreases as saturation is approached and its lipid storage is replenished. This strategy is likely employed to maximize reproductive output. In the case of *L. conchilega*, the experiment revealed that larvae are able to regulate their lipid C content (homeostasis) regardless of the fatty acid (FA) availability in their diet. Although lipid C assimilation was low in the larvae (meroplanktonic organisms are usually lipid-poor), the individuals selectively accumulated certain FAs and biosynthesized others (from dietary FA) which were not available in the diet.

Concomitant changes in several ecosystem parameters make it difficult to precisely evaluate how the zooplankton will react to changing food regimes in marine environments and to comprehend the full scale of the effects and feedback mechanisms of ecosystem changes. Nevertheless this thesis presents a robust contribution towards a better understanding of how zooplanktonic organisms might be affected by changes in the quality of their prey. Although the work for this thesis refers to organisms from a very specific location, the ubiquitousness of the species and of the forcing (of environmental change) studied enables the comparison of the results with those from several marine ecosystems.



ZUSAMMENFASSUNG

In marinen Ökosystemen hat das Zooplankton eine Schlüsselrolle – in vielen Regionen ist die Biomasse der Zooplankter hoch und sie sind ein Bindeglied zwischen Primärproduzenten und höheren trophischen Ebenen. Viele Zooplanktonorganismen reagieren sensitiv auf Fluktuationen in ihrer Nahrungsversorgung, und daher wird ihre Populationsdynamik direkt von Veränderungen der Verfügbarkeit und der Qualität der Nahrung, z. B. Phytoplankton, beeinflusst. Die Qualität der Algen als Nahrung für herbivore Organismen wiederum wird von Umweltparametern wie z.B. Wassertemperatur und Nährstoffkonzentrationen beeinflusst. Änderungen an der Basis des Nahrungsnetzes betreffen folglich alle trophischen Ebenen. Wie gut sich Zooplanktonorganismen an ein sich veränderndes Nahrungsregime anpassen können, wird u.a. dadurch bestimmt, wie sich die Nahrungsqualität auf die Physiologie der Tiere auswirkt. Zur Zeit liegen allerdings nur wenige Informationen dazu vor, wie die Zooplankter auf Nahrung von geringer Qualität reagieren.

In der südlichen Nordsee wird die Zooplanktongemeinschaft generell von calanoiden Copepoden dominiert, aber auch meroplanktische Larven, wie z.B. Polychaetenlarven, können saisonal hohe Abundanzen erreichen. Im Rahmen dieser Dissertation wurde untersucht, welchen Einfluss die Nahrungsqualität auf die Leistungsfähigkeit und die biochemische Zusammensetzung der Weibchen des calanoiden Copepoden *Temora longicornis* und der meroplanktonischen Larven des Polychaeten *Lanice conchilega* hat. Außerdem wurde untersucht, wie viel und wie schnell Kohlenstoff (C) unter unterschiedlichen Nahrungsbedingungen assimiliert und umgesetzt wird.

Die Zooplanktonproben wurden in Mai und Juni 2016 in der südlichen Nordsee nahe der deutschen Insel Helgoland gesammelt. Direkt nach dem Fang wurden die Copepoden und die Polychaetenlarven lebend aus den Fängen sortiert und dann für fünf Tage im Labor inkubiert. Mit *T. longicornis* wurden zwei Versuche und mit Larven von *L. conchilega* wurde ein Versuch durchgeführt. Dabei wurden die Tiere mit Diatomeen (*Conticribra weissflogii*) und Dinoflagellaten (*Oxyrrhis marina*) gefüttert. Die Algen wurden dabei entweder unter hohen Nährstoffkonzentrationen oder unter limitiertem Angebot an Stickstoff (N) kultiviert. Um der Aufnahmen von C in Form von Lipiden zu folgen, wurden die Beuteorganismen mit  $^{13}\text{C}$  angereichert, bevor sie den Copepoden und Polychaetenlarven als Futter angeboten wurden. In zusätzlichen Experimenten mit *T. longicornis* wurden täglich die Ingestions-, Respiration- und

## ZUSAMMENFASSUNG

---

Exkretionsraten der Tiere gemessen. Außerdem wurden die Eier und die Anzahl Kotballen, die Weibchen innerhalb von 24 h produzierten gezählt. Diese Daten wurden dazu verwendet, Kohlen- und Stickstoffbudgets zu berechnen und die Aufnahme und Nutzung von Energie aus der Nahrung in Abhängigkeit von der Nahrungsqualität zu bewerten.

Die Ergebnisse zeigen, dass Copepoden, deren Nahrung ein molares C:N-Verhältnis hatte, welches ihrem eigenen ähnelte, den meisten Kohlenstoff und Stickstoff in somatisches Wachstum und Reproduktion investierten. Diese Tiere gaben jedoch gleichzeitig am meisten C und N über ihre Ausscheidungsprodukte und den Katabolismus ab. Diese Ergebnisse waren eigentlich bei den Copepoden zu erwarten, denen Beuteorganismen mit dem höchsten molaren C: N-Verhältnis angeboten wurden. Ein wesentliches Ergebnis dieser Untersuchungen ist, dass die Dinoflagellaten eine Nahrungsquelle von höherer Qualität waren als die Diatomeen, die unter nährstoffreichen bzw. N-limitierenden Bedingungen kultiviert wurden. Hier ist jedoch zu bedenken, dass dies zunächst nur für die Temperatur gelten kann, die während der Laborexperimente herrschte, da generell die Temperatur einen erheblichen Einfluss auf den Metabolismus der Copepoden und ihren Bedarf an Kohlenstoff hat. Darüber hinaus zeigten die Experimente mit *T. longicornis*, dass überschüssiger Kohlenstoff vor allem über die Egestion abgegeben wird und dass eine geringe Nahrungsqualität sowohl die Atmung als auch die Intensität und Geschwindigkeit, mit der DOC aus den Kotballen austritt, beeinflussen kann.

Die Experimente, die die Assimilation und den Umsatz von Kohlenstoff bei *T. longicornis* untersuchten, zeigten, dass Tiere, deren Nahrung ein molares C:N Verhältnis aufwies, welches sich am stärksten von ihrem eigenen unterschied, mehr Kohlenstoff über Lipide assimilierten. Die relative Assimilation von Lipidkohlenstoff (in % des über die Nahrung aufgenommenen Kohlenstoffs) war jedoch bei allen Copepoden trotz Fütterung mit unterschiedlicher Nahrung ähnlich. Dieses Ergebnis legt nahe, dass bei *T. longicornis* die biochemische Zusammensetzung der Beuteorganismen keine Auswirkungen auf die Assimilation von Lipid-C hatte. Die hohen C-Turnover-Raten deuten auf eine opportunistische Strategie hin. Es ist zu vermuten, dass die Weibchen, Kohlenstoff aus der Nahrung zunächst schnell akkumulieren, und die Geschwindigkeit abnimmt, wenn die Lipidspeicher aufgefüllt sind. Diese Strategie könnte dazu dienen, die Fortpflanzungsleistung zu maximieren.

Im Fall von *L. conchilega* zeigte das Experiment, dass die Larven dazu in der Lage sind, ihren Lipid C-Gehalt unabhängig von Verfügbarkeiten von Fettsäuren (FA) in ihrer Nahrung zu regulieren (Homöostase). Obwohl die Assimilation von Kohlenstoff in den Lipiden der Larven niedrig war (meroplanktonische Organismen sind gewöhnlich lipidarm), akkumulierten die Individuen selektiv bestimmte FAs und biosynthetisierten andere, die nicht in der Nahrung verfügbar waren.

Gleichzeitige Veränderungen mehrerer Ökosystemparametern machen es schwierig, genau vorherzusagen, wie das Zooplankton auf Veränderungen im Nahrungsregime reagieren wird. Damit ist es schwierig, Auswirkungen und Rückkopplungsmechanismen von ökosystemaren Veränderungen vollständig zu erfassen. Nichtsdestotrotz liefert diese Arbeit einen Beitrag zu einem besseren Verständnis darüber, wie Veränderungen der Qualität der Nahrungsgrundlage Zooplanktonorganismen beeinflussen können.



**SAMENVATTING**

Zooplankton is een onmisbare schakel in mariene ecosystemen vanwege hun hoge biomassa en de connectie tussen primaire producenten. Zooplankton zijn echter kwetsbaar voor schommelingen in hun voedselvoorzieningen. De populatiedynamiek van zooplankton wordt rechtstreeks beïnvloed door veranderingen in de beschikbaarheid en de voedingskwaliteit van fytoplankton. Veranderingen in parameters zoals watertemperatuur en nutriënten concentraties beïnvloeden de voedingskwaliteit, de samenstelling van de gemeenschap en de dominantie van fytoplankton, die een voedselbron zijn voor herbivore zooplankton. Veranderingen aan de basis van het voedselweb zullen bijgevolg alle trofische niveaus beïnvloeden. Het aanpassingsvermogen van zooplankton aan veranderende voedselregimes zal worden bepaald door de effecten van de voedselkwaliteit op hun fysiologie. Het begrijpen van dergelijke effecten is een dringende kwestie, gezien de kloof in de literatuur over hoe soorten omgaan met ontoereikende (lage kwaliteit) voedselvoorzieningen. In de zuidelijke Noordzee wordt de gemeenschap van zooplankton gedomineerd door calanoïde copepoden, met een belangrijke bijdragen van polychaete larven. Dit doctoraatsproject onderzoekt hoe voedselkwaliteit de prestaties, de elementaire en biochemische samenstellingen en de assimilatie en turnover van koolstof (C) beïnvloedde bij de vrouwelijke calanoïde copepoden *Temora longicornis* en in de meroplanktonische larven van de polychaete *Lanice conchilega*.

Zooplankton stalen werden verzameld tussen mei en juni 2016 nabij het Duitse eiland Helgoland, in de zuidelijke Noordzee. Copepoden en polychaetes werden geselecteerd uit deze stalen voor vijfdaagse laboratorium voedingsexperimenten. Twee experimenten werden uitgevoerd met *T. longicornis* en één met *L. conchilega*, waarbij de individuen werden gevoed met diëten van verschillende kwaliteit; diatomeeën (*Conticribra weissflogii*) en dinoflagellaten (*Oxyrrhis marina*) werden gekweekt in nutriënten-verzadigde en in stikstof (N) arme voedings media. Om supplementaire lipide C-assimilatie te kunnen volgen, werden prooi-items verrijkt met <sup>13</sup>C voordat ze als voedsel aan copepoden en polychaetes werden aangeboden. Productiesnelheden voor graas, ademhaling, excretie, egestie en faecale pellet productie werden gemeten in de incubatie tanken tijdens de experimenten met de copepoden. Deze snelheden werden gebruikt om C- en N-budgetten te construeren en om de inname en gebruik van energie in de copepoden

## SAMENVATTING

---

te evalueren. De algehele prestaties van de copepoden, in termen van de efficiëntie waarmee deze processen plaatsvonden, worden ook in deze thesis besproken.

Copepoden die zich voedden met prooien met een molaire C:N verhouding die leek op hun eigen samenstelling vertoonden de hoogste investering in somatische en reproductieve groei. Ze vertoonden echter ook de hoogste energetische uitgaven aan eier productie en katabolisme, deze resultaten werden echter verwacht voor copepoden die zich voeden met prooien met de hoogste molaire C:N verhouding. De belangrijkste uitkomst van de copepod prestaties, in betrekking tot de voedselkwaliteit, was dat de dinoflagellaten een voedingsbron zijn van superieure of vergelijkbare kwaliteit wanneer deze werden vergeleken bij diatomeeën onder voedings verzadigde of onder N arme omstandigheden, respectievelijk. Dit resultaat is echter beperkt tot de geteste temperatuur tijdens de laboratoriumexperimenten, omdat verschillende temperaturen uiteraard het copepod metabolisme en de bijgaande C-vereisten zullen beïnvloeden. De de experimenten met *T. longicornis* onthulden tevens dat egestie een belangrijk pad is voor het elimineren van een overmaat aan C, en dat een lage voedselkwaliteit de ademhaling en de intensiteit en snelheid van DOC-lekkage uit fecale pellets kan beïnvloeden.

De experimenten die de C turnover en assimilatie in *T. longicornis* onderzochten, lieten een hogere absolute lipide C-assimilatie zien in copepoden die gevoed werden met prooien waarvan de molaire C: N-verhouding het meest verschillend was van die van henzelf. De relatieve lipide C-assimilatie (als % ingenomen voedsel) was echter vergelijkbaar tussen copepoden die werden gevoed met verschillende diëten. Dit resultaat suggereert dat de biochemische samenstelling van de prooi geen effect heeft op lipide C-assimilatie voor *T. longicornis*, alleen voor ingestiecijfers. De hoge C turnover snelheden duiden op een opportunistische voedingsstrategie voor *T. longicornis*, waarbij C wordt geaccumuleerd wanneer het gemakkelijk beschikbaar is met een snelheid die afneemt naarmate de verzadiging wordt benaderd en de opslag van lipiden wordt aangevuld. Deze strategie wordt waarschijnlijk gebruikt om het reproductievermogen te maximaliseren. In het geval van *L. conchilega* onthulde mijn experiment dat larven hun lipide C-gehalte (homeostase) kunnen reguleren ongeacht de beschikbaarheid van vetzuren (FA) in hun dieet. Hoewel lipide C-assimilatie laag was in de larven (meroplanktonische organismen zijn meestal lipidearm), accumuleerden de individuen selectief bepaalde FA's en biosynthetisereerde ze andere (uit FA-dieet) die niet beschikbaar waren in het dieet.

Gelijktijdige veranderingen in verschillende ecosysteemparameters maken het moeilijk om nauwkeurig te evalueren hoe zooplankton zal reageren op veranderende voedselregimes in mariene omgevingen. Dit maakt het niet triviaal om de volledige schaal van effecten en feedbackmechanismen van ecosysteemveranderingen te vatten. Niettemin presenteert dit proefschrift een krachtige bijdrage, en een beter begrip aan de manier waarop zooplanktonische organismen kunnen worden beïnvloed door veranderingen in de voedingskwaliteit van hun prooi. Hoewel het werk voor dit proefschrift verwijst naar organismen van een zeer specifieke locatie, maakt de alomtegenwoordigheid van deze soorten en de focus op verandering in nutriënten concentraties de resultaten van deze thesis vergelijkbaar met die van verschillende mariene ecosystemen.





**ACKNOWLEDGEMENTS**

I would like to acknowledge and thank the Marine Ecosystem Health and Conservation (MARES) Joint Doctoral Training Programme and the European Union for the scholarship which enabled the completion of this PhD. I am forever grateful for being granted the opportunity to work with the great researchers mentioned below, and at the renowned institutions I spent the last years at. I have learnt a great deal through this PhD degree, knowledge and lessons which make me a better professional, though I finish with more questions that I started with.

I would like to thank my supervisors, PD Dr Barbara Niehoff, Dr Marleen De Troch, and Dr Holger Auel, for their guidance, patience, and support during this thesis work. Barbara deserves a special thank you, not only for the professional support (which even included abstract translation into German and many skype and phone calls between Bremerhaven and wherever I was), but also for the personal support whenever it was necessary, and for the understanding that came with it. I can honestly say that, had it not been for Barbara, I most likely would not have finished this project. She also enabled me to fulfill an old dream of mine, by ensuring my participation in an expedition onboard the RV Polarstern, and was responsible for introducing me to a place I now hold dear in my heart, Helgoland. Thank you, Barbara. I would also like to thank my co-authors Dr Martin Graeve, Dr Cedric Meunier, and Dr Maarten Boersma, without whose advice and expertise I would have had a much harder time trying to accomplish my PhD goals. Another special thank you goes to Dr Lauris Boissonnot, without whose support my work with lipids work would have been that much harder.

A well-deserved thank you goes to the technicians at both the BAH and the Polar Biological and Ecological Chemical sections at the AWI in Bremerhaven. Thank you Saskia, Julia, and Ursula for helping me out with finding my bearings at the BAH and with some of the equipment and analysis I needed for my experiments. Thank you Erika for the diatom cultures and Sandra and Cristiane, who, despite the occasional language barrier, did their utmost to help me with processing my samples back in Bremerhaven. Last but not least, thank you Claudia and Jana for the help with the excretion samples, thank you Ruth for putting up with my never-ending needs while processing my lipid samples, and thank you Valeria, whose presence during lipid extractions always made bright days of the grim Bremerhaven weather.

## **ACKNOWLEDGEMENTS**

---

To the few friends who were with me for this journey, thank you for the love and support. To Henrieke during the first half of the PhD, thank you for the fun and for letting me bother you all the time with silly conversations. To Nicole, thank you for showing me the ropes around the lab and for the time we shared the office. To Dr Xuxu, Rafa, and Ole, thank you so much for keeping me sane and making life a little more colorful, cozier, and warmer – and for the countless meals at the Portuguese restaurant! And a thank you the size of my heart (which is by no means proportional to my body size) to my three European sisters, Julia, Joana, and Anouk, who embraced the concept of being my home away from home and have ever since that late September 7 years ago been some of the people I rely the most on. Knowing you were close by and that I'd get to see you soon was always enough.

To my family, what can be said? Thank you for putting my needs over yours, and for not only letting me go, but for encouraging me to, through both the good and the bad. Thank you for listening carefully and attentively when I discussed theories and problems and solutions. I'm sorry for making you having to understand about yet more tiny animals, and I am thankful and filled with joy to see you were as interested in it as always. Thank you for being my rock, even though I'm as far away as I can be, and for most of the time. Thank you for helping me be a little closer nonetheless. Mãe, papis, Lindinha, Clara, vô e Liba, por vocês eu tenho um amor maior que o mundo, e devo tudo o que sou na vida a vocês. A big thank you also to my in-laws, who took me in as their own and who have filled my life with joy and happiness ever since.

And thank you, Eric. You are one of the greatest surprises life had in store for me. I have learnt so much with you, and I keep on learning every single day. Thank you for being my home, and for making me your home. I look forward to all the professional and personal adventures we are still about to embark on in our lives, knowing there's never a bad time when we're together. For you, I would cross oceans.

## **1 SCIENTIFIC BACKGROUND AND OBJECTIVES**

### 1.1 ZOOPLANKTON COMMUNITIES

The zooplankton is a key group in biological and abiotic processes in marine ecosystems, especially since its estimated biomass is larger than that from all other consumers (Conover, 1978). They are the most important link in the energy transfer between the lower and higher trophic levels, and contribute to the cycling of organic matter and to the CO<sub>2</sub> (carbon dioxide) export from the surface to deeper water via the carbon (C) pump (Castellani and Altunbas, 2014; Jónasdóttir et al., 1998; Juul-Pedersen et al., 2006). Zooplankton are not often limited by food quantity in coastal regions (Huntley and Boyd, 1984), but some food sources will be of lower nutritional value (Sterner and Schulz, 1998). Elemental composition, digestion resistance, and biochemical composition all are important factors determining the nutritional value / quality of food particles (Sterner and Schulz, 1998).

The experiments performed during this PhD used zooplanktonic species sampled around the German island of Helgoland, located in the southern North Sea. In this area the zooplankton community is dominated by calanoid copepods, both in terms of density and abundance, but polychaete larvae, harpacticoid copepods, and appendicularians have also been found in great numbers between June and September (Fransz et al., 1991; Hickel, 1975).

#### 1.1.1 Zooplankton species investigated


##### *Temora longicornis*

The small (approximately 1 mm prosome length) calanoid copepod *Temora longicornis* (Müller, 1785) (Fig. 1) is one of the dominant species in the coastal zooplankton community in the North Atlantic and North Sea, reaching peak densities ( $\sim 3 \mu\text{g C L}^{-1}$ ) during the spring, early summer, and early fall (Castellani and Altunbas, 2014; Hickel, 1975). Off Helgoland, this abundant species (up to 600 individuals m<sup>3</sup>) reproduces all year round (Wesche et al., 2006), and its grazing may have a substantial impact on the phytoplankton standing stock (Gentsch et al., 2009; Maar et al., 2004). They are, thus, an important link in the energy transfer between primary producers and higher trophic levels. Copepods have adapted to different environmental conditions by developing several life history traits, such as high reproduction rates, opportunistic

## SCIENTIFIC BACKGROUND AND OBJECTIVES

---

feeding modes, and efficient energy utilization (Kiørboe et al., 1985, Niehoff et al., 2015). *T. longicornis* is omnivorous (Marshall, 1973), though with weak prey selectivity (DeMott, 1988), and has high metabolic turnover rates (Helland et al., 2003; Kreibich et al., 2008). It is, however, unable to accumulate significant amounts of energy reserves (Båmstedt, 1986; Evjemo and Olsen, 1997; Kreibich et al., 2008), being thus dependent on a constant availability of prey and vulnerable to fluctuations in food supply (Helland et al., 2003; Kiørboe et al., 1985, Kreibich et al., 2008). This can be particularly problematic in dynamic systems such as the North Sea, where plankton community composition can change rapidly (Kiørboe and Nielsen, 1994). The availability of prey for this copepod, both in qualitative and quantitative terms, can be further aggravated by the decrease in nutrient loading around Helgoland (Boersma et al., 2015). Efficient food utilization is, thus, of paramount importance for the survival of *T. longicornis*.



**Figure 1:** *Temora longicornis*. Confocal image (10x) showing both auto fluorescence and Congo Red Fluorescence (© Jan Michels, Christian-Albrechts-Universität zu Kiel, Germany).

### *Lanice conchilega*


The tube-dwelling polychaete *Lanice conchilega* (Pallas, 1766) has a worldwide distribution (Holthe, 1986). It is commonly found in European waters, especially in the North Sea (Godet et al., 2008; Ropert and Dauvin, 2000, Van Hoey et al., 2008), where it can amount to 15% of the zooplankton biomass between July and September when water temperatures are above 13°C (Hickel, 1975). It is a non-selective suspension-deposit feeder (Braeckman et al., 2012; Bühr,

## **SCIENTIFIC BACKGROUND AND OBJECTIVES**

1976; Bühr and Winter, 1977) which derives its diet from the organic matter available in the water column and in the sediment. Common food items include diatoms, bacteria and microphytobenthos, and the species is able to switch between food sources (Braeckman et al., 2012; Bühr and Winter, 1977; Dubois et al., 2007). *L. conchilega* forms reefs (Fig. 2a) capable of structuring the surrounding habitat and creating a complex and heterogeneous environment (Zühlke et al., 1998; Zühlke, 2001). This, in turn, provides refuge against predation and physical and chemical stresses, but also nursery and feeding grounds to several other species, fish and birds included (Alves et al., 2017; Braber and De Groot, 1973; De Smet et al., 2013; Godet et al., 2008, and references therein; Petersen and Exo, 1999; Rabaut et al., 2007). By modifying its habitat into a more favorable one and with better living conditions, *L. conchilega* positively affects the benthic community living in it, favoring species richness and composition and faunal abundance (Callaway, 2006; Rabaut et al., 2007; Zühlke et al., 1998; Zühlke, 2001). Its reproductive peak in the southern North Sea occurs in spring, and is followed by smaller peaks during the summer and autumn (Van Hoey, 2006), though larval supply can vary between years (Strasser and Pieloth, 2001). *L. conchilega* larvae evolve from a short planktonic to a benthic phase and, after a few days, to a second planktonic stage, the aulophore larvae (Bhaud, 1988) (Fig. 2b). At this stage larvae already present a (transparent) tube, and display morphological features of a juvenile (Rabaut, 2009). They are able to feed in the water column, where they can remain up to 60 days before succeeding to an intermediate benthic stage (Bhaud and Cazaux, 1990). In comparison with adult organisms, few studies have been conducted on the basic ecology of *L. conchilega* larvae, addressing topics such as juvenile settlement and buoyancy (Bhaud and Cazaux, 1990; Callaway, 2003; Ropert and Dauvin, 2000; Strasser and Pieloth, 2001). There is a considerable gap in the literature regarding the feeding ecology of the larvae.

## SCIENTIFIC BACKGROUND AND OBJECTIVES

---



**Figure 2:** *Lanice conchilega* aggregations, or reefs (a), and aulophore larva stage (b).


### 1.1.2 Common prey items for the studied zooplankton species

Diatoms and dinoflagellates are common prey for *T. longicornis* and for *L. conchilega* larvae (Bakker and van Rijswijk, 1987; Evans, 1981; Larink and Westheide, 2011). It has been claimed that diatoms, although a stable food source, are of low quality for calanoid copepods and only make up a minor part of their diet, whereas dinoflagellates are consumed in higher proportions and have a higher nutritional value (Gentsch et al., 2009; Jónasdóttir et al., 1998, 2009; Jones and Flynn, 2005; Löder et al., 2011; Saiz and Calbet, 2011).

#### *Diatoms*

Diatoms are silicified members of the phytoplankton. They can be responsible for up to 40% of the carbon fixed in the ocean, which characterizes them as key players in the biological pump (Smetacek, 1999). *Conticribra weissflogii* (Grunow) (Stachura-Suchoples and Williams, 2009) (Fig. 3a) is a planktonic, centric diatom that can be found as single cells or in groups in marine environments in many parts of the world. It varies in size from 4 to 32  $\mu\text{m}$  diameter (larger in winter, 15  $\mu\text{m}$ , and smaller in summer, 5  $\mu\text{m}$ ) (TW1200 Reed Mariculture Inc. Retrieved 2011-11-18). Great controversy involves the topic of diatoms being good or bad quality diet items for copepods, given that their defense mechanisms against predation (production of compounds toxic to zooplankton, for example) and nutritional quality vary interspecifically (Jónasdóttir et al., 1998; Paffenhöfer et al., 2005). Some studies have further pointed out the suitability of diatoms as food source for the production of eggs in copepods but its reduced success for egg

hatching (Miralto et al., 1999; Ban et al., 1997). Nevertheless they have been successfully used in laboratory feeding experiments with *T. longicornis* (Arendt et al., 2005; Boersma et al., 2016; Niehoff et al., 2015; van Someren Gréve et al., 2017).



**Figure 3:** Prey species used during feeding experiments with copepods and polychaete: (a) *Conticribra weissflogii* (© Friedel Hinz, AWI, Germany); (b) *Oxyrrhis marina* (© Liliana Gomez Luna, CNEA, Cuba).

### *Dinoflagellates*

Marine dinoflagellates often display mixotrophy and are able to feed on both phytoplankton and heterotrophically. Some species are toxic and can produce harmful algal blooms (HABs), which can impact water quality (Smayda, 1997). *Oxyrrhis marina* (Dujardin, 1841) (Fig. 3b) is a common heterotrophic dinoflagellate said to be representative of many marine planktonic protists (Roberts et al., 2011). It is considered a cosmopolitan species, though the lack of studies on its geographic may overlook genetic differences between locations (Watts et al., 2011 and references therein). This dinoflagellate is omnivorous and feeds by engulfing its prey, which range from dissolved matter to bacteria, small algae and organisms as large as itself (Hantzsche and Boersma, 2010). It has been shown that, although food quality does not influence dietary uptake in *O. marina*, it can negatively affect growth in cases of phosphorus (P) limitation (Hantzsche and Boersma, 2010). *Oxyrrhis marina* has been routinely used in laboratory experiments for over a century (Montagnes et al., 2011), including for feeding experiments with copepods (Boersma et al., 2016).

## ***SCIENTIFIC BACKGROUND AND OBJECTIVES***

---

### 1.2 NUTRITIONAL ECOLOGY

The transfer of energy from one trophic level to the next is the most basic process in food web dynamics (Lindeman, 1942). At first, studies on the interactions between the environment and the nutrition of an organism were conducted based on single currencies, such as energy and its transfer through trophic levels in ecosystems (Lindeman, 1942). With the development of the fields of trophic and nutritional ecology, it became clear that individual fitness and ecosystem dynamics, a. o., can be better explained by considering the transfer of multiple elements and nutrients between prey and predator (Sperfeld et al., 2017).

Stoichiometry is the branch of science concerned with the application of the Law of definite proportions and conservation of mass, i.e., with the patterns that can be observed in the proportions of elements or molecules in the reactants and products of reactions. Chemical stoichiometry defines and limits quantitatively how reactants will combine and how the combinations interact with one another. Ecological stoichiometry (ES) sets similar restrictions to interacting biological systems, which are further combined with limitations imposed by the conservation of matter (Sterner and Elser, 2002). In doing so, it provides a framework under which ecologists can establish links between food webs, ecosystem metabolism, and biogeochemistry. There are several other fields for which ES can have practical applications, such as biotechnology (quantifying biomass production and yield), waste removal systems, and livestock production (determining diet for optimum growth) (Sterner and Elser, 2002).

The availability of nutrients, together with light and seawater temperature, can dictate processes within the phytoplankton on the individual (in terms of elemental and biochemical composition) and community (composition, abundance, and dynamics) levels (Harrison et al., 1990; Klein Breteler et al., 2005; Reitan et al., 1994; Wiltshire and Manly, 2004; Wiltshire et al., 2008). The quality and abundance of the phytoplankton will, in turn, influence the developmental time and adult size of several animal species (Evans, 1981). It will also affect mesozooplankton production (Klein Breteler et al., 2005; Malzahn et al., 2010; Nejstgaard et al., 2001; Thor et al., 2007), which itself can determine the recruitment success of organisms in higher trophic levels (Beaugrand et al., 2003; Black et al., 2016; Malzahn et al., 2007).



### 1.2.1 Effects of nutrient limitation on prey quality

The nutritional value of food particles can be determined by several factors, such as its elemental and biochemical compositions and digestion resistance (Sterner and Schulz, 1998). The availability of limiting nutrients to algal growth can influence the transition between exponential (non-nutrient limited) and stationary (nutrient limited) growth phases. Under nutrient-limiting conditions, green algae will show an increase in cell wall thickness, a defensive mechanism against herbivore grazing when algal growth rates are low and which renders them a poor quality food source for predators (Van Donk and Hessen, 1993, 1995).

When algae have all the resources necessary for photosynthesis, C is allocated to lipid synthesis, growth and cell division (Kattner et al., 1983; Parrish and Wangersky, 1990). Lipids have the highest energetic content ( $39 \text{ kJ g}^{-1}$ ) among the major organic compounds in organisms ( $\sim 24$  and  $17 \text{ kJ g}^{-1}$  for carbohydrates and proteins, respectively) (Gentsch et al., 2009). They have several functions, such as energy storage (triacylglycerols (TAG), wax esters, phospholipids, and diacylglycerol ethers), components of biomembranes (cholesterol, phospholipids), as regulatory hormones (ecdysone), antioxidants (tocopherol, pigments), and buoyancy (neutral lipids) (Lee et al., 2006). Under exponential growth, over 50% of the total lipid content may be (n-3) polyunsaturated fatty acids (PUFA) (Falk-Petersen et al., 1998). Under stationary growth, however, algae may become deficient in FAs (Weers and Gulati, 1997), but will most likely accumulate C as TAG (Bromke et al., 2015; Henderson et al., 1998; Kattner et al., 1983; Parrish and Wangersky, 1990), which are richer in saturated (SFA) and monounsaturated (MUFA) fatty acids (Reitan et al., 1994; Roessler, 1990). TAGs, which are formed by a glycerol backbone esterified with 3 FAs, are the main type of storage lipids in animals, and provide the energy necessary for locomotion and vertical migration, feeding-related processes (or survival through starvation), and reproduction (Lee et al., 2006).

### 1.2.2 Effects of prey quality on predator performance

Ecological stoichiometry attempts to explain the balance of energy and materials between prey and predator in terms of their elemental composition, which can also be considered as a proxy for biochemical composition (Sterner and Elser, 2002). As heterotrophs cannot synthesize or transform elements, they must obtain them from their diet in the necessary quantity in order to

## **SCIENTIFIC BACKGROUND AND OBJECTIVES**

---

meet their own metabolic requirements (Anderson et al., 2004). Biomass formation and energy supply are usually demands met by C, whereas nutrient elements primarily fulfill a structural role (Anderson et al., 2004). ES often relates the content of an element to that of C, and predicts that, if present in limiting quantities, this element will be used with the highest efficiency (Kuijper et al., 2004; Anderson et al., 2005). It also characterizes consumers in terms of their ability (or lack thereof) to handle nutrient limitation in their prey (homeostasis) and to buffer such effects for higher trophic levels (Sterner and Elser, 2002). In general, autotrophs display a wider range of C:nutrient ratios, whereas marine zooplankton have a narrower range and must cope with excess elements in their diet (Sterner and Elser, 2002). Homeostatic regulation requires the use of extra energy, and stoichiometric imbalances between prey and predator will greatly affect the efficiency with which the latter grows (Sterner and Elser, 2002). In summary, food quantity and quality can influence the physiology of copepods, their efficiency in nutrient uptake, and their ability to produce/metabolize energy (Ambler, 1986; Guisande et al., 2000; Hessen and Anderson, 2008; Jónasdóttir et al., 1995, 2009; Møller, 2007; Peterson and Kimmerer, 1994; Thor and Wendt, 2010).

*Temora longicornis* has been reported to selectively graze according to its physiological requirements, choosing its prey by their biochemical composition and nutritional value (Cottonnec et al., 2001). Although knowledge on the feeding ecology of calanoids is broad, information on how they react to different food regimes in nature, i.e., before, during, and after blooms, is scarce (Gentsch et al., 2009). It has been pointed out that *T. longicornis* prefers to feed on *Conticriba* spp. in the field (Bakker and van Rijswijk, 1987; Evans, 1981) and that, when algae are not abundantly available, it may prey on heterotrophic species to satisfy its metabolic needs (Gentsch et al., 2009). It has been shown, however, that dinoflagellates are a food source of higher nutritional value to copepods than diatoms, even under N-limitation scenarios, a common situation during the late spring bloom (Jones and Flynn, 2005). Furthermore, certain diatom species can inhibit copepod growth and reproduction (Paffenhöfer et al., 2005), further lowering the quality of such food source.


Although certain studies have addressed the effects of food limitation (in terms of quantity) on polychaete larvae (Almeda et al., 2009; Hansen, 1999; Pedersen et al., 2010), a gap exists in the literature concerning how these meroplanktonic organisms are affected by food quality. No

information is available on the effect of either food quantity or quality for the larvae of *Lanice conchilega*.

### 1.2.3 Measuring predator performance

#### *Estimation of vital rates*

The measurement of vital rates can provide information on the efficiency with which food is utilized and the costs of maintenance of an individual (Almeda et al., 2011). According to the First Law of Thermodynamics, all the energy that is ingested by an organism must equal the sum of energy that is absorbed or lost to the environment. The biological processes which define how energy is absorbed or lost are species-specific and also vary with sex and life stage of an organism. In the case of adult females of copepod species, it would be possible to devise an energy budget by measuring ingestion, respiration, excretion, faecal pellet (FP) production, somatic growth, and egg production rates (Fig. 4). Although performance entails the amount of energy ingested and used by an individual for all these vital activities, the development and continuity of a population is initially constrained by its ability to transform dietary intake into viable offspring (Wendt and Thor, 2015). Performance is, thus, ultimately defined by what is invested in reproduction and how successful this process is. Each of these processes has been extensively documented in copepod species (e.g., Castellani and Altunbas, 2014; Hirst and Bunker, 2003; Ikeda et al., 2001; Jónasdóttir et al., 1995, 1998, 2009; Mayzaud, 1976; Miller and Glibert, 1998; Møller et al., 2003; Thor, 2003), but on few occasions have ingestion, respiration, excretion, egestion, and reproduction been estimated altogether (Debs, 1984; Kiørboe et al., 1985; Lee and Yan, 1994) or in consideration of nutrient unbalances of their food source (Malzahn and Boersma, 2012; Nobili et al., 2013; Siuda and Dam, 2010). Far few studies on the topic have been conducted with polychaetes (Bühr, 1976; De Smet et al., 2016; McHugh, 1993; Ropert and Gouletquer, 2000), and to date no information is available for the larval stage of *L. conchilega*.



**Figure 4:** A simplified diagram with the source (food ingestion) and sinks (egestion, respiration, excretion, reproduction and somatic growth) of energy in an individual.

#### *Estimation of carbon assimilation*

Feeding experiments and gut content analyses, although widely used to study trophic ecology in marine ecosystems, can only reveal information on short-term feeding patterns of organisms (Dalsgaard et al., 2003; Gentsch et al., 2009 and references therein). A long-term record of feeding can, for example, be accessed by investigating the patterns in lipid deposition within consumer tissues (Dalsgaard et al., 2003). Fatty acid trophic markers (FATM) are compounds that would ideally (a) have a distinctive origin, which can be easily traced; (b) be inert and harmless to organisms; (c) not be selectively taken up and incorporated; and (d) be metabolically stable, thus being qualitatively and quantitatively transferred to the next trophic level (Dalsgaard et al., 2003). FATM are usually assimilated by consumers in a conservative manner (Graeve et al., 1994; Lee et al., 1971, 2006), though this assumption has only been investigated in holoplanktonic organisms with intermediate to high lipid contents. FATM can also provide information on the feeding ecology of organisms on short and long time scales (Dalsgaard et al., 2003). Although no FA can be uniquely found in any single species, some phytoplankton taxonomic classes can be identified based on common features and combinations (presence and/or absence) of certain FA groups (Ackman et al., 1968). 16:1(n-7), 16:2(n-4), 16:3(n-4), and 20:5(n-3) (EPA), for example, are considered to be diatom FATM, whereas 18:4(n-3) and DHA

## ***SCIENTIFIC BACKGROUND AND OBJECTIVES***

---

are considered dinoflagellate FATM (Boissonnot et al., 2016; Braeckman et al., 2012; Dalsgaard et al., 2003; Falk-Petersen et al., 1998; Graeve et al., 2005).

Essential qualitative and quantitative information on pathways of energy flow between prey and predator can be obtained via feeding experiments which combine the use of trophic markers and stable isotopes. Whereas bulk stable isotope analysis (BSIA) can only determine the isotopic composition of a complex mixture, compound-specific stable isotope analysis (CSIA) is a methodology that is able to measure the isotopic composition of individual compounds within a mixture, such as fatty acids (FA) and amino acids (AA) (for a review on CSIA see Meier-Augenstein, 2002). CSIA is a powerful tool for tracing organic matter origin and fate and to investigate biochemical processes and patterns in individuals and ecosystems (e.g., Guilini et al., 2010; Rix et al., 2018; Wegener et al., 2012), and has been applied in research on a variety of fields such as archeology, ecology, and pollution (Evershed et al., 2007). It first became available with the commercial production of the gas chromatograph/combustion/isotope ratio mass spectrometer (GC-c-IRMS) in the early 90's, and has since been recognized by its potential applications (Evershed et al., 2007; Lichtfouse, 2000; Newton, 2010). FAs have been used over the past 40 years as an ecological tool to study topics that range from individual nutrition and metabolism to ecosystem structure and trophic ecology scale (Budge et al., 2006). When FA determination is coupled to measurements of isotopic composition, quantitative estimates can be obtained for the transfer of dietary C from prey to predator and for FA-specific assimilation and potential biosynthesis and bioconversion (Boissonnot et al., 2016; De Troch et al., 2012; Graeve et al., 2005; Kluijver et al., 2013; Middelburg et al., 2000). Unfortunately, FA dynamics in marine invertebrates have been mostly characterized for larger, often herbivorous, calanoid copepods, whereas few studies are available on the lipid biochemistry of smaller copepods (Boissonnot et al., 2016; Dalsgaard et al., 2003; Fraser et al., 1989) or of polychaetes (Braeckman et al., 2012).

### 1.3 FOOD REGIME SHIFTS IN THE NORTH SEA AND AROUND HELGOLAND

Climate change has the potential to directly and indirectly impact entire food webs. Seawater temperature, together with light and nutrient availability, dictate the phytoplankton composition

## ***SCIENTIFIC BACKGROUND AND OBJECTIVES***

---

and dynamics, and phytoplankton production can affect the recruitment success of zooplankton. The planktonic system has been well studied in the North Sea over the past 60 years (Edwards et al., 2002; Reid et al., 2003; Beaugrand, 2004), and several plankton regime shifts have been observed in relation to temperature (Alvarez-Fernandez et al. 2012). Three water temperature shifts have been recorded in the North Sea in the past 50 years. In the late 70's a cold event was related to air-sea exchange processes, which are dependent on the atmospheric circulation in the North Atlantic (Becker & Pauly, 1996). As a result, diatom blooms were smaller and delayed in time, and individuals from species that were previously common in early blooms and/or dominant in that system, such as dinoflagellates and meroplankton, were present in lower numbers or absent altogether (Becker & Pauly, 1996; Edwards et al., 2002). Two warmer periods were then in the late 80's and early 2000s (Payne et al., 2009). The episode in the 80's caused a shift in the pelagic ecosystem of the Northeast Atlantic towards a warmer dynamic regime, which was characterized by higher phytoplankton abundance in the central North Sea, especially during winter and summer months, an extended phytoplankton season, and a change in the structure of the zooplankton community, with the takeover of warm-water copepod and fish species (Reid et al., 2001, 2003). Recent analyses of continuous plankton recorder data from the entire North Sea have also detected a decrease in dinoflagellate abundance over the last decade (Alvarez-Fernandez et al., 2012). These alterations in climate regimes have the potential to reorganize trophodynamic relationships and to induce changes in community composition and dominance (Alheit and Niquen, 2004; Alheit et al., 2005). It would seem, thus, that zooplankton organisms feeding primarily on these prey items have to cope with severe changes in timing of occurrence (match/mismatch), and in quantity and quality of their food items.

Considerable change in the physical and biological components of ecosystems in the southern North Sea have also been recorded in the past decades (e.g., Alheit et al., 2005; Beaugrand et al., 2003; Edwards et al., 2002; Reid et al., 2003; Wiltshire et al., 2008). The long-term monitoring data from Helgoland has shown (a) increases in water temperature ( $\sim 1.2^{\circ}\text{C}$  between 1979 and 2011), light penetration, and wind speeds; (b) decreases in P and nitrogen (N) loading (from 25.32 to 13.06 mol N L<sup>-1</sup> and from 0.83 to 0.56 mol P L<sup>-1</sup> between 1979 and 2011) and in dinoflagellate abundances (3.82 to 2.14 \*10<sup>9</sup> μm<sup>3</sup> L<sup>-1</sup> from 1979 until 2011); and (c) a temperature-related change in copepod phenology (Boersma et al., 2015; Schlüter et al., 2010; Wiltshire and Boersma, 2016). Nutrient availability in the North Sea will become even more

## ***SCIENTIFIC BACKGROUND AND OBJECTIVES***

---

dependent of influx from the North Atlantic in the future (Boersma et al., 2015), and will thus have an even greater impact on the system dynamics than it currently does. Changes in all of these variables make it difficult to evaluate how the zooplankton will react to future prey composition changes in Helgoland. If the availability (or lack thereof) of nutrients for algae leads to changing phytoplankton elemental composition and community structure, copepod populations and production could also be affected, and major consequences could be expected for food webs, ranging from nutrient regeneration to fisheries production (Beaugrand et al., 2003; Boersma et al., 2015; Jones and Flynn, 2005; Malzahn and Boersma, 2012; Nobili et al., 2013). The scale of the effects and feedback mechanisms of climate change in the southern North Sea are yet to be clarified and merit further investigation. The mechanisms which regulate zooplankton community diversity and secondary production can only be understood in light of zooplankton physiology, and how it is directly and indirectly affected by environmental conditions and changes therein (Castellani and Altunbas, 2014).

### 1.4 OBJECTIVES AND THESIS OUTLINE

Climate change has become a major topic of discussion in present society given its impact on ecosystems and economies. Changes have already been recorded for several biotic and abiotic parameters of marine ecosystems, such as water temperature, nutrient loading, light penetration, wind patterns, and phenologies of phytoplankton and zooplankton. It would be difficult enough to predict the overall effect to an ecosystem of a change in one of these parameters, let alone understanding the synergetic reactions caused by changes in a suite of parameters. This multiple change scenario is the case in the area monitored by the Helgoland Roads time-series, and is most likely also the case elsewhere. Nevertheless local studies can help to understand these reactions by separately investigating components of an ecosystem and testing possible change scenarios. These, in turn, can be used to understand, predict, and possibly remedy, overall ecosystem response in the face of climate change.

The zooplankton is responsible for the transfer of energy originated from the photosynthesis of unicellular algae to higher trophic levels. They are the foremost consumers of phytoplankton, and serve as food for invertebrates, fish, and baleen whales, which makes them an important

## **SCIENTIFIC BACKGROUND AND OBJECTIVES**

---

trophic level on the food web. They also play a role in the biogeochemical cycles of C, N, and P, though their part in it has yet to be better understood. Within the zooplankton, copepods are one of the most significant secondary producers, given their worldwide distribution, abundance, and dominance. Due to their rapid biomass turnover and low capacity to accumulate lipid reserves, most copepods rely on continuous food supply (Helland et al., 2003; Kreibich et al., 2008, 2011). Their growth and reproductive activity is, therefore, closely linked to bloom events (Ianora, 1998; Halsband and Hirche, 2001; Halsband-Lenk et al., 2004). Meroplanktonic larvae may occur in high abundances on a seasonal basis (Fransz et al., 1991), and they also have restricted capacity to sustain food limitation. Changes in phytoplankton availability and nutritional quality thus directly influence zooplankton metabolism and lipid dynamics.

The rate with which physiological processes such as ingestion and respiration occur in zooplankton organisms will vary according to parameters such as temperature, but also in relation to how their nutritional requirements are met by the food items ingested. Furthermore, the various members of the zooplankton have different lipid requirements, and will display varying ability to store lipids as energy reserves. A change in the food available to zooplankters could, thus, hinder their ability to fulfill their nutritional requirements, and the adaptive capacity of these organisms to a shifting food regime in the North Sea will likely be determined by the effects of food quality (a.o.) on their physiology. Understanding such effects is a pressing issue given the gap in the literature regarding how species cope with inadequate food supply, especially during reproductive periods (Lee et al., 2006). In an attempt to fill this gap, this thesis consists of studies on the effects of prey quality on the performance of two members of the zooplankton community at Helgoland, adult females of the calanoid copepod *Temora longicornis* and meroplanktonic larvae from the polychaete *Lanice conchilega*. The former represents holoplanktonic organisms which have intermediate levels of total lipid contents and the latter is a proxy for lipid-poor meroplanktonic organisms, which likely have different lipid requirements and, thus, assimilation strategies. The decrease in nutrient loading in the area surrounding Helgoland was reproduced in the experiments as a limitation in the availability of N for the prey species with which the study species were fed. Although N is a limiting nutrient for primary production in the marine environment (Anderson et al., 2017; Litchman et al., 2007) it does not receive much attention as a limiting nutrient to zooplankton (Sterner and Schulz, 2002). In the context of this PhD, the effects of N-limitation were considered worthy of investigation in the



experiments due to the high N content (and thus demand for) in adult copepods (Hessen, 1992). Prey choice for feeding the copepod and polychaete larvae took into account species which they feed on in their natural habitat which can also be easily cultured.

The major working hypothesis investigated in this thesis is that predators (copepods or polychaetes) will exhibit the best possible physiological performance when feeding on a diet whose elemental and biochemical compositions more closely resemble their own. Chapter-specific hypotheses and how they were addressed in this PhD are described below.

**Chapter 1: copepods feeding on a diet with a molar C:N ratio close to their own body composition will have the highest possible growth and reproduction rates and the lowest possible egestion, respiration and excretion rates.**

The Ecological Stoichiometry framework proposes that the efficiency with which a consumer utilizes the food it ingests is higher when it feeds on prey with an elemental composition similar to its own. A vast body of literature is available on the functional responses of calanoid copepods to diet quality, but few address all important physiological rates at once. A comprehensive approach should concomitantly estimate metabolism, feeding, growth and reproduction and, thus, assess the energy allocation to different processes in relation to food quality. Chapter 1 presents rates of ingestion, respiration, excretion, egestion, growth, and reproduction for adult females of the copepod *Temora longicornis* fed with different diets, and an analysis of its overall performance in relation to food quality. Quality was herein defined by the elemental composition of prey species, i.e., their molar C:N ratio, and by prey type (diatoms and dinoflagellates).

**Chapter 2: how does prey nutritional quality affect the biochemical composition, the dynamics of lipid utilization and assimilation, and the reproductive output of copepods?**

A gap in knowledge still exists on the role and importance of lipids in marine zooplankton, and on how species cope with inadequate food supply, especially during reproductive periods (Lee, Hagen, and Kattner, 2006). FA dynamics in marine invertebrates have been mostly characterized for larger calanoid copepods from polar regions, whereas few studies are available on the lipid

## **SCIENTIFIC BACKGROUND AND OBJECTIVES**

---

biochemistry of smaller copepod species (Boissonnot et al., 2016; Dalsgaard et al., 2003; Fraser et al., 1989). Chapter II aimed at describing qualitatively and quantitatively how the nutritional quality (biochemical and elemental composition) of prey grown under different nutrient regimes affects the biochemical composition, the dynamics of lipid utilization and assimilation, and the reproductive output by *T. longicornis* females. In order to obtain data on lipid dynamics and on reproductive output, copepods were fed with prey cultures previously labelled with  $^{13}\text{C}$ , which enabled the calculation of egg production rates, the monitoring of lipid C transfer between prey and predator, and the calculation of FA-specific C assimilation and turnover rates.


### **Chapter 3: how is lipid C assimilated in the larvae of polychaetes? Can the concept of FATM be applied to the study of lipid dynamics in these lipid-poor meroplanktonic larvae?**

The first goal of Chapter 3 was to obtain basic knowledge on the. Experiments investigating FA-related C assimilation have been conducted mostly for holoplanktonic organisms with intermediate to high lipid contents, whose FA profile tends to reflect that of the food items it ingests. Little is known, however, about how meroplanktonic organisms, which are usually lipid-poor, incorporate dietary FAs. It is possible that the lipid content and, thus, requirement, of an organism can dictate whether it will assimilate dietary FAs in an unmodified manner or not. One of the objectives of Chapter 3 was thus to test the hypothesis that the concept of FATM can also be applied in a feeding study with lipid-poor meroplanktonic larvae. Basic knowledge on the feeding ecology of *L. conchilega* larvae is scarcely available, so the other objective of Chapter 3 was to provide information on lipid C assimilation in these organisms.

## **2 MATERIALS AND METHODS**

### **2.1 SAMPLING AREA – THE NORTH SEA AND THE ISLAND OF HELGOLAND**


The North Sea (Fig. 5) is bordered by England, France, Belgium, Holland, Germany, Denmark, Sweden, and Norway. It receives low-salinity discharges from rivers and from the Kattegat sea, and opens broadly to the North Atlantic (Ducrotoy et al., 2000). This large (750 000 km<sup>2</sup>), semi-enclosed body of water supports one of the world’s most active fisheries (IMM, 1997). The North Sea can be divided into four key areas. The northern North Sea (0-500 m depth) is strongly influenced by oceanic inflow and is mostly stratified in the summer. The southern North Sea (0-50 m depth) receives a strong riverine discharge and has a well-mixed depth layer (ICES, 2016). The English Channel is adjacent to the southern North Sea and opens to the Atlantic Ocean. The Skagerrak and Kattegat region links the North Sea to the Baltic Sea (ICES, 2016).



**Figure 5:** Location of Helgoland, a German island in the southern part of the North Sea. The dot between Helgoland and Düne, named “Kabeltonne”, indicates the area where sampling for the Helgoland Roads time-series programme takes place. Source: Pizzetti et al., 2011.

**MATERIALS AND METHODS**

The North Sea is an area with intensive human activities, such as fishing (Fig. 6), shipping, wind farms, aggregate (sand and gravel) extraction and oil and gas production (ICES, 2016). Its food web is characterized by high primary production, which in turn is consumed by the zooplankton and benthos, and passed on to higher trophic levels such as birds, fish, and mammals (ICES, 2016; Fig. 6). It can be currently considered as perturbed, as the larger fish species which used to be the major predators have been depleted by the fishing industry and are now absent or present in reduced numbers in the North Sea (ICES, 2016).



**Figure 6:** Major components of the North Sea food web and significant interactions with fishing fleets, as modelled with statistical tGAMs. Source: modified from ICES, 2016.

The island of Helgoland is located in the German Bight, in the southern North Sea (Fig. 5). It lies approximately 60km from the mainland and is at the mouth of the River Elbe. Being home to a

rich fauna and flora, Helgoland caught the attention of German biologists in the mid-19<sup>th</sup> century. The area surrounding Helgoland has been monitored since 1873 (Wiltshire and Manly, 2004), and in 1892 a marine laboratory (formerly Biologische Anstalt Helgoland (BAH), currently a part of the Alfred-Wegener-Institute für Polar- und Meeresforschung (AWI)) was established in the island (Fischer, 2013). It was only in 1962 that a long term monitoring programme was established to record pelagic data on phytoplankton species, salinity, nutrients, and water temperature and transparency (Hickel et al., 1993). The Helgoland Roads time-series, as it is known, is one of the longest datasets in the North Sea, and the one with the shortest time intervals, and is kept up by the BAH (Hickel et al., 1993; Wiltshire et al., 2015). Zooplankton, rocky shore macroalgae, macrozoobenthos and pelagic bacteria are also included in the current sampling scheme (Wiltshire et al., 2010 and references therein). Samples are collected at the site denominated Kabeltonne (54°11' N, 07°54' E), in the narrow channel between the main island of Helgoland and the small, sandy island of Düne nearby (Fig.5).

### 2.2 ZOOPLANKTON SAMPLING

Zooplankton samples were collected for experiments with *T. longicornis* females and with *L. conchilega* larvae by horizontal hauls with a 500- $\mu$ m mesh-size CalCOFI net. Tows were conducted for 15 minutes at 5m depth at the Kabeltonne site off the German island of Helgoland (54°11'N, 07°54'E). The samples were then taken to the laboratory, where intact and active individuals were immediately sorted under an Olympus SZX16 stereoscopic microscope.

The quantity of *T. longicornis* females needed for the full experimental design required that the experiment be split into two parts. Samplings were performed as close in time as environmental conditions allowed on 17 (Experiment I) and 30 (Experiment II) May 2016. A total of 1300 females were sorted at each date ( $t_{0h}$ ), 1080 for the experiments, 30 for analysis of *in situ* body C and N contents, 150 for determination of *in situ* lipid composition and FA-specific <sup>13</sup>C isotopic enrichment, and 40 were fixed in 4% formalin buffered with hexamethylenetetramine for measurement of prosome length (PL) under a Leica M205C. Adult male copepods were also sorted and added to the experiments for continuous fertilization of females.

## **MATERIALS AND METHODS**

---

*Lanice conchilega* larvae were sorted after the sampling on 6 June 2016. A total of 390 individuals were sorted, 30 for determination of *in situ* body C and N contents, 150 for determination of *in situ* lipid composition and FA-specific  $^{13}\text{C}$  isotopic enrichment, and 210 for the feeding experiment.

### 2.3 PREY CULTURE


Three species were cultured in the laboratory, the diatom *C. weissflogii*, the heterotrophic dinoflagellate *O. marina*, and the cryptophycean *Rhodomonas salina* (Wislouch) (Hill and Wetherbee, 1989). The latter was used solely as food for *O. marina*, whereas the diatoms and dinoflagellates were used to feed copepods and polychaetes. Detailed descriptions on how these species were cultured and sampled are available in Chapter II.

A stock solution was maintained for each of the three species. New batch cultures were created daily for five consecutive days by diluting part of the stock solution with fresh nutrient-replete f/2 medium (after Guillard, 1975) or fresh N-depleted medium (f/2 without the addition of nitrate). Silicate was only added to the medium used for diatom culture. Diatom and cryptophycean cultures were directly labelled with  $^{13}\text{C}$  by adding  $^{13}\text{C}$ -enriched sodium bicarbonate ( $\text{NaH}^{13}\text{CO}_3$ ) to the medium. *Oxyrrhis marina* was indirectly labelled with  $^{13}\text{C}$  via daily feeding on labelled *R. salina*. Cultures were kept in a temperature-controlled room at 18°C. *R. salina* and *C. weissflogii* were provided constant light, while *O. marina* was kept in the dark. Cultures were grown for 5 days and then used as food suspension for copepods during experiments. Cell densities were determined with a BD Accuri C6 Flow Cytometer. *C. weissflogii* and *O. marina* cultures were sampled (filtered) daily for determination of cell C, N, and lipid contents and of FA-specific  $^{13}\text{C}$  isotopic enrichment. Filters for determination of prey C and N content were dried at 60°C for 48 h, folded inside aluminum foil, and stored in a desiccator until analysis. Samples for FA analyses were placed into pre-combusted lipid vials and stored at -80°C. The remaining volume of the cultures was then used in the feeding experiments at concentrations of  $8 \cdot 10^3$  and  $2 \cdot 10^3$  cells  $\text{mL}^{-1}$  for diatom and dinoflagellate diets, respectively.

2.4 EXPERIMENTAL DESIGNS

2.4.1 Feeding experiments with *T. longicornis* females

Experiments were initiated right after field sampling. Figure 7 depicts the distribution of different diet treatments, replication and the sampling scheme. Copepods from Experiment I were fed with diatoms cultured in nutrient-replete (Diat+) and N-depleted (Diat-) medium. In Experiment II the copepods were fed with dinoflagellates cultured with nutrient-replete (Dino+) and N-depleted (Dino-) food (*R. salina*, as described above). Prey were offered to copepods ad libitum ( $>350 \mu\text{g C L}^{-1}$ ,  $8 \cdot 10^3$  diatom and  $2 \cdot 10^3$  dinoflagellate cells  $\text{mL}^{-1}$ ) during the entire experiment. The only exceptions were the respiration and excretion incubations (described below), which were performed without any food.



**Figure 7:** Experimental design used in the feeding experiments with females of the copepod *Temora longicornis* (Experiments I and II) and with larvae of the polychaete *Lanice conchilega* (Experiment III). All experiments took place immediately after zooplankton sampling. The experimental design was almost equal between Experiments I and II, except that the former was conducted with diet treatments (names in bold) consisting of diatoms cultured in nutrient-rich (Diat+) and N-limited (Diat-) conditions, whereas the latter was conducted with dinoflagellates fed with nutrient-rich (Dino+) and N-limited (Dino-) prey. Experiment III was conducted only with nutrient-rich diatom (Diat+) and dinoflagellate (Dino+) cultures. The timing of experimental phases in Experiments I and II (acclimation period, between  $t_{0h}$  and  $t_{48h}$ ; incubations, between  $t_{48h}$  and  $t_{72h}$ ; and remaining feeding time, between  $t_{48h}$  and  $t_{120h}$ ) is shown in the left column. Experiment III consisted only of feeding time (between  $t_{0h}$  and  $t_{120h}$ ). Copepods and

## ***MATERIALS AND METHODS***

---

polychaete larvae were kept in triplicates. Sampling of copepods and polychaete larvae for carbon (C) and nitrogen (N) contents and for lipid composition, and of copepods for prosome length (PL) measurement, are included in the diagram, as are the estimations of vital rates for copepods. Source: modified from Franco-Santos et al., 2018 (Chapter I).

Copepods were incubated in groups of 180 females in partially filled 3 L plastic beakers (75 females L<sup>-1</sup>) fitted with a 300 µm meshed-bottom cylinder to keep the copepods from feeding on eggs and fecal pellets (FP). Although the in situ temperature was different between Experiments I and II (10 and 12°C, respectively), all experiments were conducted in the same temperature, and copepods were kept in a dark temperature-controlled room at 10 ± 0.3°C. The water was gently stirred three times a day for food resuspension. Partial water exchanges (66%) were performed daily in order to remove eggs and FP and to renew copepod food, which was done immediately after exchanges. The copepods were acclimated under these conditions for 48 h. Subsequently, 60 females were sampled from each replicate, 10 individuals to determine body C and N contents and 50 for lipid composition and FA-specific <sup>13</sup>C isotopic enrichment (samples referred to as *t*<sub>48h</sub>). Some of the remaining individuals were distributed among four experimental incubations in order to separately measure grazing, respiration, excretion and egg and FP production rates (methodologies described in more detail in Chapter I). These were then used to construct C and N budgets and to calculate C and N turnover and efficiencies for copepods. Experimental units were kept in the same temperature-controlled room as the beakers. The remaining copepods were kept in their respective beakers, under the same conditions as during the acclimation, and fed another three days before being sampled (referred to as *t*<sub>120h</sub>) to determine body C and N contents, lipid composition, and FA-specific <sup>13</sup>C isotopic enrichment. Copepods were gently washed in distilled water, placed into either pre-weighed tin cartridges for body C and N content determination or pre-combusted lipid vials for FA analyses, and stored at -80°C until further analysis. Tin capsules with copepods were dried and stored in a desiccator until analysis.

### *Grazing*

Ingestion rates were estimated based on changes in prey cell concentrations in the absence and presence of copepods (Frost, 1972). The prey cell concentrations at the beginning and end (after 24 h) of the grazing incubation were measured from bottles containing prey or prey+copepods.



At the end of the incubation copepods were sieved and sampled to measure body C and N contents; the food suspension was fixed immediately with 4% buffered formalin for cell densities determination with a BD Accuri C6 Flow Cytometer. Cell densities were used to calculate grazing rates as number of prey cells ingested per female and per day.

### *Respiration*

The sealed chamber method was applied to measure oxygen consumption rates (Harris et al., 2000). The O<sub>2</sub> saturation of water was measured at the beginning and end (after 24 h) of the respiration incubation from bottles which contained aerated artificial seawater (ASW) or ASW+copepods. After the end of the incubation the copepods were sampled to measure body C and N contents. O<sub>2</sub> consumption rates were obtained per female and per day.

### *Excretion*

The sealed chamber method was also used to calculate daily, individual excretion rates. Bottles with ASW and ASW+copepods were incubated for 24 h, after which their contents were filtered and immediately frozen at -20°C. The copepods were trapped inside the syringe filters and could not be sampled for C and N content analysis. Since both the excretion and the respiration incubations were conducted under similar conditions, the C and N contents of the copepods from the latter were also used for calculating the excretion rates. The equation from Miller and Glibert (1998) was used to calculate the total dissolved nitrogen (TDN) and the non-purgeable organic carbon (NPOC) content of filtrate samples.

### *Egg and faecal pellet production*

For each diet treatment, females were placed individually in 12-well cell culture plates filled with filtered seawater and food suspension. The production of eggs and FP was checked under a binocular 1, 6, 12, 18 and 24 h after the beginning of the experiment. After eggs and feces were counted, they were carefully pipetted out of the well, at times being collected for determination of their C and N contents. Food was immediately replenished until the next count, and plates were left undisturbed. Copepods were sampled after 24 h for determination of body C and N contents. Individual egg production rates (EPR) were calculated as a daily estimate based on the production observed over 24 h. FP production rates (FPR) were also observed during 24 h, but

## ***MATERIALS AND METHODS***

---

due to problems during the last 6 h period, only the data from the first 18 h of incubation were used to estimate daily rates. Filters with eggs and FPs were dried at 60°C for 48 h, folded inside aluminum foil, and stored in a desiccator until analysis.

### *Carbon and nitrogen budgets, turnover rates, and efficiencies*

The amount of energy ingested by an organism should equal the sum of the amounts of energy egested and used for growth and metabolism. In that sense, the vital rates of an individual can also be expressed as a balanced equation, such that  $I = G + R + U + E + F$ , where  $I$  is ingestion,  $G$  is somatic growth,  $R$  is respiration,  $U$  is excretion,  $E$  is egg production, and  $F$  is FP production. In order to do so, the vital rates were converted to C and N units ( $\mu\text{g C or N female}^{-1} \text{ day}^{-1}$ ), as described in detail in Chapter I.

The budget data were further standardized to daily C and N turnover rates (% body C or N  $\text{day}^{-1}$ ), as described in Chapter I. An arcsine square root transformation was then applied to the calculated percentage values. Given that there were negative values in the N growth term, and that the transformation is not possible for negative numbers, a fixed, minimum value was added to all treatments in that category to ensure values above zero.

Different physiological efficiencies were calculated to evaluate copepod performance in relation to food quality: absorption efficiency ( $AE_C$  and  $AE_N$ ), the percentage of ingested C or N (respectively) absorbed in the gut of copepods; and net somatic growth efficiency ( $NSGE_C$  and  $NSGE_N$ ) and net egg production efficiency ( $NEPE_C$  and  $NEPE_N$ ), the percentage of absorbed C and N (respectively) that was converted into copepod somatic growth and into reproduction, respectively. The equations and a more detailed description of how efficiencies were calculated are given in Chapter I.

### 2.4.2 Feeding experiment with *L. conchilega* larvae

Experiments were initiated right after field sampling (Fig. 7). Polychaete larvae were kept for five days in triplicate 500 mL glass beakers fitted with a 300  $\mu\text{m}$  meshed-bottom cylinder (140  $\text{ind L}^{-1}$ ). Only nutrient-replete cultures of *C. weissflogii* and of *O. marina* were used for this feeding experiment. The diatom and dinoflagellate food suspensions were provided on a daily

basis at a concentration of  $8 \times 10^3$  and  $2 \times 10^3$  cells  $\text{mL}^{-1}$ , respectively. Individuals were kept in a dark temperature-controlled room at the temperature recorded when they were sampled in the field,  $13.5 \pm 0.3^\circ\text{C}$ . The beakers were gently stirred three times a day for food resuspension in the water. Partial water exchanges (66%) were performed daily and followed by the addition of new food suspension. At the end of the experiment the larvae were sampled from each replicate for body C and N content determination and for analysis of FA composition and FA-specific  $^{13}\text{C}$  isotopic enrichment (10 and 50 individuals per sample, respectively). Individuals were gently washed in distilled water, placed into either pre-weighed tin cartridges (body C and N content determination) or pre-combusted lipid vials (FA analyses), and stored at  $-80^\circ\text{C}$  until further analysis. Tin cartridges with larvae were dried at  $60^\circ\text{C}$  for 48 h and stored in a desiccator until analysis.

## 2.5 ANALYTICAL WORK

### 2.5.1 Carbon, nitrogen, and FA content analyses

The C and N contents of all copepod, prey culture, egg, and FP samples were measured with an elemental analyzer (detection limit:  $2 \mu\text{g C} / 0.5 \mu\text{g N}$ ; maximum error:  $\pm 3\%$ , Euro EA 3000, EuroVector S.P.A., Milan, Italy) using acetanilide as a standard.

The extraction and determination of fatty acid methyl esters (FAMES) in the samples (according to Boissonnot et al., 2016) is described in detail in Chapter III. The chromatograms generated for the FAMES were evaluated with the ChemStation software from Agilent. The A:B(n-X) shorthand notation was used to refer to FAs, where A is the number of carbon atoms, B is the number of double bonds, and (n-X) gives the position of the double bond closest to the terminal methyl group. Total lipid contents of zooplankton and prey cultures were calculated by adding the mass of all FAs in a sample.

### 2.5.2 Compound-specific stable isotope analysis (CSIA)

The FA-specific stable isotope composition of carbon in FAMES extracted was obtained according to Boissonnot et al. (2016) with a Thermo gas chromatography combustion-isotope ratio mass spectrometry (GC-c-IRMS) system, equipped with a Trace GC Ultra gas

## ***MATERIALS AND METHODS***

---

chromatograph, a GC Isolink, and a Delta V Plus isotope ratio mass spectrometer connected via a Conflo IV interface (Thermo Scientific Corporation, Bremen, Germany). The chromatograms containing peak areas and C isotope ratios were obtained with the (instrument-specific) software Isodat 3.0. The 14:0 and 18:0 FAME reference standards (Iowa University) with known  $\delta$ -values were used for further calculations.

The equations used to calculate carbon assimilation (following Boissonnot et al., 2016) are described in detail in Chapter III.

### 2.5.3 Statistical analyses

Statistical analyses applied to results from the feeding experiments with *T. longicornis* females are described in Chapters I and II, and Chapter III contains the analyses regarding results from the feeding experiment with *L. conchilega* larvae.

Problems with the isotopic enrichment of nutrient-replete *O. marina* cultures in all the feeding experiments hindered the use of the data collected from these treatments for statistical analyses. In the case of *T. longicornis*, data for females fed with the Dino+ diet are presented, but are not included in some analyses, as described in Chapter II. For *L. conchilega*, analyses were only performed on data from the diatom cultures and from larvae fed with them, as described in Chapter III.

### **3 PUBLICATIONS**

The following overview encompasses a publication as first author and another as co-author, and two additional chapters, to be published as first author. These constitute the main part of my PhD thesis. The minor part of the thesis consisted of similar experiments with crab zoeae in Helgoland and with copepods in the Arctic, and of isotopic labelling experiments with the three algae species. The project under which this PhD was developed was first idealized by PD Dr. Barbara Niehoff, Dr. Marleen De Troch, and PD. Dr. Holger Auel for the Doctoral Programme on Marine Ecosystem Health and Conservation - MARES. I conducted the entire laboratory work for chapters I, II, and III, and one of the experiments for chapter IV, at the BAH, in Helgoland. I processed all of my samples at the BAH and at the AWI in Bremerhaven (at the Polar Biological Oceanography and Ecological Chemistry sections). I conducted the data analyses for chapters I, II, and III at the AWI in Bremerhaven and at UGhent in Ghent, Belgium. My contribution (% total workload) for the elaboration of each chapter is also stated below.

#### CHAPTER I

Franco-Santos RM, Auel H, Boersma M, De Troch M, Meunier CL, Niehoff B

#### **Bioenergetics of the copepod *Temora longicornis* under different nutrient regimes**

Experimental concept and design: 70% - the experiments were mainly designed by me and BN, with valuable input from the co-authors.

Acquisition of experimental data: 100% - I gathered all the experimental data during laboratory experiments I performed at the BAH during April-June 2016.

Data analysis and interpretation: 75% - I processed and analyzed the experimental at the BAH, AWI, and UGhent, and BN and HA helped with interpretation.

Preparation of figures and tables: 90% - I prepared all figures and tables for the paper and all co-authors provided feedback and valuable input and suggestions.

## ***PUBLICATIONS***

---

Drafting of the manuscript: 75% - I wrote the manuscript and received scientific and editorial advice from all co-authors.

Journal of Plankton Research, volume 40(4), doi:10.1093/plankt/fby016, published in May 2018.

### CHAPTER II

Franco-Santos RM, Auel H, Boersma M, De Troch M, Graeve M, Meunier CL, Niehoff B

#### **Life history strategies of the copepod *Temora longicornis* as inferred from lipid carbon assimilation experiments**

Experimental concept and design: 70% - the experiments were mainly designed by me and BN, with valuable input from the co-authors.

Acquisition of experimental data: 100% - I gathered all the experimental data during laboratory experiments I performed at the BAH during April-June 2016.

Data analysis and interpretation: 70% - I processed and analyzed the experimental at the BAH, AWI, and UGhent. All co-authors helped with interpretation.

Preparation of figures and tables: 90% - I prepared all figures and tables for the paper and all co-authors provided feedback and valuable input and suggestions.

Drafting of the manuscript: 75% - I wrote the manuscript and received scientific and editorial advice from all co-authors.

The manuscript is under preparation for submission to the journal Functional Ecology.

### CHAPTER III

Franco-Santos RM, Auel H, Boersma M, De Troch M, Graeve M, Meunier CL, Niehoff B

#### **Fatty acid bioconversion and homeostasis in the larvae of the sand mason worm *Lanice conchilega***

Experimental concept and design: 90% - the experiments were designed by me and BN.

Acquisition of experimental data: 100% - I gathered all the experimental data during laboratory experiments I performed at the BAH during April-June 2016.

Data analysis and interpretation: 80% - I processed and analyzed the experimental at the BAH, AWI, and UGhent, and HA helped with interpretation.

Preparation of figures and tables: 90% - I prepared all figures and tables for the paper and all co-authors provided feedback and valuable input and suggestions.

Drafting of the manuscript: 85% - I wrote the manuscript and received scientific and editorial advice from all co-authors.

The manuscript was submitted to PLoS ONE on October 31<sup>st</sup> 2018.

#### CHAPTER IV

Boersma M, Mathew KA, Niehoff B, Schoo KL, Franco-Santos RM, Meunier CL

#### **Temperature driven changes in the diet preference of omnivorous copepods: no more meat when it's hot?**

Experimental concept and design: 0% - the experiments were conceived by MB, BN, and KLS.

Acquisition of experimental data: 10% - I carried out one of the experiments of the study.

Preparation of figures and tables: 10% - I provided feedback on the tables and figures that were prepared for the paper.

Drafting of the manuscript: 10% - I provided scientific and editorial feedback to the drafts.

Ecology Letters, volume 19, doi: 10.1111/ele.12541, published in November 2015.





## CHAPTER I

### **Bioenergetics of the copepod *Temora longicornis* under different nutrient regimes**

Franco-Santos RM, Auel H, Boersma M, De Troch M, Meunier CL, Niehoff B

Published in Journal of Plankton Research, 40(4), doi:10.1093/plankt/fby016, in May 2018.





*J. Plankton Res.* (2018) 40(4): 420–435. First published online May 21, 2018 doi:10.1093/plankt/fby016

# Bioenergetics of the copepod *Temora longicornis* under different nutrient regimes

R. M. FRANCO-SANTOS<sup>1,2,4\*</sup>, H. AUEL<sup>2</sup>, M. BOERSMA<sup>2,3</sup>, M. DE TROCH<sup>4</sup>, C. L. MEUNIER<sup>3</sup> AND B. NIEHOFF<sup>1\*</sup>

<sup>1</sup>POLAR BIOLOGICAL OCEANOGRAPHY SECTION, ALFRED WEGENER INSTITUTE HELMHOLTZ CENTRE FOR POLAR AND MARINE RESEARCH, AM HANDELSHAFFEN 12, 27570 BREMERHAVEN, GERMANY, <sup>2</sup>UNIVERSITY OF BREMEN, MARINE ZOOLOGY, BREMEN, GERMANY, <sup>3</sup>ALFRED WEGENER INSTITUTE FOR POLAR AND MARINE RESEARCH, BIOLOGISCHE ANSTALT HELGOLAND, HELGOLAND, GERMANY AND <sup>4</sup>GHENT UNIVERSITY, MARINE BIOLOGY, GENT, BELGIUM

\*CORRESPONDING AUTHOR: rsantos@awi.de, ritinhamf@hotmail.com (R.M. Franco-Santos); barbara.niehoff@awi.de (B. Niehoff)

Received January 3, 2018; editorial decision May 2, 2018; accepted May 8, 2018

Corresponding Editor: Xabier Irigoien

The copepod *Temora longicornis* depends on constant prey availability, but its performance also depends on how efficiently it utilizes its food sources. Our research goal was to understand copepod energy allocation in relation to diet quality. The working hypothesis was that *Temora* performs better on the diet whose elemental ratio is closest to its own. Diatoms (Diat) and dinoflagellates (Dino) cultured in nutrient-replete (+) and nitrogen-depleted (–) conditions were fed to the copepods. Ingestion, respiration, excretion and egg and fecal pellet production rates were measured. Carbon (C) and nitrogen (N) budgets were built to investigate differences in dietary C and N partitioning. Copepods fed nitrogen-depleted diatoms (Diat–), which had the most different C:N ratio to that of *Temora longicornis*, had high metabolic losses and low growth. Copepods fed nitrogen-rich dinoflagellates (Dino+) with a more similar C:N ratio to their own also had high metabolic losses, but displayed the highest investment into somatic growth and egg production. The results indicate that dinoflagellates are a better food source for *T. longicornis*. Furthermore, consumption of low-quality food leads to higher respiration rates and faster leakage of dissolved organic carbon from copepod fecal pellets; and egestion is a main pathway in copepods for eliminating unabsorbed and non-metabolized carbon.

KEYWORDS: bioenergetics; budget; carbon; nitrogen; stoichiometry

## INTRODUCTION

Copepods are an important link in the energy transfer between primary producers and higher trophic levels,

and contribute to the cycling of organic matter (Juul-Pedersen *et al.*, 2006; Castellani and Altunbas, 2014). The small calanoid copepod *Temora longicornis* (Müller, 1785) is one of the dominant species in the coastal

zooplankton community in the North Atlantic and North Sea, reaching peak densities during the spring and early summer (Hickel, 1975; Castellani and Altunbas, 2014). Off Helgoland, this omnivorous species is abundant all year round and its grazing may have a substantial impact on the phytoplankton standing stock (Gentsch *et al.*, 2009; Maar *et al.*, 2004). *Temora longicornis* has high metabolic turnover rates, but is unable to accumulate significant amounts of energy reserves (Kreibich *et al.*, 2008, 2011). The species is, thus, dependent on a constant availability of prey and is vulnerable to fluctuations in food supply (Helland *et al.*, 2003; Kreibich *et al.*, 2008, 2011). This can be particularly problematic in systems such as the North Sea, where plankton community composition can change rapidly (Kiørboe and Nielsen, 1994). As *T. longicornis* inhabits dynamic systems, it must be able to quickly react to changes in trophic conditions (Gentsch *et al.*, 2009; Kreibich *et al.*, 2008), and efficient food utilization is of paramount importance for its survival.

Food quantity and quality influence the physiology of copepods, their efficiency in nutrient uptake, and their ability to both convert food into energy and channel stored energy into reproduction (Møller, 2007; Hessen and Anderson, 2008; Jónasdóttir *et al.*, 2009), and changes in these prey characteristics could impact not only copepod populations but also the recruitment of their predators (Boersma *et al.*, 2015). Zooplankton are often limited by food quantity in coastal regions (Hirst and Bunker, 2003), and some food sources are also of lower nutritional value (Sterner and Schulz, 1998). It has been suggested, for example, that protozoans are qualitatively important to copepod diet (Stoecker and Capuzzo, 1990), and that egg hatching success is dependent upon the ingestion of essential fatty acids (Broglio *et al.*, 2003). Elemental composition, digestion resistance and biochemical composition are important factors determining the nutritional value of food particles (Sterner and Schulz, 1998).

A vast body of literature is available on the functional responses of calanoids to diet quality (e.g. Dam and Lopes, 2003; Arendt *et al.*, 2005; Jónasdóttir *et al.*, 2009; Nobili *et al.*, 2013), but few address all important vital rates at once (e.g. Abou Debs, 1984). This comprehensive approach would allow for the assessment of energy allocation to different processes in copepods, and is necessary to come to a better understanding of the potential responses of *T. longicornis* to climate change-induced food regime shifts. The present work investigated metabolism, feeding, growth and reproduction in *T. longicornis* females in light of different prey elemental compositions/limitations. Based on the concepts of ecological stoichiometry, homeostasis and trophic upgrading

(Klein Breteler *et al.*, 1999; Sterner and Elser, 2002; Malzahn *et al.*, 2010), specifically on how prey elemental composition affects consumer performance and energy utilization, the research goal was to investigate the partitioning of dietary carbon (C) and nitrogen (N) in relation to food quality. The working hypothesis was that copepods feeding on a diet with a C:N ratio close to their own body composition would perform best, i.e. would have the highest possible growth and reproduction rates and the lowest possible egestion, respiration and excretion rates. The opposite pattern would thus be observed for copepods feeding on prey with a C:N ratio as different from their own as possible (higher or lower).

## METHOD


### Field sampling

Zooplankton were collected by horizontal hauls with a 500- $\mu$ m mesh-size CalCOFI net, which were conducted for 15 minutes at 5 m depth off the German island of Helgoland (54°11'N, 07°54'E), in the southern North Sea. The quantity of females needed for the full experimental design required that the experiment be split into two parts. Samplings were performed as close in time as environmental conditions allowed on 17 (Experiment I) and 30 (Experiment II) May 2016. The samples were taken to the laboratory, where intact and active adult females of *T. longicornis* were immediately sorted under an Olympus SZX16 stereoscopic microscope. A total of 1150 females were sorted at each date ( $t_{0h}$ ), 1080 for the experiments, 30 for determination of *in situ* body C and N contents and another 40 were fixed in 4% formalin buffered with hexamethylenetetramine for measurement of prosome length (PL) under a Leica M205C (Fig. 1).

### Prey culture

Three prey species were cultured in the laboratory, the cryptophycean *Rhodomonas salina* (Wislouch) (Hill and Wetherbee, 1989), the heterotrophic dinoflagellate *Oxymnis marina* (Dujardin, 1841) and the diatom *Conticribra weissflogii* (Grunow) (Stachura-Suchoples and Williams, 2009). *Rhodomonas salina* was used solely as food for *O. marina*, whereas the diatoms and dinoflagellates, which are both common prey for *T. longicornis* (Evans, 1981), were used to feed copepods.

A stock solution was maintained for each of the species. New cultures were created daily by diluting part of the stock solution with fresh medium. Two types of medium were used: nutrient-replete *f/2* (after Guillard, 1975) and N-depleted (*f/2* without the addition of



**Fig. 1.** Experimental design used in the present study with females of the copepod *Temora longicornis*. The two zooplankton sampling dates in May 2016 are indicated, after which Experiments I and II immediately took place. The experimental design was almost equal between these, except that the former was conducted with diet treatments (names in bold) consisting of diatoms cultured in nutrient-rich (Diat+) and N-limited (Diat-) conditions, whereas the latter was conducted with dinoflagellates fed with nutrient-rich (Dino+) and N-limited (Dino-) prey. The timing of experimental phases (acclimation period, between  $t_{0h}$  and  $t_{48h}$ , and incubations, between  $t_{48h}$  and  $t_{72h}$ ) is shown in the left column. Copepods were acclimated and incubated in triplicates. Sampling of copepods for C and N analysis and PL measurement are included in the diagram, as are the estimations of vital rates.

nitrate). Silicate was only added to the medium used for diatoms. Cultures were kept in a temperature-controlled room at 18°C. *Rhodomonas salina* and *C. weissflogii* were provided constant light, while *O. marina* was kept in the dark. Aeration was provided to *R. salina*, and *C. weissflogii* and *O. marina* cultures were stirred twice a day to keep cells suspended. Cultures were grown for 5 days and then used as food suspension for copepods during experiments. Daily feeding of *O. marina* with *R. salina* was planned such that prey cells were depleted by the dinoflagellate on Day 5. Cell densities were determined with a BD Accuri C6 Flow Cytometer. *Conticribra weissflogii* and *O. marina* cultures were sampled daily for the determination of cell C and N contents by filtering known cell concentrations through Whatman GF/F filters. The remaining volume was then used to feed copepods.

### Experimental design

Experiments were initiated right after field sampling. Figure 1 depicts the distribution of different diet treatments, replication and the sampling scheme. Copepods from Experiment I were fed diatoms cultured in nutrient-replete (Diat+) and N-depleted (Diat-) medium. In Experiment II, the copepods were fed with dinoflagellates cultured with nutrient-replete (Dino+) and N-depleted (Dino-) food. The effects of food

quality on predators are more evident when food is abundant than when prey quantity is low (Sterner, 1997). In order to better observe the effects of food quality on the copepods, and to be able to distinguish them from the effects of food quantity, prey were offered to copepods *ad libitum* ( $>350 \mu\text{g C L}^{-1}$ ,  $8 \times 10^3$  diatom and  $2 \times 10^3$  dinoflagellate cells  $\text{mL}^{-1}$ ) during the entire experiment. The only exceptions were the respiration and excretion incubations, which were performed without any food, as described below. Copepod density varied in the different experimental units for practical reasons, such as the amount of copepods available for performing the experiments and the volumes of the different experimental units themselves. Visual observations of female behavior during the experiments and the results obtained indicate that these differences in density did not affect copepod feeding.

Copepods were incubated in groups of 180 females in partially filled 3 L plastic beakers ( $75 \text{ females L}^{-1}$ ) fitted with a 300  $\mu\text{m}$  meshed-bottom cylinder to keep the copepods from feeding on eggs and fecal pellets (FP). Although the *in situ* temperature was different between Experiments I and II (10 and 12°C, respectively), all experiments were conducted in the same temperature, and copepods were kept in a dark temperature-controlled room at  $10 \pm 0.3^\circ\text{C}$ . The water was gently stirred three times a day for food resuspension. Partial

water exchanges (66%) were performed daily in order to remove eggs and FP and to renew copepod food. The copepods were acclimated under these conditions for 48 h. Subsequently, 10 females were sampled from each replicate to determine body C and N contents (samples referred to as  $t_{48h}$ ). The remaining individuals were distributed among four experimental incubations in order to separately measure grazing, respiration, excretion and egg and FP production rates. Experimental units were kept in the same temperature-controlled room as the beakers.

### Grazing

Ingestion rates were estimated based on changes in prey cell concentrations in the absence and presence of copepods (Frost, 1972). The prey cell concentrations at the beginning and end of the grazing incubation were measured from triplicate 1 L glass bottles containing filtered seawater (FSW) + the respective food suspension (“start bottles”) and FSW + food suspension + 10 copepods (“grazed bottles”), respectively. The start bottles were fixed immediately with 4% buffered formalin. Prey growth during the incubation was accounted for by creating another set of three 1 L glass bottles with FSW + food suspension (“control bottles”). The control and grazed bottles were attached onto a plankton wheel rotating with speed between 0.5 and 1 rpm for 24 h, after which their contents were fixed with formalin as described above. The copepods were retained in 300  $\mu$ m mesh-sized sieves prior to fixation and sampled to measure body C and N contents. Cell densities of the preserved food suspension were determined within 6 to 8 days of fixation with a BD Accuri C6 Flow Cytometer. These were used to calculate grazing rates as number of prey cells ingested per female and per day.

### Respiration

The sealed chamber method was applied to measure oxygen consumption rates (Harris *et al.*, 2000) with a NTH oxygen microsensor (PreSens GmbH, Regensburg, Germany) connected to a 4-channel oxygen meter (Microx TX3, PreSens GmbH). Calibration of the microsensor was performed with aerated artificial seawater (ASW, salinity 32) as the 100% O<sub>2</sub> reference and with a saturated Na<sub>2</sub>SO<sub>3</sub> solution as the 0% O<sub>2</sub> reference. In order to avoid complicated corrections for O<sub>2</sub> production or consumption by prey (Ikeda, 1976), this incubation was conducted without prey. For each treatment, ASW was added to six 60-mL Winkler bottles. Three of these were replicate control bottles and contained only ASW and the other three were replicates with ASW and 10 copepods each. Females were placed inside the bottles and allowed to acclimate for an hour with the lid open.

The O<sub>2</sub> saturation of water was then measured, and the lid was closed. After 24 h, it was measured again, and the copepods were sampled to measure body C and N contents. O<sub>2</sub> consumption rates (under starvation conditions) were obtained per female and per day. The use of control bottles allowed for correction for potential oxygen consumption or production by microbes during incubation. Air pressure values for Helgoland were obtained from [www.wetter.com](http://www.wetter.com).

### Excretion

Daily, individual excretion rates were also calculated following the sealed chamber method, and the equation from Miller and Glibert (1998) was used to calculate the total dissolved nitrogen (TDN) and the non-purgeable organic carbon (NPOC) content of filtrate samples. Non-feeding conditions were also employed in this incubation in order to create a low background against which to measure and compare changes in dissolved organic carbon (DOC) and nitrogen (DON). For each treatment, triplicate 100-mL (pre-combusted) glass bottles were prepared with only ASW (controls) and with ASW + ca. 15 copepods. After 24 h of incubation, ASW and copepods were filtered through pre-cleaned (10% HCl) syringes fitted with Whatman GF/F w/GMF (0.7  $\mu$ m pore size, 25 mm diameter) syringe filters. Filtrates were immediately frozen at  $-20^{\circ}\text{C}$  in pre-cleaned high-density polyethylene bottles. DOC and TDN in the filtrate were determined by high temperature catalytic oxidation (HTCO) and subsequent non-dispersive infrared spectroscopy and chemiluminescence detection using a Shimadzu TOC-VCPN analyzer. In the autosampler, the samples (6.5 mL) were acidified with HCl and sparged with oxygen (100 mL min<sup>-1</sup>) for 5 min to remove inorganic carbon. A 50- $\mu$ L sample volume was injected directly on the catalyst (heated to 680°C). Final DOC concentrations were averaged values of triplicate measurements. If the standard variation or the coefficient of variation exceeded 0.1  $\mu$ M or 1%, respectively, up to two additional analyses were performed and outliers were eliminated. After each batch of six samples, one DSR (Deep Sea Water Reference Material, Hansell Research Lab, University of Miami, US), one Milli-Q blank, and one potassium hydrogen phthalate standard were measured. The limit of quantification was 7  $\mu$ M for DOC and 11  $\mu$ M for TDN, and the accuracy was  $\pm 5\%$ . The copepods were trapped inside the syringe filters and could not be sampled for C and N content analysis. Since both the excretion and the respiration incubations were conducted under similar conditions, the C and N contents of the copepods from the latter were also used for calculating the excretion rates.

### Egg and FP production

For each diet treatment, females were placed individually in triplicate 12-well cell culture plates (12 females per replicate) filled with FSW and food suspension to 4 mL volume. The production of eggs and FP was checked under a binocular 1, 6, 12, 18 and 24 h after the beginning of the experiment. After eggs and feces were counted, they were carefully pipetted out of the well, at times being collected in filters for determination of their C and N contents. Due to the high amount of eggs and feces required for C and N analysis, only one pooled sample was taken from each of the different treatments, with a minimum of 400 eggs and 379 FP. Due to time limitations, eggs and FP were not washed in distilled water before sampling, but the presence of algal cells in the samples was unlikely, since most of the food had been eaten by copepods at the time of collection. Food was immediately replenished until the next count, and plates were left undisturbed. Copepods were sampled after 24 h for determination of body C and N contents. Individual egg production rates (EPRs) were calculated as a daily estimate based on the production observed over 24 h. FP production rates (FPRs) were also observed during 24 h, but due to problems during the last 6 h period, only the data from the first 18 h of incubation were used to estimate daily rates.

### C and N content analysis

Copepods, prey cultures, eggs and FP were sampled for the determination of their C and N contents. Copepods were gently washed in distilled water, placed into pre-weighed tin cartridges and stored at  $-80^{\circ}\text{C}$  until further analysis. *In situ* water samples and prey cultures were filtered onto pre-combusted ( $500^{\circ}\text{C}$  for 24 h) Whatman GF/F filters (0.7  $\mu\text{m}$  pore size, 25 mm diameter). Eggs and FP were counted and pipetted onto pre-combusted Whatman GF/F filters. Both tin cartridges and filters with samples were dried at  $60^{\circ}\text{C}$  for 48 h, folded (filters inside aluminum foil), and stored in a desiccator. The C and N contents of all samples were later measured with an elemental analyzer (detection limit: 2  $\mu\text{g}$  C / 0.5  $\mu\text{g}$  N; maximum error:  $\pm 3\%$ , Euro EA 3000, EuroVector S.P.A., Milan, Italy) using acetanilide as a standard.

### C and N budgets

The amount of energy ingested by an organism should equal the sum of the amounts of energy egested and used for growth and metabolism. In that sense, the vital rates of an individual can also be expressed as a balanced equation, such that  $I = G + R + U + E + F$ , where  $I$  is ingestion,  $G$  is somatic growth,  $R$  is respiration,  $U$  is

excretion,  $E$  is egg production and  $F$  is FP production. In order to do so, the vital rates were converted to C and N units. The number of prey cells ingested and of eggs and FP produced were multiplied by their respective C and N contents. Respiratory quotients (RQs), which ranged from 0.74 to 0.76, were estimated for each replicate (as described by McConnaughey, 1978) and used to convert the  $\text{O}_2$  consumption into C-equivalent respiration (Harris *et al.*, 2000). The excretion incubations did not allow for the differentiation between C properly excreted (in the form of urea and amino acids, for example) and C leaked from the FP before absorption by the copepods. The majority of the DOC measured was likely leaked from FP, and thus added to the calculation of  $F$ . The  $G$  term was obtained by the formula  $G = (\text{X}_{t_{48\text{h}}} - \text{X}_{t_{0\text{h}}}) / (\mathcal{N} * \Delta t)$ , where  $\text{X}_{t_{48\text{h}}}$  and  $\text{X}_{t_{0\text{h}}}$  are the C or the N content of the copepod samples from  $t_{48\text{h}}$  and  $t_{0\text{h}}$ , respectively,  $\mathcal{N}$  is the number of females in the sample and  $\Delta t$  is the time in days between  $t_{48\text{h}}$  and  $t_{0\text{h}}$ . Values were compared in the format  $\mu\text{g}$  C or N female $^{-1}$  day $^{-1}$ . Some organisms have a limited ability to store C relative to other elements, which is especially true for many copepod species (Meunier *et al.*, 2014), and will be in excess of this element when they are supplied a diet with more C than they can absorb and/or utilize (Hessen and Anderson, 2008). Therefore, the discussion on the C budget also approaches the topic of excess C and the pathways for returning this excess C to the environment (i.e. eliminating it).

### Turnover rates

The budget data were further standardized to daily C and N turnover rates (% body C or N day $^{-1}$ ). This was achieved by dividing the budget values by the median C or N contents of the copepods sampled at  $t_{48\text{h}}$  and at the end of each incubation (to account for possible weight losses during the incubations). Somatic growth was derived from weight differences between  $t_{0\text{h}}$  and  $t_{48\text{h}}$ , so the budget values for this specific turnover rate were divided by the median C or N contents of the copepods sampled at  $t_{0\text{h}}$  and  $t_{48\text{h}}$  (to account for possible weight gain during the acclimation). An arcsine square root transformation was then applied to the calculated percentage values. Given that there were negative values in the  $\mathcal{N}$  growth term, and that the transformation is not possible for negative numbers, a fixed, minimum value was added to all treatments in that category to ensure values above zero.

### Efficiencies

Different physiological efficiencies were calculated to evaluate copepod performance in relation to food quality. Absorption efficiency (AE), the percentage of ingested

carbon absorbed in the gut of copepods, was calculated as  $AE = 1 - (F/I)$ , where  $F$  is the egestion and  $I$  is the ingestion (Harris *et al.*, 2000). The DOC excreted by copepods was added to the  $F$  term in the carbon AE to account for leakage from FP. The net somatic growth efficiency (NSGE) and the net egg production efficiency (NEPE) indicated the percentage of absorbed food converted into somatic growth and egg production, respectively. The formulas were adapted from Kjørboe *et al.* (1985) and Wendt and Thor (2015) as  $NSGE = G/(G + E + R + U)$  and  $NEPE = E/(G + E + R + U)$ , where  $G$  is the somatic growth,  $E$  is the egg production,  $R$  is the respiration and  $U$  is the excretion.

### Statistical analysis

Differences in seston and in copepod *in situ* elemental composition and PL were verified with  $t$ -tests. Differences in vital rates, turnover rates and efficiency calculations were tested with one-way analysis of variance (ANOVA). Differences in copepod elemental composition during the experiments were tested with a two-way ANOVA, in order to account for interactions between diet treatment and sampling time. When these results were significant, the Tukey HSD *post hoc* test was used at 95% confidence limits to further identify origin of differences. Prior to the ANOVA, the data were tested for normality and homogeneity of variances with Shapiro–Wilk and Bartlett tests, respectively. If the data were non-normal and/or heteroscedastic, they were analyzed with a Kruskal–Wallis test and with the *post hoc* Nemenyi test (with  $P$ -value being determined with the Tukey method). Analyses were performed using R ver. 3.2.5 (Ihaka and Gentleman, 1993).

## RESULTS

### C and N contents

#### Prey culture

The C and N contents and molar C:N ratio of prey cultures are presented in Table I. The average C content

was always above  $500 \mu\text{g C L}^{-1}$ , confirming the *ad libitum* feeding condition. The average N content was always above  $100 \mu\text{g N L}^{-1}$ , with the exception of the Diat– treatment, which was clearly N-limited and contained less than half this amount. The molar C:N ratio was also similar between all treatments with the exception of the Diat–, such that the approximate values were 6 and 19 for the former and the latter, respectively. Cultures were thus significantly different from one another (Supplementary Tables S1 and S2).

#### Copepods

The PL of *in situ* females was significantly greater in Experiment II (Table II, Supplementary Table S1), but their elemental compositions were not statistically different.

Copepod C and N contents varied between 13 and  $21 \mu\text{g C copepod}^{-1}$  and between 3 and  $5 \mu\text{g N copepod}^{-1}$ , and molar C:N ratios ranged from 4.0 to 5.6 (Table II). Although no significant differences were found between the  $t_{0h}$  samples from Experiments I and II, the C and N contents and the molar C:N ratio were significantly different when comparing  $t_{0h}$  with  $t_{48h}$  and  $t_{72h}$ , both within and between experiments (Supplementary Tables 1 and 2). A significant increase in C and N contents was observed between  $t_{0h}$  and  $t_{48h}$ , except in the Diat– treatment, for which the N content remained the same throughout the experiment. C and N contents of copepods sampled between  $t_{48h}$  and the end of the incubations remained constant in the grazing incubation (Fig. 2, Table II, Supplementary Table S2), for which surplus conditions were maintained. Some level of food limitation was present in the egg and FP production rate incubations, given the small volume of the experimental unit, and copepods in the respiration incubation were food-deprived. In both incubations, the C and N contents of copepods decreased (Table II, Supplementary Table S2). The highest increase in molar C:N ratio was observed for the Diat– treatment (from 4.7 at  $t_{0h}$  to 5.6 at the end of the grazing incubation), which was always significantly different from all other treatments (Fig. 2, Table II, Supplementary Table S2).

Table I: C and N contents and molar C:N ratio of the prey cultures (nutrient-replete and N-depleted *C. weissflogii*, Diat+ and Diat–, and *O. marina*, Dino+ and Dino–, respectively). Values are mean  $\pm$  standard deviation from the three batches (temporal replicates) used for feeding in each experiment

Experiment	Treatment	Prey culture				
		$\mu\text{g C L}^{-1}$	$\text{pg C cell}^{-1}$	$\mu\text{g N L}^{-1}$	$\text{pg N cell}^{-1}$	C:N ratio
I	Diat+	$572 \pm 55$	$72 \pm 7$	$105 \pm 18$	$13 \pm 2$	$6.4 \pm 0.5$
	Diat–	$696 \pm 46$	$87 \pm 6$	$43 \pm 4$	$5 \pm 1$	$19.1 \pm 2.2$
II	Dino+	$906 \pm 179$	$453 \pm 90$	$179 \pm 26$	$89 \pm 13$	$5.9 \pm 0.3$
	Dino–	$534 \pm 44$	$267 \pm 22$	$108 \pm 4$	$54 \pm 2$	$5.8 \pm 0.4$



**Table II:** C and N contents (in  $\mu\text{g C copepod}^{-1}$  and  $\mu\text{g N copepod}^{-1}$ ), and molar C:N ratio of *T. longicornis*. Values presented for samples obtained at  $t_{0h}$  (in situ condition),  $t_{48h}$  (after 48h of acclimation) and after the experimental incubations on Day 3 (grazing = G; respiration = R; and egg and FP production = EF). PL (in mm) are shown for  $t_{0h}$  copepods. Diet treatments: single cultures of *C. weissflogii* (Diat) and *O. marina* (Dino) in nutrient-replete (+) and N-depleted (–) conditions. Values presented are mean  $\pm$  standard deviation of triplicate samples containing between 10 and 12 copepods each (except for PLs, which had 40 replicates each)

Treatment	Carbon				Nitrogen				C:N				PL $t_{0h}$			
	$t_{0h}$	$t_{48h}$	G	R	EF	$t_{0h}$	$t_{48h}$	G	R	EF	$t_{0h}$	$t_{48h}$		G	R	EF
Diat+	16 $\pm$ 1	20 $\pm$ 0	20 $\pm$ 1	18 $\pm$ 1	18 $\pm$ 1	4 $\pm$ 0	5 $\pm$ 0	5 $\pm$ 0	4 $\pm$ 0	4 $\pm$ 0	4.7 $\pm$ 0.2	4.9 $\pm$ 0.1	5.0 $\pm$ 0.0	4.8 $\pm$ 0.1	4.8 $\pm$ 0.0	0.99 $\pm$ 0.1
Diat–	15 $\pm$ 0	19 $\pm$ 1	20 $\pm$ 0	17 $\pm$ 1	17 $\pm$ 2	4 $\pm$ 0	4 $\pm$ 0	4 $\pm$ 0	4 $\pm$ 0	4 $\pm$ 0	4.4 $\pm$ 0.1	5.4 $\pm$ 0.2	5.6 $\pm$ 0.1	5.1 $\pm$ 0.1	5.5 $\pm$ 0.1	1.05 $\pm$ 0.1
Dino+	15 $\pm$ 0	20 $\pm$ 0	20 $\pm$ 1	15 $\pm$ 2	18 $\pm$ 2	4 $\pm$ 0	5 $\pm$ 0	5 $\pm$ 0	4 $\pm$ 1	4 $\pm$ 0	4.4 $\pm$ 0.1	4.8 $\pm$ 0.0	4.8 $\pm$ 0.0	4.6 $\pm$ 0.0	4.9 $\pm$ 0.0	4.6 $\pm$ 0.1
Dino–	15 $\pm$ 0	18 $\pm$ 1	18 $\pm$ 1	15 $\pm$ 1	15 $\pm$ 1	4 $\pm$ 0	5 $\pm$ 0	5 $\pm$ 0	4 $\pm$ 0	4 $\pm$ 0	4.4 $\pm$ 0.1	4.6 $\pm$ 0.1	4.7 $\pm$ 0.0	4.6 $\pm$ 0.1	4.6 $\pm$ 0.0	4.6 $\pm$ 0.0

**Eggs and FP**


Diatom-fed copepods produced eggs with the highest C content and the highest and the lowest N content (72 and 69 ng C  $\text{egg}^{-1}$  and 13 and 9 ng N  $\text{egg}^{-1}$  for nutrient-replete and N-depleted treatments, respectively). Dinoflagellate-fed copepods produced eggs with similar C and N contents (60 ng C  $\text{egg}^{-1}$  and 12 and 11 ng N  $\text{egg}^{-1}$  for Dino+ and Dino–, respectively) The molar C:N ratio of eggs was similar between the Diat+ (6.5) and Dino– (6.4) diets, and lower and higher in the Dino+ (5.8) and Diat– (8.9) treatments, respectively.

The FP from diatom-fed copepods had the lowest C and N contents (26 and 20 ng C  $\text{FP}^{-1}$  and 4 and 2 ng N  $\text{FP}^{-1}$  for nutrient-replete and N-depleted treatments, respectively). The C and N contents of the FP from dinoflagellate-fed copepods were almost twice as high (52 and 48 ng C  $\text{FP}^{-1}$  and 9 and 7 ng N  $\text{FP}^{-1}$  for Dino+ and Dino–, respectively). The molar C:N ratio of FP was lower in the Dino+ (6.7), Diat+ (7.6) and Dino– (8.0) diets than in the Diat– treatment (11.7). It was not possible to measure FP sizes, however, visual observations indicate that FPs from copepods fed with diatoms were slightly longer and thicker than those produced by copepods fed with dinoflagellates.

**Vital rates and turnover rates**

Diatom-fed copepods had significantly higher average ingestion rates ( $8.6 \pm 0.5 \times 10^4$  and  $10.1 \pm 1.2 \times 10^4$  cells  $\text{female}^{-1} \text{day}^{-1}$  for Diat+ and Diat–, respectively) than the dinoflagellate-fed copepods ( $2.4 \pm 0.8 \times 10^4$  and  $1.6 \pm 0.5 \times 10^4$  cells  $\text{female}^{-1} \text{day}^{-1}$  for Dino+ and Dino–, respectively) (Supplementary Tables S1 and S3). C ingestion, which ranged from 17 to 86% body C  $\text{day}^{-1}$ , was significantly higher in the Dino+ treatment than in the Dino–, whereas N ingestion, which ranged from 12 to 66% body N  $\text{day}^{-1}$  (Table III), was significantly higher in the Dino+ diet than in the N-depleted treatments (Supplementary Tables S1 and S3).

The average oxygen consumption rates were lower in the diatom treatments ( $3.4 \pm 0.3$  and  $4.3 \pm 0.1 \text{ mL O}_2 \text{ female}^{-1} \text{day}^{-1}$  for Diat+ and Diat–, respectively) than in the dinoflagellate treatments ( $6.0 \pm 1.3$  and  $4.2 \pm 0.4 \text{ mL O}_2 \text{ female}^{-1} \text{day}^{-1}$  for Dino+ and Dino–, respectively). They corresponded to a C-equivalent respiration of 6–18% body C  $\text{day}^{-1}$  (Table III). A significant difference in  $\text{O}_2$  consumption and turnover of respired C was only found between the nutrient-replete treatments, with higher values in the Dino+ diet (Supplementary Tables S1 and S3 and Table III). Surprisingly, the average N excretion values were higher in the N-depleted treatments ( $1.0 \pm 0.3$  and  $0.6 \pm 0.1 \mu\text{g N}$



**Fig. 2.** C and N contents and molar C:N ratio of *T. longicornis* females during Experiments I and II. Experiment I tested diet treatments consisting of diatoms cultured in nutrient-rich (Diat+, gray squares and dotted lines) and N-depleted (Diat-, black squares and full lines) conditions, whereas Experiment II tested diets consisting of dinoflagellates fed with nutrient-rich (Dino+, gray triangle and dotted lines) and N-limited (Dino-, black triangle and dotted lines) prey. Samples were taken of *in situ* conditions ( $t_{0h}$ , empty symbols), after 48h of acclimation ( $t_{48h}$ ), and after 24h of incubations ( $t_{72h}$ ). Bars represent mean values  $\pm$  one standard deviation.

female<sup>-1</sup> day<sup>-1</sup> for the Diat- and Dino-, respectively) than in the nutrient-replete diets ( $0.2 \pm 0.2$  and  $0.3 \pm 0.1$  µg N female<sup>-1</sup> day<sup>-1</sup> for the Diat+ and Dino+, respectively). They corresponded to 5–32% body N day<sup>-1</sup> (Table III) and were significantly lower in the Diat- treatment (Supplementary Tables S1 and S3).

The average FP production rates were not significantly different between the Diat+ ( $89 \pm 4$  pellets female<sup>-1</sup> day<sup>-1</sup>) and Diat- ( $86 \pm 8$  pellets female<sup>-1</sup> day<sup>-1</sup>) treatments. FP production rates recorded for copepods fed with dinoflagellates were lower,  $72 \pm 5$  and  $49 \pm 1$  pellets female<sup>-1</sup> day<sup>-1</sup> for the Dino+ and Dino- treatments, respectively (Supplementary Tables S1 and S3). The average amount of DOC leaked from FP, which was significantly different between all treatments (Supplementary Table S1), was higher in the N-depleted diets ( $1.6 \pm 0.2$  and  $1.1 \pm 0.1$  µg C female<sup>-1</sup> day<sup>-1</sup> for Diat- and Dino-, respectively) than in the nutrient-replete treatments ( $0.2 \pm 0.1$  and  $0.7 \pm 0.1$  µg C female<sup>-1</sup> day<sup>-1</sup> for Diat+ and Dino+, respectively). The C and N turnover rates for FP production, which varied between 12 and 24% body C day<sup>-1</sup> and between 4 and 14% body N day<sup>-1</sup> (Table III), were significantly different between almost all treatment pairs (Supplementary Tables S1 and S3).

Somatic growth was significantly different between treatments in terms of both C and N (Supplementary Table S1). It ranged from 4 to 14% body C day<sup>-1</sup> and from 0 to 11% body N day<sup>-1</sup> (Table III), with the lowest values recorded for the N-limited diatom treatment (Supplementary Table S3). The average EPRs were significantly higher in the Dino+ treatment ( $61 \pm 4$  eggs female<sup>-1</sup> day<sup>-1</sup>) than for copepods fed with the other diets ( $39 \pm 9$ ,  $36 \pm 4$ , and  $41 \pm 5$  eggs female<sup>-1</sup> day<sup>-1</sup> for Diat+, Diat- and Dino-, respectively) (Supplementary Tables


S1 and S3). The egg C turnover ( $13$ – $21\%$  body C day<sup>-1</sup>) was similar between treatments (Table III), but the N turnover ( $7$ – $16\%$  body N day<sup>-1</sup>) was significantly different between the Dino+ and Diat- treatments (Supplementary Tables S1 and S3).

### C and N budgets

The C and N budgets obtained for each of the treatments are represented in Fig. 3. They were mostly unbalanced, which is represented in the figure by the equation symbols “-” and “+”. The “*T*” values are represented on the left side of the copepod antenna, whereas the right side represents the sum of “*G* + *R* + *U* + *E* + *P*”. When the gain of energy via feeding was higher than growth and energy expenditure (e.g. Fig 3B and G), the “+” and “-” signs are placed on the left and right sides of the antenna, respectively. When the gain of energy was lower than growth and energy expenditure (e.g. Fig 3D and F), the “+” and “-” signs are placed on the right and left sides of the antenna, respectively. The symbol “=” was used for the C budget which was nearly balanced (Fig. 3C). To the right of these symbols is a percentage, which indicates the percentage of energy gained via feeding that was recorded as being used. The budgets recorded for the Dino+ treatment seem to have the best estimates (percentages close to 100%). The budgets for the other treatments mostly point to higher growth and energy expenditure. The Dino- treatment had the most deviating results, with expenses amounting to twice as much the recorded energy ingested, a pattern visually represented by the use of double equation symbols (Fig. 3D and H). The Diat+ budgets also show higher expenses than energy gain (Fig. 3A and E), whereas the Diat- budgets show a mixed response, with an

**Table III:** C and N turnover rates and AE, NSGE and NEPE efficiencies from experimental incubations with *T. longicornis*. Diet treatments: single cultures of *C. weissflogii* (Diat) and *O. marina* (Dino) in nutrient-replete (+) and N-depleted (–) conditions. Mass-specific values for ingestion (I), growth (G), carbon-equivalent respiration (R), excretion (U), egg production (E) and FP production (F) turnover rates are expressed as % body C or N day<sup>-1</sup>, and efficiencies as %. Values presented are mean ± standard deviation.

Treatment	Carbon										Nitrogen									
	I	G	R	E	F	AE	NSGE	NEPE	I	G	U	E	F	AE	NSGE	NEPE				
Diat+	32 ± 1	11 ± 1	7 ± 1	15 ± 4	13 ± 1	60 ± 3	33 ± 4	45 ± 5	24 ± 1	9 ± 1	5 ± 9	11 ± 3	8 ± 0	68 ± 1	36 ± 11	48 ± 6				
Diat–	49 ± 6	6 ± 2	10 ± 0	14 ± 1	19 ± 2	64 ± 7	20 ± 5	47 ± 2	14 ± 2	0 ± 1	27 ± 5	8 ± 1	5 ± 0	67 ± 7	–1 ± 4	23 ± 4				
Dino+	65 ± 23	13 ± 1	14 ± 3	19 ± 1	23 ± 1	64 ± 12	27 ± 2	44 ± 4	50 ± 17	10 ± 1	8 ± 4	15 ± 1	13 ± 1	73 ± 9	29 ± 3	48 ± 7				
Dino–	27 ± 10	10 ± 2	10 ± 1	14 ± 1	20 ± 1	36 ± 5	29 ± 5	42 ± 2	21 ± 8	9 ± 2	15 ± 3	11 ± 1	8 ± 0	60 ± 12	25 ± 6	32 ± 3				



**Fig. 3.** Carbon (A–D) and nitrogen (E–H) budgets obtained for *T. longicornis* females feeding on diatoms cultured in nutrient-rich (Diat+) and N-depleted (Diat–) conditions and on dinoflagellates fed with nutrient-rich (Dino+) and N-limited (Dino–) prey. Values given represent daily ingestion, egestion, respiration, excretion, reproduction and somatic growth rates for one individual copepod, and are visually represented by proportionately sized arrows. The comparison between the amount of C and N ingested by copepods (“I” values on the right side of the copepod antenna) and the amount used by them (the sum of “G + R + U + E + F” values on the left side of the copepod antenna) is represented in the figure by the equation symbols “+”, “–” and “=”. When the gain of energy via feeding was higher than growth and energy expenditure (B and G), the “+” and “–” signs are placed on the left and right sides of the antenna, respectively. When the gain of energy was lower than growth and energy expenditure (A, D, E, F and H), the “+” and “–” signs are placed on the right and left sides of the antenna, respectively. The symbol “=” was used for the C budget which was nearly balanced (C). To the right of these symbols is a percentage, which indicates the percentage of energy gained via feeding that was recorded as being used. The closer to 100%, the more balanced the measurements; values below and above 100% indicated higher energy gain from feeding than energy expenditure and higher energy expenditure than energy gain from ingestion, respectively. The DOC and FP components of the egestion term of the C budget (percentage of total “P”) are indicated in gray and white colors (respectively) in the corresponding arrows.

accurate estimate for the C budget (Fig. 3B) but an inaccurate estimate for the N budget (Fig. 3F). The DOC and FP components of the egestion term of the

C budget (percentage of total “*F*”) are indicated in gray and white colors (respectively) in the corresponding arrows in Fig. 3.

### Efficiencies

The AEs varied between 33% and 75% and between 46% and 81% of C and N ingested, respectively (Table III), with the carbon AE for Dino– being significantly lower than that of Diat– and Dino+ (Supplementary Tables S1 and S3). The carbon NSGEs varied between 15% and 36%, and were significantly different between the diatom treatments, while the nitrogen NSGEs varied from –4% to 48% and were significantly lower in Diat– (Table III, Supplementary Tables S1 and S3). No significant differences were found between the carbon NEPE, which ranged from 39% to 50%, whereas the nitrogen NEPEs, which varied from 19% to 56% (Table III), were significantly higher in the nutrient-replete treatments (Supplementary Tables S1 and S3).

## DISCUSSION

To our knowledge, this is the first study of energy budgets of *T. longicornis* females feeding on prey items of different qualities, which enabled the comparison of copepod performance under different food regimes. Our working hypothesis was that the highest growth and reproduction rates and the lowest catabolic rates would be recorded for copepods feeding on a diet with a C:N ratio as close as possible to their own body composition. Our study was able to partially confirm the working hypothesis. The highest investment in somatic and reproductive growth was indeed shown by copepods fed with a food source of C:N ratio close to their own (Dino+), but so were the highest expenditures with egestion and catabolism, contrary to what had been postulated. Performance entails the amount of energy ingested and used by an individual for different vital activities, but is ultimately defined by what is invested in reproduction and how successful this process is. Thus, the major finding of this study is that dinoflagellates are a food source for copepods of superior or similar quality to diatoms under nutrient-replete or N-depleted conditions, respectively. This is, however, only valid for the temperatures investigated in the present study, as individual metabolism and C requirements are affected by temperature (e.g. Boersma *et al.*, 2016). Results also revealed that egestion is a major pathway for *T. longicornis* females to eliminate the excess C; and that low food quality can influence copepod respiration (regardless of its C-to-nutrient ratio) and the intensity and speed with which DOC leaks from

FP. The fact that results only partially agreed with predictions from stoichiometric theory indicates the need for further investigations into copepod ecophysiology and adaptive capacity to shifting food regimes.

The level of N-limitation achieved in the Diat– treatment was not expected for the Dino– diet, as it has been shown that *O. marina* can regulate its body composition to incorporate the stoichiometric imbalances of its prey in an attenuated form (Malzahn *et al.*, 2010; Meunier *et al.*, 2014). Although the elemental composition of the Dino– treatment was similar to that of the Dino+, its fatty acid profile was not (unpublished data). This could explain why performance was different for copepods fed with these two diets, and why the former constituted a food source of lower quality.

### Copepod feeding

The C ingestion rates reported by Arendt *et al.* (2005) and Jónasdóttir *et al.* (2009) for *T. longicornis* feeding on *C. weissflogii* are twice as high as those reported herein, despite their use of lower prey concentrations (Table IV). The values from our study might be underestimated, as indicated by the comparison of the amount of energy ingested and used by copepods (symbols on the top left and right sides of the copepod antenna in Fig. 3, respectively). Despite measurements being recorded in the same fashion for all treatments, some show a nearly balanced budget (e.g. Fig. 3C) while others reveal unbalanced budgets (e.g. Fig. 3F). We speculate that the ingestion rates are underestimated and that this might be due to copepods feeding on food sources that were unaccounted for, which were ingested in quantities inversely proportional to the quality of the diet treatments. Coprophagy and filial cannibalism are known to occur among copepods, regardless of availability of alternative food, and can reach values of up to 50% of produced FP and 60 eggs female<sup>–1</sup> day<sup>–1</sup> (e.g. Lampitt *et al.*, 1990; Noji *et al.*, 1991; Dam and Lopes, 2003; Boersma *et al.*, 2014) in starved or food-limited conditions. The glass bottles used for the grazing incubation in the present study did not allow for the separation of copepods from eggs and FP. The eggs had a similar or higher nutritional quality (in terms of elemental composition/molar C:N ratio) than the prey items, and the microbiota associated with the peritrophic membrane of the FPs might be a valuable food source for copepods (as suggested by Lampitt *et al.*, 1990), so by feeding on its own eggs and FPs, copepods could have complemented nutritionally inadequate diets. This response would be a form of compensatory behavior for this copepod, and has been identified in other studies (Augustin and Boersma, 2006; Siuda and Dam, 2010) as

**Table IV: Comparative values between prey species and quantity used in laboratory experiments and EPR of *T. longicornis* recorded for that prey in the present study and elsewhere in the literature**

Copepod	Prey		Ingestion		μg C ind <sup>-1</sup> day <sup>-1</sup>	% body C day <sup>-1</sup>	EPR Eggs ind <sup>-1</sup> day <sup>-1</sup>	Reference
	Type	μg C L <sup>-1</sup>	Cells ind <sup>-1</sup> day <sup>-1</sup>	1–4 <sup>a</sup>				
<i>Temora longicornis</i>	Natural prey assemblage	200–800 <sup>a</sup>					2–55	Dam and Peterson (1991) and Peterson and Dam (1996)
	Phytoplankton						50	Castellani and Altunbas (2014)
Dinoflagellates <i>Conticribra weissflogii</i>		349	204 000 ± 70 000	15.6 ± 4.8	124	27.8 ± 7.3	3	Jansen <i>et al.</i> (2006)
		349		6.5	70		11	Arendt <i>et al.</i> (2005)
		225–276			68		5–19	Dam and Lopes (2003)
<i>Oxyrrhis marina</i>		274		2.2 ± 0.2	25		20–60	Jónasdóttir <i>et al.</i> (2009)
		572 ± 55	86 000 ± 500	6.3 ± 0.2	32 ± 1		39 ± 9	Niehoff <i>et al.</i> (2015)
		286		4.7 ± 0.4	51		20–80	This study
		906 ± 179	24 000 ± 800	13.1 ± 4.6	65 ± 23		61 ± 4	Niehoff <i>et al.</i> (2015) van Someren Gréve (2013) This study

<sup>a</sup>Using a carbon:chlorophyll  $\alpha$  ratio of 50, according to Dam and Peterson (1991).

a mechanism for other small calanoids to overcome the elemental limitation of their prey.

### Copepod waste production (or catabolic activities)

Few studies are available on the metabolic activities of small copepods such as *T. longicornis* (e.g. Berner, 1962; Dam and Peterson, 1993; Nobili *et al.*, 2013; Castellani and Altunbas, 2014), all reporting lower O<sub>2</sub> consumption and C-equivalent respiration than the ones described herein (1–2 μL O<sub>2</sub> ind<sup>-1</sup> day<sup>-1</sup> and 1% body C day<sup>-1</sup>). Our respiration and excretion incubations were performed without feeding the copepods, and probably underestimate the rates of fed copepods (Miller and Glibert, 1998; Thor, 2002a, 2002b; Nobili *et al.*, 2013). High respiration rates have been suggested as a mechanism for removal of excess C in unbalanced nutrient conditions (Anderson *et al.*, 2005; Hessen and Anderson, 2008), even though the mechanism itself is yet to be described (Malzahn *et al.*, 2010). The results of the present study partially agree with this assumption. The second highest respiration rates were reported for copepods fed with the N-poor diets, but one of them was enriched in C in relation to N (Diat–) and the other had a C:N ratio close to that of the copepods (Dino–). Furthermore, the highest rates were measured for another one of the diets whose C:N more closely resembled that of the copepods (Dino+). The type of substrate catabolized by starved copepods in order to obtain energy influences respiration (Castellani and Altunbas, 2014) and excretion (Anderson, 1992) rates. Proteins are the main substrates catabolized in small calanoids (Thor, 2002). The RQ of proteins is higher than that of lipids (Gnaiger, 1983) and would, thus, generate higher C-equivalent respiration rates in starved individuals that were previously fed N-rich diets, as observed for *O. marina* by Meunier *et al.* (2012). Our results thus suggest that O<sub>2</sub> consumption increases with decreasing food quality, regardless of an excess of C or not. If prey is of high nutritional quality, then O<sub>2</sub> consumption seems to be modulated by the amount of food ingested.

The preferential utilization of proteins by copepods previously fed with a N-rich diet should also result in higher ammonia excretion rates than those of copepods previously fed N-poor diets, but the opposite was recorded. It is possible that the high food C:N ratio of 19 in the Diat– treatment led to an increase in ingestion rates and a decrease in gut passage time. This, in turn, reduced the efficiency with which nutrients were absorbed and resulted in higher excretion rates (Plath and Boersma, 2001).

FP production by copepods is an important source of POM and DOC in the epipelagic (Møller *et al.*, 2003;

Thor *et al.*, 2003), contributing to microbial production with recycled nutrients (Smetacek, 1980; Strom *et al.*, 1997). Although it has been claimed that undisturbed FP do not release DOC (Strom *et al.*, 1997), it has been shown that 50% of the total C content of FP is released as DOC at the onset of defecation (Thor *et al.*, 2003). DOC can be generated by copepods before absorption, as leakage from egested FP, and after absorption, as excretion in the form of nitrogenous organic compounds (Frangoulis *et al.*, 2004). The methodology did not allow for the separation of both components. Even though some DOC may have been excreted, it was considered that the majority of DOC measured was leaked from FP, which is why DOC values were added to the C content of FP to calculate the egestion term (“*F*”) of the C budget. In our study, the percentage of total C egested (“*F*”) that corresponded to DOC varied widely between treatments, going from approximately 8% in Diat+ to 16% in Dino+, 31% in Dino– and 50% in Diat–. It would seem, thus, that the level of DOC leakage from FP is directly or indirectly influenced by food quality.

Small calanoids such as *T. longicornis* are unable to store significant amounts of energy reserves, and must return to the environment any C consumed in excess of its needs and ability to absorb. Our results suggest that the primary pathway for *T. longicornis* to eliminate excess C was egestion, which removed twice as much C as that burnt via respiration (Fig. 3A–D). Furthermore, if all C had been assimilated and metabolized and no excess was left, assuming that all N was also assimilated, the molar C:N ratio of FP would not have differed from that of the diet treatments (Checkley, 1980). The fact that it was higher in the FP (except for Diat–) indicates that N was used constantly and efficiently for production and that the excess C was egested (Checkley, 1980). This pattern is further supported by the statistically similar nitrogen AE in the treatments, contrary to what stoichiometric theory would have predicted. The carbon AE, on the other hand, was significantly lower in the Dino– treatment. It could be that the resulting biochemical composition of the Dino– diet indirectly caused an inefficient absorption of C in copepods and led to an egestion similar to that of the other treatments despite a lower ingestion rate. Our estimates for the carbon AE were greater than the assimilation efficiency estimates for another small copepod (*A. tonsa*) feeding on the same prey species (Besiktepe and Dam, 2002).

### Copepod growth and reproduction (or anabolic activities)

Most studies assume negligible increases in body weight in adult copepods which do not store energy reserves and only investigate reproductive investment as a

measure of growth (Hirst and McKinnon, 2001). The C and N somatic growth rates observed in the present study, though lower than the EPRs, contest this assumption, as do other studies (Hirst and McKinnon, 2001; Dam and Lopes, 2003). A comparison between the net efficiencies for somatic growth (NSGE) and egg production (NEPE) emphasizes how the majority of assimilated energy is used in reproduction, but also that the portion destined to somatic growth is not negligible. The only absence of somatic growth was observed in terms of N for the Diat– treatment, as expected.

The EPR recorded for *T. longicornis* in this study were within the range of those obtained when this copepod was fed with natural plankton (Peterson and Dam, 1996), similar prey species (Niehoff *et al.*, 2015) and with water from the spring phytoplankton bloom (Castellani and Altunbas, 2014; Peterson and Kimmerer, 1994), but higher than when it was fed water from a dinoflagellate bloom (Jansen *et al.*, 2006) (Table IV). The highest EPR in our study was obtained with the nutrient-rich dinoflagellate diet, as opposed to results from other studies (Turner *et al.*, 2001; Dam and Lopes, 2003; Jónasdóttir *et al.*, 2009). It must be noted that great controversy involves the topic of diatoms being good or bad quality diet items (for reviews, see Jónasdóttir *et al.*, 1998; Paffenhöfer *et al.*, 2005), with many suggesting its suitability for egg production but reduced success for hatching (Miralto *et al.*, 1999; Ban *et al.*, 1997).

The quantity of body C invested into reproduction was similar across treatments, but body N investment was significantly different between the Diat– and the Dino+ treatments. This indicates N-limitation in egg production, as also suggested by Nobili *et al.* (2013) for *T. longicornis*. The nitrogen NEPE observed in our study further indicates a clear limitation of this element in copepods fed with the N-depleted diets. This contrasts the assumption that nutrients in limited amounts should be used with higher efficiency. Other studies have reported similar trends, with lower egg production efficiencies associated with higher food C:N ratios and with P-limitation (Anderson *et al.*, 2005; Nobili *et al.*, 2013). This pattern might be due to increased maintenance requirements for copepods fed with nutrient-limited diets (Anderson *et al.*, 2005; Wendt and Thor, 2015). Even so, the NEPE values reported herein seem to be mostly higher than those obtained from the literature as either egg production efficiency or gross growth efficiency (e.g. Dam and Lopes, 2003; Thor *et al.*, 2007; Nobili *et al.*, 2013; Wendt and Thor, 2015). Few studies have approached the effect of N-limitation on EPR, which has been said to be both deleterious (Koski *et al.*, 2006; Nobili *et al.*, 2013) and advantageous (Augustin and Boersma, 2006). EPR does not provide information

on egg viability (hatching success), so it would be necessary to conduct further investigations in order to understand the importance of diet quality for secondary production by *T. longicornis*.

### Climate change and marine C and N cycles

Long-term monitoring data from Helgoland have shown a drastic decrease in calanoid copepod densities since 1985, a decline most likely caused by the decrease in nutrient (N and P) loading coupled to the increase in light penetration in the region (Boersma *et al.*, 2015). These would have resulted in phytoplankton with higher C:nutrient ratios and, thus, of lower nutritional value for zooplankton (Boersma *et al.*, 2015). Recent studies show controversial results regarding the relative dominance in recent years of diatoms and dinoflagellates in the southern North Sea (Wiltshire *et al.*, 2010; Alvarez-Fernandes *et al.* 2012; Hinder *et al.*, 2012; Boersma *et al.*, 2015). Changes in the abundance of these prey species can affect secondary production, and a further increase in their C:N ratio could have major consequences for food web processes (Jones and Flynn, 2005; Malzahn and Boersma, 2011; Nobili *et al.*, 2013; Boersma *et al.*, 2015). On the other hand, increasing local temperatures (Wiltshire *et al.*, 2010) could result in higher metabolic costs for copepods (Castellani and Altunbas, 2014), which can only be met by an increased C consumption (Boersma *et al.*, 2016; Malzahn *et al.*, 2016). In that scenario, increasing temperatures would dampen the effects of prey with higher C:nutrient ratios (Boersma *et al.*, 2016; Malzahn *et al.*, 2016). In N-limitation conditions (e.g. late spring bloom), dinoflagellates are still of better nutritional value than diatoms (Jones and Flynn, 2005; this study), but at higher temperatures, copepods might need to preferentially consume autotrophs to supply metabolic demand, as observed for *T. longicornis* (Boersma *et al.*, 2016) and *A. tonsa* (Malzahn *et al.*, 2016). All of these variables make it difficult to say at this point how *T. longicornis* (and possibly other calanoid copepods) will react to future changes in prey composition, even though its omnivorous strategy and related adaptations allow it to switch food sources depending on their availability (Daan *et al.*, 1988; Gentsch *et al.*, 2009).

### CONCLUSIONS

The results obtained in the present study indicate that, under nutrient-replete conditions, dinoflagellates such as *O. marina* are a better food source for *T. longicornis* than diatoms. This major finding is, however, dependent upon environmental conditions. Furthermore, low-quality food

also leads to higher respiration rates, regardless of its C-to-nutrient ratio, and to faster leakage of DOC from copepod FP. In addition, egestion seems to be the main pathway for eliminating excess C, contrary to the common belief of excess C being mostly respired. The vital rates measured herein for *T. longicornis* provide important information on this species' food utilization efficiency and maintenance costs in relation to diet quality. The budget approach used herein is not free of flaws, and future research in the field should investigate more complex scenarios (mixed diets, different temperatures, varying food concentrations, include all life stages).

### SUPPLEMENTARY DATA

Supplementary data are available at *Journal of Plankton Research* online.

### ACKNOWLEDGEMENTS

We kindly thank the crew of the Aade research vessel (AWI-BAH) for providing us with zooplankton samples and the technicians at the AWI, BAH and Claudia Burau for the help with sample analyses.

### FUNDING

R.M.F.-S. was supported by a doctoral research grant from the Doctoral Programme on Marine Ecosystem Health and Conservation (MARES, Framework Agreement Number 2011-0016). The work was further supported by Ghent University (BOF-GOA project 01G02617). C.L.M. was supported by the Bundesministerium für Bildung und Forschung (BMBF grant no. 01LN1702A).

### DATA ARCHIVING

The laboratory experimental raw data and metadata for this study are available through the PANGAEA repository under <https://doi.pangaea.de/10.1594/PANGAEA.886050>.

### Ethical standards

The experiments described in the present study comply with the current laws of the country in which they were performed.

## Conflict of interest

The authors declare that they have no conflict of interest.

## REFERENCES

- Abou Debs, C. (1984) Carbon and nitrogen budget of the calanoid copepod *Temora stylifera*: effect of concentration and composition of food. *Mar. Ecol. Prog. Ser.*, **15**, 213–223.
- Alvarez-Fernandes, S., Lindeboom, H. and Meesters, E. (2012) Temporal changes in plankton of the North Sea: community shifts and environmental drivers. *Mar. Ecol. Prog. Ser.*, **462**, 21–38.
- Anderson, T. R. (1992) Modelling the influence of food C:N ratio, and respiration on growth and nitrogen excretion in marine zooplankton and bacteria. *J. Plankton Res.*, **14**, 1645–1671.
- Anderson, T. R., Hessen, D. O., Elser, J. J. and Urabe, J. (2005) Metabolic stoichiometry and the fate of excess carbon and nutrients in consumers. *Am. Nat.*, **165**, 1–15.
- Arendt, K. E., Jónasdóttir, S. H., Hansen, P. J. and Gärtner, S. (2005) Effects of dietary fatty acids on the reproductive success of the calanoid copepod *Temora longicornis*. *Mar. Biol.*, **146**, 513–530.
- Augustin, C. B. and Boersma, M. (2006) Effects of nitrogen stressed algae on different *Acartia* species. *J. Plankton Res.*, **28**, 429–436.
- Ban, S., Burns, C., Castel, J., Chaudron, Y., Christou, E., Escribano, R., Umani, S. F., Gasparini, S. et al. (1997) The paradox of diatom-copepod interactions. *Mar. Ecol. Prog. Ser.*, **157**, 287–293.
- Berner, A. (1962) Feeding and respiration in the copepod *Temora longicornis* (Müller). *J. Mar. Biol. Assoc. U. K.*, **42**, 625–640.
- Besiktepe, S. and Dam, H. G. (2002) Coupling of ingestion and defecation as a function of diet in the calanoid copepod *Acartia tonsa*. *Mar. Ecol. Prog. Ser.*, **229**, 151–164.
- Boersma, M., Mathew, K. A., Niehoff, B., Schoo, K. L., Franco-Santos, R. M. and Meunier, C. L. (2016) Temperature driven changes in the diet preference of omnivorous copepods: no more meat when it's hot? *Ecol. Lett.*, **19**, 45–53.
- Boersma, M., Wesche, A. and Hirche, H. J. (2014) Predation of calanoid copepods on their own and other copepods' offspring. *Mar. Biol.*, **161**, 733–743.
- Boersma, M., Wiltshire, K. H., Kong, S. M., Greve, W. and Renz, J. (2015) Long-term change in the copepod community in the southern German Bight. *J. Sea Res.*, **101**, 41–50.
- Broglio, E., Jónasdóttir, S. H., Calbet, A., Jakobsen, H. H. and Saiz, E. (2003) Effect of heterotrophic versus autotrophic food on feeding and reproduction of the calanoid copepod *Acartia tonsa*: relationship with prey fatty acid composition. *Aquat. Microb. Ecol.*, **31**, 267–278.
- Castellani, C. and Altunbas, Y. (2014) Seasonal change in acclimatised respiration rate of *Temora longicornis*. *Mar. Ecol. Prog. Ser.*, **500**, 83–101.
- Checkley, D. M. (1980) The egg production of a marine planktonic copepod in relation to its food supply: laboratory studies. *Limnol. Oceanogr.*, **25**, 430–446.
- Daan, R., Gonzalez, S. R. and Klein Breteler, W. C. M. (1988) Cannibalism in omnivorous calanoid copepods. *Mar. Ecol. Prog. Ser.*, **47**, 45–54.
- Dam, H. G. and Lopes, R. M. (2003) Omnivory in the calanoid copepod *Temora longicornis*: feeding, egg production and egg hatching rates. *J. Exp. Mar. Biol. Ecol.*, **292**, 119–137.
- Dam, H. G. and Peterson, W. T. (1991) In situ feeding behavior of the copepod *Temora longicornis*: effects of seasonal changes in chlorophyll size fractions and female size. *Mar. Ecol. Prog. Ser.*, **71**, 113–123.
- Dam, H. G. and Peterson, W. T. (1993) Seasonal contrasts in the diel vertical distribution, feeding behavior, and grazing impact of the copepod *Temora longicornis* in Long Island Sound. *J. Mar. Res.*, **51**, 561–594.
- Evans, F. (1981) An investigation into the relationship of sea temperature and food supply to the size of the planktonic copepod *Temora longicornis* Müller in the North Sea. *Estuar. Coast. Shelf Sci.*, **13**, 145–158.
- Frangoulis, C., Christou, E. D. and Hecq, J. H. (2004) Comparison of marine copepod outfluxes: nature, rate, fate and role in the carbon and nitrogen cycles. *Adv. Mar. Biol.*, **47**, 254–309.
- Frost, B. W. (1972) Effects of size and concentration of food particles on the feeding behavior of the marine planktonic copepod *Calanus pacificus*. *Limnol. Oceanogr.*, **17**, 805–815.
- Gentsch, E., Kreibich, T., Hagen, W. and Niehoff, B. (2009) Dietary shifts in the copepod *Temora longicornis* during spring: evidence from stable isotope signatures, fatty acid biomarkers and feeding experiments. *J. Plankton Res.*, **31**, 45–60.
- Gnaiger, E. (1983) Appendix C—calculation of energetic and biochemical equivalents of respiratory oxygen consumption. In Gnaiger, E. and Forstner, H. (eds), *Polarographic Oxygen Sensors: Aquatic and Physiological Applications*. Springer, Berlin, pp. 337–345.
- Guillard, R. R. L. (1975) Culture of phytoplankton for feeding marine invertebrates. In Smith, W. L. and Chanley, M. H. (eds), *Culture of Marine Invertebrate Animals*. Springer, US, pp. 29–60.
- Harris, R., Wiebe, P., Lenz, J., Skjoldal, H. R. and Huntley, M. (eds) (2000) *ICES Zooplankton Methodology Manual*. Academic Press, London.
- Helland, S., Terjesen, B. F. and Berg, L. (2003) Free amino acid and protein content in the planktonic copepod *Temora longicornis* compared to *Artemia franciscana*. *Aquaculture*, **215**, 213–228.
- Hessen, D. O. and Anderson, T. R. (2008) Excess carbon in aquatic organisms and ecosystems: physiological, ecological, and evolutionary implications. *Limnol. Oceanogr.*, **53**, 1685–1696.
- Hickel, W. (1975) The mesozooplankton in the Wadden Sea of Sylt (North Sea). *Helgol. Wiss. Meeres.*, **27**, 254–262.
- Hinder, S. L., Hays, G. C., Edwards, M., Roberts, E. C., Walne, A. W. and Gravenor, M. B. (2012) Changes in marine dinoflagellate and diatom abundance under climate change. *Nat. Clim. Change*, **2**, 271–275.
- Hirst, A. G. and Bunker, A. J. (2003) Growth of marine planktonic copepods: global rates and patterns in relation to chlorophyll  $\alpha$ , temperature, and body weight. *Limnol. Oceanogr.*, **48**, 1988–2010.
- Hirst, A. G. and McKinnon, A. D. (2001) Does egg production represent adult female copepod growth? A call to account for body weight changes. *Mar. Ecol. Prog. Ser.*, **223**, 179–199.
- Ihaka, R. and Gentleman, R. (1993) R (Version 3.2.5) [software]. Retrieved from <https://cran.r-project.org/bin/windows/base/old/3.2.4/>.
- Ikeda, T. (1976) The effect of laboratory conditions on the extrapolation of experimental measurements to the ecology of marine



- zooplankton I. Effect of feeding condition on the respiration rate. *Bull. Plankton Soc. Jpn.*, **23**, 51–60.
- Jansen, S., Riser, C. W., Wassmann, P. and Bathmann, U. (2006) Copepod feeding behaviour and egg production during a dinoflagellate bloom in the North Sea. *Harmful Algae*, **5**, 102–112.
- Jones, R. H. and Flynn, K. J. (2005) Nutritional status and diet composition affect the value of diatoms as copepod prey. *Science*, **307**, 1457–1459.
- Juul-Pedersen, T., Nielsen, T. G., Michel, C., Møller, E. F., Tiselius, P., Thor, P., Olesen, M., Selander, E. *et al.* (2006) Sedimentation following the spring bloom in Disko Bay, West Greenland, with special emphasis on the role of copepods. *Mar. Ecol. Prog. Ser.*, **314**, 239–255.
- Jónasdóttir, S. H., Kiørboe, T., Tang, K. W., St John, M., Visser, A. W., Saiz, E. and Dam, H. G. (1998) Role of diatoms in copepod production: good, harmless or toxic? *Mar. Ecol. Prog. Ser.*, **172**, 305–308.
- Jónasdóttir, S. H., Visser, A. W. and Jespersen, C. (2009) Assessing the role of food quality in the production and hatching of *Temora longicornis* eggs. *Mar. Ecol. Prog. Ser.*, **382**, 139–150.
- Kiørboe, T., Møhlenberg, F. and Hamburger, K. (1985) Bioenergetics of the planktonic copepod *Acartia tonsa*: relation between feeding, egg production and respiration, and composition of specific dynamic action. *Mar. Ecol. Prog. Ser.*, **26**, 85–97.
- Kiørboe, T. and Nielsen, T. G. (1994) Regulation of zooplankton biomass and production in a temperate, coastal ecosystem. 1. Copepods. *Limnol. Oceanogr.*, **39**, 493–507.
- Klein Breteler, W. C. M., Schogt, N., Baas, M., Schouten, S. and Kraay, G. W. (1999) Trophic upgrading of food quality by protozoans enhancing copepod growth: role of essential lipids. *Mar. Biol.*, **135**, 191–198.
- Koski, M., Klein Breteler, W., Schogt, N., Gonzalez, S. and Jakobsen, H. H. (2006) Life-stage-specific differences in exploitation of food mixtures: diet mixing enhances copepod egg production but not juvenile development. *J. Plankton Res.*, **28**, 919–936.
- Kreibich, T., Saborowski, R., Hagen, W. and Niehoff, B. (2008) Short-term variation of nutritive and metabolic parameters in *Temora longicornis* females (Crustacea, Copepoda) as a response to diet shift and starvation. *Helgol. Mar. Res.*, **62**, 241–249.
- Kreibich, T., Saborowski, R., Hagen, W. and Niehoff, B. (2011) Influence of short-term nutritional variations on digestive enzyme and fatty acid patterns of the calanoid copepod *Temora longicornis*. *J. Exp. Mar. Biol. Ecol.*, **407**, 182–189.
- Lampitt, R. S., Noji, T. and Von Bodungen, B. (1990) What happens to zooplankton faecal pellets? Implications for material flux. *Mar. Biol.*, **104**, 15–23.
- Maar, M., Nielsen, T. G., Gooding, S., Tønnesson, K., Tiselius, P., Zervoudaki, S., Christou, E., Sell, A. *et al.* (2004) Trophodynamic function of copepods, appendicularians and protozooplankton in the late summer zooplankton community in the Skagerrak. *Mar. Biol.*, **144**, 917–933.
- Malzahn, A. M. and Boersma, M. (2011) Effects of poor food quality on copepod growth are dose dependent and non-reversible. *Oikos*, **121**, 1408–1416.
- Malzahn, A. M., Doerfler, D. and Boersma, M. (2016) Junk food gets healthier when it's warm. *Limnol. Oceanogr.*, **61**, 1677–1685.
- Malzahn, A. M., Hantzsche, F., Schoo, K. L., Boersma, M. and Aberle, N. (2010) Differential effects of nutrient-limited primary production on primary, secondary or tertiary consumers. *Oecologia*, **162**, 35–48.
- McConnaughey, T. (1978) Ecosystems naturally labelled with Carbon-13: applications to the study of consumer food-webs. Thesis, University of Alaska in Fairbanks, Alaska.
- Meunier, C. L., Haafke, J., Oppermann, B., Boersma, M. and Malzahn, A. M. (2012) Dynamic stoichiometric response to food quality fluctuations in the heterotrophic dinoflagellate *Oxyrrhis marina*. *Mar. Biol.*, **159**, 2241–2248.
- Meunier, C. L., Malzahn, A. M. and Boersma, M. (2014) A new approach to homeostatic regulation: towards a unified view of physiological and ecological concepts. *PLoS ONE*, **9**, e107737.
- Miller, C. A. and Glibert, P. M. (1998) Nitrogen excretion by the calanoid copepod *Acartia tonsa*: results of mesocosm experiments. *J. Plankton Res.*, **20**, 1767–1780.
- Miralto, A., Barone, G., Romano, G., Poulet, S. A., Ianora, A., Russo, G. L., Buttino, I., Mazzarella, G. *et al.* (1999) The insidious effect of diatoms on copepod reproduction. *Nature*, **402**, 173–176.
- Møller, E. F. (2007) Production of dissolved organic carbon by sloppy feeding in the copepods *Acartia tonsa*, *Centropages typicus*, and *Temora longicornis*. *Limnol. Oceanogr.*, **52**, 79–84.
- Møller, E. F., Thor, P. and Nielsen, T. G. (2003) Production of DOC by *Calanus finmarchicus*, *C. glacialis* and *C. hyperboreus* through sloppy feeding and leakage from fecal pellets. *Mar. Ecol. Prog. Ser.*, **262**, 185–191.
- Niehoff, B., Kreibich, T., Saborowski, R. and Hagen, W. (2015) Feeding history can influence physiological responses of copepods: an experimental study comparing different cohorts of *Temora longicornis* from the Southern North Sea. *J. Exp. Mar. Biol. Ecol.*, **469**, 143–149.
- Nobili, R., Robinson, C., Buitenhuis, E. and Castellani, C. (2013) Food quality regulates the metabolism and reproduction of *Temora longicornis*. *Biogeosci. Discuss.*, **10**, 3203–3239.
- Noji, T. T., Estep, K. W., Macintyre, F. and Norrbin, F. (1991) Image analysis of faecal material grazed upon by three species of copepods: evidence for coprorhexy, coprophagy and coprochaly. *J. Mar. Biol. Ass. UK*, **71**, 465–480.
- Paffenhöfer, G. A., Ianora, A., Miralto, A., Turner, J. T., Kleppel, G. S., Ribeira d'Alcalá, M., Casotti, R., Caldwell, G. S. *et al.* (2005) Colloquium on diatom-copepod interactions. *Mar. Ecol. Prog. Ser.*, **286**, 293–305.
- Peterson, W. K. and Kimmerer, W. J. (1994) Processes controlling recruitment of the marine calanoid copepod *Temora longicornis* in Long Island Sound: Egg production, egg mortality, and cohort survival rates. *Limnol. Oceanogr.*, **39**, 1594–1605.
- Peterson, W. T. and Dam, H. G. (1996) Pigment ingestion and egg production rates of the calanoid copepod *Temora longicornis*: implications for gut pigment loss and omnivorous feeding. *J. Plankton Res.*, **18**, 855–861.
- Plath, K. and Boersma, M. (2001) Mineral limitation of zooplankton: stoichiometric constraints and optimal foraging. *Ecology*, **82**, 1260–1269.
- Siuda, A. N. S. and Dam, H. G. (2010) Effects of omnivory and predator-prey elemental stoichiometry on planktonic trophic interactions. *Limnol. Oceanogr.*, **55**, 2107–2116.
- Smetacek, V. S. (1980) Zooplankton standing stock, copepod faecal pellets and particulate detritus in Kiel Bight. *Estuar. Coast. Mar. Sci.*, **11**, 477–490.

- Sterner, R. W. (1997) Modelling interactions of food quality and quantity in homeostatic consumers. *Freshwater Biol.*, **38**, 473–482.
- Sterner, R. W. and Elser, J. J. (2002) *Ecological Stoichiometry: The Biology of Elements from Molecules to the Biosphere*. Princeton University Press, Princeton, NJ.
- Sterner, R. W. and Schulz, K. L. (1998) Zooplankton nutrition: recent progress and a reality check. *Aquat. Ecol.*, **32**, 261–279.
- Stoecker, D. K. and Capuzzo, J. M. (1990) Predation on protozoa: its importance to zooplankton. *J. Plankton Res.*, **12**, 891–908.
- Strom, S. L., Benner, R., Ziegler, S. and Dagg, M. J. (1997) Planktonic grazers are a potentially important source of marine dissolved organic carbon. *Limnol. Oceanogr.*, **42**, 1364–1374.
- Thor, P. (2002) Specific dynamic action and carbon incorporation in *Calanus finmarchicus* copepodites and females. *J. Exp. Mar. Biol. Ecol.*, **272**, 159–169.
- Thor, P., Cervetto, G., Besiktepe, S., Ribera-Maygas, E., Tang, K. W. and Dam, H. G. (2002) Influence of two different green algal diets on specific dynamic action and incorporation of carbon into biochemical fractions in the copepod *Acartia tonsa*. *J. Plankton Res.*, **24**, 293–300.
- Thor, P., Dam, H. G. and Rogers, D. R. (2003) Fate of organic carbon released from decomposing copepod fecal pellets in relation to bacterial production and ectoenzymatic activity. *Mar. Ecol. Prog. Ser.*, **33**, 279–288.
- Thor, P., Koski, M., Tang, K. W. and Jónasdóttir, S. H. (2007) Supplemental effects of diet mixing on absorption of ingested organic carbon in the marine copepod *Acartia tonsa*. *Mar. Ecol. Prog. Ser.*, **331**, 131–138.
- Turner, J. T., Ianora, A., Miralto, A., Laabir, M. and Esposito, F. (2001) Decoupling of copepod grazing rates, fecundity and egg-hatching success on mixed and alternating diatom and dinoflagellate diets. *Mar. Ecol. Prog. Ser.*, **220**, 187–199.
- van Someren Gréve, H. (2013) Feeding performance of the marine calanoid copepod *Temora longicornis*. Thesis, Utrecht University, Netherlands.
- Wendt, I. and Thor, P. (2015) Influence of prey species and concentration on egg production efficiency and hatching success in *Acartia tonsa* Dana (Copepoda, Calanoidea). *Crustaceana*, **88**, 675–687.
- Wiltshire, K. H., Kraberg, A., Bartsch, I., Boersma, M., Franke, H.-D., Freund, J., Gebühr, C., Gerds, G. *et al.* (2010) Helgoland Roads, North sea: 45 years of change. *Estuar. Coast.*, **33**, 295–310.

## CHAPTER II

### **Life history strategies of the copepod *Temora longicornis* as inferred from lipid carbon assimilation experiments**

Franco-Santos RM, Auel H, Boersma M, De Troch M, Graeve M, Meunier CL, Niehoff B

To be submitted for publication to the journal Functional Ecology



## Life cycle strategies of *Temora longicornis*

### 1 Life history strategies of the copepod *Temora longicornis* as inferred from lipid carbon 2 assimilation experiments

3

4 Franco-Santos RM<sup>a,b,d,\*</sup>, Auel H<sup>b</sup>, Boersma M<sup>b,c</sup>, De Troch M<sup>d</sup>, Graeve M<sup>a</sup>, Meunier CL<sup>c</sup>, Niehoff  
5 B<sup>a</sup>

6

7 <sup>a</sup> Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Am Handelshafen 12,  
8 27570, Bremerhaven, Germany

9 <sup>b</sup> University of Bremen, Marine Zoology, Bremen, Germany

10 <sup>c</sup> Alfred-Wegener-Institut, Helmholtz-Zentrum für Polar- und Meeresforschung, Biologische Anstalt  
11 Helgoland, Helgoland, Germany

12 <sup>d</sup> Ghent University, Marine Biology, Gent, Belgium

13

14 \* Corresponding authors, Rita M. Franco-Santos and Barbara Niehoff

15 [rsantos@awi.de](mailto:rsantos@awi.de), [ritinhamf@hotmail.com](mailto:ritinhamf@hotmail.com) +61 0451 644 313

16 [barbara.niehoff@awi.de](mailto:barbara.niehoff@awi.de)

17

### 18 **Abstract**

19 1. The calanoid copepod *T. longicornis* needs to cope with rapidly changing plankton community  
20 composition in the North Sea. These changes can be further aggravated by recent and ongoing  
21 changes in the nutrient regime in the area, so efficient food utilization by copepods is of paramount  
22 importance for their survival.

23 2. The present work aimed at describing qualitatively and quantitatively how differences in the  
24 biochemical composition of prey grown under different nutrient regimes affects the dynamics of  
25 lipid utilization and assimilation and the reproductive output by *T. longicornis* females.

26 3. Copepods were fed prey cultures isotopically enriched with  $^{13}\text{C}$ ; the transfer of lipid C from prey  
27 to consumer was followed; and FA-specific C assimilation and turnover rates were calculated.

28 4. Absolute C assimilation was higher in copepods fed prey whose molar C:N ratio was the most  
29 different (higher) from their own but, due to differences in ingestion rates, relative C assimilation  
30 (as % food ingested) was similar between copepods fed with different diets.

31 5. The assimilation of lipid C by copepods was inversely related to their C investment in egg  
32 production. The copepods which showed highest egg production rates also displayed the highest  
33 levels of the FAs 18:3(n-3) and DHA.

34 6. High lipid C turnover rates indicate an opportunistic feeding strategy for *T. longicornis*, which  
35 would accumulate readily available dietary C with a speed that decreases as saturation is  
36 approached and its lipid storage is replenished. This strategy would enable them to cope with  
37 environmental changes in feeding conditions and to invest in reproduction year round.

38 7. These results are an important addition to a small, yet growing, body of literature which estimates  
39 the transfer of lipid C between prey and predator, and to the field of copepod feeding ecology.

40

41 **Keywords:** assimilation, carbon, compound-specific stable isotope analysis, copepod, fatty acid,  
42 feeding ecology, life history strategies, turnover

43

44

## 45 INTRODUCTION

46 *Temora longicornis* (Müller, 1785) is a small calanoid copepod (approx. 1 mm prosome  
47 length) abundant all year round in the southern North Sea off the German island of Helgoland,  
48 where its grazing can have a substantial impact on the phytoplankton standing stock (Gentsch,  
49 Kreibich, Hagen, and Niehoff, 2009; Maar et al., 2004). It is one of the dominant copepod species  
50 in the North Sea, and plays a pivotal role in the energy transfer from primary producers to higher  
51 trophic levels. Because *T. longicornis* has high metabolic turnover rates and is unable to accumulate

52 significant amounts of energy reserves, it is dependent on a constant availability of prey and  
53 vulnerable to fluctuations in food supply (Helland, Terjesen, and Berg, 2003; Kreibich, Saborowski,  
54 Hagen, and Niehoff, 2008, 2011). Efficient food utilization is, thus, of paramount importance for  
55 the survival of *T. longicornis*, as this species needs to cope with rapidly changing plankton  
56 community composition in the North Sea (Gentsch, Kreibich, Hagen, and Niehoff, 2009; Kiørboe  
57 and Nielsen, 1994; Kreibich, Saborowski, Hagen, and Niehoff, 2008), which can be further  
58 aggravated by recent changes in the nutrient regime in the area (Boersma, Wiltshire, Kong, Greve,  
59 and Renz, 2015).

60 Several studies have been conducted on how diet quality can influence copepod physiology,  
61 reproduction success, nutrient uptake efficiency, and ability to produce and metabolize energy  
62 (Hessen and Anderson, 2008; Jónasdóttir, Visser, and Jespersen, 2009; Klein Breteler, Schogt,  
63 Baas, Schouten, and Kraay, 1999; Møller, 2007). *Temora longicornis* has been reported to  
64 selectively graze according to its physiological requirements, choosing its prey by their biochemical  
65 composition and nutritional value (Cottonnec, Brunet, Sautour, and Thoumelin, 2001). The  
66 nutritional value of a food source can be defined, a.o., by its elemental and biochemical  
67 composition (Sterner and Schulz, 1998).

68 A gap in knowledge still exists on the role and importance of lipids in marine zooplankton,  
69 and on how species cope with inadequate food supply, especially during reproductive periods (Lee,  
70 Hagen, and Kattner, 2006). Studies which apply isotope labelling to specific compounds such as  
71 lipids or proteins (compound-specific stable isotope analysis, CSIA) can obtain rates of carbon (C)  
72 transfer, accumulation, and turnover and information on lipid biosynthesis (Boissonnot, Niehoff,  
73 Hagen, Søreide, and Graeve, 2016; Dalsgaard, John, Kattner, Müller-Navarra, and Hagen, 2003; De  
74 Troch et al., 2012; Graeve, Albers, and Kattner, 2005). FA dynamics in marine invertebrates have  
75 been mostly characterized for larger, often herbivorous, calanoid copepods from polar regions,  
76 whereas few studies are available on the lipid biochemistry of smaller copepod species (Boissonnot,

77 Niehoff, Hagen, Søreide, and Graeve, 2016; Dalsgaard, John, Kattner, Müller-Navarra, and Hagen,  
78 2003; Fraser, Sargent, and Gamble, 1989).

79 The present work thus aimed at describing qualitatively and quantitatively how the  
80 biochemical composition of prey (diatoms and dinoflagellates) grown under different nutrient  
81 regimes affects the dynamics of lipid utilization and assimilation and the reproductive output by *T.*  
82 *longicornis* females. We assumed that lipid C assimilation would be inversely related to  
83 reproductive output and that these are, in turn, affected by the nutritional quality (elemental and  
84 biochemical composition) of the diet. In order to obtain data on lipid dynamics and on reproductive  
85 output, copepods were fed with prey cultures previously labelled with  $^{13}\text{C}$ , which enabled the  
86 calculation of egg production rates, the monitoring of lipid C transfer between prey and predator,  
87 and the calculation of FA-specific C assimilation and turnover rates.

88

89

## 90 **METHODS**

### 91 **Field sampling**

92 Zooplankton were sampled with a 500  $\mu\text{m}$  mesh-size CalCOFI net towed horizontally for 15  
93 minutes at 5 m depth off the German island of Helgoland (54°11'N, 07°54'E), in the southern North  
94 Sea. Samples were taken to the laboratory and intact and active adult females of *T. longicornis* were  
95 immediately sorted under an Olympus SZX16 stereoscopic microscope. A total of 1260 females  
96 were sorted during each sampling date, 1080 for feeding experiments (with diets of varying  
97 nutritional qualities) and 180 for determination of *in situ* body C, N, and lipid contents, FA  
98 composition, and FA-specific  $^{13}\text{C}$  isotopic enrichment. Samples for *in situ* conditions were collected  
99 in triplicates of 10 and 50 individuals each for elemental and biochemical analyses, respectively.  
100 Due to an insufficient number of females during the first sampling date, the sorting procedure was  
101 performed twice, on May 17<sup>th</sup> (Experiment I) and 30<sup>th</sup> (Experiment II), 2016. The effects of the  
102 nutrient-replete and N-depleted diatom treatments (Diat+ and Diat-, respectively) on copepod



103 elemental and biochemical composition were investigated in Experiment I, and the effects of the  
104 nutrient-replete and N-depleted dinoflagellate treatments (Dino+ and Dino-, respectively) were  
105 tested in Experiment II.

106

## 107 **Prey culture**

108 Common prey species for *T. longicornis* (Evans, 1981) were used in the feeding  
109 experiments, the diatom *Conticribra weissflogii* (Grunow) (Stachura-Suchoples and Williams,  
110 2009) and the dinoflagellate *Oxyrrhis marina* (Dujardin, 1841). Batch cultures were created daily  
111 for *C. weissflogii* and *O. marina* for five consecutive days under two nutrient treatments. The  
112 cryptophycean *Rhodomonas salina* (Wislouch) (Hill and Wetherbee, 1989) was also cultured in  
113 batches (created on a daily basis) for five consecutive days to serve as food for the dinoflagellates.  
114 *Conticribra weissflogii* and *R. salina* batches were created by diluting stock solutions with fresh  
115 nutrient replete (f/2) medium and with fresh N-depleted medium (f/2 without nitrate). The two  
116 different types of medium were created (after Guillard, 1975) with seawater filtered in cellulose  
117 acetate filters (0.2 µm pore size). Diatom and cryptophycean cultures were labelled by adding <sup>13</sup>C-  
118 enriched sodium bicarbonate (NaH<sup>13</sup>CO<sub>3</sub>) to the medium at concentrations of 4 and 12 mg L<sup>-1</sup>,  
119 respectively. *Oxyrrhis marina* was labelled via daily feeding on <sup>13</sup>C-labelled *R. salina*. Cell  
120 quantities were adjusted so that cryptophycean cells were depleted by the dinoflagellate on day 5.  
121 Cultures were grown for five days (details can be found in Franco-Santos et al., 2018) and then used  
122 as food suspension for copepods at a concentration of 8 and 2 \*10<sup>3</sup> cells mL<sup>-1</sup> for diatom and  
123 dinoflagellate treatments, respectively.

124 *Conticribra weissflogii* and *O. marina* cultures were sampled daily for determination of cell  
125 C, N, and FA content and of FA-specific <sup>13</sup>C isotopic enrichment by filtering subsamples of known  
126 cell concentrations through pre-combusted (500°C for 24h) Whatman GF/F filters (0.7 µm pore  
127 size, 25 mm diameter). Samples for FA analyses were placed into pre-combusted lipid vials and  
128 stored at -80°C. Filters for determination of prey C and N content were dried at 60°C for 48 h,

129 folded inside aluminum foil, and stored in a desiccator until analysis. Cell densities were  
130 determined with a BD Accuri C6 Flow Cytometer.

131

## 132 **Experimental design**

133 Feeding experiments with *T. longicornis* were initiated immediately after sorting of the  
134 copepods. Copepods were fed each diet for five days in triplicate plastic beakers fitted with a 300  
135  $\mu\text{m}$  meshed-bottom cylinder (75 females  $\text{L}^{-1}$ ). Food was provided on a daily basis after partial water  
136 exchanges (66%). Females from Experiment I were fed with the Diat+ and Diat- diets, and  
137 copepods from Experiment II were fed with the Dino+ and Dino- diets. The copepods were kept at  
138  $10 \pm 0.3^\circ\text{C}$  in a dark temperature-controlled room. This temperature was similar to that recorded in  
139 the field (surface water) during sampling for Experiment I but was approximately  $2^\circ\text{C}$  lower than  
140 that recorded *in situ* during sampling for Experiment II. The beakers were gently stirred three times  
141 a day for food resuspension in the water.

142 Copepods were sampled from each replicate of each treatment for body C and N content  
143 determination and for FA profiling and FA-specific  $^{13}\text{C}$  isotopic enrichment (10 and 50 individuals  
144 per sample, respectively). Sampling was conducted before (day 1, *in situ* condition,  $t_{0\text{h}}$ ), during (day  
145 3,  $t_{48\text{h}}$ ), and at the end (day 6,  $t_{120\text{h}}$ ) of the experiment. Copepods were gently washed in distilled  
146 water, placed into either pre-weighed tin capsules (5\*9 mm, IVA Analysentechnik) for body C and  
147 N content determination or pre-combusted lipid vials for FA analyses, and stored at  $-80^\circ\text{C}$  until  
148 further analysis. Tin capsules with copepods were dried and stored in a desiccator until analysis.

149

## 150 **Carbon, nitrogen and fatty acid content analyses**

151 The C and N contents of all copepod and prey culture samples were measured with an  
152 elemental analyzer (detection limit:  $2 \mu\text{g C} / 0.5 \mu\text{g N}$ ; maximum error:  $\pm 3\%$ , Euro EA 3000,  
153 EuroVector S.P.A., Milan, Italy) using acetanilide as a standard.

154 Fatty acid methyl esters (FAMES) were extracted and identified as described by Boissonnot,  
155 Niehoff, Hagen, Søreide, and Graeve (2016). Samples were homogenized in a  
156 dichloromethane:methanol (2:1, v:v) solution, from which total lipids were extracted. A known  
157 amount of an internal standard, the tricosanoic acid methyl ester (23:0), was added to each sample.  
158 Potassium chloride (0.88% solution) was added to create a biphasic system and aid in extraction.  
159 Lipid extracts were transesterified by heating samples with 3% sulfuric acid in methanol at 80°C  
160 under nitrogen atmosphere for 4 hours. FAMES were then extracted with cyclohexane and  
161 determined with a gas chromatograph (HP 6890 N, Agilent Technologies Deutschland GmbH &  
162 Co. KG) equipped with a 60m × 0.25mm i.d. wall-coated open tubular capillary column (film  
163 thickness: 0.25 µm; liquid phase: DB-FFAP), a split/splitless injector (250°C) and a flame  
164 ionization detector (280°C). The chromatograms generated were evaluated with the ChemStation  
165 software (Agilent). The A:B(n-X) shorthand notation was used to refer to FAs, where A is the  
166 number of carbon atoms, B is the number of double bonds, and (n-X) gives the position of the  
167 double bond closest to the terminal methyl group. *T. longicornis* does not have major storage  
168 reserves and exhibits TAG as its primary neutral lipids (Fraser, Sargent, and Gamble, 1989);  
169 therefore lipid classes were not separated in the present study. Total lipid contents of copepods and  
170 prey cultures were calculated as the sum of the mass from of all FAs.

171

## 172 **Compound-specific stable isotope analysis (CSIA)**

173 The FA-specific stable isotope composition of carbon in FAMES extracted was obtained  
174 according to Boissonnot, Niehoff, Hagen, Søreide, and Graeve (2016) with a Thermo gas  
175 chromatography combustion-isotope ratio mass spectrometry (GC-c-IRMS) system, equipped with  
176 a Trace GC Ultra gas chromatograph, a GC Isolink, and a Delta V Plus isotope ratio mass  
177 spectrometer connected via a Conflo IV interface (Thermo Scientific Corporation, Bremen,  
178 Germany). The chromatograms containing peak areas and C isotope ratios were obtained with the

179 (instrument-specific) Isodat 3.0 software. The 14:0 and 18:0 FAME reference standards (Iowa  
180 University) with known  $\delta$ -values were used for further calculations.

181 The equations used by Boissonnot, Niehoff, Hagen, Søreide, and Graeve (2016) to calculate  
182 carbon assimilation were also applied in the present study, and are:

183

184 Isotopic ratios of FAs

$$185 \quad \delta^{13}\text{C} (\text{‰}) = \left[ \left( \frac{R_{\text{sample}}}{R_{\text{standard}}} \right) - 1 \right] * 1000, \quad (1)$$

186 where  $R_{\text{sample}}$  and  $R_{\text{standard}}$  are the ratio of  $^{13}\text{C}/^{12}\text{C}$  in the sample and reference standard, respectively;

187

188 Atom percent (atom%), converted  $\delta$ -values which express isotope data in terms of concentrations to  
189 inform on the  $^{13}\text{C}$  enrichment in each FA

$$190 \quad \text{atom}\% = \left( \frac{R_{\text{sample}}}{(R_{\text{sample}} + 1)} \right) * 100, \quad (2)$$

191 where R is the ratio  $^{13}\text{C}/^{12}\text{C}$  in the sample;

192

193 Atom percent excess (APE), the isotopic enrichment in experimental copepods (in comparison to  
194 field individuals) due to the assimilation of  $^{13}\text{C}$ -enriched prey cultures

$$195 \quad \text{APE} = \text{atom}\%_{\text{experimental copepods}} - \text{atom}\%_{\text{field copepods}} \quad (3)$$

196

197 Carbon mass of each FA ( $C_{\text{mass}}$ , in  $\mu\text{g C ind}^{-1}$ )

$$198 \quad C_{\text{mass}} = \frac{\{(\text{atom}\% / 100 * A_{13\text{C}}) + [(100 - \text{atom}\%) / 100 * A_{12\text{C}}]\} * C_{\text{FA}} * \text{FA}_{\text{mass}}}{\{[(\text{atom}\% / 100 * A_{13\text{C}}) + ((100 - \text{atom}\%) / 100 * A_{12\text{C}})] * C_{\text{FA}}\} + A_{12\text{C}} + (\text{H}_{\text{FAME}} * A_{\text{H}}) + (\text{O}_{\text{FAME}} * A_{\text{O}})}, \quad (4)$$

199 where  $A_{12\text{C}}$ ,  $A_{13\text{C}}$ ,  $A_{\text{H}}$ , and  $A_{\text{O}}$  are the atomic masses of  $^{12}\text{C}$ ,  $^{13}\text{C}$ , H and O, respectively,  $C_{\text{FA}}$  is the  
200 number of carbon atoms in the FA,  $\text{H}_{\text{FAME}}$  and  $\text{O}_{\text{FAME}}$  are the number of hydrogen and oxygen  
201 atoms in the FAME, and  $\text{FA}_{\text{mass}}$  is the mass (in  $\mu\text{g ind}^{-1}$ ) of the FA;

202

203 Proportion of carbon assimilated (PA)

204 
$$PA = \frac{APE}{L}, \quad (5)$$

205 where L is the atom% averaged for all prey FAs except for 18:0. The averaging is to account for the  
206 elongation and/or desaturation of small amounts of dietary FAs when they are assimilated by the  
207 copepods (Dalsgaard, John, Kattner, Müller-Navarra, and Hagen, 2003), and the exclusion of 18:0  
208 is due to it not labelling as well as all other FAs in prey cultures (present study; Boissonnot,  
209 Niehoff, Hagen, Søreide, and Graeve, 2016 and references therein). When calculating the PA for  
210 FA 18:0 in the copepod, L was the average of labeling in prey FA 18:0. For copepods sampled at  
211  $t_{48h}$ , L was calculated from the first two batches of prey cultures used for feeding, whereas for  
212 copepods sampled at  $t_{120h}$ , L was calculated from all prey culture batches;

213

214 FA-specific carbon assimilation ( $C_{\text{assim}}$ , in  $\mu\text{g } ^{13}\text{C ind}^{-1}$ )

215 
$$C_{\text{assim}} = C_{\text{mass}} * PA \quad (6)$$

216 Total C assimilation into lipids was obtained by summing the  $C_{\text{assim}}$  values for all FAs, and C  
217 assimilation as % of food ingested was obtained with the grazing rates reported in Franco-Santos et  
218 al. (2018). Assimilation rates were obtained by dividing  $C_{\text{assim}}$  by the number of experimental days  
219 passed when sample was collected;

220

221 Carbon turnover rates ( $C_T$  in  $\% \text{ day}^{-1}$ )

222 
$$C_T = \frac{\frac{C_{\text{assim}}(t)}{B(t)}}{\Delta t}, \quad (7)$$

223 where  $(t)$  indicates the sampling time for which the values of  $C_{\text{assim}}$  and B should be used and  $\Delta t$  the  
224 number of experimental days when the sample was collected.

225

## 226 **Statistical analysis**

227 The statistical significance of differences in C, N, and total lipid contents of prey culture,  
228 their C:N molar ratio and atom% was investigated with one-way analyses of variance (ANOVA).

229 Differences in atom% were tested (a) between the different diets; and (b) within each diet between  
230 the different sampling days. When the results were significant, the origin of differences was  
231 identified by applying the Tukey HSD (Honestly Significant Difference) post-hoc test with a 95%  
232 confidence limit. Prior to the ANOVAs, the data were tested for normality and homogeneity of  
233 variances with Shapiro-Wilk and Bartlett tests, respectively. Non-normal and/or heteroscedastic  
234 data were either analyzed with ANOVA and Tukey HSD after normalization/homogenization or  
235 with the non-parametric test Kruskal–Wallis, followed by the post-hoc Nemenyi test (with P-value  
236 being determined with the Tukey method).

237 Differences in copepod C, N, and total lipid contents and C:N molar ratio were investigated  
238 with two-way repeated measures ANOVA. When the results regarding the diet treatment factor  
239 were significant, the origin of differences was identified by applying one-way repeated measures  
240 ANOVA for each treatment, followed by pairwise Tukey HSD comparisons with a Bonferroni  
241 correction (*p*-value significance threshold was set at 0.0167). When the results regarding the  
242 sampling day factor were significant, the origin of differences was identified by applying one-way  
243 ANOVA for each day, followed by a Tukey HSD test. Because the *in situ* samples were shared  
244 between the treatments of Experiment I and Experiment II, these were investigated for differences  
245 with a t-test rather than with the Tukey HSD. Repeated measures ANOVA data were tested for  
246 normality with graphical methods and residuals analysis and for sphericity with Mauchly's test.

247 Differences between copepods with regard to <sup>13</sup>C labelling (average APE) and rates of C  
248 assimilation and turnover within each sampling period ( $t_{48h-t_{0h}}$  and  $t_{120h-t_{0h}}$ ) were tested with one-  
249 way ANOVAs, and differences between treatments were identified with a Tukey HSD test.  
250 Differences in C assimilation and turnover for SFA, MUFA and PUFA were also investigated  
251 between the different diets within each sampling period with one-way ANOVAs. Differences  
252 within each diet between sampling days in C assimilation and turnover rates for SFA, MUFA and  
253 PUFA were analyzed with paired t-tests, with data previously checked for normality and  
254 homoscedasticity with Shapiro-Wilk and F tests, respectively. Problems in the labelling of the

255 Dino+ diet affected the isotopic enrichment (and, consequently, the calculation of C assimilation  
256 and turnover rates) in copepods fed with it, and only the data from copepods fed with the Diat+,  
257 Diat- and Dino- diets was used for the statistical tests. All univariate analyses were performed  
258 using R ver. 3.4.4 (Ihaka and Gentleman, 1993).

259         Multivariate analysis of FA composition (untransformed relative FA concentration data for  
260 FAs accounting for > 1% total fatty acids, TFA) for both prey cultures and copepods was performed  
261 with a non-metric multidimensional scaling (nMDS) method generated from a Bray-Curtis  
262 similarity matrix. A one-way analysis of similarity (ANOSIM) was conducted in order to identify  
263 differences in the FA profiles of the diets offered to the copepods, and followed by a similarity  
264 percentages (SIMPER) analysis to identify the specific FAs contributing to the difference.  
265 Differences in the FA profile of *T. longicornis* across factors were investigated with a permutational  
266 multivariate analysis of variance (PERMANOVA) with Monte Carlo simulations to account for  
267 small sample sizes. Significant results were followed by a test of homogeneity of dispersions  
268 (PERMDISP), the multivariate data equivalent of a sphericity test for repeated measures design.  
269 Although not a requirement for PERMANOVA, PERMDIST enables, in cases of a significant  
270 effects result, the differentiation between differences caused by the treatment and differences  
271 caused by the variation within replicates among time points due to their non-independence in  
272 repeated measures designs (Anderson et al., 2008). All multivariate analyses were performed with  
273 PRIMER 7.0 software (Clarke and Gorley, 2015) with the PERMANOVA+ add-on (Anderson et  
274 al., 2008).

275

276

## 277 **RESULTS**

### 278 **Prey culture composition and labelling**

279         Although the Dino+ culture had significantly higher C content, all diets used for feeding  
280 copepods contained >500  $\mu\text{g C L}^{-1}$ , confirming the *ad libitum* feeding condition (Table 1). The

281 approximate N content of diet cultures was 200  $\mu\text{g N L}^{-1}$  for Dino +, 100  $\mu\text{g N L}^{-1}$  for Diat+ and  
 282 Dino-, and 50  $\mu\text{g N L}^{-1}$  for Diat-. The molar C:N ratio was also similar between all treatments  
 283 (approx. 6) with the exception of the Diat-, which was almost three times greater. Total lipid  
 284 content in the diets was approximately 100  $\mu\text{g L}^{-1}$  except for Dino+, which had double that amount.

285

286 **Table 1** Carbon (C), nitrogen (N), and total lipid contents, molar C:N ratios (C:N), and average  $^{13}\text{C}$   
 287 enrichment (of all FAs but 18:0, atom%) of the prey cultures used in the feeding experiment  
 288 (nutrient-replete and N-depleted *Conticribra weissflogii*, Diat+ and Diat-, and *Oxyrrhis marina*,  
 289 Dino+ and Dino-, respectively). Values are mean  $\pm$  standard deviation from four to five batches  
 290 (temporal replicates).

Experiment	Treatment	pg C cell <sup>-1</sup>	pg N cell <sup>-1</sup>	C:N	pg lipid cell <sup>-1</sup>	atom%
I	Diat+	71 $\pm$ 5	13 $\pm$ 2	6.2 $\pm$ 0.6	13 $\pm$ 1	2.73 $\pm$ 0.11
	Diat-	79 $\pm$ 12	6 $\pm$ 1	17.0 $\pm$ 3.4	13 $\pm$ 2	2.81 $\pm$ 0.09
	Dino+	517 $\pm$ 110	101 $\pm$ 18	6.0 $\pm$ 0.3	107 $\pm$ 21	1.18 $\pm$ 0.05
II	Dino-	273 $\pm$ 18	51 $\pm$ 4	6.3 $\pm$ 0.8	54 $\pm$ 9	2.03 $\pm$ 0.14

291

292 The FA composition of all prey cultures (Table 2) was mostly characterized by  
 293 polyunsaturated fatty acids (PUFAs), which accounted for 44 to 67% of the total fatty acid (TFA)  
 294 concentration. Saturated (SFA) and monounsaturated (MUFA) fatty acids were equally present in  
 295 the Diat+ (20% TFA) and Diat- (30% TFA) diets, but in the Dino+ and Dino- diets SFAs were  
 296 present in higher relative concentrations (26 and 41% TFA, respectively) than MUFAs (7 and 15%  
 297 TFA, respectively). The most important FAs in the Diat+ and Diat- cultures were 20:5(n-3)  
 298 (eicosapentaenoic acid, EPA, with 21 and 18% TFA, respectively), 16:1(n-7) (16 and 26% TFA,  
 299 respectively), 16:3(n-4) (20 and 12% TFA, respectively), and 16:0 (14 and 19% TFA, respectively).  
 300 16:0 and 22:6(n-3) (docosahexaenoic acid, DHA) were the dominant FAs in dinoflagellate cultures  
 301 (19 and 29% TFA in Dino+ and 31 and 30% TFA in Dino-, respectively), with a high importance  
 302 also of 18:3(n-3) in Dino+ (15% TFA). The FAs 16:2(n-4) and 16:3(n-4) were only detected in the



303 diatom cultures. The ANOSIM (global R: 0.974,  $p = 0.001$ ) indicated differences in FA  
304 composition between the prey cultures, which the nMDS (Fig. 1, stress 0.06) and SIMPER  
305 (Supplementary Table S1) showed to be a clear separation between the diatom and dinoflagellate  
306 cultures. The Diat+ culture was mostly different from the Dino cultures in terms of the FAs DHA  
307 (16-18%) and 16:3(n-4) (15-16%), whereas the Diat- culture was mostly different from the Dino  
308 cultures due to the FAs 16:1(n-7) (20-22%) and DHA (19-21%). The nMDS and SIMPER results  
309 also indicated that the nutrient-replete and N-depleted treatments were similar within the diatom  
310 and dinoflagellate cultures, mostly because of the FAs 16:1(n-7) (23%) and 16:3(n-4) (23%) for the  
311 former and to the FAs 16:0 (23%), 18:4(n-3) (20%), and 18:3(n-3) (14%) for the latter. The FAs  
312 showed significantly higher  $^{13}\text{C}$  enrichment in the diatom cultures ( $\sim 3$  atom%) and lower in the  
313 Dino+ diet (1.18 atom%). Labelling was significantly higher in the first batch cultures (i.e., days),  
314 with the exception for the Diat+ batch from day 5 (Fig. 2 and Supplementary Tables ST1 and ST2).  
315 FAs within the same diet had similar  $^{13}\text{C}$  enrichment, with the exception of 18:0, which varied from  
316 55 to 93% of the average atom% of all FAs in a given diet. Nevertheless, EPA, DHA, 18:3(n-3) and  
317 18:4(n-3) were the FAs with better labelling in all diets but Dino+ (Table2). Cultures were  
318 significantly different from one another in all of these variables (Supplementary Tables S1 and S2).


## Life cycle strategies of *Temora longicornis*

319 **Table 2** Prey culture absolute (Mass, in  $\mu\text{g L}^{-1}$ ) and relative (% total fatty acid, TFA) fatty acid (FA) compositions ( $> 1\%$  TFA) and  $^{13}\text{C}$  enrichment  
 320 (atom%). *Conticribra weissflogii* and *Oxyrrhis marina* were cultured in nutrient replete (Diat+ and Dino+, respectively) and nitrogen depleted (Diat-  
 321 and Dino-, respectively) conditions. Values are means  $\pm$  standard deviation from prey cultures and from the sum of saturated (SFA), monounsaturated  
 322 (MUFA) and polyunsaturated (PUFA) fatty acids (n = 4 or 5). - : not detected.

FA	Diat+			Diat-			Dino+			Dino-		
	Mass	% TFA	atom%	Mass	% TFA	atom%	Mass	% TFA	atom%	Mass	% TFA	atom%
14:0	6.6 $\pm$ 0.6	6.7 $\pm$ 1.0	2.8 $\pm$ 0.1	6.3 $\pm$ 1.5	5.9 $\pm$ 0.6	2.9 $\pm$ 0.1	9.3 $\pm$ 2.1	4.4 $\pm$ 0.4	1.2 $\pm$ 0.1	5.0 $\pm$ 2.6	4.2 $\pm$ 1.1	2.0 $\pm$ 0.2
i 15:0	0.7 $\pm$ 0.1	0.8 $\pm$ 0.2	2.7 $\pm$ 0.1	0.3 $\pm$ 0.1	0.3 $\pm$ 0.2	2.7 $\pm$ 0.1	1.5 $\pm$ 0.4	0.7 $\pm$ 0.2	1.2 $\pm$ 0.1	1.4 $\pm$ 0.2	1.2 $\pm$ 0.2	2.0 $\pm$ 0.3
15:0	1.0 $\pm$ 0.1	1.0 $\pm$ 0.1	2.8 $\pm$ 0.1	1.8 $\pm$ 0.5	1.7 $\pm$ 0.2	2.9 $\pm$ 0.1	1.5 $\pm$ 0.2	0.8 $\pm$ 0.2	1.2 $\pm$ 0.1	0.5 $\pm$ 0.1	0.5 $\pm$ 0.0	1.9 $\pm$ 0.1
16:0	10.8 $\pm$ 7.2	10.4 $\pm$ 6.9	2.8 $\pm$ 0.1	19.4 $\pm$ 2.7	18.5 $\pm$ 0.5	2.9 $\pm$ 0.1	39.8 $\pm$ 6.4	19.0 $\pm$ 2.7	1.2 $\pm$ 0.1	34.3 $\pm$ 7.9	30.2 $\pm$ 0.8	2.0 $\pm$ 0.1
16:1n7	15.9 $\pm$ 0.5	16.2 $\pm$ 1.5	2.8 $\pm$ 0.1	27.3 $\pm$ 6.8	25.7 $\pm$ 2.3	2.9 $\pm$ 0.1	1.6 $\pm$ 0.4	0.8 $\pm$ 0.1	1.2 $\pm$ 0.1	1.9 $\pm$ 1.8	1.5 $\pm$ 1.0	1.7 $\pm$ 0.1
16:2n4	9.1 $\pm$ 1.6	9.1 $\pm$ 0.9	2.8 $\pm$ 0.1	4.2 $\pm$ 0.3	4.0 $\pm$ 0.3	2.9 $\pm$ 0.1	-	-	-	-	-	-
16:3n4	19.7 $\pm$ 2.4	20.2 $\pm$ 5.0	2.8 $\pm$ 0.1	11.2 $\pm$ 0.7	10.8 $\pm$ 1.9	2.9 $\pm$ 0.1	-	-	-	-	-	-
18:0	0.6 $\pm$ 0.1	0.6 $\pm$ 0.1	1.5 $\pm$ 0.2	1.0 $\pm$ 0.3	1.0 $\pm$ 0.2	1.9 $\pm$ 0.2	1.7 $\pm$ 0.5	0.8 $\pm$ 0.2	1.1 $\pm$ 0.0	3.0 $\pm$ 2.3	2.4 $\pm$ 1.2	1.2 $\pm$ 0.0
18:1n7	0.7 $\pm$ 0.1	0.7 $\pm$ 0.2	2.5 $\pm$ 0.2	0.6 $\pm$ 0.1	0.6 $\pm$ 0.2	2.3 $\pm$ 0.1	11.3 $\pm$ 2.0	5.3 $\pm$ 0.3	1.2 $\pm$ 0.1	5.8 $\pm$ 1.0	5.1 $\pm$ 0.4	1.9 $\pm$ 0.1
18:1n9	0.5 $\pm$ 0.1	0.5 $\pm$ 0.1	2.6 $\pm$ 0.1	1.2 $\pm$ 0.5	1.1 $\pm$ 0.3	2.9 $\pm$ 0.1	2.2 $\pm$ 1.0	1.0 $\pm$ 0.3	1.2 $\pm$ 0.1	8.5 $\pm$ 4.2	7.1 $\pm$ 1.8	2.1 $\pm$ 0.2
18:2n6	1.0 $\pm$ 0.1	1.0 $\pm$ 0.1	2.7 $\pm$ 0.1	0.9 $\pm$ 0.1	0.9 $\pm$ 0.2	2.9 $\pm$ 0.1	8.3 $\pm$ 2.7	3.8 $\pm$ 0.6	1.2 $\pm$ 0.0	6.8 $\pm$ 3.3	5.8 $\pm$ 1.4	2.1 $\pm$ 0.2
18:3n3	0.2 $\pm$ 0.0	0.2 $\pm$ 0.0	2.8 $\pm$ 0.1	0.3 $\pm$ 0.0	0.2 $\pm$ 0.1	2.9 $\pm$ 0.1	32.9 $\pm$ 11.6	15.1 $\pm$ 3.0	1.1 $\pm$ 0.0	9.7 $\pm$ 4.8	8.2 $\pm$ 2.2	2.2 $\pm$ 0.1
18:4n3	3.7 $\pm$ 1.3	3.6 $\pm$ 1.0	2.8 $\pm$ 0.1	4.9 $\pm$ 1.5	4.6 $\pm$ 0.7	2.9 $\pm$ 0.1	28.2 $\pm$ 9.9	12.9 $\pm$ 2.5	1.2 $\pm$ 0.1	3.4 $\pm$ 2.2	2.8 $\pm$ 1.1	2.2 $\pm$ 0.1

20:4n3	0.2 ± 0.0	0.3 ± 0.1	2.5 ± 0.1	0.5 ± 0.1	0.5 ± 0.1	2.8 ± 0.1	1.3 ± 0.2	0.6 ± 0.1	1.2 ± 0.0	1.0 ± 0.9	1.1 ± 1.0	2.1 ± 0.0
20:5n3	21.1 ± 2.3	21.2 ± 0.6	2.8 ± 0.1	19.2 ± 2.5	18.3 ± 0.7	2.9 ± 0.1	10.7 ± 2.4	5.0 ± 0.3	1.2 ± 0.1	3.5 ± 0.4	3.1 ± 0.4	2.1 ± 0.1
22:6n3	5.6 ± 0.5	5.6 ± 0.5	2.8 ± 0.1	4.7 ± 0.2	4.5 ± 0.5	2.9 ± 0.1	59.7 ± 5.9	28.6 ± 3.4	1.2 ± 0.1	28.7 ± 2.6	26.6 ± 7.8	2.1 ± 0.1
ΣSFA	19.7 ± 7.2	19.4 ± 5.9	-	28.9 ± 4.8	27.5 ± 0.4	-	55.1 ± 9.1	26.2 ± 3.5	-	44.7 ± 13.2	40.6 ± 5.1	-
ΣMUFA	18.3 ± 0.6	18.6 ± 1.8	-	30.0 ± 7.1	28.3 ± 2.3	-	15.1 ± 3.2	7.1 ± 0.2	-	16.2 ± 6.9	14.5 ± 3.6	-
ΣPUFA	61.2 ± 3.1	62.0 ± 4.1	-	46.3 ± 4.3	44.3 ± 2.6	-	142.7 ± 32.2	66.7 ± 3.5	-	47.6 ± 4.4	44.9 ± 8.6	-


323



324

325 **Figure 1:** Non-metric multidimensional scaling (nMDS) generated with a Bray-Curtis similarity on untransformed, relative (>1% TFA) concentrations of FAs for field (Experiments  
326 I and II) and experimental (Temora (Diat+), Temora (Diat-), Temora (Dino+) and Temora (Dino-)) copepods and for the prey cultures (Diat+, Diat-, Dino+ and Dino-). Numbers  
327 represent the day the copepods and prey cultures were sampled during the experiments. Temora: *Temora longicornis*; Diat+: nutrient-replete *Conticribra weissflogii*; Diat-:  
328 Nitrogen-depleted *C. weissflogii*; Dino+: nutrient-replete *Oxyrrhis marina*; Dino-: Nitrogen-depleted *O. marina*.

## Life cycle strategies of *Temora longicornis*



329

330 **Figure 2:** <sup>13</sup>C enrichment (atom%) for each prey batch culture grown for the feeding experiments. Batches were created  
331 every day consecutively for five days, and let grow for five days before being used to feed copepods. Prey used were  
332 the diatom *Conticribra weissflogii* cultured in nutrient-replete and nitrogen-depleted conditions (Diat+ and Diat-,  
333 respectively) and the dinoflagellate *Oxyrrhis marina*, fed daily with prey (*Rhodomonas salina*) cultured in nutrient-  
334 replete and nitrogen-depleted conditions (Dino+ and Dino-, respectively).

335

### 336 Copepods

#### 337 Composition

338 The average C, N, and total lipid contents and molar C:N ratio of copepods at each sampling  
339 time are shown in Table 3. All values increased significantly in the first two days of the experiment  
340 (Supplementary Tables S1 and S2). Copepod C content varied between 15 and 20  $\mu\text{g C copepod}^{-1}$   
341 and was significantly higher in the copepods fed with Diat+ than in those fed with N-depleted prey  
342 at  $t_{48\text{h}}$  (Supplementary Tables S1 and S2). N content, which varied between 4 and 5  $\mu\text{g N copepod}^{-1}$ ,  
343 was significantly lower in the Diat- treatment at  $t_{48\text{h}}$  (Supplementary Tables S1 and S2). Lipid  
344 content varied between 2 and 4  $\mu\text{g copepod}^{-1}$  and was significantly higher in the copepods feeding  
345 on Diat- (Supplementary Tables S1 and S2). The lipid C content increased throughout the  
346 experiment, from 8-9% total body C in field copepods to approximately 10 and 12% total body C at

347  $t_{48h}$  and  $t_{120h}$ , respectively, in the Diat+ and dinoflagellate diets. Copepods fed with Diat- had the  
 348 highest relative lipid C content, 13 and 15% total body C at  $t_{48h}$  and  $t_{120h}$ , respectively. Molar C:N  
 349 ratios, which ranged from 4.4 to 5.3, were significantly higher in the Diat- copepods at  $t_{48h}$ , and  
 350 significantly lower in Diat+ and Dino- in comparison to Diat- and Dino+ at  $t_{120h}$  (Supplementary  
 351 Tables S1 and S2).

352

353 **Table 3** Carbon (C), nitrogen (N), and total lipid C contents (in  $\mu\text{g}$  C, N, and lipids copepod<sup>-1</sup>,  
 354 respectively), molar C:N ratios, and <sup>13</sup>C enrichment (APE) of *Temora longicornis* females. Samples  
 355 were obtained at  $t_{0h}$  (*in situ* condition),  $t_{48h}$  (after 48h of experiment), and  $t_{120h}$  (at the end of the  
 356 experiment). Diet treatments: single cultures of *Conticribra weissflogii* (Diat) and *Oxyrrhis marina*  
 357 (Dino) in nutrient-replete (+) and N-depleted (-) conditions. Values presented are mean  $\pm$  standard  
 358 deviation of triplicates.

Diets	C			N			molar C:N ratio			Lipid			APE%	
	$t_{0h}$	$t_{48h}$	$t_{120h}$	$t_{0h}$	$t_{48h}$	$t_{120h}$	$t_{0h}$	$t_{48h}$	$t_{120h}$	$t_{0h}$	$t_{48h}$	$t_{120h}$	$t_{48h}$	$t_{120h}$
Diat+	16 $\pm$ 1	20 $\pm$ 0	20 $\pm$ 1	4 $\pm$ 0	5 $\pm$ 0	5 $\pm$ 0	4.7 $\pm$ 0.1	4.9 $\pm$ 0.1	4.7 $\pm$ 0.1	2 $\pm$ 0	3 $\pm$ 0	3 $\pm$ 0	0.74 $\pm$ 0.03	0.94 $\pm$ 0.03
Diat-	16 $\pm$ 1	18 $\pm$ 1	18 $\pm$ 2	4 $\pm$ 0	4 $\pm$ 0	4 $\pm$ 0	4.7 $\pm$ 0.1	5.3 $\pm$ 0.2	5.0 $\pm$ 0.1	2 $\pm$ 0	3 $\pm$ 0	4 $\pm$ 0	0.73 $\pm$ 0.01	1.02 $\pm$ 0.02
Dino+	15 $\pm$ 0	20 $\pm$ 0	20 $\pm$ 2	4 $\pm$ 0	5 $\pm$ 0	5 $\pm$ 0	4.4 $\pm$ 0.1	4.8 $\pm$ 0.0	5.0 $\pm$ 0.1	2 $\pm$ 0	3 $\pm$ 0	3 $\pm$ 0	0.10 $\pm$ 0.01	0.08 $\pm$ 0.00
Dino-	15 $\pm$ 0	18 $\pm$ 1	18 $\pm$ 1	4 $\pm$ 0	5 $\pm$ 0	4 $\pm$ 0	4.4 $\pm$ 0.1	4.6 $\pm$ 0.1	4.8 $\pm$ 0.0	2 $\pm$ 0	2 $\pm$ 0	3 $\pm$ 0	0.52 $\pm$ 0.03	0.70 $\pm$ 0.03

359

360 The relative FA composition of *T. longicornis* changed when copepods were introduced to  
 361 experimental diets and reflected that of their prey, though *in situ* composition seemed to indicate  
 362 feeding on diatoms in the field (Table 4, Figs 1 and 3). In copepods from the field, SFA, MUFA and  
 363 PUFA accounted for approximately 25, 10, and 65% TFA. Feeding on the Diat+ and dinoflagellate  
 364 diets provided a similar partition of SFA, MUFA and PUFA, but feeding on Diat- increased the  
 365 relative MUFA content (21% TFA) and decreased the relative PUFA content (55% TFA) in  
 366 copepods. EPA, DHA and 16:0 were the major FA in *T. longicornis* throughout the experiment.  
 367 EPA was the major FA constituent in diatom-fed copepods, whereas DHA was the major FA in

368 dinoflagellate-fed copepods. Together, EPA and DHA accounted for 54-72% TFA in copepods fed  
369 with diatoms, and approximately 70-75% TFA in copepods fed with dinoflagellates. The FA  
370 16:1(n-7) accounted for another 8-16% TFA in diatom-fed copepods, and the FAs 18:3(n-3) and  
371 18:4(n-3) together made up 11-16% TFA in the copepods fed with the Dino+ diet. The FAs 16:2(n-  
372 4) and 16:3(n-4) were only detected in copepods fed with diatoms. The difference in relative  
373 abundance for the previously mentioned major FAs between the beginning and the end of the  
374 experiments ( $t_{120h}-t_{0h}$ ) can be visualized in Fig. 3, and the difference in absolute composition (for  
375 FA > 1% TFA) in Fig. 4. The relative abundance of the FAs 16:0 and 18:4(n-3) increased in all  
376 treatments, and decreased for EPA. The diatom FATMs (16:1(n-7), 16:2(n-4), and 16:3(n-4))  
377 increased in diatom-fed copepods and decreased in dinoflagellate-fed females, whereas the  
378 dinoflagellate FATMs (18:3(n-3) and DHA) decreased in the former and increased in the latter (Fig.  
379 3). The nMDS (Fig. 1) suggests a clear separation between diet treatments and between sampling  
380 days within some diets, showing a continuous change in FA composition through the experimental  
381 time. According to the PERMANOVA results (main effects and pairwise comparisons) all  
382 treatments are different from one another in all sampling days, and all sampling days within a  
383 treatment are different from one another, except for Diat+ at  $t_{48h}$  and  $t_{120h}$  (Supplementary Table  
384 S3). It was not possible to confirm that these differences are solely due to treatment effects, as the  
385 PERMDIST result was significant ( $F = 704.82$ ,  $df1 = 2$ ,  $df2 = 33$ ,  $P(\text{perm}) = 0.001$ ).

### Life cycle strategies of *Temora longicornis*


386 **Table 4** Absolute (in ng C ind<sup>-1</sup>) and relative (% total fatty acid, TFA, in parenthesis) FA composition (>1% total FAs) of *Temora longicornis* females  
 387 at the beginning (t<sub>0h</sub>, *in situ* condition), middle (t<sub>48h</sub>, after 48h of experiment), and end (t<sub>120h</sub>) of the feeding experiment. Diet treatments: single cultures  
 388 of *Conticribra weissflogii* (Diat) and *Oxyrrhis marina* (Dino) in nutrient-replete (+) and N-depleted (-) conditions. Values presented are mean ±  
 389 standard deviation of triplicates. Values are mean ± standard deviation (n = 3 each) from each FA and from the sum of all (TFA), saturated (SFA),  
 390 monounsaturated (MUFA), and polyunsaturated (PUFA) fatty acids (n = 4 or 5). – : not detected.

FA	Diat+		Diat-		Dino+		Dino-		
	t <sub>0h</sub>	t <sub>120h</sub>	t <sub>0h</sub>	t <sub>120h</sub>	t <sub>0h</sub>	t <sub>120h</sub>	t <sub>0h</sub>	t <sub>120h</sub>	
TFA	1509 ± 92	2129 ± 116	2105 ± 129	2707 ± 365	1249 ± 170	2025 ± 111	2509 ± 91	1794 ± 226	2117 ± 146
14:0	65 ± 17 (5 ± 1)	89 ± 11 (5 ± 0)	70 ± 9 (4 ± 0)	104 ± 19 (4 ± 0)	46 ± 12 (4 ± 0)	52 ± 1 (3 ± 0)	67 ± 2 (3 ± 0)	43 ± 4 (3 ± 0)	48 ± 10 (2 ± 0)
16:0	207 ± 13 (14 ± 0)	322 ± 16 (16 ± 0)	320 ± 14 (16 ± 1)	424 ± 57 (16 ± 0)	186 ± 27 (16 ± 0)	369 ± 12 (19 ± 1)	452 ± 14 (19 ± 0)	369 ± 44 (22 ± 0)	473 ± 37 (23 ± 0)
16:1n7	78 ± 9 (5 ± 0)	162 ± 6 (8 ± 0)	155 ± 15 (8 ± 0)	326 ± 36 (14 ± 0)	29 ± 6 (2 ± 0)	18 ± 2 (1 ± 0)	13 ± 2 (1 ± 0)	20 ± 4 (1 ± 0)	18 ± 3 (1 ± 0)
16:2n4	11 ± 1 (0 ± 0)	83 ± 2 (4 ± 0)	92 ± 12 (5 ± 0)	80 ± 13 (3 ± 0)	6 ± 1 (1 ± 0)	–	–	–	–
16:3n4	6 ± 1 (0 ± 0)	88 ± 67 (4 ± 3)	156 ± 24 (8 ± 1)	158 ± 30 (6 ± 0)	5 ± 1 (0 ± 0)	–	–	–	–
18:0	47 ± 4 (3 ± 0)	45 ± 2 (2 ± 0)	45 ± 8 (2 ± 1)	55 ± 7 (2 ± 0)	53 ± 14 (4 ± 1)	48 ± 4 (3 ± 0)	39 ± 1 (2 ± 0)	49 ± 7 (3 ± 0)	33 ± 2 (2 ± 0)
18:1n7	52 ± 2 (4 ± 0)	67 ± 3 (3 ± 0)	69 ± 3 (3 ± 0)	87 ± 11 (3 ± 0)	37 ± 4 (3 ± 0)	77 ± 5 (4 ± 0)	108 ± 5 (4 ± 0)	61 ± 7 (4 ± 0)	78 ± 6 (4 ± 0)

18:1n9	23 ± 3 (2 ± 0)	17 ± 2 (1 ± 0)	19 ± 8 (1 ± 0)	23 ± 3 (2 ± 0)	32 ± 9 (1 ± 0)	29 ± 4 (1 ± 0)	13 ± 4 (1 ± 0)	11 ± 3 (1 ± 0)	16 ± 1 (1 ± 0)	13 ± 4 (1 ± 0)	47 ± 11 (3 ± 1)	84 ± 7 (4 ± 0)
18:2n6	12 ± 1 (1 ± 0)	14 ± 1 (1 ± 0)	13 ± 1 (1 ± 0)	12 ± 1 (1 ± 0)	17 ± 3 (1 ± 0)	17 ± 2 (1 ± 0)	7 ± 2 (1 ± 0)	31 ± 3 (2 ± 0)	59 ± 4 (2 ± 0)	7 ± 2 (1 ± 0)	40 ± 7 (2 ± 0)	83 ± 6 (4 ± 0)
18:3n3	8 ± 1 (1 ± 0)	7 ± 1 (0 ± 0)	6 ± 2 (0 ± 0)	8 ± 1 (1 ± 0)	10 ± 2 (0 ± 0)	9 ± 2 (0 ± 0)	7 ± 2 (1 ± 0)	122 ± 8 (6 ± 0)	234 ± 15 (9 ± 0)	7 ± 2 (1 ± 0)	60 ± 9 (3 ± 0)	121 ± 10 (6 ± 0)
18:4n3	18 ± 1 (1 ± 0)	64 ± 1 (3 ± 0)	59 ± 7 (3 ± 0)	18 ± 1 (1 ± 0)	72 ± 7 (3 ± 0)	92 ± 15 (3 ± 0)	16 ± 3 (1 ± 0)	97 ± 6 (5 ± 0)	159 ± 11 (6 ± 0)	16 ± 3 (1 ± 0)	25 ± 4 (1 ± 0)	41 ± 4 (2 ± 0)
20:5n3	589 ± 34 (38 ± 0)	717 ± 26 (33 ± 1)	686 ± 42 (32 ± 1)	589 ± 34 (38 ± 0)	761 ± 61 (30 ± 1)	752 ± 89 (27 ± 0)	457 ± 41 (36 ± 2)	337 ± 21 (16 ± 0)	270 ± 9 (11 ± 0)	457 ± 41 (36 ± 2)	353 ± 55 (19 ± 1)	249 ± 25 (12 ± 1)
22:6n3	299 ± 19 (19 ± 1)	338 ± 16 (15 ± 1)	317 ± 13 (15 ± 1)	299 ± 19 (19 ± 1)	336 ± 19 (13 ± 1)	330 ± 33 (12 ± 0)	305 ± 43 (24 ± 0)	758 ± 41 (36 ± 0)	973 ± 31 (38 ± 0)	305 ± 43 (24 ± 0)	639 ± 64 (35 ± 1)	805 ± 37 (37 ± 1)
ΣSFA	334 ± 30 (23 ± 1)	483 ± 26 (24 ± 1)	457 ± 17 (23 ± 1)	334 ± 30 (23 ± 1)	593 ± 59 (25 ± 1)	624 ± 88 (24 ± 0)	301 ± 56 (25 ± 1)	505 ± 9 (26 ± 1)	600 ± 17 (25 ± 0)	301 ± 56 (25 ± 1)	483 ± 57 (28 ± 0)	578 ± 50 (29 ± 1)
ΣMUFA	188 ± 15 (13 ± 0)	287 ± 16 (14 ± 1)	280 ± 26 (14 ± 1)	188 ± 15 (13 ± 0)	484 ± 60 (20 ± 1)	589 ± 86 (22 ± 0)	103 ± 18 (8 ± 0)	125 ± 17 (6 ± 1)	156 ± 7 (6 ± 0)	103 ± 18 (8 ± 0)	153 ± 31 (9 ± 1)	199 ± 16 (10 ± 0)
ΣPUFA	987 ± 52 (64 ± 1)	1360 ± 92 (62 ± 2)	1368 ± 95 (64 ± 1)	987 ± 52 (64 ± 1)	1391 ± 111 (55 ± 1)	1494 ± 191 (54 ± 0)	845 ± 97 (66 ± 0)	1131 ± 414 (168 ± 21)	1754 ± 70 (69 ± 0)	845 ± 97 (66 ± 0)	1158 ± 143 (63 ± 1)	1340 ± 81 (62 ± 1)




## Life cycle strategies of *Temora longicornis*



391

392 **Figure 3:** Difference in *Temora longicornis* fatty acid (FA) relative abundance (% total FA) between the beginning and  
 393 end of the feeding experiments ( $t_{120h} - t_{0h}$ ). Color legend represents the main FAs in the copepods, whereas fill pattern  
 394 legend indicates the feeding treatment applied. Diat+ and Diat-: nutrient-replete and Nitrogen-depleted *Conticribra*  
 395 *weissflogii*; respectively. Dino+ and Dino-: nutrient- replete and Nitrogen-depleted *Oxyrrhis marina*. Values are mean  
 396  $\pm$  standard deviation of triplicate samples.



397

398 **Figure 4:** Change in *Temora longicornis* main fatty acids (FA,  $>1\%$  total FA) absolute abundance ( $\mu\text{g FA ind}^{-1}$ )  
 399 between the beginning ( $t_{0h}$ ), middle ( $t_{48h}$ ), and end ( $t_{120h}$ ) of the feeding experiments in relation to the diet treatments  
 400 they received. Diets were made of monocultures of diatoms (*Conticribra weissflogii*) and dinoflagellates (*Oxyrrhis*  
 401 *marina*) cultured in nutrient-replete (Diat+ and Dino+, respectively) and in nitrogen-depleted (Diat- and Dino-,  
 402 respectively) conditions.

403 *Labelling*

404 The average  $^{13}\text{C}$  enrichment of copepod FAs, which varied between 0.1 and 1.0 APE (Table  
 405 3, FA-specific APE in Supplementary Table S4), increased significantly from day 3 to day 6 in all  
 406 treatments but Dino+, for which it significantly decreased (Supplementary Tables S1 and S2). The  
 407 APE values were lowest for the Dino+ treatment. Within the same sampling day, the APE of  
 408 diatom-fed copepods was significantly higher than that of copepods fed with Dino-, and those fed  
 409 with Diat- had the highest APE at the end of the experiment (Supplementary Tables S1 and S2).  
 410 The FA-specific  $^{13}\text{C}$  enrichment in copepods on days 3 and 6 is shown in Table 5. The diatom  
 411 FATMs 16:2(n-4) and 16:3(n-4) had some of the better labelling results in copepods fed with these  
 412 diets, whereas the dinoflagellate FATMs 18:3(n-3) and 18:4(n-3) showed the highest labelling in  
 413 Dino-. The FA 18:4(n-3) was also well labelled in the Diat- diet. FAs were homogeneously  
 414 labelled in copepods fed with the Dino+ diet, though DHA seemed to label slightly better.

415

416 **Table 5** Total (TFA) and fatty acid (FA)-specific lipid carbon (C) assimilation (as ng C ind<sup>-1</sup> for  
 417 TFA and as % TFA assimilated for specific FAs) between  $t_{0h-t_{48h}}$  and between  $t_{0h-t_{120h}}$  for *Temora*  
 418 *longicornis* females. Value in parenthesis for TFA indicates lipid C assimilation as % C ingested  
 419 reported by Franco-Santos et al. (2018) between  $t_{0h-t_{48h}}$ . Diet treatments: single cultures of  
 420 *Conticribra weissflogii* (Diat) and *Oxyrrhis marina* (Dino) in nutrient-replete (+) and N-depleted (-  
 421 ) conditions. SFA: saturated FA; MUFA: monounsaturated FA; PUFA: polyunsaturated FA;  
 422 PUFA<sub>3</sub>: set of FAs 18:3(n-3), 20:5(n-3), and 22:6(n-3). Values are mean ± standard deviation (n = 3  
 423 each). - : not detected.


C assimilation								
FA	Diat+		Diat-		Dino+		Dino-	
	$t_{0h-t_{48h}}$	$t_{0h-t_{120h}}$	$t_{0h-t_{48h}}$	$t_{0h-t_{120h}}$	$t_{0h-t_{48h}}$	$t_{0h-t_{120h}}$	$t_{0h-t_{48h}}$	$t_{0h-t_{120h}}$
TFA	657 ± 46 (5 ± 0)	901 ± 68	755 ± 69 (4 ± 1)	1200 ± 165	201 ± 10 (1 ± 0)	187 ± 4	498 ± 52 (6 ± 2)	842 ± 47
14:0	4 ± 0	3 ± 0	4 ± 0	4 ± 0	2 ± 0	2 ± 0	2 ± 0	2 ± 0

i-15:0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	1 ± 0	1 ± 0	1 ± 0	1 ± 0
15:0	1 ± 0	1 ± 0	2 ± 0	1 ± 0	1 ± 0	1 ± 0	0 ± 0	0 ± 0
16:0	16 ± 1	16 ± 0	18 ± 0	17 ± 0	17 ± 0	16 ± 0	24 ± 0	23 ± 0
16:1n5	1 ± 0	1 ± 0	1 ± 0	1 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
16:1n7	11 ± 1	9 ± 0	23 ± 0	21 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
16:2n4	7 ± 0	6 ± 0	3 ± 0	4 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
16:3n4	8 ± 6	11 ± 1	7 ± 0	8 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
16:4n1	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
17:0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	1 ± 0	1 ± 0	0 ± 0	0 ± 0
18:0	1 ± 0	2 ± 0	1 ± 0	2 ± 0	1 ± 0	1 ± 0	2 ± 0	1 ± 0
18:1n7	3 ± 0	3 ± 0	3 ± 0	3 ± 0	3 ± 0	3 ± 0	3 ± 0	3 ± 0
18:1n9	0 ± 0	0 ± 0	1 ± 0	1 ± 0	0 ± 0	0 ± 0	4 ± 0	5 ± 0
18:2n6	1 ± 0	1 ± 0	1 ± 0	1 ± 0	1 ± 0	2 ± 0	4 ± 0	5 ± 0
18:3n3	0 ± 0	0 ± 0	0 ± 0	0 ± 0	5 ± 0	5 ± 0	5 ± 3	7 ± 0
18:3n6	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	2 ± 3	0 ± 0
18:4n3	5 ± 0	4 ± 0	5 ± 0	5 ± 0	5 ± 0	4 ± 0	1 ± 1	2 ± 0
20:1n7	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
20:1n9	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	1 ± 0
20:2n6	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	1 ± 0	1 ± 0	1 ± 0
20:4n3	0 ± 0	0 ± 0	1 ± 0	1 ± 0	1 ± 0	1 ± 0	1 ± 0	0 ± 0
20:4n6	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
20:5n3	31 ± 2	32 ± 1	24 ± 1	24 ± 1	8 ± 0	8 ± 0	6 ± 0	6 ± 0
22:1n7	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
22:1n9	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
22:5n3	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
22:6n3	8 ± 1	10 ± 1	6 ± 0	7 ± 0	53 ± 0	54 ± 0	44 ± 1	41 ± 1
ΣSFA	22 ± 1	20 ± 0	25 ± 1	24 ± 0	22 ± 1	21 ± 0	29 ± 0	28 ± 0
ΣMUFA	15 ± 1	13 ± 0	28 ± 0	26 ± 0	4 ± 0	5 ± 0	8 ± 0	9 ± 0
ΣPUFA	63 ± 2	65 ± 0	47 ± 1	50 ± 1	74 ± 0	75 ± 0	63 ± 0	63 ± 1

426 *Lipid C assimilation*


427 Total assimilation of C into copepod lipids, which varied between 0.2 and 1.2  $\mu\text{g C ind}^{-1}$  at  
428 the end of the experiment, was higher in copepods fed with diatoms than in those fed with  
429 dinoflagellates (Table 5). C assimilation into SFAs was higher for copepods fed with the N-depleted  
430 diets (both in absolute and relative terms), whereas assimilation into MUFAs and PUFAs was  
431 higher for copepods fed with diatom diets (both in absolute and relative terms) (Table 5, Fig. 5).  
432 Absolute C assimilation into PUFAs and, more specifically, into the FAs 18:3(n-3), EPA and DHA,  
433 was higher for copepods fed with the Diat+ prey, but relative assimilation was higher for copepods  
434 fed with dinoflagellates (Table 5, Fig. 5). Copepods assimilated a higher amount of PUFAs than  
435 MUFAs and SFAs in all diet treatments (Fig. 5). The C assimilation rates were significantly higher  
436 for all treatments (but for Diat-) in the first two days than for the entire duration of the experiment  
437 (Supplementary Table S1). At  $t_{48\text{h}}$  the rates were significantly higher for copepods on the diatom  
438 diets than for those feeding on Dino-, but by the end of the experiment significantly higher rates  
439 were recorded for the copepods fed with Diat- (Supplementary Tables ST1 and ST2). C  
440 assimilation in FATM on days 3 and 6 was also different between copepods fed with diatoms and  
441 dinoflagellates, and reflected the FA composition of their respective diets (Table 5, Fig. 6). For  
442 Diat+ and Diat- copepods, most of the total lipid C assimilated was into EPA (~30 and 25%,  
443 respectively), followed by 16:0 (~15%), 16:1(n-7) (~10 and 20%, respectively), and 16:3(n-4) and  
444 DHA (~10% each). For Dino+ and Dino- copepods, most of the total lipid C assimilated was into  
445 DHA (~55 and 42%, respectively), followed by 16:0 (16 and 24%, respectively), EPA (8 and 6%,  
446 respectively), and 18:3(n-3) (~5%). Lipid C assimilation between  $t_{0\text{h}}$  and  $t_{48\text{h}}$  accounted for between  
447 1 and 6% of the food ingested (in terms of C content) at the same time period (Table 5), and was not  
448 significantly different between copepods fed with diatom diets and Dino-.

449



450

451 **Figure 5:** Carbon (C) assimilation (ng C ind<sup>-1</sup>) and turnover rate (%) recorded for *Temora longicornis* at the beginning  
 452 ( $t_{0h}$ ), middle ( $t_{48h}$ ), and end ( $t_{120h}$ ) of the feeding experiments in relation to the diet treatments. Values are shown for total  
 453 lipids (sum of all fatty acids identified) and for saturated (SFA), monounsaturated (MUFA), and polyunsaturated  
 454 (PUFA) fatty acids. Diet treatments consisted of monocultures of diatoms (*Conticribra weissflogii*) and dinoflagellates  
 455 (*Oxyrrhis marina*) cultured in nutrient-replete (Diat+ and Dino+, respectively) and in nitrogen-depleted (Diat- and  
 456 Dino-, respectively) conditions. Values are mean  $\pm$  standard deviation of triplicate samples.



457

458 **Figure 6:** Carbon (C) assimilation ( $\text{ng C ind}^{-1}$ ) and turnover rate (%) recorded for *Temora longicornis* at the beginning  
 459 ( $t_{0h}$ ), middle ( $t_{48h}$ ), and end ( $t_{120h}$ ) of the feeding experiments in relation to the diet treatments. Values are shown for  
 460 diatom (16:1(n-7), 16:3(n-4), and 20:5(n-3)) and dinoflagellate (18:3(n-3), 18:4(n-3), and 22:6(n-3)) fatty acid trophic  
 461 markers (FATM). Diet treatments consisted of monocultures of diatoms (*Conticribra weissflogii*) and dinoflagellates  
 462 (*Oxyrrhis marina*) cultured in nutrient-replete (Diat+ and Dino+, respectively) and in nitrogen-depleted (Diat- and  
 463 Dino-, respectively) conditions. Values are mean  $\pm$  standard deviation of triplicate samples.

464

465 *Lipid C turnover*

466 Lipid C turnover rates, which ranged from 1 to 15%  $\text{ind}^{-1} \text{day}^{-1}$  (Table 6), were significantly  
 467 higher during the first 2 days of experiment than for the total experiment duration (Supplementary  
 468 Tables S1 and S2). No differences were found between the diatom and Dino- treatments within the  
 469 same sampling day though. C turnover of SFA, MUFA and PUFA for these three treatments was  
 470 also similar, though they were usually higher in copepods fed with Diat+ and Diat- (Fig. 5, Table  
 471 6). Overall, approximately 40-45% of all lipid C was replaced with labelled dietary lipid by the end  
 472 of the experiment, except for copepods fed with the Dino+ diet, for which only approximately 10%  
 473 of lipid C was substituted (Fig. 5). Approximately half (40-45%) of the internal SFA, MUFA, and

474 PUFA content was also replaced with dietary FAs by the end of the experiment for the diatom and  
 475 Dino- diets and ~ 10% for Dino+ (Fig. 5). The diatom FATM 16:1(n-7) and 16:3(n-4) showed high  
 476 C turnover (~ 50 to 60% at the end of the experiment) in diatom-fed copepods and low in  
 477 dinoflagellate-fed copepods. Even though the diatom diets were relatively poor in the FAs 18:3(n-3)  
 478 and 18:4(n-3) (< 1% and <5% TFA, respectively), which are dinoflagellate FATM, these FAs had  
 479 high C turnover (~ 30% and 60% at the end of the experiment, respectively, Fig. 6, Table 6) in  
 480 copepods fed with Diat+ and Diat-. High C turnover of these dinoflagellate FATMs were also  
 481 recorded for copepods fed with Dino- (Fig. 6, Table 6). The replacement of internal by dietary EPA  
 482 and DHA also followed the availability of each of these essential FA in the respective diet.  
 483 Copepods fed with Diat+ and Diat- replaced 40-45% and 25-30% of their original EPA and DHA  
 484 by the end of the experiment, respectively, whereas those fed with Dino- substituted 20% and 45%  
 485 of their original EPA and DHA by the end of the experiment, respectively (Fig. 6). C turnover for  
 486 copepods fed with Dino+ were usually well below 10%, with maximum rates recorded for DHA (~  
 487 10% at the end of the experiment, Fig. 6, Table 6).

488

489 **Table 6** Total (TFA) and fatty acid (FA)-specific lipid carbon (C) turnover rate (as % ind<sup>-1</sup> day<sup>-1</sup>)  
 490 between  $t_{0h}$ - $t_{48h}$  and between  $t_{0h}$ - $t_{120h}$  for *Temora longicornis* females. Diet treatments: single  
 491 cultures of *Conticribra weissflogii* (Diat) and *Oxyrrhis marina* (Dino) in nutrient-replete (+) and N-  
 492 depleted (-) conditions. SFA: saturated FA; MUFA: monounsaturated FA; PUFA: polyunsaturated  
 493 FA; PUFA<sub>3</sub>: set of FAs 18:3(n-3), 20:5(n-3), and 22:6(n-3). Values are mean ± standard deviation  
 494 (n = 3 each). - : not detected.

C turnover									
FA	Diat+		Diat-		Dino+		Dino-		
	$t_{0h} - t_{48h}$	$t_{0h} - t_{120h}$	$t_{0h} - t_{48h}$	$t_{0h} - t_{120h}$	$t_{0h} - t_{48h}$	$t_{0h} - t_{120h}$	$t_{0h} - t_{48h}$	$t_{0h} - t_{120h}$	
TFA	15 ± 1	9 ± 0	15 ± 0	9 ± 0	5 ± 0	1 ± 1	14 ± 0	8 ± 0	
14:0	15 ± 1	9 ± 0	14 ± 0	9 ± 0	3 ± 0	1 ± 0	10 ± 1	7 ± 0	

15:0	21 ± 0	10 ± 0	24 ± 1	11 ± 0	7 ± 0	2 ± 0	10 ± 1	7 ± 0
16:0	16 ± 0	9 ± 0	17 ± 0	9 ± 0	5 ± 0	1 ± 0	16 ± 0	8 ± 0
16:1n5	26 ± 1	11 ± 0	22 ± 0	11 ± 0	-	-	-	-
16:1n7	22 ± 0	10 ± 0	27 ± 0	12 ± 0	1 ± 0	1 ± 0	1 ± 0	4 ± 1
16:2n4	29 ± 0	12 ± 0	27 ± 0	12 ± 0	-	-	-	-
16:3n4	31 ± 0	12 ± 0	30 ± 0	12 ± 0	-	-	-	-
16:4n1	8 ± 0	8 ± 0	4 ± 0	4 ± 1	-	-	-	-
17:0	2 ± 1	2 ± 0	8 ± 1	5 ± 0	6 ± 0	2 ± 0	10 ± 0	7 ± 0
18:0	10 ± 0	7 ± 1	9 ± 1	8 ± 0	2 ± 0	1 ± 0	8 ± 0	7 ± 0
18:1n7	13 ± 0	8 ± 1	14 ± 0	9 ± 0	4 ± 0	1 ± 0	14 ± 1	7 ± 0
18:1n9	4 ± 0	1 ± 1	13 ± 1	7 ± 0	3 ± 1	1 ± 0	21 ± 4	10 ± 0
18:2n6	15 ± 1	8 ± 0	12 ± 1	8 ± 0	5 ± 0	1 ± 0	25 ± 0	10 ± 0
18:3n3	8 ± 1	5 ± 2	9 ± 0	7 ± 1	4 ± 0	1 ± 0	20 ± 11	10 ± 0
18:3n6	24 ± 1	11 ± 0	19 ± 0	10 ± 0	4 ± 1	1 ± 0	16 ± 0	8 ± 0
18:4n3	27 ± 0	11 ± 0	26 ± 0	12 ± 0	5 ± 0	1 ± 0	13 ± 11	10 ± 0
20:1n7	6 ± 2	4 ± 2	6 ± 1	4 ± 1	3 ± 1	1 ± 1	0 ± 1	-
20:1n9	2 ± 0	2 ± 0	5 ± 0	4 ± 1	0 ± 1	1 ± 0	15 ± 1	9 ± 0
20:2n6	5 ± 2	5 ± 0	4 ± 1	3 ± 1	4 ± 1	1 ± 0	15 ± 3	9 ± 0
20:4n3	15 ± 2	8 ± 1	18 ± 1	10 ± 0	4 ± 0	1 ± 0	16 ± 2	9 ± 0
20:4n6	7 ± 1	5 ± 1	4 ± 0	4 ± 0	3 ± 1	1 ± 0	5 ± 2	6 ± 0
20:5n3	14 ± 0	9 ± 0	12 ± 0	8 ± 0	2 ± 0	1 ± 0	4 ± 0	4 ± 0
22:1n7	3 ± 3	4 ± 0	4 ± 1	3 ± 0	4 ± 1	1 ± 0	3 ± 1	2 ± 0
22:1n9	0 ± 1	1 ± 2	-2 ± 1	0 ± 1	-	-	-	-
22:5n3	6 ± 0	5 ± 0	8 ± 0	6 ± 0	4 ± 0	1 ± 0	8 ± 0	0 ± 0
22:6n3	8 ± 0	6 ± 0	6 ± 0	5 ± 0	7 ± 0	2 ± 0	17 ± 0	9 ± 0
ΣSFA	15 ± 0	9 ± 0	16 ± 0	9 ± 0	4 ± 0	1 ± 0	15 ± 0	8 ± 0
ΣMUFA	17 ± 1	9 ± 0	22 ± 1	11 ± 0	3 ± 0	1 ± 0	13 ± 1	8 ± 0
ΣPUFA	15 ± 1	9 ± 0	13 ± 0	8 ± 0	5 ± 0	2 ± 0	14 ± 0	8 ± 0



## 495 **DISCUSSION**

496

### 497 **Lipid C assimilation**

498           The results obtained in the present study indicate that feeding upon N-depleted food can  
499 influence the metabolism of *T. longicornis* females, as the copepods which were fed with the Diat-  
500 culture showed the highest lipid C content and assimilation. This is in agreement with results for  
501 other calanoid copepods such as *Acartia tonsa*, which accumulated more lipids when feeding on N-  
502 depleted in comparison to nutrient-replete *R. salina* (Burian, Grosse, Winder, and Boschker, 2018).  
503 However, when total C assimilation is calculated as the % dietary C that was ingested by copepods,  
504 there were no significant differences between the treatments. Although the quality of a food source  
505 affects the rate at which it is ingested by copepods (Franco-Santos et al., 2018), it would seem that  
506 it does not influence its assimilability. A possible explanation is provided by Mitra and Flynn  
507 (2005) who suggest that, where nutrient limitation significantly affects the ingestion of prey by  
508 zooplankton, it may be unimportant at the stage of food assimilation. Furthermore, it has been  
509 shown that, although phosphorus (P) assimilation efficiency differs for *Daphnia pulex* feeding  
510 on prey of different P (and C) contents, it does not affect its C assimilation efficiency (Urabe,  
511 Shimizu, and Yamaguchi, 2017), which could also explain the results obtained herein. Indeed, C  
512 was not the limiting element in the prey offered to copepods, so its assimilation from dietary  
513 material should not be affected by prey quality, as it was freely available for ingestion.

514

### 515 **Copepod life history strategies**

516           The differences in absolute C assimilation were most likely caused by varying copepod  
517 ingestion rates, as observed by Franco-Santos et al. (2018). If copepods ingested different amounts  
518 of dietary C but assimilated similar quantities, then it follows that they also had different C  
519 investment into reproduction. This assumption is confirmed when the C assimilation results of the  
520 present study are compared with the EPR reported by Franco-Santos et al. (2018), which were

521 higher for copepods fed with the Dino+ culture, followed by those fed with Dino-, Diat+, and Diat-  
522 (61, 41, 39, and 36 eggs ind<sup>-1</sup> day<sup>-1</sup>). As previously assumed, reproductive investment was inversely  
523 related to the C assimilation rates of the copepods, and both of these factors were affected by the  
524 elemental and biochemical compositions of the prey items.

525 All experimental diets allowed for lipid accumulation in *T. longicornis*, confirming both an  
526 active uptake of dietary FAs and that the quantity of food ingested was more than enough to cover  
527 for the metabolic costs of living, even though the latter are specific for each copepod-diet  
528 combination (Franco-Santos et al., 2018). The copepods fed with nutrient-replete diatoms did not  
529 shown an increase in lipid C between  $t_{48h}$  and  $t_{120h}$ , such that their metabolic cost during that time  
530 period was equal to lipid C turnover, 4% day<sup>-1</sup>. It should be noted that the lipid C turnover rates  
531 decreased by ½ to ¼ from the first 2 days to the last 3 days of the experiments. This could indicate  
532 an opportunistic strategy of immediate accumulation of dietary energy when it is readily available,  
533 for which speed decreases as saturation is approached and depots are replenished. High efficiency  
534 in lipid assimilation is usually regarded as an adaptation of herbivorous copepods from higher  
535 latitudes to the strong (short) seasonal character of food supply (for a review on the topic see Hagen  
536 and Auel, 2001). Smaller species with a more flexible feeding mode which have high metabolic  
537 needs, on the other hand, generally do not depend on the accumulation of large lipid reserves for  
538 guaranteed survival and reproduction. The lipid C turnover rates reported by Boissonnot, Niehoff,  
539 Hagen, Søreide, and Graeve (2016) and Graeve, Albers, and Kattner (2005) for either of these life  
540 strategies (*Calanus finmarchicus*, 2.7% day<sup>-1</sup>; *C. glacialis*, 3.0 % day<sup>-1</sup>; *Pseudocalanus minutus*,  
541 2.6% day<sup>-1</sup>; *Oithona similis*, 0.5% day<sup>-1</sup>) are lower than the ones obtained in the present study. This  
542 is true even for our lowest recorded rates, 4-6% day<sup>-1</sup>, during the last three days of the experiments.  
543 This shows how well adapted *T. longicornis* is to feeding on these diatom and dinoflagellate  
544 species. *Temora longicornis* is believed to actively grow and reproduce during the winter in the  
545 North Sea (Wesche et al., 2007), and stored dietary FAs could be mobilized for demanding  
546 metabolic processes or during events of food scarcity, thus enabling copepods to cope with

547 environmental changes in feeding conditions and to invest in reproductive output year round  
548 (Boissonnot, Niehoff, Hagen, Søreide, and Graeve, 2016; Gentsch, Kreibich, Hagen, and Niehoff,  
549 2009). This is of utmost importance for the survival of *T. longicornis* inhabiting the North Sea, an  
550 area where environmental changes (a decrease in nutrient loading coupled to an increase in light  
551 penetration) in the past three decades have affected prey quality (C:nutrient ratios) and drastically  
552 reduced the densities of calanoid copepod populations (Boersma, Wiltshire, Kong, Greve, and  
553 Renz, 2015).

554

### 555 **Relating prey biochemical composition and egg production in copepods**

556 Several studies have suggested the existence of (multiple) correlations between various  
557 biochemicals (in special the level of (n-3) PUFAs) and different indices of copepod reproduction  
558 (Anderson and Pond, 2000; Jónasdóttir, Fields, and Pantoja, 1995; Jónasdóttir, Visser, and  
559 Jespersen, 2009; Peters, Dutz, and Hagen, 2007; Sterner and Schulz, 1998). EPA is a precursor to  
560 several molecules involved in the reproductive processes, although the pathway for its use in egg  
561 production is not fully understood in copepods; DHA is essential for neural function; and both have  
562 an important role in cellular physiology regulating membrane fluidity (Brett & Müller-Navarra,  
563 1997; Jónasdóttir, Visser, and Jespersen, 2009; Sterner and Schulz, 1998). In agreement with the  
564 literature, the absolute content of certain dietary PUFAs seemed to positively affect EPR. Dino+  
565 provided copepods with only half the amount of EPA delivered by diatom diets, but it supplied per  
566 volume of culture from 100-150x more 18:3(n-3) and 10x more DHA than the latter. It also  
567 contained 2-3x more of these PUFAs than its N-limited counterpart. Dino- also showed higher  
568 18:3(n-3) and DHA and lower EPA than the diatom diets, and yielded the second highest EPR. The  
569 combination of the results from this study and from Franco-Santos et al. (2018) suggests that the  
570 FAs 18:3(n-3) and DHA are more important than EPA for egg production in *T. longicornis* females.  
571 This is contrary to the findings by Jónasdóttir, Visser, and Jespersen (2009), which highlight the  
572 importance of EPA and total C for EPR and relate 18:3(n-3) and DHA to egg hatching success in *T.*

573 *longicornis*. There is contradictory information as to whether EPR can also be influenced by  
574 maternal contribution of energy reserves or not (Anderson and Pond, 2000; Graeve, Albers, and  
575 Kattner, 2005; Jónasdóttir, Visser, and Jespersen, 2009; Peters, Dutz, and Hagen, 2007), but it is  
576 possible that dinoflagellate-fed copepods used field-accumulated EPA reserves to fuel higher EPR.

577

## 578 **Methodological constrains of results**

579 The low level of labelling transfer from *R. salina* to *O. marina* in the Dino+ cultures made it  
580 difficult to follow the transfer of labelled FAs from the latter into the copepods they were fed to. It  
581 is not possible to affirm with certainty whether the low APE and C assimilation and turnover values  
582 obtained for these copepods were due to methodological problems, biological processes, or both.  
583 There is indication, however, that it was a methodology problem. The C assimilation of these  
584 copepods accounted for only 15-26% of the recorded increase in their total lipid C, suggesting that  
585 C assimilation was underestimated. Henceforth we refer to the increase in total lipid C between  $t_{0h}$ -  
586  $t_{48h}$  and  $t_{0h}$ - $t_{120h}$  as a proxy or minimum value for C assimilation (adjusted to  $0.4 \pm 0.1$  and  $0.3 \pm 0.0$   
587  $\mu\text{g C ind}^{-1} \text{ day}^{-1}$ , respectively, and to  $3 \pm 1$  % food ingested) and turnover (adjusted to  $19 \pm 4$  and  $10$   
588  $\pm 1$  %  $\text{day}^{-1}$ , respectively) in the copepods fed with the Dino+ diet. FA-specific data on C  
589 assimilation and turnover for these copepods, however, are not included in the discussion unless in  
590 a qualitative manner.

591

592

## 593 **CONCLUSIONS**

594 Three major results can be described from the data reported herein and are discussed in the  
595 following sections. Absolute C assimilation in copepods was inversely proportional to their C  
596 investment in reproductive output, and these were influenced by the chemical and elemental  
597 composition (nutritional quality) of the prey copepods were fed with. Furthermore, relative C  
598 assimilation (as % food ingested) was similar between copepods fed with different diets, suggesting

599 that differences in reproductive output were related to copepod C ingestion rates. The high C  
600 turnover rates indicate an opportunistic feeding strategy for *T. longicornis*, which would accumulate  
601 dietary C as soon as it is readily available with a speed that decreases as saturation is approached  
602 and its lipid storage is replenished. This strategy would enable them to cope with environmental  
603 changes in feeding conditions and to invest in reproduction year round. Egg production rates were  
604 directly proportional to the copepod content of the FAs 18:3(n-3) and DHA. These results are an  
605 important addition to a small, yet growing, body of literature which estimates the transfer of lipid C  
606 between prey and predator, and to the field of copepod feeding ecology.

607

608

#### 609 **AUTHOR CONTRIBUTIONS**

610 All authors contributed to conceiving the ideas and to design the methodology;

611 RMFS collected and analyzed the data and led the writing of the manuscript;

612 All authors contributed critically to the drafts and gave final approval for publication.

613

614

#### 615 **SUPPLEMENTARY DATA**

616 Supplementary tables are available at Functional Ecology online.

617

618

#### 619 **ACKNOWLEDGEMENTS**

620 We kindly thank the crew of the Aade research vessel (AWI-BAH) for providing us with  
621 zooplankton samples and Dr Lauris Boissonnot for the insightful discussions.

622

623

#### 624 **FUNDING**

625 MARES Doctoral Programme on Marine Ecosystem Health and Conservation (Framework  
626 Agreement Number 2011-0016), to RMF-S. The work was further supported by Ghent University  
627 (BOF-GOA project 01G02617). CLM was supported by the Bundesministerium für Bildung und  
628 Forschung (BMBF grant no. 01LN1702A).

629

630

### 631 **DATA ARCHIVING**

632 The laboratory experimental data and metadata for this study are available through the  
633 PANGEA repository under the doi number (to be included once submission to PANGEA is  
634 completed).

635

636

### 637 **INTEGRITY OF RESEARCH AND REPORTING**

638

#### 639 *Ethical standards*

640 The experiments described in the present study comply with the current laws of the country in  
641 which they were performed.

642

#### 643 *Conflict of interest*

644 The authors declare that they have no conflict of interest.

645

646

### 647 **REFERENCES**

648 Anderson, M. J., Gorley, R. N., and Clarke, K. R. (2008) PERMANOVA+ for PRIMER: guide to  
649 software and statistical methods. PRIMER-E Ltd.

- 650 Anderson, T. R., & Pond, D. W. (2000). Stoichiometric theory extended to micronutrients:  
651 Comparison of the roles of essential fatty acids, carbon, and nitrogen in the nutrition of  
652 marine copepods. *Limnology and Oceanography*, 45(5), 1162–1167. doi:  
653 10.4319/lo.2000.45.5.1162
- 654 Boersma, M., Wiltshire, K. H., Kong, S. M., Greve, W., & Renz, J. (2015). Long-term change in  
655 the copepod community in the southern German Bight. *Journal of Sea Research*, 101, 41–50.  
656 doi: 10.1016/j.seares.2014.12.004
- 657 Boissonnot, L., Niehoff, B., Hagen, W., Søreide, J. E., & Graeve, M. (2016). Lipid turnover reflects  
658 life-cycle strategies of small-sized Arctic copepods. *Journal of Plankton Research*, 38(6),  
659 1420–1432. doi: 10.1093/plankt/fbw076
- 660 Brett, M. T., & Müller-Navarra, D. C. (1997). The role of highly unsaturated fatty acids in aquatic  
661 foodweb processes. *Freshwater Biology*, 38(3), 483–499. doi: 10.1046/j.1365-  
662 2427.1997.00220.x
- 663 Burian, A., Grosse, J., Winder, M., & Boschker, H. T. S. (2018). Nutrient deficiencies and the  
664 restriction of compensatory mechanisms in copepods. *Functional Ecology*, 32(3), 636–647.  
665 doi: 10.1111/1365-2435.13016
- 666 Clarke, K. R., and Gorley, R. N. (2015) PRIMER v7: User Manual/Tutorial. PRIMER-E, Plymouth.
- 667 Cotonnec, G., Brunet, C., Sautour, B. & Thoumelin, G. (2001). Nutritive value and selection of  
668 food particles by copepods during a spring bloom of *Phaeocystis* sp. in the English Channel,  
669 as determined by pigment and fatty acid analyses. *Journal of Plankton Research*, 23(7), 693–  
670 703. doi: 10.1093/plankt/23.7.693
- 671 Dalsgaard, J., John, M. S., Kattner, G., Müller-Navarra, D., & Hagen, W. (2003). Fatty acid trophic  
672 markers in the pelagic marine environment. *Advances in Marine Biology*, 46, 225–340. doi:  
673 10.1016/S0065-2881(03)46005-7
- 674 De Troch, M., Boeckx, P., Cnudde, C., Van Gansbeke, D., Vanreusel, A., Vincx, M., & Caramujo,  
675 M. J. (2012). Bioconversion of fatty acids at the basis of marine food webs: insights from a

676 compound-specific stable isotope analysis. *Marine Ecology Progress Series*, 465, 53–67.  
677 doi: 10.3354/meps09920

678 Evans, F. (1981). An investigation into the relationship of sea temperature and food supply to the  
679 size of the planktonic copepod *Temora longicornis* Müller in the North Sea. *Estuarine,  
680 Coastal and Shelf Science*, 13(2), 145–158. doi: 10.1016/S0302-3524(81)80072-2

681 Franco-Santos, R. M., Auel, H., Boersma, M., De Troch, M., Meunier, C. L., & Niehoff, B. (2018).  
682 Bioenergetics of the copepod *Temora longicornis* under different nutrient regimes. *Journal  
683 of Plankton Research*, 40(4), 420–435. doi:10.1093/plankt/fby016.

684 Fraser, A. J., Sargent, J. R., & Gamble, J. C. (1989). Lipid class and fatty acid composition of  
685 *Calanus finmarchicus* (Gunnerus), *Pseudocalanus* sp. and *Temora longicornis* Muller from  
686 a nutrient-enriched seawater enclosure. *Journal of Experimental Marine Biology and  
687 Ecology*, 130(1), 81–92. doi: 10.1016/0022-0981(89)90020-8

688 Gentsch, E., Kreibich, T., Hagen, W., & Niehoff, B. (2009). Dietary shifts in the copepod *Temora  
689 longicornis* during spring: evidence from stable isotope signatures, fatty acid biomarkers and  
690 feeding experiments. *Journal of Plankton Research*, 31(1), 45–60. doi:  
691 10.1093/plankt/fbn097

692 Graeve, M., Albers, C., & Kattner, G. (2005). Assimilation and biosynthesis of lipids in Arctic  
693 *Calanus* species based on feeding experiments with a <sup>13</sup>C labelled diatom. *Journal of  
694 Experimental Marine Biology and Ecology*, 317(1), 109–125. doi:  
695 10.1016/j.jembe.2004.11.016

696 Guillard, R. R. L. (1975). Culture of phytoplankton for feeding marine invertebrates. In Smith, W.  
697 L., and Chanley, M. H. (Eds.), *Culture of marine invertebrate animals* (pp. 29–60). US:  
698 Springer. doi: 10.1007/978-1-4615-8714-9\_3

699 Hagen, W., & Auel, H. (2001). Seasonal adaptations and the role of lipids in oceanic zooplankton.  
700 *Zoology*, 104(3-4), 313–326. doi: 10.1078/0944-2006-00037



- 701 Helland, S., Terjesen, B. F., & Berg, L. (2003). Free amino acid and protein content in the  
702 planktonic copepod *Temora longicornis* compared to *Artemia franciscana*. *Aquaculture*,  
703 215(1-4), 213–228. doi: 10.1016/S0044-8486(02)00043-1
- 704 Hessen, D. O., & Anderson, T. R. (2008). Excess carbon in aquatic organisms and ecosystems:  
705 physiological, ecological, and evolutionary implications. *Limnology and Oceanography*,  
706 53(4), 1685–96. doi: 10.4319/lo.2008.53.4.1685
- 707 Ihaka, R., & Gentleman, R. (1993). R (Version 3.4.4) [software]. Retrieved from [https://cran.r-](https://cran.r-project.org/bin/windows/base/old/3.2.4/)  
708 [project.org/bin/windows/base/old/3.2.4/](https://cran.r-project.org/bin/windows/base/old/3.2.4/).
- 709 Jónasdóttir, S. H., Fields, D., & Pantoja, S. (1995). Copepod egg production in Long Island Sound,  
710 USA, as a function of the chemical composition of seston. *Marine Ecology Progress Series*,  
711 119, 87–98. doi: 10.3354/meps119087
- 712 Jónasdóttir, S. H., Visser, A. W., & Jespersen, C. (2009). Assessing the role of food quality in the  
713 production and hatching of *Temora longicornis* eggs. *Marine Ecology Progress Series*, 382,  
714 139–150. doi: 10.3354/meps07985
- 715 Kiørboe, T., & Nielsen, T. G. (1994). Regulation of zooplankton biomass and production in a  
716 temperate, coastal ecosystem. 1. Copepods. *Limnology and Oceanography*, 39(3), 493–507.  
717 doi: 10.4319/lo.1994.39.3.0493
- 718 Klein Breteler, W. C. M., Schogt, N., Baas, M., Schouten, S., & Kraay, G. W. (1999). Trophic  
719 upgrading of food quality by protozoans enhancing copepod growth: role of essential lipids.  
720 *Marine Biology*, 135(1), 191–198. doi: 10.1007/s002270050616
- 721 Kreibich, T., Saborowski, R., Hagen, W., & Niehoff, B. (2008). Short-term variation of nutritive  
722 and metabolic parameters in *Temora longicornis* females (Crustacea, Copepoda) as a  
723 response to diet shift and starvation. *Helgoland Marine Research*, 62(3), 241–249. doi:  
724 10.1007/s10152-008-0112-0
- 725 Kreibich, T., Saborowski, R., Hagen, W., & Niehoff, B. (2011). Influence of short-term nutritional  
726 variations on digestive enzyme and fatty acid patterns of the calanoid copepod *Temora*

727 *longicornis*. Journal of Experimental Marine Biology and Ecology, 407(2), 182–189. doi:  
728 10.1016/j.jembe.2011.06.013

729 Lee, R. F., Hagen, W., & Kattner, G. (2006). Lipid storage in marine zooplankton. Marine Ecology  
730 Progress Series, 307, 273–306. doi: 10.3354/meps307273

731 Löder, M. G. J., Meunier, C. L., Wiltshire, K. H., Boersma, M., & Aberle, N. (2011). The role of  
732 ciliates, heterotrophic dinoflagellates and copepods in structuring spring plankton  
733 communities at Helgoland Roads, North Sea. Marine Biology, 158(7), 1551–1580. doi:  
734 10.1007/s00227-011-1670-2

735 Maar, M., Nielsen, T. G., Gooding, S., Tönnesson, K., Tiselius, P., Zervoudaki, S., ... Richardson,  
736 K. (2004). Trophodynamic function of copepods, appendicularians and protozooplankton in  
737 the late summer zooplankton community in the Skagerrak. Marine Biology, 144(5), 917–  
738 933. doi: 10.1007/s00227-003-1263-9

739 Mitra, A., & Flynn, K. J. (2005). Predator–prey interactions: is ‘ecological stoichiometry’ sufficient  
740 when good food goes bad? Journal of Plankton Research, 27(5), 393–399. doi:  
741 10.1093/plankt/fbi022

742 Møller, E. F. (2007). Production of dissolved organic carbon by sloppy feeding in the copepods  
743 *Acartia tonsa*, *Centropages typicus*, and *Temora longicornis*. Limnology and Oceanography,  
744 52(1), 79–84. doi: 10.4319/lo.2007.52.1.0079

745 Peters, J., Dutz, J., & Hagen, W. (2007). Role of essential fatty acids on the reproductive success of  
746 the copepod *Temora longicornis* in the North Sea. Marine Ecology Progress Series, 341,  
747 153–163. doi: 10.3354/meps341153

748 Peters, J., Dutz, J., & Hagen, W. (2013). Trophodynamics and life-cycle strategies of the copepods  
749 *Temora longicornis* and *Acartia longiremis* in the Central Baltic Sea. Journal of Plankton  
750 Research, 35(3), 595–609. doi: 10.1093/plankt/fbt004

751 Sterner, R. W., & Elser, J. J. (2002). Ecological stoichiometry: the biology of elements from  
752 molecules to the biosphere. Princeton University Press. doi:

- 753 Sterner, R. W., & Schulz, K. L. (1998). Zooplankton nutrition: recent progress and a reality check.  
754 *Aquatic Ecology*, 32(4), 261–279. doi: 10.1023/A:1009949400573
- 755 Urabe, J., Shimizu, Y., & Yamaguchi, T. (2018). Understanding the stoichiometric limitation of  
756 herbivore growth: the importance of feeding and assimilation flexibilities. *Ecology Letters*,  
757 21(2), 197–206. doi: 10.1111/ele.12882
- 758 Weers, P. M. M., & Gulati, R. D. (1997). Growth and reproduction of *Daphnia galeata* in response  
759 to changes in fatty acids, phosphorus, and nitrogen in *Chlamydomonas reinhardtii*.  
760 *Limnology and Oceanography*, 42(7), 1584–1589. doi: 10.4319/lo.1997.42.7.1584
- 761 Wesche, A., Wiltshire, K. H., & Hirche, H. J. (2007). Overwintering strategies of dominant  
762 calanoid copepods in the German Bight, southern North Sea. *Marine Biology*, 151, 1309–  
763 1320. doi: 10.1007/s00227-006-0560-5

764

765

## 766 SUPPLEMENTARY MATERIAL

767

768 **Supplementary table S1** Post hoc test results. Tukey HSD results ( $p$ -value) for differences in (1)  
769 prey culture: carbon (C), nitrogen (N), and total lipid contents, molar C:N ratios, and  $^{13}\text{C}$  labelling  
770 (atom%); and (2) *Temora longicornis* females: C, N, and total lipid contents, molar C:N ratios,  $^{13}\text{C}$   
771 enrichment (APE), C assimilation rate ( $C_{\text{assim}}$ , as  $\mu\text{g C ind}^{-1} \text{ day}^{-1}$ ), and C turnover rate ( $C_{\text{T}}$ ,  $\% \text{ day}^{-1}$ ).  
772 Similarity percentages (SIMPER) results (% average dissimilarity) for differences in FA  
773 composition of prey cultures. Sampling days:  $t_{0\text{h}}$  (*in situ*),  $t_{48\text{h}}$  (after 48h of experiment),  $t_{120\text{h}}$  (at the  
774 end of the experiment). Diet treatments: single cultures in nutrient-replete and N-depleted  
775 conditions of *Conticribra weissflogii* (Diat+ and Diat–, respectively) and *Oxyrrhis marina* (Dino+  
776 and Dino–, respectively). Significant values: \* ( $p < 0.05$ ), \*\* ( $p < 0.01$ ); \*\*\* ( $p < 0.001$ ).  $p$ -values  
777 for tests with an adjusted  $p$ -value of 0.0167 are given in number format.

778

Effect	Factor	Variable	Treatment pair	Result		
			Dino+/Diat-	**		
		C content	Dino+/Diat+; Dino+/Dino-	***		
		N content	Diat-/Diat+; Diat-/Dino+; Diat-/Dino-; Dino+/Diat+; Dino+/Dino-	***		
		C:N	Diat-/Diat+; Diat-/Dino+; Diat-/Dino-	***		
		Lipid content	Dino+/Dino-	**		
		atom%	Dino+/Diat+; Dino+/Diat-; Dino+/Dino-; Dino-/Diat+; Dino-/Diat-	***		
Prey cultures	Day	Diat+	Day 1/Day 5	**		
		Diat+	Day 4/Day 1; Day 4/Day 3; Day 4/Day 5	***		
		Diat-	Day 5/Day 1; Day 5/Day 4	**		
		Diat-	Day 5/Day 2	***		
		Dino+	Day 3/Day 5	*		
		Dino+	Day 1/Day 3; Day 1/Day 4; Day 1/Day 5; Day 2/Day 3; Day 2/Day 4; Day 2/Day 5	***		
		Dino-	Day 4/Day 5	**		
		Dino-	Day 5/Day 1; Day 5/Day 3	***		
				SIMPER	Diat+/Diat-	21%
					Diat+/Dino+	65%
			Diat+/Dino-	68%		
			Diat-/Dino+	58%		
			Diat-/Dino-	60%		
			Dino+/Dino-	25%		
Copepod		APE	Dino+/Diat+; Dino+/Diat-; Dino+/Dino-; Dino-/Diat+; Dino-/Diat-	***		
			Diat+/Diat-	*		
	Diet	C <sub>assim</sub>	Diat+/Dino-; Diat-/Dino-	**		
			Dino+/Diat-; Dino+/Dino-	***		
		C <sub>T</sub>	Diat-/Dino-	**		
			Dino+/Diat+; Dino+/Diat-; Dino+/Dino-	***		
	Day	APE	$t_{48h}/t_{120h}$	***		
		C <sub>assim</sub>	$t_{48h}/t_{120h}$	***		
		C <sub>T</sub>	$t_{48h}/t_{120h}$	***		
			$t_{0h}/t_{48h}$	0.009		
		C content	$t_{0h}/t_{120h}$	0.014		
			$t_{0h}/t_{48h}$	0.008		
	Diet:		$t_{0h}/t_{120h}$	0.009		
	Day	Dino-	$t_{0h}/t_{48h}$	0.013		
		Diat+	$t_{0h}/t_{120h}$	0.0163		
		Diat-	$t_{0h}/t_{48h}$	0.011		
		Dino+	$t_{0h}/t_{120h}$	0.002		
		Dino-	$t_{0h}/t_{120h}$	0.004		

	Diat+		0.007
	Diat-	$t_{0h}/t_{48h}$	0.014
	Dino+		0.002
	Diat+	Lipid content	0.008
	Dino+		0.0003
	Dino-	$t_{0h}/t_{120h}$	0.006
	Dino-		0.009
$t_{48h}$	C content	Diat+/Diat-; Diat+/Dino-	*
$t_{48h}$	N content	Diat-/Diat+; Diat-/Dino+; Diat-/Dino-	***
$t_{48h}$	C:N	Diat-/Diat+; Diat+/Dino-	**
		Diat-/Dino+; Diat-/Dino-	***
$t_{120h}$		Dino-/Diat-; Dino-/Dino+	*
$t_{120h}$		Diat+/Diat-; Diat+/Dino+	**
$t_{48h}$	Lipid content	Diat-/Dino+	*
		Diat-/Dino-	**
$t_{120h}$		Diat-/Diat+; Diat-/Dino-	*
$t_{48h}$	APE	Dino-/Diat+; Dino-/Diat-	***
		Diat+/Diat-	*
$t_{120h}$		Dino-/Diat+; Dino-/Diat-	***
$t_{48h}$	$C_{assim}$	Diat+/Dino-	*
		Diat-/Dino-	**
$t_{120h}$		Diat-/Diat+; Diat-/Dino-	*
$t_{48h}$	$C_{assim}$ SFA	Diat-/Diat+; Diat-/Dino-	*
$t_{120h}$		Diat+/Diat-	*
$t_{48h}$	$C_{assim}$ MUFA	Diat+/Diat-; Diat+/Dino-; Diat-/Dino-	***
		Diat+/Dino-	**
$t_{120h}$		Diat-/Diat+; Diat-/Dino-	***
$t_{48h}$	$C_{assim}$ PUFA	Diat+/Dino-	*
		Diat-/Dino-	*
$t_{120h}$	$C_T$ SFA	Diat-/Diat+	*
		Diat-/Dino-	**
$t_{48h}$		Diat+/Diat-; Diat+/Dino-; Diat-/Dino-	***
$t_{120h}$	$C_T$ MUFA	Diat+/Dino-	*
		Diat-/Diat+; Diat-/Dino-	***
$t_{48h}$	$C_T$ PUFA	Diat+/Dino-	*
		Diat+/Diat-	**
$t_{120h}$		Diat+/Diat-	*
		Diat+/Dino-	**

779 **Supplementary table S2** One-way Analysis of Variance (ANOVA) and one- and two-way repeated  
780 measures ANOVA (RM ANOVA) results (F-statistics, degrees of freedom (df), and *p*-values) for  
781 statistical analyses of (1) prey culture: carbon (C), nitrogen (N), and total lipid contents, molar C:N  
782 ratios, and <sup>13</sup>C labelling of fatty acids (atom%); and (2) *Temora longicornis* females: C, N, and lipid  
783 content, molar C:N ratios, <sup>13</sup>C enrichment (APE), total and FA-specific C assimilation rate (*C*<sub>assim</sub>,  
784 as μg C ind<sup>-1</sup> day<sup>-1</sup>), and total and FA-specific C turnover rate (*C*<sub>T</sub> % day<sup>-1</sup>). T-test results (t-  
785 statistics, df, *p*-values) for within treatment comparisons of copepod APE, *C*<sub>assim</sub> and *C*<sub>T</sub> are also  
786 given. Non-significant values are omitted. Significant values: \* (*p* < 0.05), \*\* (*p* < 0.01); \*\*\* (*p* <  
787 0.001). FA denominations: saturated (SFA), monounsaturated (MUFA) and polyunsaturated  
788 (PUFA) fatty acids.

	Test	Factor	Response variable	F-statistics	df	<i>p</i> -value			
Prey cultures	1-way ANOVA		C content	19.99	3, 16	***			
			N content	121.1	3, 16	***			
			molar C:N ratio	74.31	3, 16	***			
			Lipid content	22.73	3, 14	***			
			atom%	362.3	3, 14	***			
			Diat+		21.79	3, 71	***		
			Diat-	atom%	5.824	4, 89	***		
			Dino+		30.32	4, 76	***		
			Dino-		9.901	3, 52	***		
Copepods	2-way RM ANOVA	All	C content	11.38	3	**			
			N content	17.76	3	***			
			molar C:N ratio	36.52	3	***			
			Total lipid content	14.26	3	**			
	Diet treatments	1-way RM ANOVA	Diat+		20.69	2,4	**		
				Dino+	C content	12.64	2,4	*	
			Dino-	23.93		2,4	**		
			Diat+		14.61	2,4	*		
				Dino+	N content	8.888	2,4	*	
			Dino-	15.01		2,4	*		
			Diat-		15.61	2,4	*		
				Dino+	molar C:N ratio	35.67	2,4	**	
			Dino-	28.05		2,4	**		
			Diat+		25.46	2,4	**		
				Diat-	Total lipid content	15.54	2,4	*	
			Dino+	116.4		2,4	***		
			Dino-	21.77		2,4	**		
				t-test	Diat+		-11.977	2	**
					Diat-	APE	-17.397	2	**
	Dino+	4.5826	2		*				

		Dino-		-9.2307	2	*
		Diat+		8.1848	2	*
		Dino-	$C_{\text{assim}}$	4.3235	2	*
		Diat+	$C_{\text{assim}}$ SFA	22.258	2	**
		Dino-		4.7035	2	*
		Diat+	$C_{\text{assim}}$ MUFA	20.524	2	**
		Diat+		5.5807	2	*
		Dino-	$C_{\text{assim}}$ PUFA	4.5783	2	*
		Diat+		14.637	2	**
		Diat-		113.28	2	***
		Dino+	$C_T$	501.43	2	***
		Dino-		35.43	2	***
		Diat+		22.942	2	**
		Diat-	$C_T$ SFA	35.203	2	***
		Dino-		31.262	2	**
		Diat+		11.75	2	**
		Diat-	$C_T$ MUFA	43.64	2	***
		Dino-		11.426	2	**
		Diat+		7.1813	2	*
		Diat-	$C_T$ PUFA	54.049	2	***
		Dino-		43.25	2	***
			C content	43.448	2	***
			N content	23.345	2	***
			molar C:N ratio	42.315	2	***
			Total lipid content	94.937	2	***
Days sampled	2-way RM ANOVA	$t_{48h}$	C content	7.265	3,8	*
		$t_{48h}$	N content	29.43	3,8	***
		$t_{48h}$	molar C:N ratio	45.52	3,8	***
		$t_{100h}$		11.74	3,8	**
	$t_{48h}$	Total lipid content	8	3,8	**	
	$t_{100h}$		6.101	3,8	*	
	$t_{48h}$	APE	60.7	2,6	***	
	$t_{100h}$		124.6	2,6	***	
	$t_{48h}$	$C_{\text{ass}}$	15.88	2,6	**	
	$t_{100h}$		9.81	2,6	*	
	$t_{48h}$	$C_{\text{assim}}$ SFA	11.09	2,6	**	
	$t_{100h}$		9.391	2,6	*	
	$t_{48h}$	$C_{\text{assim}}$ MUFA	237.5	2,6	***	
	$t_{100h}$		130.8	2,6	***	
	$t_{48h}$	$C_{\text{assim}}$ PUFA	5.34	2,6	*	
	$t_{48h}$	$C_T$ SFA	6.333	2,6	*	
	$t_{100h}$		21.44	2,6	**	
	$t_{48h}$	$C_T$ MUFA	162.1	2,6	***	
$t_{100h}$		135.1	2,6	***		
$t_{48h}$	$C_T$ PUFA	16.78	2,6	**		
$t_{100h}$		16.8	2,6	**		
Diet treatment : Day sampled		All	molar C:N ratio	9.693	6	***

789 **Supplementary table S3** Permutational Multivariate Analysis of Variance (PERMANOVA) results  
790 (pseudo F-statistics or t-statistics, *p*-values for standard permutations (p(perm)) and for Monte Carlo  
791 simulations (p(MC)), unique number of permutations (UNP)) for *Temora longicornis* fatty acid  
792 (FA) profiles (>1% total FAs). Diet treatments: nutrient-replete and N-depleted *Conticribra*  
793 *weissflogii* (Diat+ and Diat-, respectively) and *Oxyrrhis marina* (Dino+ and Dino-, respectively).  
794 Non-significant values are omitted. Significant values: \* (*p* < 0.05), \*\* (*p* < 0.01); \*\*\* (*p* < 0.001).

Main effects	F-statistics	p(perm)	p(MC)	UNP
Treatment	968.73	***	***	971
Day sampled	360.93	***	***	996
Treatment : Day sampled	120.01	***	***	997
Pairwise comparisons	t-statistics	p(perm)	p(MC)	UNP
Diat+ x Diat-	8.7585		***	10
Diat+ x Dino+	32.515		***	798
Diat+ x Dino-	30.096		***	10
Diat- x Dino+	52.824		***	799
Diat- x Dino-	45.824		***	10
Dino+ x Dino-	8.8395		***	800
<i>t</i> <sub>0h</sub> X <i>t</i> <sub>48h</sub>	16.796	***	***	998
<i>t</i> <sub>0h</sub> X <i>t</i> <sub>120h</sub>	26.109	***	***	801
<i>t</i> <sub>48h</sub> X <i>t</i> <sub>120h</sub>	7.7516	***	***	998
<i>t</i> <sub>0h</sub> X <i>t</i> <sub>48h</sub>	5.2527		*	38
Diat+ <i>t</i> <sub>0h</sub> X <i>t</i> <sub>120h</sub>	8.8608		**	38
<i>t</i> <sub>48h</sub> X <i>t</i> <sub>120h</sub>	1.7474			38
<i>t</i> <sub>0h</sub> X <i>t</i> <sub>48h</sub>	10.718		**	38
Diat- <i>t</i> <sub>0h</sub> X <i>t</i> <sub>120h</sub>	14.355		**	38
<i>t</i> <sub>48h</sub> X <i>t</i> <sub>120h</sub>	5.6138		*	38
<i>t</i> <sub>0h</sub> X <i>t</i> <sub>48h</sub>	22.247		***	38
Dino+ <i>t</i> <sub>0h</sub> X <i>t</i> <sub>120h</sub>	28.433		**	38
<i>t</i> <sub>48h</sub> X <i>t</i> <sub>120h</sub>	Negative			38
<i>t</i> <sub>0h</sub> X <i>t</i> <sub>48h</sub>	15.496		**	38
Dino- <i>t</i> <sub>0h</sub> X <i>t</i> <sub>120h</sub>	24.23		***	38
<i>t</i> <sub>48h</sub> X <i>t</i> <sub>120h</sub>	13.578		**	38
Diat+ x Diat-	4.4046		**	10
Diat+ x Dino+	18.338		***	809
<i>t</i> <sub>48h</sub> Diat+ x Dino-	16.359		***	10
Diat- x Dino+	35.394		***	810
Diat- x Dino-	26.979		***	10
Dino+ x Dino-	8.9636		***	811
Diat+ x Diat-	7.4701		***	10
Diat+ x Dino+	30.698		***	812
<i>t</i> <sub>120h</sub> Diat+ x Dino-	28.165		***	10
Diat- x Dino+	57.689		***	813
Diat- x Dino-	45.74		***	10
Dino+ x Dino-	12.907		***	814



815 **Supplementary table S4** Total (TFA) and fatty acid-specific <sup>13</sup>C enrichment (APE) between  $t_{0h}$ - $t_{48h}$   
816 and between  $t_{0h}$ - $t_{120h}$  for *Temora longicornis* females. Diet treatments: single cultures of *Conticribra*  
817 *weissflogii* (Diat) and *Oxyrrhis marina* (Dino) in nutrient-replete (+) and N-depleted (-) conditions.  
818 Values are mean  $\pm$  standard deviation (n = 3 each). – : not detected.

APE								
FA	Diat+		Diat-		Dino+		Dino-	
	$t_{48h}$	$t_{120h}$	$t_{48h}$	$t_{120h}$	$t_{48h}$	$t_{120h}$	$t_{48h}$	$t_{120h}$
TFA	0.7 $\pm$ 0.0	0.9 $\pm$ 0.0	0.7 $\pm$ 0.0	1.0 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.5 $\pm$ 0.0	0.7 $\pm$ 0.0
14:0	0.8 $\pm$ 0.0	1.2 $\pm$ 0.0	0.8 $\pm$ 0.0	1.2 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.4 $\pm$ 0.0	0.7 $\pm$ 0.1
i-15:0	0.4 $\pm$ 0.1	0.6 $\pm$ 0.0	0.3 $\pm$ 0.0	0.6 $\pm$ 0.0	0.2 $\pm$ 0.0	0.1 $\pm$ 0.0	0.9 $\pm$ 0.0	0.9 $\pm$ 0.0
15:0	1.2 $\pm$ 0.0	1.4 $\pm$ 0.0	1.4 $\pm$ 0.1	1.6 $\pm$ 0.0	0.2 $\pm$ 0.0	0.1 $\pm$ 0.0	0.4 $\pm$ 0.0	0.7 $\pm$ 0.0
16:0	0.9 $\pm$ 0.0	1.2 $\pm$ 0.0	1.0 $\pm$ 0.0	1.3 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.7 $\pm$ 0.0	0.8 $\pm$ 0.0
16:1n5	1.5 $\pm$ 0.0	1.5 $\pm$ 0.0	1.3 $\pm$ 0.0	1.6 $\pm$ 0.0	–	–	–	–
16:1n7	1.3 $\pm$ 0.0	1.4 $\pm$ 0.0	1.5 $\pm$ 0.0	1.7 $\pm$ 0.0	0.0 $\pm$ 0.0	0.0 $\pm$ 0.0	0.1 $\pm$ 0.0	0.4 $\pm$ 0.1
16:2n4	1.7 $\pm$ 0.0	1.6 $\pm$ 0.0	1.5 $\pm$ 0.0	1.7 $\pm$ 0.0	–	–	–	–
16:3n4	1.7 $\pm$ 0.0	1.7 $\pm$ 0.0	1.8 $\pm$ 0.0	1.8 $\pm$ 0.0	–	–	–	–
16:4n1	0.5 $\pm$ 0.0	1.1 $\pm$ 0.0	0.2 $\pm$ 0.0	0.6 $\pm$ 0.1	–	–	–	–
17:0	0.1 $\pm$ 0.0	0.3 $\pm$ 0.1	0.5 $\pm$ 0.1	0.7 $\pm$ 0.1	0.2 $\pm$ 0.0	0.1 $\pm$ 0.0	0.4 $\pm$ 0.0	0.7 $\pm$ 0.0
18:0	0.3 $\pm$ 0.0	0.5 $\pm$ 0.1	0.4 $\pm$ 0.0	0.7 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.2 $\pm$ 0.0	0.4 $\pm$ 0.0
18:1n7	0.8 $\pm$ 0.0	1.1 $\pm$ 0.1	0.8 $\pm$ 0.0	1.2 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.6 $\pm$ 0.1	0.7 $\pm$ 0.0
18:1n9	0.2 $\pm$ 0.0	0.2 $\pm$ 0.1	0.7 $\pm$ 0.1	1.0 $\pm$ 0.1	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.9 $\pm$ 0.2	1.0 $\pm$ 0.0
18:2n6	0.9 $\pm$ 0.0	1.1 $\pm$ 0.0	0.7 $\pm$ 0.0	1.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	1.1 $\pm$ 0.0	1.0 $\pm$ 0.0
18:3n3	0.5 $\pm$ 0.0	0.6 $\pm$ 0.3	0.5 $\pm$ 0.0	0.9 $\pm$ 0.1	0.1 $\pm$ 0.0	0.0 $\pm$ 0.0	1.1 $\pm$ 0.1	1.0 $\pm$ 0.0
18:3n6	1.3 $\pm$ 0.1	1.5 $\pm$ 0.0	1.1 $\pm$ 0.0	1.5 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.7 $\pm$ 0.0	0.8 $\pm$ 0.0
18:4n3	1.5 $\pm$ 0.0	1.6 $\pm$ 0.0	1.5 $\pm$ 0.0	1.7 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.8 $\pm$ 0.0	1.0 $\pm$ 0.0
20:1n7	0.4 $\pm$ 0.1	0.6 $\pm$ 0.3	0.4 $\pm$ 0.1	0.6 $\pm$ 0.1	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.0 $\pm$ 0.0	–
20:1n9	0.1 $\pm$ 0.0	0.3 $\pm$ 0.0	0.3 $\pm$ 0.0	0.5 $\pm$ 0.1	0.0 $\pm$ 0.0	0.1 $\pm$ 0.0	0.6 $\pm$ 0.0	0.9 $\pm$ 0.0
20:2n6	0.3 $\pm$ 0.1	0.7 $\pm$ 0.0	0.2 $\pm$ 0.0	0.5 $\pm$ 0.1	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.7 $\pm$ 0.1	0.9 $\pm$ 0.0
20:4n3	0.8 $\pm$ 0.1	1.0 $\pm$ 0.1	1.1 $\pm$ 0.1	1.4 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.7 $\pm$ 0.1	0.9 $\pm$ 0.0
20:4n6	0.4 $\pm$ 0.1	0.7 $\pm$ 0.2	0.3 $\pm$ 0.0	0.6 $\pm$ 0.1	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.2 $\pm$ 0.1	0.6 $\pm$ 0.0

20:5n3	$0.8 \pm 0.0$	$1.2 \pm 0.0$	$0.7 \pm 0.0$	$1.1 \pm 0.0$	$0.1 \pm 0.0$	$0.1 \pm 0.0$	$0.2 \pm 0.0$	$0.4 \pm 0.0$
22:1n7	$0.2 \pm 0.1$	$0.6 \pm 0.0$	$0.2 \pm 0.0$	$0.5 \pm 0.1$	$0.1 \pm 0.0$	$0.1 \pm 0.0$	$0.1 \pm 0.0$	$0.2 \pm 0.0$
22:1n9	–	$0.1 \pm 0.2$	$-0.1 \pm 0.1$	$0.0 \pm 0.1$	–	–	$0.1 \pm 0.0$	–
22:5n3	$0.4 \pm 0.0$	$0.8 \pm 0.0$	$0.4 \pm 0.0$	$0.8 \pm 0.0$	$0.1 \pm 0.0$	$0.1 \pm 0.0$	$0.4 \pm 0.0$	$0.0 \pm 0.0$
22:6n3	$0.5 \pm 0.0$	$0.8 \pm 0.0$	$0.4 \pm 0.0$	$0.7 \pm 0.0$	$0.2 \pm 0.0$	$0.1 \pm 0.0$	$0.7 \pm 0.0$	$0.9 \pm 0.0$

## CHAPTER III

### **Fatty acid bioconversion and homeostasis in the larvae of the sand mason worm *Lanice conchilega***

Franco-Santos RM, Auel H, Boersma M, De Troch M, Graeve M, Meunier CL, Niehoff B

Submitted for publication to PLoS ONE on October 30<sup>th</sup> 2018.



# PLOS ONE

## You are not always what you eat - Fatty acid bioconversion and lipid homeostasis in the larvae of the sand mason worm *Lanice conchilega*

--Manuscript Draft--

<b>Manuscript Number:</b>	
<b>Article Type:</b>	Research Article
<b>Full Title:</b>	You are not always what you eat - Fatty acid bioconversion and lipid homeostasis in the larvae of the sand mason worm <i>Lanice conchilega</i>
<b>Short Title:</b>	FA bioconversion and lipid homeostasis in <i>Lanice conchilega</i>
<b>Corresponding Author:</b>	Rita Melo Franco-Santos, M.Sc. Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung Bremerhaven, GERMANY
<b>Keywords:</b>	assimilation; bioconversion; compound-specific stable isotope analysis; fatty acids; larvae; polychaete
<b>Abstract:</b>	The meroplanktonic larvae of benthic adults are an important seasonal component of the zooplankton in temperate coastal waters. The larvae of the reef-building polychaete <i>Lanice conchilega</i> can make up to 15% of the summer zooplankton biomass in the North Sea. Despite their importance for reef maintenance (which positively affects the benthic community), little is known of the trophic ecology of these meroplanktonic larvae. Qualitative and quantitative estimates of carbon (C) transfer between trophic levels and of FA-specific assimilation, biosynthesis, and bioconversion can be obtained by compound-specific stable isotope analysis of fatty acids (FA). The present work tested the hypothesis that the concept of fatty acid trophic markers (FATM), widely used for studies on holoplankton with intermediate to high lipid contents, is also applicable to lipid-poor organisms such as meroplanktonic larvae. The incorporation of isotopically-enriched dietary C by <i>L. conchilega</i> larvae was traced, and lipid assimilation did not follow FA-specific relative availabilities in the diet. Furthermore, FAs that were unavailable in the diet, such as 22:5(n-3), were recorded in <i>L. conchilega</i> , suggesting their bioconversion by the larvae. The results indicate that <i>L. conchilega</i> larvae preferentially assimilate certain FAs and regulate their FA composition (lipid homeostasis) independently of that of their diet. Their quasi-homeostatic response to dietary FA availability could imply that the concept of FATM has limited application in lipid-poor organisms such as <i>L. conchilega</i> larvae.
<b>Order of Authors:</b>	Rita M Franco-Santos, M.Sc. Holger Auel Boersma Maarten Marleen De Troch Martin Graeve Cédric Leo Meunier Barbara Niehoff
<b>Opposed Reviewers:</b>	
<b>Additional Information:</b>	
<b>Question</b>	<b>Response</b>
<b>Financial Disclosure</b>  Enter a financial disclosure statement that describes the sources of funding for the work included in this submission. Review the <a href="#">submission guidelines</a> for detailed	RMF-S received a scholarship from the MARES Doctoral Programme on Marine Ecosystem Health and Conservation ( <a href="http://www.mares-eu.org">www.mares-eu.org</a> ) (Framework Agreement Number 2011-0016). MDT was supported by Ghent University ( <a href="http://www.ugent.be">www.ugent.be</a> ) (BOF-GOA project 01G02617). CLM was supported by the Bundesministerium für Bildung und Forschung ( <a href="http://www.bmbf.de">www.bmbf.de</a> ) (BMBF grant no. 01LN1702A). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the

<p>requirements. View published research articles from <a href="#">PLOS ONE</a> for specific examples.</p> <p>This statement is required for submission and <b>will appear in the published article</b> if the submission is accepted. Please make sure it is accurate.</p> <p><b>Unfunded studies</b> Enter: <i>The author(s) received no specific funding for this work.</i></p> <p><b>Funded studies</b> Enter a statement with the following details:</p> <ul style="list-style-type: none"> <li>• Initials of the authors who received each award</li> <li>• Grant numbers awarded to each author</li> <li>• The full name of each funder</li> <li>• URL of each funder website</li> <li>• Did the sponsors or funders play any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript?</li> <li>• <b>NO</b> - Include this sentence at the end of your statement: <i>The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.</i></li> <li>• <b>YES</b> - Specify the role(s) played.</li> </ul> <p>* typeset</p>	<p>manuscript.</p>
<p><b>Competing Interests</b></p> <p>Use the instructions below to enter a competing interest statement for this submission. On behalf of all authors, disclose any <a href="#">competing interests</a> that could be perceived to bias this work—acknowledging all financial support and any other relevant financial or non-financial competing interests.</p> <p>This statement <b>will appear in the published article</b> if the submission is accepted. Please make sure it is accurate. View published research articles from <a href="#">PLOS ONE</a> for specific examples.</p>	<p>The authors have declared that no competing interests exist.</p>

**NO authors have competing interests**

Enter: *The authors have declared that no competing interests exist.*

**Authors with competing interests**

Enter competing interest details beginning with this statement:

*I have read the journal's policy and the authors of this manuscript have the following competing interests: [insert competing interests here]*

\* typeset

**Ethics Statement**

Enter an ethics statement for this submission. This statement is required if the study involved:

- Human participants
- Human specimens or tissue
- Vertebrate animals or cephalopods
- Vertebrate embryos or tissues
- Field research

Write "N/A" if the submission does not require an ethics statement.

General guidance is provided below.

Consult the [submission guidelines](#) for detailed instructions. **Make sure that all information entered here is included in the Methods section of the manuscript.**

N/A

### Format for specific study types

#### Human Subject Research (involving human participants and/or tissue)

- Give the name of the institutional review board or ethics committee that approved the study
- Include the approval number and/or a statement indicating approval of this research
- Indicate the form of consent obtained (written/oral) or the reason that consent was not obtained (e.g. the data were analyzed anonymously)

#### Animal Research (involving vertebrate animals, embryos or tissues)

- Provide the name of the Institutional Animal Care and Use Committee (IACUC) or other relevant ethics board that reviewed the study protocol, and indicate whether they approved this research or granted a formal waiver of ethical approval
- Include an approval number if one was obtained
- If the study involved *non-human primates*, add *additional details* about animal welfare and steps taken to ameliorate suffering
- If anesthesia, euthanasia, or any kind of animal sacrifice is part of the study, include briefly which substances and/or methods were applied

#### Field Research

Include the following details if this study involves the collection of plant, animal, or other materials from a natural setting:

- Field permit number
- Name of the institution or relevant body that granted permission

#### Data Availability

Authors are required to make all data underlying the findings described fully available, without restriction, and from the time of publication. PLOS allows rare exceptions to address legal and ethical concerns. See the [PLOS Data Policy](#) and [FAQ](#) for detailed information.

Yes - all data are fully available without restriction



<p>A Data Availability Statement describing where the data can be found is required at submission. Your answers to this question constitute the Data Availability Statement and <b>will be published in the article</b>, if accepted.</p> <p><b>Important:</b> Stating 'data available on request from the author' is not sufficient. If your data are only available upon request, select 'No' for the first question and explain your exceptional situation in the text box.</p> <p>Do the authors confirm that all data underlying the findings described in their manuscript are fully available without restriction?</p>	
<p><b>Describe where the data may be found in full sentences. If you are copying our sample text, replace any instances of XXX with the appropriate details.</b></p> <ul style="list-style-type: none"> <li>• If the data are <b>held or will be held in a public repository</b>, include URLs, accession numbers or DOIs. If this information will only be available after acceptance, indicate this by ticking the box below. For example: <i>All XXX files are available from the XXX database (accession number(s) XXX, XXX).</i></li> <li>• If the data are all contained <b>within the manuscript and/or Supporting Information files</b>, enter the following: <i>All relevant data are within the manuscript and its Supporting Information files.</i></li> <li>• If neither of these applies but you are able to provide <b>details of access elsewhere</b>, with or without limitations, please do so. For example:</li> </ul> <p><i>Data cannot be shared publicly because of [XXX]. Data are available from the XXX Institutional Data Access / Ethics Committee (contact via XXX) for researchers who meet the criteria for access to confidential data.</i></p> <p><i>The data underlying the results presented in the study are available from (include the name of the third party</i></p>	<p>The laboratory experimental data and metadata for this study will be available through the PANGAEA repository under the doi number (to be included once submission to PANGAEA is completed).</p>

<p><i>and contact information or URL).</i></p> <ul style="list-style-type: none"><li>• This text is appropriate if the data are owned by a third party and authors do not have permission to share the data.</li></ul> <p>* typeset</p>	
Additional data availability information:	Tick here if the URLs/accession numbers/DOIs will be available only after acceptance of the manuscript for publication so that we can ensure their inclusion before publication.

Dear Editor-in-Chief Dr. Joerg Heber,

October 30<sup>th</sup>, 2018

Enclosed you will find the manuscript entitled “**You are not always what you eat - Fatty acid bioconversion and lipid homeostasis in the larvae of the sand mason worm *Lanice conchilega***”, that I wish to submit for publication in PLoS ONE. Given that the topic of this manuscript encompasses zooplankton ecology, I would like to suggest that it be handled by Dr. Jian-Shiou Hwang. This is the first submission of this manuscript, and albeit we do not oppose any reviewers, we prefer reviewers with ecological backgrounds.

This manuscript is a research article with two main goals. The first was to provide data on the feeding ecology of *L. conchilega* larvae, which is scarcely found in the literature. The second was to investigate whether the concept of fatty acid trophic markers (FATM), commonly applied in tracer studies with holoplankton of intermediate to high total lipid contents, can be applied in feeding studies with lipid-poor meroplanktonic larvae. In order to do so, we followed the assimilation of <sup>13</sup>C-enriched dietary lipid into the larvae.

To the best of our knowledge, this study is the first to present data on carbon assimilation and turnover and on fatty acid (FA) bioconversion in any meroplanktonic organism. The results show that *L. conchilega* larvae do not assimilate dietary FAs in an unmodified manner, which is one of the assumptions for FATMs. Instead they regulate (maintain) their lipid composition (regardless of that of their prey) via preferential assimilation and bioconversion of dietary FAs. This behavior limits the application of the FATM concept in feeding studies with *L. conchilega* larvae and, probably, with other lipid-poor meroplanktonic larvae. The combination of our results with published work reporting on the total lipid and wax ester contents of several zooplankton species led to our suggestion of the hypothesis that 3 different patterns in lipid homeostasis (regulation) are found among meroplanktonic and holoplanktonic organisms in relation to their lipid content.

I look forward to hearing from you at your earliest convenience regarding the submission of this manuscript.

Yours sincerely,

Rita M. Franco-Santos

[ritinhamf@hotmail.com](mailto:ritinhamf@hotmail.com)

Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research  
Am Handelshafen 12, 27570, Bremerhaven, Germany

## FA bioconversion and lipid homeostasis in *Lanice conchilega*

### 1 You are not always what you eat - Fatty acid bioconversion and lipid homeostasis in the 2 larvae of the sand mason worm *Lanice conchilega*

3  
4 Franco-Santos RM<sup>a,b,d,\*</sup>, Auel H<sup>b</sup>, Boersma M<sup>b,c</sup>, De Troch M<sup>d</sup>, Graeve M<sup>a</sup>, Meunier CL<sup>c</sup>, Niehoff  
5 B<sup>a</sup>

6  
7 <sup>a</sup> Alfred-Wegener-Institut, Helmholtz-Zentrum für Polar- und Meeresforschung, Am Handelshafen  
8 12, 27570, Bremerhaven, Germany

9 <sup>b</sup> University of Bremen, Marine Zoology, Bremen, Germany

10 <sup>c</sup> Alfred-Wegener-Institut, Helmholtz-Zentrum für Polar- und Meeresforschung, Biologische Anstalt  
11 Helgoland, Helgoland, Germany

12 <sup>d</sup> Ghent University, Marine Biology, Gent, Belgium

13  
14 \* Corresponding authors, Rita M. Franco-Santos and Barbara Niehoff

15 [rsantos@awi.de](mailto:rsantos@awi.de), [ritinhamf@hotmail.com](mailto:ritinhamf@hotmail.com) +61 0451 644 313

16 [barbara.niehoff@awi.de](mailto:barbara.niehoff@awi.de)

17

### 18 **Abstract**

19 The meroplanktonic larvae of benthic adults are an important seasonal component of the  
20 zooplankton in temperate coastal waters. The larvae of the reef-building polychaete *Lanice*  
21 *conchilega* can make up to 15% of the summer zooplankton biomass in the North Sea. Despite their  
22 importance for reef maintenance (which positively affects the benthic community), little is known  
23 of the trophic ecology of these meroplanktonic larvae. Qualitative and quantitative estimates of  
24 carbon (C) transfer between trophic levels and of FA-specific assimilation, biosynthesis, and  
25 bioconversion can be obtained by compound-specific stable isotope analysis of fatty acids (FA).  
26 The present work tested the hypothesis that the concept of fatty acid trophic markers (FATM),  
27 widely used for studies on holoplankton with intermediate to high lipid contents, is also applicable  
28 to lipid-poor organisms such as meroplanktonic larvae. The incorporation of isotopically-enriched  
29 dietary C by *L. conchilega* larvae was traced, and lipid assimilation did not follow FA-specific  
30 relative availabilities in the diet. Furthermore, FAs that were unavailable in the diet, such as 22:5(n-  
31 3), were recorded in *L. conchilega*, suggesting their bioconversion by the larvae. The results  
32 indicate that *L. conchilega* larvae preferentially assimilate certain FAs and regulate their FA  
33 composition (lipid homeostasis) independently of that of their diet. Their quasi-homeostatic

34 response to dietary FA availability could imply that the concept of FATM has limited application in  
35 lipid-poor organisms such as *L. conchilega* larvae.

36

37 **Keywords:** assimilation, bioconversion, compound-specific stable isotope analysis, fatty acids,  
38 larvae, polychaete

39

40

## 41 INTRODUCTION

42 The meroplanktonic larvae of macrobenthic organisms are an important seasonal component  
43 of the zooplankton in temperate coastal waters [1]. In the North Sea, the larvae of the sand mason  
44 worm *Lanice conchilega* (Pallas, 1766) can make up to 15% of the zooplankton biomass between  
45 July and September [2], when water temperatures are above 13°C. The reproductive peak of *L.*  
46 *conchilega* in the southern North Sea occurs in spring, and is followed by smaller peaks during the  
47 summer and autumn [3], though larval supply can vary between years [4]. The larvae of this  
48 polychaete evolve from a short planktonic to a benthic phase and, after a few days, to a second  
49 planktonic stage, the aulophore larvae [5]. At this stage larvae already present a (transparent) tube,  
50 and display morphological features of a juvenile [6]. They are able to feed in the water column,  
51 where they can remain up to 60 days before succeeding to a benthic stage [7].

52 The benthic, tube-dwelling adults of *L. conchilega* are non-selective suspension-deposit  
53 feeders [8, 9, 10], which derive their diet from the organic matter available in the water column and  
54 in the sediment. Common food items include diatoms, bacteria and microphytobenthos, and the  
55 species is able to switch between food sources [8, 10, 11]. *Lanice. conchilega* forms reefs capable of  
56 structuring the surrounding habitat and creating a complex and heterogeneous environment [12, 13].  
57 The reefs provide refuge against predation and physical and chemical stresses, attract a variety of  
58 other organisms, and serve as nursery and feeding grounds to several species, including fish and  
59 birds [14, 15, 16, 17, 18, 19]. *L. conchilega* is an ecosystem engineer whose presence favors species  
60 richness and faunal abundance and positively affects the benthic community [12, 13, 19, 20]. In  
61 comparison with adult organisms, few studies have been conducted on the ecology of *L. conchilega*

62 larvae, addressing topics such as juvenile settlement and buoyancy [4, 7, 21, 22]. The survival and  
63 development of these meroplanktonic larvae and, thus, the maintenance of *L. conchilega* reefs,  
64 depend, a.o., on their feeding success. To the best of our knowledge there are no studies published  
65 on the feeding ecology of the larvae, a gap which needs to be addressed if the larval stage of this  
66 species is to be better understood.

67 Information on the feeding ecology of an organism on short and long time scales and on the  
68 pathways of energy flow between prey and predator can be obtained from the analysis of fatty acid  
69 trophic markers (FATM) [23] in combination with the use of stable isotopes. FATMs are usually  
70 assimilated in a conservative manner by consumers [24, 25]. The FAs 16:1(n-7), 16:2(n-4), 16:3(n-  
71 4), and 20:5(n-3) (EPA) are characteristic of diatoms, whereas 18:4(n-3) and 22:6(n-3) (DHA) are  
72 considered dinoflagellate markers [8, 23, 26, 27]. Compound-specific stable isotope analysis (CSIA)  
73 is a powerful tool for tracing organic matter origin and fate and to investigate biochemical processes  
74 and patterns in individuals and ecosystems [e.g., 28, 29, 30]. When FA determination is coupled to  
75 measurements of isotopic composition, quantitative estimates can be obtained for the transfer of  
76 dietary carbon (C) from prey to predator and for FA-specific assimilation and potential biosynthesis  
77 and bioconversion [26, 27, 31, 32].

78 The present work had two objectives. The first was to start filling the gap of knowledge on  
79 the feeding ecology of *L. conchilega* larvae. Experiments investigating FA-related C assimilation  
80 have been conducted mostly for holoplanktonic organisms with intermediate to high lipid contents,  
81 whose FA profile tends to reflect that of the food items it ingests. Little is known, however, about  
82 how meroplanktonic organisms, which are usually lipid-poor, incorporate dietary FAs. It is possible  
83 that the lipid content and, thus, requirement, of an organism can dictate whether it will assimilate  
84 dietary FAs in an unmodified manner or not. We assume that the larvae of *L. conchilega* are lipid-  
85 poor, so our second objective was to test the hypothesis that the concept of FATM can also be  
86 applied in a feeding study with lipid-poor meroplanktonic larvae. In order to do so, the polychaete  
87 larvae were fed with a diatom culture previously labelled with  $^{13}\text{C}$  and the incorporation of

88 isotopically enriched FA-specific lipid C into the consumer was followed. This allowed for the  
89 recording of C assimilation and turnover in the organism, and also for the investigation of possible  
90 bioconversion pathways.

91

92

## 93 **METHOD**

### 94 **Field sampling**

95 Zooplankton samples for the feeding experiment were collected on June 6<sup>th</sup> 2016 with a 500  
96 µm mesh-size CalCOFI net, which was towed horizontally for 15 minutes at 5 m depth off the  
97 German island of Helgoland (54°11'N, 07°54'E), in the southern North Sea. Samples were taken to  
98 the laboratory and intact and active aulophore larvae of the polychaete *L. conchilega* were  
99 immediately sorted under an Olympus SZX16 stereoscopic microscope. A total of 390 larvae were  
100 sorted, 30 for determination of *in situ* body C and nitrogen (N) content, 150 for analyzing the *in situ*  
101 FA content and composition, and 210 for a feeding experiment. The planktonic diatom *Conticribra*  
102 *weissflogii* (Grunow) (Stachura-Suchoples and Williams, 2009) was used as prey in the feeding  
103 experiment with *L. conchilega* larvae.

104

### 105 **Algae culture**

106 Batch cultures were created daily for *C. weissflogii* for five consecutive days by diluting a  
107 stock solution with fresh *f/2* medium [after 33]. Diatoms were labelled with <sup>13</sup>C by adding <sup>13</sup>C-  
108 enriched sodium bicarbonate (NaH<sup>13</sup>CO<sub>3</sub>) to the medium at a concentration of 4 mg L<sup>-1</sup>. Cultures  
109 were kept in constant light inside a temperature-controlled room at 18°C, and stirred twice a day in  
110 order to keep cells suspended. Algae were grown for five days and then used as food suspension  
111 (exponential growth phase) for the polychaete larvae during the experiment. *C. weissflogii* cultures  
112 were sampled daily during the experiment for determination of cell C, N, and FA content and of  
113 FA-specific <sup>13</sup>C isotopic enrichment. This was done by filtering subsamples of known cell

114 concentrations through pre-combusted (500°C for 24 h) Whatman GF/F filters (0.7 µm pore size, 25  
115 mm diameter). Filters for determination of prey C and N content were dried at 60°C for 48 h, folded  
116 inside aluminum foil, and stored in a desiccator until analysis. Samples for FA analyses were placed  
117 into pre-combusted lipid vials and stored at -80°C. The remaining volume of the cultures was then  
118 used to feed polychaete larvae. Cell densities were determined with a BD Accuri C6 Flow  
119 Cytometer.

120

## 121 **Experimental design**

122 Feeding experiments with *L. conchilega* were initiated immediately after sorting of the  
123 larvae. Individuals were kept for five days in triplicate 500 mL glass beakers fitted with a 300 µm  
124 meshed-bottom cylinder (140 ind L<sup>-1</sup>). The diatom suspension was provided on a daily basis at a  
125 concentration of 8000 cells mL<sup>-1</sup>. The polychaete larvae were kept in a dark temperature-controlled  
126 room at the temperature recorded when they were sampled in the field, i.e., 13.5 ± 0.3°C. The  
127 beakers were gently stirred three times a day for food resuspension in the water. A partial water  
128 exchange (66%) was performed daily and followed by the addition of new algae culture, with a  
129 final diatom concentration > 8000 cells mL<sup>-1</sup>. At the end of the experiment the larvae were sampled  
130 from each replicate for body C and N content determination and for analysis of FA composition and  
131 FA-specific <sup>13</sup>C isotopic enrichment (10 and 50 individuals per sample, respectively). Individuals  
132 (inside their tubes) were gently washed in distilled water, placed into pre-weighed tin capsules (5\*9  
133 mm, IVA Analysentechnik) for body C and N content determination or into pre-combusted lipid  
134 vials for FA analyses, and stored at -80°C until further analysis. Tin capsules with larvae were dried  
135 at 60°C for 48 h, weighed with an ultra micro-balance (detection limit: 0.1 µg; XP6U Ultra Micro  
136 Balance, Mettler Toledo, Germany) and stored in a desiccator until analysis.

137

## 138 **Carbon, nitrogen and fatty acid content analyses**



139 The C and N contents of all larvae and algae samples were measured with an elemental  
140 analyzer (detection limit: 2  $\mu\text{g C}$  / 0.5  $\mu\text{g N}$ ; maximum error:  $\pm 3\%$ , Euro EA 3000, EuroVector  
141 S.P.A., Milan, Italy) using acetanilide as a standard.

142 Lipids were extracted and FAs identified as described by [26]. Samples were homogenized  
143 in a dichloromethane:methanol (2:1, v:v) solution, from which total lipids were extracted. A known  
144 amount of an internal standard, the tricosanoic acid methyl ester (23:0), was added to each sample.  
145 Potassium chloride (0.88% solution) was added to create a biphasic system and aid in extraction.  
146 Lipid extracts were transesterified by heating samples with 3% sulfuric acid in methanol at 80°C  
147 under nitrogen atmosphere for 4 hours. Fatty acid methyl esters (FAMES) were then extracted with  
148 cyclohexane, and determined and quantified with a gas chromatograph (HP 6890 N, Agilent  
149 Technologies Deutschland GmbH & Co. KG) equipped with a 60m  $\times$  0.25mm i.d. wall-coated open  
150 tubular capillary column (film thickness: 0.25  $\mu\text{m}$ ; liquid phase: DB-FFAP), a split/splitless injector  
151 (250°C) and a flame ionization detector (280°C). The chromatograms generated were evaluated  
152 with the ChemStation software from Agilent. The A:B(n-X) shorthand notation was used to refer to  
153 FAs, where A is the number of carbon atoms, B is the number of double bonds, and (n-X) gives the  
154 position of the double bond closest to the terminal methyl group. Total lipid contents of larvae and  
155 algae cultures were calculated by adding the mass of all FAs.

156

### 157 **Compound specific stable isotope analysis (CSIA)**

158 The FA-specific stable isotope composition of carbon in FAMES extracted was obtained  
159 according to [26] with a Thermo gas chromatography combustion-isotope ratio mass spectrometry  
160 (GC-c-IRMS) system, equipped with a Trace GC Ultra gas chromatograph, a GC Isolink, and a  
161 Delta V Plus isotope ratio mass spectrometer connected via a Conflo IV interface (Thermo  
162 Scientific Corporation, Bremen, Germany). The chromatograms containing peak areas and C  
163 isotope ratios were analyzed with the IRMS software Isodat 3.0. The 14:0 and 18:0 FAME  
164 reference standards (Iowa University) with known  $\delta$ -values were used for further calculations.

165 The equations used by [26] to calculate carbon assimilation were also applied in the present  
166 study, and were:

167

168 Isotopic ratios of FAs

$$169 \quad \delta^{13}\text{C} (\text{‰}) = \left[ \left( \frac{R_{\text{sample}}}{R_{\text{standard}}} \right) - 1 \right] * 1000, \quad (1)$$

170 where  $R_{\text{sample}}$  and  $R_{\text{standard}}$  are the ratio of  $^{13}\text{C}/^{12}\text{C}$  in the sample and reference standard, respectively;

171

172 Atom percent (atom%), which are converted  $\delta$ -values and express isotope data in terms of isotope  
173 concentrations to inform on the  $^{13}\text{C}$  enrichment in each FA

$$174 \quad \text{atom}\% = \left( \frac{R_{\text{sample}}}{(R_{\text{sample}} + 1)} \right) * 100,$$

175 (2)

176 where R is the ratio  $^{13}\text{C}/^{12}\text{C}$  in the sample;

177

178 Atom percent excess (APE), which considers the isotopic enrichment in experimental larvae (in  
179 comparison to field individuals) due to the assimilation of algae enriched in  $^{13}\text{C}$

$$180 \quad \text{APE} = \text{atom}\%_{\text{experimental polychaetes}} - \text{atom}\%_{\text{field polychaetes}}$$

181 (3)

182

183 Carbon mass of each FA ( $C_{\text{mass}}$ , in  $\mu\text{g C ind}^{-1}$ ), which divides the FA mass by the FAME mass

$$184 \quad C_{\text{mass}} = \frac{\{(\text{atom}\% / 100 * A_{13\text{C}}) + [(100 - \text{atom}\%) / 100 * A_{12\text{C}}]\} * C_{\text{FA}} * \text{FA}_{\text{mass}}}{\{[(\text{atom}\% / 100 * A_{13\text{C}}) + ((100 - \text{atom}\%) / 100 * A_{12\text{C}})] * C_{\text{FA}}\} + A_{12\text{C}} + (\text{H}_{\text{FAME}} * A_{\text{H}}) + (\text{O}_{\text{FAME}} * A_{\text{O}})}, \quad (4)$$

185 where  $A_{12\text{C}}$ ,  $A_{13\text{C}}$ ,  $A_{\text{H}}$ , and  $A_{\text{O}}$  are the atomic masses of  $^{12}\text{C}$ ,  $^{13}\text{C}$ , H and O, respectively,  $C_{\text{FA}}$  is the  
186 number of carbon atoms in the FA,  $\text{H}_{\text{FAME}}$  and  $\text{O}_{\text{FAME}}$  are the number of hydrogen and oxygen  
187 atoms in the FAME, and  $\text{FA}_{\text{mass}}$  is the mass (in  $\mu\text{g ind}^{-1}$ ) of the FA;

188

189 Proportion of carbon assimilated (PA)

$$190 \quad PA = \frac{APE}{L}, \quad (5)$$

191 where L is the average labeling (in atom%) in all algae FAs but 18:0 (L = 3.62%). The averaging is  
192 to account for the elongation and/or desaturation of small amounts of dietary FAs when they are  
193 assimilated by the larvae [23], and the exclusion of 18:0 is due to its poor labelling in comparison to  
194 other FAs in prey cultures [26, 34]. When calculating the PA in the larvae FA 18:0, L was the  
195 average of labeling in algae FA 18:0 ( $L_{FA\ 18:0} = 1.87\%$ );

196

197 Carbon assimilation ( $C_{assim}$ , in  $\mu\text{g C ind}^{-1}$ ), calculated for each FA

$$198 \quad C_{assim} = C_{mass} * PA \quad (6)$$

199 Total C assimilation was obtained by summing the  $C_{assim}$  values for all FAs. Assimilation rates were  
200 obtained by dividing  $C_{assim}$  by the number of experimental days when sample was collected;

201

202 Carbon turnover rates ( $C_T$  in  $\% \text{ day}^{-1}$ )

$$203 \quad C_T = \frac{\frac{C_{assim}(t)}{C_{mass}(t)}}{\Delta t}, \quad (7)$$

204 where (t) indicates the specific sampling time for which the values of  $C_{assim}$  and  $C_{mass}$  should be used  
205 and  $\Delta t$  the number of experimental days when sample was collected.

206

## 207 **Statistical analysis**

208 Differences in atom% for the different batches of the diatom culture were investigated with  
209 one-way analysis of variance (ANOVA), and the origin of differences was identified by applying  
210 the Tukey HSD (Honestly Significant Difference) post-hoc test with a 95% confidence limit. Prior  
211 to the ANOVAs, the data were tested for normality and homogeneity of variances with Shapiro-  
212 Wilk and Bartlett tests, respectively. Differences between *in situ* and laboratory-fed polychaete

213 larvae in dry mass (DM), C, N, and total lipid contents and C:N molar ratio were investigated with  
214 t-tests. All univariate analyses were performed using R ver. 3.4.4 [35].

215 Multivariate analysis of FA composition for *L. conchilega* individuals was performed with a  
216 dendrogram, which clustered individuals with similar FA profiles (based on their group average  
217 linkage clustering). The dendrogram was generated from a Bray-Curtis similarity matrix obtained  
218 from absolute and percentage (with logit transformation, as suggested by 36) data on FAs with  
219 relative content > 1% total fatty acids (TFA). All multivariate analyses were performed with  
220 PRIMER 7.0 software [37].

221

222

## 223 RESULTS

### 224 Algal composition and labelling

225 The average C and N contents of the *C. weissflogii* cultures were 580 and 120  $\mu\text{g L}^{-1}$ ,  
226 respectively, resulting in an average molar C:N ratio of 5.8 (Table 1). The average lipid content was  
227 120  $\mu\text{g L}^{-1}$  (Table 1). The FA profile of the diatom (Table 1, Fig. 1) was dominated by  
228 polyunsaturated fatty acids (PUFAs), which comprised 61% of total fatty acids (TFA). Saturated  
229 (SFA) and monounsaturated (MUFA) fatty acids were present in similar amounts, 22 and 17%  
230 TFA, respectively. The most important FAs in the *C. weissflogii* batches were 20:5(n-3)  
231 (eicosapentaenoic acid, EPA), 16:3(n-4), 16:1(n-7), and 16:0 (22, 19, 15 and 13% TFA,  
232 respectively). The FA 22:6(n-3) (docosahexaenoic acid, DHA) was present in a smaller amount,  
233 i.e., 5% of TFA.

234 Fig 1. Fatty acid absolute and relative concentrations in diatoms (*Conticribra weissflogii*)  
235 and in *Lanice conchilega* larvae.

236 Absolute and relative (concentrations are given in  $\text{ng C ind}^{-1}$  and in % total FA (%TFA),  
237 respectively. Data for larvae are differentiated between organisms sampled from the field and  
238 sampled after 5 days of feeding on the diatom.

239 Table 1 Dry mass (DM), carbon (C), nitrogen (N) and total lipid contents, molar C:N ratio,  
 240 and FA-specific absolute and relative composition of *Lanice conchilega* larvae and its diet, the  
 241 diatom *Conticribra weissflogii*. Total lipid C content is given by the  $\Sigma$ FAs term. A distinction is  
 242 made between the internal (non-labelled, from *in situ* origin) and the internal + dietary (labelled,  
 243 assimilated from the diet) FA contents in *L. conchilega* at the end of the feeding experiment. DM is  
 244 reported in  $\mu\text{g ind}^{-1}$ . C, N, and lipid contents are reported in  $\text{ng ind}^{-1}$  for larvae samples and in  $\mu\text{g L}^{-1}$   
 245  $^1$  and in  $\text{pg cell}^{-1}$  (in parenthesis) for algae samples. Absolute concentration of FAs are reported in  
 246  $\text{ng C ind}^{-1}$  and in  $\text{ng C L}^{-1}$  for larvae and algae samples, respectively; and relative concentration of  
 247 FAs are shown in % total FA (TFA, in parenthesis) for all samples. Only FAs > 1% TFA are  
 248 indicated. Values are means  $\pm$  standard deviation from samples and from the sum of saturated  
 249 (SFA), monounsaturated (MUFA) and polyunsaturated (PUFA) fatty acids (n = 3 or 4). – : not  
 250 detected.

	<i>L. conchilega</i>				<i>C. weissflogii</i>
	<i>in situ</i>	Experiment			
		Internal and dietary C	Internal C		
DM	55 $\pm$ 6	52 $\pm$ 13			
C content	7668 $\pm$ 2858	8511 $\pm$ 966		577 $\pm$ 44 (72 $\pm$ 6)	
N content	1929 $\pm$ 628	2170 $\pm$ 270		117 $\pm$ 11 (15 $\pm$ 1)	
Molar C:N ratio	4.6 $\pm$ 0.3	4.6 $\pm$ 0.1		5.8 $\pm$ 0.5	
Total lipid content	632 $\pm$ 116	845 $\pm$ 11		116 $\pm$ 13 (15 $\pm$ 2)	
$\Sigma$ FAs	465 $\pm$ 85	622 $\pm$ 8	423 $\pm$ 26	84920 $\pm$ 9311	
14:0	28 $\pm$ 11 (6 $\pm$ 1)	21 $\pm$ 1 (4 $\pm$ 0)	16 $\pm$ 2	5214 $\pm$ 1086 (6 $\pm$ 1)	
16:0	91 $\pm$ 18 (20 $\pm$ 0)	132 $\pm$ 4 (22 $\pm$ 1)	88 $\pm$ 8	11166 $\pm$ 1591 (13 $\pm$ 0)	
16:1n7	20 $\pm$ 5 (4 $\pm$ 0)	29 $\pm$ 2 (5 $\pm$ 0)	16 $\pm$ 0	12155 $\pm$ 2206 (15 $\pm$ 1)	
16:1n5	–	–	–	930 $\pm$ 149 (1 $\pm$ 0)	
16:2n4	–	8 $\pm$ 1 (1.3 $\pm$ 0)	4 $\pm$ 0	6860 $\pm$ 736 (8 $\pm$ 0)	
16:3n4	–	4 $\pm$ 1 (0.7 $\pm$ 0)	2 $\pm$ 0	16110 $\pm$ 1781 (19 $\pm$ 2)	
18:0	49 $\pm$ 4 (11 $\pm$ 1)	57 $\pm$ 3 (9 $\pm$ 1)	37 $\pm$ 6	475 $\pm$ 85 (1 $\pm$ 0)	
18:1n11	18 $\pm$ 3 (4 $\pm$ 0)	31 $\pm$ 2 (5 $\pm$ 0)	26 $\pm$ 3	–	

18:1n9	6 ± 3 (1.2 ± 0)	–	1 ± 2	348 ± 46 (0.4 ± 0)
18:1n7	26 ± 5 (6 ± 0)	33 ± 1 (5 ± 0)	24 ± 1	474 ± 81 (1 ± 0)
18:2n6	–	–	–	862 ± 52 (1 ± 0)
18:4n3	–	3 ± 0 (0.5 ± 0)	1 ± 1	3833 ± 995 (4 ± 1)
20:1n11	11 ± 2 (2 ± 0)	17 ± 1 (3 ± 0)	15 ± 2	–
20:5n3	103 ± 12 (22 ± 1)	119 ± 2 (19 ± 0)	77 ± 3	19363 ± 1661 (22 ± 1)
22:5n3	73 ± 8 (15 ± 1)	99 ± 5 (15 ± 1)	74 ± 6	–
22:6n3	31 ± 4 (6 ± 0)	53 ± 1 (8 ± 0)	32 ± 1	4839 ± 216 (5 ± 1)
ΣSFA	173 ± 34 (38 ± 1)	219 ± 8 (36 ± 1)	148 ± 15	18140 ± 2743 (22 ± 1)
ΣMUFA	83 ± 22 (18 ± 1)	112 ± 1 (18 ± 0)	81 ± 5	13907 ± 2329 (17 ± 1)
ΣPUFA	209 ± 29 (44 ± 2)	291 ± 8 (45 ± 1)	194 ± 8	52873 ± 4641 (61 ± 2)

251

252 The <sup>13</sup>C isotopic enrichment in all FAs (but 18:0) varied from 2.88 ± 0.21 atom% to 4.00 ±  
253 0.19 atom% between the different batches, averaging 3.62 ± 0.50 atom% for all days (Table 2, Fig.  
254 2). For the FA 18:0, isotopic enrichment averaged 1.87 ± 0.30 atom% between the different batches  
255 (Table 2, Fig. 2). Labelling was significantly different between batches (ANOVA, F = 21.66, df =  
256 3,67, p = \*\*\*), being higher on days 1 and 5 than on day 4 and on day 2 (Tukey, p = \* and p = \*\*\*,  
257 respectively), and higher on day 4 than on day 2 (Tukey, p = \*\*\*). The major FAs cited above (or  
258 PUFAs in general) showed the highest labelling values within the FAs (Table 2).

259 Fig 2: <sup>13</sup>C isotopic enrichment of *Conticribra weissflogii* batch cultures fed to polychaete  
260 larvae during the experiment.

261 <sup>13</sup>C isotopic enrichment is presented in atom%. Straight lines represent atom% values for the fatty  
262 acid (FA) 18:0, whereas boxplots and open circles represent average atom% values from all the  
263 other FAs.

264 Table 2 Total fatty acid (TFA) and FA-specific (> 1% TFA) <sup>13</sup>C isotopic labelling (APE)  
265 and carbon assimilation (C<sub>assim</sub>, as ng C ind<sup>-1</sup> and as % total C assimilated, in parenthesis) and  
266 turnover rate (C<sub>turn</sub>, as % day<sup>-1</sup>) for *Lanice conchilega*. <sup>13</sup>C isotopic labelling is also shown for  
267 *Conticribra weissflogii* (atom% - the atom% of TFA does not include values for the FA 18:0). SFA:

268 saturated FA; MUFA: monounsaturated FA; PUFA: polyunsaturated FA. Values are mean  $\pm$   
 269 standard deviation (n = 3 or 4 each). – : not detected.

FA	<i>C. weissflogii</i>		<i>L. conchilega</i>	
	atom%	APE	C <sub>assim</sub>	C <sub>turn</sub>
TFA	3.62 $\pm$ 0.50	1.16 $\pm$ 0.48	199 $\pm$ 21	6 $\pm$ 1
14:0	3.72 $\pm$ 0.53	0.93 $\pm$ 0.13	6 $\pm$ 1 (3 $\pm$ 0)	5 $\pm$ 1
16:0	3.62 $\pm$ 0.52	1.21 $\pm$ 0.14	44 $\pm$ 4 (22 $\pm$ 0)	7 $\pm$ 1
16:1n7	3.72 $\pm$ 0.54	1.67 $\pm$ 0.09	14 $\pm$ 1 (7 $\pm$ 0)	9 $\pm$ 0
16:1n5	3.48 $\pm$ 0.52	-	-	-
16:2n4	3.73 $\pm$ 0.52	1.94 $\pm$ 0.02	4 $\pm$ 0 (2 $\pm$ 0)	11 $\pm$ 0
16:3n4	3.73 $\pm$ 0.52	2.01 $\pm$ 0.06	2 $\pm$ 1 (1 $\pm$ 0)	11 $\pm$ 0
18:0	1.87 $\pm$ 0.30	0.66 $\pm$ 0.11	20 $\pm$ 2 (10 $\pm$ 0)	7 $\pm$ 1
18:1n11	-	0.67 $\pm$ 0.10	6 $\pm$ 1 (3 $\pm$ 0)	4 $\pm$ 1
18:1n9	3.19 $\pm$ 0.55	-	-	-
18:1n7	3.26 $\pm$ 0.45	0.97 $\pm$ 0.11	9 $\pm$ 1 (4 $\pm$ 0)	5 $\pm$ 1
18:2n6	3.68 $\pm$ 0.51	-	-	-
18:4n3	3.76 $\pm$ 0.54	1.91 $\pm$ 0.09	2 $\pm$ 0 (1 $\pm$ 0)	11 $\pm$ 0
20:1n11	-	0.49 $\pm$ 0.08	2 $\pm$ 0 (1 $\pm$ 0)	3 $\pm$ 0
20:5n3	3.74 $\pm$ 0.53	1.26 $\pm$ 0.12	41 $\pm$ 4 (21 $\pm$ 0)	7 $\pm$ 1
22:5n3	-	0.94 $\pm$ 0.11	25 $\pm$ 2 (13 $\pm$ 0)	5 $\pm$ 1
22:6n3	3.76 $\pm$ 0.53	1.42 $\pm$ 0.10	21 $\pm$ 2 (11 $\pm$ 0)	8 $\pm$ 1
$\Sigma$ SFA	3.30 $\pm$ 0.90	0.95 $\pm$ 0.24	72 $\pm$ 7 (36 $\pm$ 0)	7 $\pm$ 1
$\Sigma$ MUFA	3.41 $\pm$ 0.51	0.90 $\pm$ 0.49	31 $\pm$ 3 (15 $\pm$ 0)	5 $\pm$ 1
$\Sigma$ PUFA	3.70 $\pm$ 0.45	1.46 $\pm$ 0.44	97 $\pm$ 11 (49 $\pm$ 0)	7 $\pm$ 1

270

### 271 Larval composition and labelling

272 Approximately 85% of the *L. conchilega* larvae survived at the end of the experiment. The  
 273 average *in situ* DM and C and N contents of larvae were 55  $\mu\text{g ind}^{-1}$  and 7668 and 1929  $\text{ng ind}^{-1}$ ,  
 274 respectively, and the average values at end of the experiment were 52  $\mu\text{g ind}^{-1}$  and 8511 and 2170

275 ng ind<sup>-1</sup>, respectively (Table 1). Changes in DM and C, N, and lipid contents between larvae  
276 sampled from the field and after the experiment were not significant, indicating that the organisms  
277 did not lose or gain weight during the five-day incubation. The molar C:N ratio of larvae remained  
278 the same throughout the experiment at an average of 4.6. The average lipid content was similar  
279 between field (632 ng ind<sup>-1</sup>) and experimental (845 ng ind<sup>-1</sup>) individuals (Table 1). The lipid C  
280 content accounted for approximately 7% of the total C content of larvae sampled from the field and  
281 at the end of the experiment. Total lipid content comprised 1.2 ± 0.3% DM for *in situ* larvae and 1.7  
282 ± 0.4% DM for larvae at the end of the experiment.

283 The major FAs in *L. conchilega* larvae were EPA, 16:0 and 22:5n3, which corresponded to  
284 22, 20, and 15% TFA at the beginning and to 19, 22, and 15% TFA at the end of the experiment,  
285 respectively (Table 1). About a quarter of the FAs identified were only present in small amounts  
286 (<1% TFA, Fig. 1). The relative FA composition of *L. conchilega* larvae fed with diatoms remained  
287 similar to that of the individuals collected *in situ* (Table 1, Fig. 1), as conveyed by the 95.5%  
288 similarity between samples shown in the Bray-Curtis dendrogram (Fig. 3a). A similar pattern was  
289 observed for the absolute FA composition of the larvae (Table 1, Fig. 1), except that one of the *in*  
290 *situ* replicates was grouped with the experimental ones at 91% similarity, and these were found to  
291 be 82% similar to the other *in situ* replicates (Fig. 3b). These results further reinforce the above  
292 mentioned statistically non-significant differences in lipid C between *in situ* larvae and those fed  
293 during the experiment. In general, changes (i.e., increase or decrease) in the absolute concentration  
294 of a FA matched the change in its relative concentration (Fig. 1). The only exception to this pattern  
295 were the FAs 18:0, 18:1(n-7) and EPA, whose relative concentrations in the larvae decreased  
296 despite an increase in their absolute concentrations (Fig. 1).

297 Fig 3: Dendrograms generated with Bray-Curtis similarity matrices comparing fatty acid  
298 profiles of *Lanice conchilega* larvae.

299 Similarity matrices present data on the FA profile of *Lanice conchilega* larvae sampled from the  
300 field and after 5 days of feeding on the diatom *Conticribra weissflogii*. Only FAs with relative



301 content > 1% TFA were used to generate the similarity matrices. a) logit transformed relative values  
302 for FA composition (% TFA). b) absolute (ng C ind<sup>-1</sup>) values for FA composition.

303 The average <sup>13</sup>C enrichment of *L. conchilega* FAs was of 1.16 ± 0.48 APE, ranging from  
304 0.49 APE in FA 20:1(n-11) to 2.01 APE for FA 16:3(n-4) (Table 2). The FAs 18:1(n-11), 20:1(n-  
305 11), and 22:5(n-3) had the lowest labelling values (together with 18:0) in *L. conchilega*, whereas the  
306 FAs 16:2(n-4), 16:3(n-4), and 16:1(n-7) (diatom fatty acid trophic markers, FATM) had the highest  
307 APE (together with 18:4(n-3)).

308

### 309 **Origin of FAs in the larvae**

310 The labelling of FAs with <sup>13</sup>C enabled for the discrimination of the source of each FA-  
311 specific lipid C content: internal (Table 1), for FAs already present in the *in situ* larvae, and dietary  
312 (Table 2), for FAs assimilated via feeding on the labelled diatoms. The absolute content of internal  
313 FAs at the end of the experiment remained the same (~98%) as recorded from *in situ* larvae for two  
314 of the three replicates, and on average represented 92% of the *in situ* internal FA content (Table 1).  
315 Given the similarity of C content of internal FA between the beginning and end of the experiment  
316 the FA-specific differences are not considerable. The internal FA which showed the greatest  
317 decrease in terms of C content during the experiment was EPA, which fell from 22% TFA in *in situ*  
318 larvae to 18% of the total internal FA in diatom-fed larvae (Table 1). SFAs also showed slight  
319 decreases in C content between the beginning and end of the experiment.

320

### 321 **Lipid C assimilation and turnover in larvae**

322 Total assimilation of C into lipids in polychaete larvae at the end of the experiment was 200  
323 ng lipid C ind<sup>-1</sup> (Table 2). The relative assimilation of lipid C into FAs (Table 2) followed the  
324 relative composition of internal FAs (Table 1). Approximately half of the lipid C was assimilated  
325 into PUFAs, a third into SFAs, and only a sixth into MUFAs (Table 2, Fig. 4). The majority of the  
326 total lipid C was assimilated into 16:0 and EPA (22 and 21%, respectively), followed by 22:5(n-3),

327 DHA, and 18:0 (13, 11 and 10%, respectively). It is worth noting that the incorporation of dietary C  
328 into specific FAs did not follow their relative availability in the diet. The absolute concentration of  
329 the FAs 17:0, 18:1(n-11), 20:1(n-11), and 22:5(n-3) increased in *L. conchilega*, even though they  
330 were absent from the diatom cultures fed to the larvae (Table 2). The absolute concentration of the  
331 FAs 18:0 and DHA also increased in the diatom-fed larvae (Table 2), even though they were  
332 available in low amounts in the diet (1 and 5% TFA in *C. weissflogii*, respectively). Furthermore,  
333 the FAs i-15:0, 16:1(n-5), 18:2(n-6), 18:3(n-6), and 20:4(n-3) were only identified in *C. weissflogii*  
334 (Fig. 1) and were either not assimilated or bioconverted by the polychaete larvae. C assimilation in  
335 FATM was low in comparison to that into other FAs, but amounted to the highest C turnover rates  
336 (Table 2).

337 Fig 4: Carbon assimilation and turnover in *Lanice conchilega* larvae after the feeding  
338 experiment.

339 Carbon assimilation and turnover are given in ng C ind<sup>-1</sup> and in %, respectively. Values are shown  
340 for the sum of all fatty acids (TFA), saturated (SFA), monounsaturated (MUFA), and  
341 polyunsaturated (PUFA) fatty acids; and for specific FAs.

342 The average lipid C turnover rate was of 6% day<sup>-1</sup>, but varied between 3 and 11% day<sup>-1</sup> for  
343 different FAs (Table 2). C turnover of SFA and PUFA was similar, 7% day<sup>-1</sup>, and slightly higher  
344 than that of MUFAs, 5% day<sup>-1</sup>. Overall, 32% of all lipid C was replaced with labelled dietary lipid  
345 C by the end of the experiment (Fig. 4).

346

347

## 348 **DISCUSSION**

349 The present study is the first to our knowledge to present data on C assimilation and  
350 turnover and on FA bioconversion in the larvae of the polychaete *L. conchilega*, and in any other  
351 meroplanktonic larvae for that matter. Although these were recorded from a short-term feeding  
352 experiment (5 days), they indicate how quickly polychaete larvae can assimilate dietary material.

353 These lipid-poor larvae seem to be able to regulate their lipid composition (homeostasis) by  
354 preferentially assimilating certain FAs and through bioconversion of dietary and internal FAs.

355

### 356 **Experimental conditions**

357 The active uptake of dietary C during the experiment was sufficient to compensate for the  
358 larval metabolic costs of living during the 5 days. This indicates that diatoms are a nutritious source  
359 of food for *L. conchilega* larvae, which readily accept this alga and utilize it as metabolically  
360 necessary. The lack of literature on feeding studies with meroplanktonic larvae prompted us to look  
361 for guidelines on food concentrations from laboratory experiments with small planktonic copepods  
362 [e.g., 38], so it is possible that individuals would have shown higher growth if they had received  
363 more food.

364 A common value of 40% DM is usually attributed for C content in marine organisms [e.g.,  
365 39]. The values obtained in the present study were almost 3 times lower, with larval C content at  
366 approximately 15% DM. As previously stated in the methodology, larvae were sampled while still  
367 inside their tubes. An attempt was made to remove the larvae, but we observed that it was not  
368 possible to do so without damaging the individuals or without the application of a sedative. The  
369 larval tube is composed of a thin organic layer, to which sand and shell fragments are attached at  
370 later stages [21]. This may have influenced the sample C and N contents and weight, and resulted on  
371 low relative values for C content in relation to DM.

372

### 373 **Lipid composition, homeostasis, and bioconversion in *L. conchilega* larvae**

374 The low total lipid content recorded for *in situ* larvae in the present study (1.2% DM) is  
375 similar to that reported for field and starving crab zoeae and bivalve larvae [40, 41, 42]. *Lanice*  
376 *conchilega* larvae thus appear to be lipid-poor organisms.

377 The relative concentration of single FAs and of SFAs, MUFAs, and PUFAs was preserved  
378 in the larvae during the feeding period. This occurred despite the provision of, and successful

379 feeding upon, a diet with a different FA composition. These results indicate that *L. conchilega*  
380 larvae can selectively accumulate FAs and/or control its FA metabolism and, thus, regulate lipid  
381 composition. A quasi-homeostatic response to variation in FA availability in the food source has  
382 also been shown for other zooplankton (daphnids) by [43]. The preferential retention of a FA over  
383 another can be easily identified if their relative proportions in the diet are similar, but their  
384 assimilation by the larvae differs. In the present study this is the case for the pairs of FAs 16:0 +  
385 16:1(n-7), EPA + 16:3(n-4), and DHA + 18:4(n-3), for which the first FA of each pair was  
386 assimilated by *L. conchilega* in quantities that were 3-20x higher than those of the second FA,  
387 indicating their preferential accumulation. EPA and DHA have an important function for marine  
388 consumers as they fuel reproductive processes and neural function and regulate cell membrane  
389 fluidity [44, 45, 46]. It has been observed that adult *L. conchilega* accumulate (n-3) FAs to store  
390 energy during gametogenesis [8], but larvae probably use EPA and DHA for growth, as tissue  
391 hormones (eicosanoids) are produced from PUFA [47]. The FA 16:0, on the other hand, can be  
392 elongated and desaturated into other necessary FAs.

393         There are three different metabolic sources from which the FAs composing the lipid reserves  
394 of a predator can be derived. FAs can be (1) assimilated unmodified from the food (dietary origin),  
395 (2) synthesized *de novo* (dietary or internal origin), or (3) bioconverted (synthesized via chain  
396 elongation and desaturation) from another FA of dietary or internal origin [48]. FA bioconversion  
397 has more often been reported for crustaceans [23, 31] than for polychaetes [49]. In the present study  
398 we report the bioconversion of the FAs 17:0, 18:1(n-11), 20:1(n-11), and 22:5(n-3). It should be  
399 noted that the synthesis of odd-chained FAs is uncommon in consumers, and that the FA 17:0 is  
400 usually considered a bacterial FATM. The presence of this FA in the lipid profile of *L. conchilega*  
401 larvae indicates that either the organisms are able to synthesize odd-chained FAs or that bacteria  
402 were present in small amounts in the experimental units and were ingested by the larvae.  
403 Bioconversion occurred for both endogenously-derived FAs, which increased in absolute content  
404 during the experiment, and for dietary-derived FAs, which were enriched in <sup>13</sup>C. FA elongation and

405 desaturation involve reactions catalyzed by enzymes and are subject to isotopic fractionation [50],  
 406 which could explain why the  $^{13}\text{C}$  enrichment was lower in the bioconverted FAs. The FA 22:5(n-3)  
 407 was bioconverted at high quantities, and made up 13% of all C assimilated by the larvae. Although  
 408 marine invertebrates are not able to biosynthesize *de novo* the FAs 18:2(n-6) and 18:3(n-3), they  
 409 have a limited ability to convert them into PUFAs via chain elongation and desaturation [23, 51]. In  
 410 the present study neither the diet nor the larvae contained 18:3(n-3), so 22:5(n-3) must have been  
 411 bioconverted from EPA, as has been suggested for brittle stars [52].

412

### 413 **Lipid homeostasis in lipid-poor and lipid-rich marine planktonic organisms**

414 The ability to regulate lipid composition does not come without an energetic cost. FA  
 415 homeostasis can be easy to maintain when an organism feeds upon a prey of similar lipid  
 416 composition, but will require that energy be allocated to bioconversion if prey have a different  
 417 biochemical make-up. Furthermore, the way in which an individual utilizes its assimilated FAs is  
 418 life-stage specific [23]. Based on our results and on the available literature on lipid content and  
 419 assimilation of mero- and holoplanktonic marine species [26, 27, 53, 54, other references shown in  
 420 Table 3, Franco-Santos et al., unpublished data], we put forward the hypothesis that 3 different  
 421 patterns in lipid homeostasis can be found among lipid-poor and lipid-rich planktonic organisms.

422 Table 3 Total lipid content (TLC, in % dry mass) and wax ester content (WE, in % total  
 423 lipid content) of several meroplanktonic and holoplanktonic species.

SPECIES	LOCATION	TLC	WE	REFERENCE
<b>Meroplankton (lipid-poor)</b>				
<b>Bivalvia</b>				
<i>Crassostrea gigas</i> larvae	Bay of Archacon, France	2-8	-	[42]
<i>Teredo navalis</i> larvae (lab-reared)	Great Harbor, MA, USA	2-4	-	[54]
<i>Bankia gouldi</i> larvae (lab-reared)	Pivers Island, NC, USA	3-8	-	[54]
<i>Crassadoma gigantean</i> larvae (lab-spawned)	?	4-8	-	[55]
<b>Bryozoa</b>				
<i>Celleporella hyalina</i> larvae (lab-spawned)	Menai Strait, UK	6-9	-	[56]
<b>Cirripedia</b>				
<i>Balanus balanoides</i>	Menai Strait, UK	13-15	-	[57]
<b>Cephalopoda</b>				
<i>Octopus vulgaris</i> hatchlings	Ría de Vigo, Spain	< 15	< 2	[58]
<b>Decapoda</b>				
<i>Campylonotus vagans</i> zoeae (lab-reared)	Beagle Channel, Argentina	7-9	-	[59]

<i>Carcinus maenas</i> zoeae	Helgoland	1-9	-	[41]
<i>Nephrops norvegicus</i> larvae	Mediterranean and Irish Seas	6-8	-	[60]
<i>Panulirus cygnus</i> larvae	Western Australia	9-13	-	[61]
<b>Vertebrata</b>				
<i>Pleuragramma antarcticum</i> (larvae)	Antarctic Peninsula	12	4	[62]
<i>Solea senegalensis</i> (lab-spawned eggs/larvae)	Bay of Cadiz, Spain	11-12	-	[63]
<b>Holoplankton (lipid-poor)</b>				
<b>Amphipoda</b>				
<i>Cyphocaris richardi</i>	Antarctic Peninsula	21	11	[62]
<i>Euprimno abyssalis</i>	Bute Inlet, Canada	26	12	[64]
<i>Hyperia galba</i>	Bute Inlet, Canada	19	8	[64]
<i>Eusirus propaperdentatus</i>	Antarctic Peninsula	22	22	[62]
<i>Parandania boeckii</i>	Antarctic Peninsula	20	33	[62]
<i>Parathemisto gaudichaudii</i>	Antarctic Peninsula	19	9	[62]
<b>Annelida</b>				
<i>Tomapteris septentrionalis</i>	Bute Inlet, Canada	22	< 0.5	[64]
<b>Copepoda</b>				
<i>Calanus helgolandicus</i>	?	12	37	[25]
<i>Euchaeta marina</i>	Andaman Sea, India	11	-	[65]
<i>Metridia gerlachei</i>	Antarctic Peninsula	21	52	[62]
<i>Rhincalanus gigas</i>	Antarctic Peninsula	8	< blank	[62]
<i>Undinula vulgaris</i>	Andaman Sea, India	9	-	[65]
<b>Chaetognatha</b>				
<i>Eukrohnia hamata</i>	Arctic	19	12	[66]
<i>Sagitta enflata</i>	Andaman Sea, India	8	-	[65]
<i>Sagitta elegans</i>	Bute Inlet, Canada	14	6	[64]
<i>Sagitta gazellae</i>	Antarctic Peninsula	17	3	[62]
<b>Coelenterata</b>				
<i>Atolla wyvillei</i>	Antarctic Peninsula	1	48	[62]
<i>Beroe cucumis</i>	Bute Inlet, Canada	13	12	[64]
<i>Diphyes antarctica</i>	Antarctic Peninsula	1	16	[62]
<i>Pleurobrachia pileus</i>	Bute Inlet, Canada	9	6	[64]
<b>Decapoda</b>				
<i>Acanthephyra sanguinea</i>	Andaman Sea, India	14	-	[65]
<i>Alpheus</i> sp.	Andaman Sea, India	13	-	[65]
<i>Lucifer hansenii</i>	Cochin estuary, India	10-16	-	[67]
<i>Pasiphaea pacifica</i>	Bute Inlet, Canada	21	4	[64]
<b>Euphasiacea</b>				
<i>Euphausia diomedae</i>	Andaman Sea, India	13	-	[65]
<i>Euphausia pacifica</i>	Bute Inlet, Canada	19	1	[64]
<b>Mysidacea</b>				
<i>Siriella</i> sp.	Andaman Sea, India	11	-	[65]
<b>Ostracoda</b>				
<i>Conchoecia elegans</i>	Bute Inlet, Canada	17	4	[64]
<i>Cypridina dentata</i>	Andaman Sea, India	11	-	[65]
<b>Tunicata</b>				
<i>Salpa thompsoni</i>	Antarctic Peninsula	24	2	[62]
<b>Holoplankton (lipid-rich)</b>				
<b>Amphipoda</b>				
<i>Eurythenes gryllus</i>	Antarctic Peninsula	55	19	[62]
<b>Copepoda</b>				
<i>Calanoides acutus</i>	Antarctic Peninsula	45	64	[62]
<i>Calanus hyperboreus</i>	Arctic	37-74	34-91	[68]

<i>Calanus finmarchicus</i>	Norway	31	71	[69]
<i>Calanus glacialis</i>	Svalbard, Norway	70	68	[70]
<i>Calanus plumchrus</i>	Bute Inlet, Canada	47	86	[64]
<i>Heterorhodus tanneri</i>	Bute Inlet, Canada	43	69	[64]
<i>Metridia longa</i>	Arctic	57	76	[66]
<i>Paraeuchaeta glacialis</i>	Arctic	43	72	[66]
<b>Decapoda</b>				
<i>Hymenodora glacialis</i>	Arctic	35-39	44-62	[66]
<b>Euphasiacea</b>				
<i>Euphausia superba</i> (subadults/adults)	Weddell and Lazarev Seas	7-40	-	[71]
<i>Thysanoessa macrura</i>	Antarctic Peninsula	60	50	[62]
<b>Pteropoda</b>				
<i>Clione limacina</i>	Bute Inlet, Canada and Arctic	19-31	4-12	[64, 66]

424

425 Lipid-poor meroplanktonic larvae from decapods, bryozoans, and vertebrates, a.o., generally  
426 display total lipid contents in the order of 5-15% DM (Table 3). According to the available  
427 literature, total lipid content is on the lower side of this range for bivalve larvae (2-8% DM) and on  
428 the higher part for barnacle larvae (13-15% DM) and octopus paralarvae (< 15% DM) (Table 3).  
429 Both lecithotrophic (which possess and feed on yolk reserves) and planktotrophic (which feed on  
430 plankton) larvae rely on obtaining enough energy to sustain metabolic functions and growth into  
431 juvenile stages, and cannot dispense of energy for storage purposes. Lipid classes which are  
432 destined for energy storage, such as wax esters and triacylglycerols, represent a small proportion of  
433 total lipid content in lipid-poor meroplankton (Table 3). These organisms have specific  
434 requirements of FAs to sustain their body functions, such as formation of biomembranes, and will  
435 thus display lipid homeostasis, even though this may require the allocation of further energy for FA  
436 bioconversion. This is the case of the meroplanktonic larvae of *L. conchilega*, as discussed in the  
437 previous section. It appears, however, that the larvae of hermatypic and soft coral are lipid-rich  
438 (total lipid content reported to range between 41-68% DM; 72, 73). These would be an exception  
439 within the lipid-poor meroplankton, and more information is necessary before we can infer on their  
440 ability to regulate lipid composition during the larval stage.

441 Holoplanktonic organisms with an intermediate total lipid content ranging from 15-25% DM  
442 include small copepods, tunicates, chaetognaths, and ostracods (Table 3). Generally speaking, the  
443 lower end of this range is observed in organisms found in tropical regions, whereas the higher end is

444 recorded for individuals from temperate and polar regions (Table 3). The wax ester content of  
445 organisms varies between different groups, and ranges from values as low as 1% total lipid content  
446 to as high as 52% total lipid content (Table 3). These holoplanktonic organisms generally have a  
447 small to modest ability for energy storage as lipid reserves, a strategy whose focus is likely the  
448 maximization of reproductive output. Such species will mostly assimilate dietary FAs in an  
449 unmodified manner (non-homeostatic).

450 Lipid-rich holoplanktonic species, on the other hand, will display total lipid contents > 30%  
451 DM (Table 3). The species for which data was available were mostly from polar regions, and  
452 represent amphipods (55% DM), large copepods (31-74% DM), krill (7-60% DM), decapods (35-  
453 39% DM), and pteropods (19-31% DM). These organisms need to store energy in the most dense  
454 and effective way to survive low feeding conditions during the winter and for buoyancy purposes,  
455 and will thus bioconvert dietary FAs into long chain MUFAs / wax esters (semi-homeostatic). This  
456 can be exemplified by the large proportion of wax esters within the total lipid content in the  
457 mentioned groups, which is approximately 50% in decapods and krill and 34-90% in copepods  
458 (Table 3). Only pteropods and amphipods have a lower proportion of wax esters, which is  
459 approximately 15-20% of the total lipid content (Table 3).

460

#### 461 **The applicability of the FATM concept**

462 In theory, FATM are assimilated by consumers in a conservative manner [24, 25]. The  
463 applicability of the FATM concept has been broadly acknowledged to depend, a.o., on the  
464 qualitative and quantitative transfer of FAs between trophic levels and on their metabolic stability  
465 and non-selective incorporation into consumer tissues [23]. In the present study the diatom FATMs  
466 were transferred between the producer and consumer trophic levels in a qualitative manner. In  
467 quantitative terms, however, the dietary FAs 16:0, EPA, and DHA were preferentially assimilated  
468 by *L. conchilega* larvae. Furthermore, we have postulated that EPA, which is a diatom FATM, was  
469 likely bioconverted (via chain elongation, as indicated in Fig. 1 of 74) into 22:5(n-3) by the



470 polychaete larvae. Unlike the trends shown by lipid-rich or lipid-accumulating organisms, the  
471 results of this study indicate that the applicability of the concept of FATM is limited in the lipid-  
472 poor larvae of *L. conchilega*. We also suggest that this will be the case for other lipid-poor  
473 meroplanktonic larvae, as they probably also display preferential assimilation and bioconversion of  
474 FAs in order to sustain body functions.

475

476

## 477 **CONCLUSIONS**

478         The present study documented preferential assimilation of FAs by *L. conchilega* larvae,  
479 which seem to be able to regulate their FA composition regardless of the FA profile of their food  
480 source. Bioconversion was also recorded in these individuals, which synthesized the PUFA 22:5(n-  
481 3) from dietary FA precursors. Our results show that dietary FAs are not transferred in a  
482 conservative (quantitative) manner, indicating that the concept of FATM is of limited applicability  
483 in the trophic study of lipid-poor *L. conchilega* larvae. This is probably also the case for other lipid-  
484 poor meroplanktonic larvae. Based on the results of this study and on data available from the  
485 literature, we propose that lipid homeostasis depends upon the lipid content of zooplanktonic  
486 organisms. Lipid-poor meroplanktonic larvae should display lipid homeostasis in order to satisfy  
487 their energetic requirements; holoplanktonic organisms with intermediate lipid levels will usually  
488 assimilate dietary lipids in a conservative manner and invest assimilated energy into reproduction;  
489 whereas lipid-rich holoplanktonic organisms will be semi-homeostatic, converting stored energy  
490 (FAs) into wax esters in order to survive starvation conditions during winter and to increase  
491 buoyancy.

492

493

## 494 **ACKNOWLEDGEMENTS**

495         We kindly thank the crew of the Aade research vessel (AWI-BAH) for providing us with  
496 zooplankton samples and the technicians from the Ecological Chemistry department at the AWI in  
497 Bremerhaven for the help with lipid analysis.

498

499

500 **REFERENCES**

- 501 [1] Umezawa Y, Tamaki A, Suzuki T, Takeuchi S, Yoshimizu C, Tayasu I. Phytoplankton as a  
502 principal diet for callianassid shrimp larvae in coastal waters, estimated from laboratory  
503 rearing and stable isotope analysis. *Mar Ecol Prog Ser.* 2018;592: 141-158.
- 504 [2] Hickel W. The mesozooplankton in the Wadden Sea of Sylt (North Sea). *Helgoland wiss Meer.*  
505 1975;27: 254-262.
- 506 [3] Van Hoey G. Spatio-temporal variability within the macrobenthic *Abra alba* community, with  
507 emphasis on the structuring role of *Lanice conchilega*. 2006. Thesis, Ghent University,  
508 Belgium.
- 509 [4] Strasser M, Pieloth U. Recolonization pattern of the polychaete *Lanice conchilega* on an  
510 intertidal sand flat following the severe winter of 1995/96. *Helgol Mar Res.* 2001;55: 176-  
511 181.
- 512 [5] Bhaud M. The two planktonic larval periods of *Lanice conchilega* (Pallas, 1766) annelida  
513 polychaeta, a peculiar example of the irreversibility of evolution. *Ophelia.* 1988;29: 141-  
514 152.
- 515 [6] Rabaut M. *Lanice conchilega*, fisheries and marine conservation – Towards an ecosystem  
516 approach to marine management. 2009. Thesis, Ghent University, Belgium.
- 517 [7] Bhaud M, Cazaux CP. Buoyancy characteristics of *Lanice conchilega* (Pallas) larvae  
518 (Terebellidae). Implications for settlement. *J Exp Mar Biol Ecol.* 1990;141: 31-45.
- 519 [8] Braeckman U, Provoost P, Sabbe K, Soetaert K, Middelburg JJ, Vincx M, et al. Temporal  
520 dynamics in the diet of two marine polychaetes as inferred from fatty acid biomarkers. *J Sea*  
521 *Res.* 2012;68: 6-19.
- 522 [9] Bühr KJ. Suspension-feeding and assimilation efficiency in *Lanice conchilega* (Polychaeta).  
523 *Mar Biol.* 1976;38: 373-383.
- 524 [10] Bühr KJ, Winter JE. Distribution and maintenance of a *Lanice conchilega* association in the  
525 Weser Estuary (FRG), with special reference to the suspension-feeding behaviour of *Lanice*  
526 *conchilega*. In: Keegan BF, Ceidigh PO, Boaden PJS, editors. *Biology of benthic organisms.*  
527 Pergamon; 1977. pp. 101-113.
- 528 [11] Dubois S, Orvain F, Marin-Leal JC, Ropert M, Lefebvre S. Small-scale spatial variability of  
529 food partitioning between cultivated oysters and associated suspension-feeding species, as  
530 revealed by stable isotopes. *Mar Ecol Prog Ser.* 2007;336: 151-160.

- 531 [12] Zühlke R, Blome D, Heinz van Bernem K, Dittmann S. Effects of the tube-building polychaete  
532 *Lanice conchilega* (Pallas) on benthic macrofauna and nematodes in an intertidal sandflat.  
533 Senck Marit. 1998;29: 131-138.
- 534 [13] Zühlke R. Polychaete tubes create ephemeral community patterns: *Lanice conchilega* (Pallas,  
535 1766) associations studied over six years. J Sea Res. 2001;46: 261-272.
- 536 [14] Alves RMS, van Colen C, Vincx M, Vanaverbeke J, de Smet B, Guarini JM, et al. A case  
537 study on the growth of *Lanice conchilega* (Pallas, 1766) aggregations and their ecosystem  
538 engineering impact on sedimentary processes. J Exp Mar Biol Ecol. 2017;489: 15-23.
- 539 [15] Braber L, De Groot SJ. The food of five flatfish species (Pleuronectiformes) in the southern  
540 North Sea. Neth. J Sea Res. 1973;6: 163-172.
- 541 [16] De Smet B, Godet L, Fournier J, Desroy N, Jaffré M, Vincx M, et al. Feeding grounds for  
542 waders in the Bay of the Mont Saint-Michel (France): the *Lanice conchilega* reef serves as  
543 an oasis in the tidal flats. Mar Biol. 2013;160: 751-761.
- 544 [17] Godet L, Toupoint N, Olivier F, Fournier J, Retière C. Considering the functional value of  
545 common marine species as a conservation stake: The case of Sandmason Worm *Lanice*  
546 *conchilega* (Pallas 1766) (Annelida, Polychaeta) beds. Ambio. 2008;37: 347-355.
- 547 [18] Petersen B, Exo KM. Predation of waders and gulls on *Lanice conchilega* tidal flats in the  
548 Wadden Sea. Mar Ecol Prog Ser. 1999;178: 229-240.
- 549 [19] Rabaut M, Guilini K, Van Hoey G, Vincx M, Degraer S. A bio-engineered soft-bottom  
550 environment: The impact of *Lanice conchilega* on the benthic species-specific densities and  
551 community structure. Estuar Coast Shelf S. 2007;75: 525-536.
- 552 [20] Callaway R. Tube worms promote community change. Mar Ecol Prog Ser. 2006;308: 49-60.
- 553 [21] Callaway R. Juveniles stick to adults: recruitment of the tube-dwelling polychaete *Lanice*  
554 *conchilega* (Pallas, 1766). Hydrobiologia. 2003;503: 121-130.
- 555 [22] Ropert M, Dauvin JC. Renewal and accumulation of a *Lanice conchilega* (Pallas) population in  
556 the Baie des Veys, western Bay of Seine. Oceanol Acta. 2000;23: 529-546.
- 557 [23] Dalsgaard J, John MS, Kattner G, Müller-Navarra D, Hagen W. Fatty acid trophic markers in  
558 the pelagic marine environment. Adv Mar Biol. 2003;46: 225-340.
- 559 [24] Graeve M, Kattner G, Hagen W. Diet-induced changes in the fatty acid composition of Arctic  
560 herbivorous copepods: Experimental evidence of trophic markers. J Exp Mar Biol Ecol.  
561 1994;182: 97-110.
- 562 [25] Lee RF, Nevenzel JC., Paffenhöfer GA. Importance of wax esters and other lipids in the  
563 marine food chain: phytoplankton and copepods. Mar Biol. 1971;9: 99-108.
- 564 [26] Boissonnot L, Niehoff B, Hagen W, Søreide JE, Graeve M. Lipid turnover reflects life-cycle  
565 strategies of small-sized Arctic copepods. J Plankton Res. 2016;38: 1420-1432.

- 566 [27] Graeve M, Albers C, Kattner G. Assimilation and biosynthesis of lipids in Arctic Calanus  
567 species based on feeding experiments with a <sup>13</sup>C labelled diatom. J Exp Mar Biol Ecol.  
568 2005;317: 109-125.
- 569 [28] Guilini K, van Oevelen D, Soetaert K, Middelburg JJ, Vanreusel A. Nutritional importance of  
570 benthic bacteria for deep-sea nematodes from the Arctic ice margin: Results of an isotope  
571 tracer experiment. Limnol Oceanogr. 2010;55: 1977-1989.
- 572 [29] Rix L, de Goeij JM, van Oevelen, D, Struck U, Al-Horani FA, Wild C, et al. Reef sponges  
573 facilitate the transfer of coral-derived organic matter to their associated fauna via the sponge  
574 loop. Mar Ecol Prog Ser. 2018;589: 85-96.
- 575 [30] Wegener G, Bausch M, Holler T, Thang NM, Mollar XP, Kellermann MY, et al. Assessing  
576 sub-seafloor microbial activity by combined stable isotope probing with deuterated water  
577 and <sup>13</sup>C-bicarbonate. Env Microbiol. 2012;14: 1517-1527.
- 578 [31] De Troch M, Boeckx P, Cnudde C, Van Gansbeke D, Vanreusel A, Vincx M, et al.  
579 Bioconversion of fatty acids at the basis of marine food webs: insights from a compound-  
580 specific stable isotope analysis. Mar Ecol Prog Ser. 2012;465: 53-67.
- 581 [32] Middelburg JJ, Barranguet C, Boschker HTS, Herman PMJ, Moens T, Heip CHR. The fate of  
582 intertidal microphytobenthos carbon: An in situ <sup>13</sup>C-labeling study. Limnol Oceanogr.  
583 2000;45: 1224-1234.
- 584 [33] Guillard RRL. Culture of phytoplankton for feeding marine invertebrates. In: Smith WL,  
585 Chanley MH, editors. Culture of marine invertebrate animals. US: Springer; 1975. pp. 29-  
586 60.
- 587 [34] Li HY, Lu Y, Zheng JW, Yang WD, Liu JS. Biochemical and genetic engineering of diatoms  
588 for polyunsaturated fatty acid biosynthesis. Mar Drugs. 2014;12: 153-166.
- 589 [35] Ihaka R, Gentleman R. R. Version 3.4.4 [software]. 1993. Available from [https://cran.r-  
590 project.org/bin/windows/base/old/3.2.4/](https://cran.r-project.org/bin/windows/base/old/3.2.4/).
- 591 [36] Warton DI, Hui FKC. The arcsine is asinine: the analysis of proportions in ecology. Ecology.  
592 2011;92: 3-10.
- 593 [37] Clarke KR, Gorley RN. PRIMER v7: User Manual/Tutorial. PRIMER-E, Plymouth; 2015.
- 594 [38] Franco-Santos RM, Auel H, Boersma M, De Troch M, Meunier CL, Niehoff B. Bioenergetics  
595 of the copepod *Temora longicornis* under different nutrient regimes. J Plankton Res.  
596 2018;40: 420-435.
- 597 [39] Kiørboe T, Møhlenberg F, Hamburger K. Bioenergetics of the planktonic copepod *Acartia  
598 tonsa*: relation between feeding, egg production and respiration, and composition of specific  
599 dynamic action. Mar Ecol Prog Ser. 1985;26: 85-97.

- 600 [40] Harms J, Anger K, Klaus S, Seeger B. Nutritional effects on ingestion rate, digestive enzyme  
601 activity, growth, and biochemical composition of *Hyas araneus* L. (Decapoda: Majidae)  
602 larvae. J Exp Mar Biol Ecol. 1991;145: 233-265.
- 603 [41] Harms J, Meyer-Harms B, Dawirs RR, Anger K. Growth and physiology of *Carcinus maenas*  
604 (Decapoda, Portunidae) larvae in the field and in laboratory experiments. Mar Ecol Prog  
605 Ser. 1994;108: 107-118.
- 606 [42] His E, Maurer D. Shell growth and gross biochemical composition of oyster larvae  
607 (*Crassostrea gigas*) in the field. Aquaculture. 1988;69: 185-194.
- 608 [43] Müller-Navarra DC. The nutritional importance of polyunsaturated fatty acids and their use as  
609 trophic markers for herbivorous zooplankton: Does it contradict? Arch Hydrobiol. 2006;167:  
610 501-513.
- 611 [44] Brett MT, Müller-Navarra DC. The role of highly unsaturated fatty acids in aquatic foodweb  
612 processes. Freshwater Biol. 1997;38: 483-499.
- 613 [45] Jónasdóttir SH, Visser AW, Jespersen C. Assessing the role of food quality in the production  
614 and hatching of *Temora longicornis* eggs. Mar Ecol Prog Ser. 2009;382: 139-150.
- 615 [46] Thor P, Koski M, Tang KW, Jónasdóttir SH. Supplemental effects of diet mixing on absorption  
616 of ingested organic carbon in the marine copepod *Acartia tonsa*. Mar Ecol Prog Ser.  
617 2007;331: 131-138.
- 618 [47] Johnston PV. Dietary fat, eicosanoids, and immunity. Adv Lipid Res. 1985;21:103-141.
- 619 [48] Budge SM, Iverson SJ, Koopman HN. Studying trophic ecology in marine ecosystems using  
620 fatty acids: a primer on analysis and interpretation. J Exp Biol. 2006;214: 3790-3800.
- 621 [49] Jeffreys RM, Wolff GA, Murty SJ. The trophic ecology of key megafaunal species at the  
622 Pakistan Margin: Evidence from stable isotopes and lipid biomarkers. Deep-Sea Res.  
623 2009;56: 1816-1833.
- 624 [50] Budge SM, Wang SW, Hollmén TE, Wooller MJ. Carbon isotopic fractionation in eider  
625 adipose tissue varies with fatty acid structure: implications for trophic studies. Mar Mammal  
626 Sci. 2011;22: 759-801.
- 627 [51] Pond DW, Allen CE, Bell MV, Van Dover CL, Fallick AE, Dixon DR, et al. Origins of long-  
628 chain polyunsaturated fatty acids in the hydrothermal vent worms *Ridgea*  
629 *piscesae* and *Protis hydrothermica*. Mar Ecol Prog Ser. 2002;225: 219-226.
- 630 [54] Lee RFL, Hagen W, Kattner G. Lipid storage in marine zooplankton. Mar Ecol Prog Ser.  
631 2006;307: 273-306.
- 632 [52] Mansour MP, Holdsworth DG, Forbes SE, Macleod CK, Volkman JK. High contents of  
633 24:6(n-3) and 20:1(n-13) fatty acids in the brittle star *Amphiura elandiformis* from  
634 Tasmanian coastal sediments. Biochem Syst Ecol. 2005;33: 659-674.

- 635 [53] Kreibich T, Saborowski R, Hagen W, Niehoff B. Influence of short-term nutritional variations  
636 on digestive enzyme and fatty acid patterns of the calanoid copepod *Temora longicornis*. J  
637 Exp Mar Biol Ecol. 2011;407: 182-189.
- 638 [54] Mann R, Gallager SM. Physiological and biochemical energetics of larvae of *Teredo nevalis* L.  
639 and *Bankia gouldi*. (Bartsch) (Bivalvia: Teredinidae). J Exp Mar Biol Ecol. 1985;85: 211-  
640 228.
- 641 [55] Whyte JNC, Bourne N, Ginther NG, Hodgson CA. Compositional changes in the larva to  
642 juvenile development of the scallop *Crassadoma gigantea* (Gray). J Exp Mar Biol Ecol.  
643 1992;163: 13-29.
- 644 [56] Orellana MC, Cancino JM, Hughes RN. Is settlement in lecithotrophic bryozoan larvae  
645 constrained by energy reserves? In: Gordon DP, Smith AM, Grant-Mackie JA, editors.  
646 Bryozoans in space and time. Proceedings of the 10<sup>th</sup> International Bryozoology Conference;  
647 Wellington, New Zealand. National Institute of Water and Atmospheric Research Ltd.,  
648 Wellington; 1996. 442p.
- 649 [57] Holland DL, Walker G. The biochemical composition of the cypris larva of the barnacle  
650 *Balanus balanoides* L. J Const Int Explor Mer. 1975;36: 162-165.
- 651 [58] Lourenço S, Roura A, Fernández-Reiriz MJ, Narciso L, González AF. Feeding relationship  
652 between *Octopus vulgaris* (Cuvier, 1797) early life-cycle stages and their prey in the  
653 Western Iberian upwelling system: Correlation of reciprocal lipid and fatty acid contents.  
654 Front Physiol. 2017;8: 1-11.
- 655 [59] Thatje S, Lovrich GA, Torres G, Hagen W, Anger K. Changes in biomass, lipid, fatty acid and  
656 elemental composition during the abbreviated larval development of the subantarctic shrimp  
657 *Campylonotus vagans*. J Exp Mar Biol Ecol. 2004;301: 159-174.
- 658 [60] Rotllant G, Anger K, Durfort M, Sardà F. Elemental and biochemical composition of *Nephrops*  
659 *norvegicus* (Linnaeus 1758) larvae from the Mediterranean and Irish Seas. Helgol Mar Res.  
660 2004;58: 206-210.
- 661 [61] Wang M, O'Rorke R, Waite AM, Beckley LE, Thompson P, Jeffs AG. Fatty acid profiles of  
662 phyllosoma larvae of western rock lobster (*Panulirus cygnus*) in cyclonic and anticyclonic  
663 eddies of the Leeuwin Current off Western Australia. Progr Oceanogr. 2014;122: 153-162.
- 664 [62] Reinhardt SB, Van Vleet ES. Lipid composition of twenty-two species of Antarctic midwater  
665 zooplankton and fish. Mar Biol. 1986;91: 149-159.
- 666 [63] Vázquez R, González S, Rodríguez A, Mourente G. Biochemical composition and fatty acid  
667 content of fertilized eggs, yolk sac stage larvae and first-feeding larvae of the Senegal sole  
668 (*Solea senegalensis* Kaup). Aquaculture. 1994;119: 273-286.

- 669 [64] Lee RF. Lipids of zooplankton from Bute Inlet, British Columbia. J Fish Res Board Can.  
670 1974b;31: 1577-1582.
- 671 [65] Goswami SC, Rao TSS, Matondkar SGP. Biochemical composition of zooplankton from the  
672 Andaman Sea. Indian J Mar Sci. 1981;10: 296-300.
- 673 [66] Lee RF. Lipids of Arctic zooplankton. Comp Biochem Physiol. 1975;51: 263-266.
- 674 [67] Antony G, Antony A. Biochemical composition of *Lucifer hanseni* from the Cochin estuary.  
675 Indian J Fish. 2001;48: 41-47.
- 676 [68] Lee RF. Lipid composition of the copepod *Calanus hyperboreas* from the Arctic Ocean.  
677 Changes with depth and season. Mar Biol. 1974a;26: 313-318.
- 678 [69] Diel S, Tande K. Does the spawning of *Calanus finmarchicus* in high latitudes follow a  
679 reproducible pattern? Mar Biol. 1992;113: 21-31.
- 680 [70] Scott CL, Kwasniewski S, Falk-Petersen S, Sargent JR. Lipids and life strategies of *Calanus*  
681 *finmarchicus*, *Calanus glacialis* and *Calanus hyperboreus* in late autumn, Kongsfjorden,  
682 Svalbard. Polar Biol. 2000;23: 510-516.
- 683 [71] Hagen W, Kattner G, Terbrüggen A, Van Vleet ES. Lipid metabolism of the Antarctic krill  
684 *Euphausia superba* and its ecological implication. Mar Biol. 2001;139: 95-104.
- 685 [72] Ben-David-Zaslow R, Benayahu Y. Biochemical composition, metabolism, and amino acid  
686 transport in planula-larvae of the soft coral *Heteroxenia fuscescens*. J Exp Zool. 2000;287:  
687 401-412.
- 688 [73] Harii S, Nadaoka K, Yamamoto M, Iwao K. Temporal changes in settlement, lipid content and  
689 lipid composition of larvae of the spawning hermatypic coral *Acropora tenuis*. Mar Ecol  
690 Prog Ser. 2007;346: 89-96.
- 691 [74] Leonard AE, Pereira SL, Sprecher H, Huang YS. Elongation of long-chain fatty acids. Prog  
692 Lipid Res. 2004;43: 36-54.












Fig 2

[Click here to access/download;Figure;Fig 2.tiff](#)









## CHAPTER IV

### **Temperature driven changes in the diet preference of omnivorous copepods: no more meat when it's hot?**

Boersma M, Mathew KA, Niehoff B, Schoo KL, Franco-Santos RM, Meunier CL

Published in Ecology Letters, 19, doi: 10.1111/ele.12541, in November 2015.



LETTER

## Temperature driven changes in the diet preference of omnivorous copepods: no more meat when it's hot?

Maarten Boersma,<sup>1,2\*</sup>  
K. Avarachen Mathew,<sup>1†</sup> Barbara  
Niehoff,<sup>3</sup> Katherina L. Schoo,<sup>1‡</sup>  
Rita M. Franco-Santos<sup>2,3</sup> and  
Cédric L. Meunier<sup>1</sup>

### Abstract

Herbivory is more prevalent in the tropics than at higher latitudes. If differences in ambient temperature are the direct cause for this phenomenon, then the same pattern should be visible in a seasonal gradient, as well as in experiments manipulating temperature. Using <sup>15</sup>N stable isotope analyses of natural populations of the copepod *Temora longicornis* we indeed observed seasonal differences in the trophic level of the copepod and a decrease in trophic level with increasing temperature. In a grazing experiment, with a mixed diet of the cryptophyte *Rhodomonas salina* and the heterotrophic dinoflagellate *Oxyrrhis marina*, *T. longicornis* preferred the cryptophyte at higher temperatures, whereas at lower temperatures it preferred the non-autotrophic prey. We explain these results by the higher relative carbon content of primary producers compared to consumers, in combination with the higher demand for metabolic carbon at higher temperatures. Thus, currently increasing temperatures may cause changes in dietary preferences of many consumers.

### Keywords

Global warming, grazing, homeostasis, metabolism, stoichiometry, zooplankton.

Ecology Letters (2016) 19: 45–53

### INTRODUCTION

For fish in particular, a clear latitudinal gradient exists with respect to the prevalence of herbivory (Floeter *et al.* 2005; Behrens & Lafferty 2007; Gonzalez-Bergonzoni *et al.* 2012), with higher incidences of herbivory in the tropics, whereas carnivory is more important at higher latitudes. Several explanations for this phenomenon have been brought forward. These were well-developed by Clements *et al.* (2009) and Gonzalez-Bergonzoni *et al.* (2012), to which we refer readers for more details, but explanations range from evolutionary constraints (colder systems are younger so herbivory has not developed as yet), through availability (in colder systems there are no plants in winter), and quality (digestibility of tropical plants tissue is higher) to a suite of directly temperature-related explanations. Digestion of plant material is considered easier in higher temperatures, so herbivory should be more prevalent (Floeter *et al.* 2005). This temperature constraint hypothesis seems to be the most widely accepted explanation for the observed gradient, and essentially argues that with declining temperature feeding rates decrease more rapidly than metabolic rates. The critical temperature required for net positive energy balance is thought to be higher for herbivores than for carnivores because animal material is a high quality food and easier to assimilate than plant material (Floeter *et al.* 2005). In their review on herbivory in marine fish Clements *et al.* (2009) argue that it is essential to also consider the nutritional content of the food and take into account nutrient intake. They state that temperature can also affect

the nutrient content of plant material, and with this, feeding rates of herbivores.

Especially the food content of nitrogen (N) and phosphorus (P) in their concentration relative to carbon (C) have received a lot of attention (Sterner & Elser 2002), and there is strong consensus in the aquatic literature that plants (algae) have higher (and more variable) carbon to nitrogen and phosphorus contents than animals (e.g. van de Waal *et al.* 2009). Hence, a consumer requiring relatively more C should select for plants, whereas a higher P or N requirement should lead to a selection for animal prey. Several studies have investigated the temperature sensitivity of nutritional demands with respect to C : N : P stoichiometry, very well summarised recently by Cross *et al.* (2015). Two competing predictions exist. The Growth Rate Hypothesis (Elser *et al.* 1996) links growth rate to phosphorus demand, and, as in poikilotherms growth rates increase with temperature, this hypothesis predicts a higher P-content and hence P-demand at higher temperatures (Dobberfuhl & Elser 2000; Persson *et al.* 2011). In a later study though, Elser *et al.* (2000) include evolutionary arguments in their reasoning, stating that as a result of shorter growing seasons, growth rates of arctic clones of the cladocerans *Daphnia pulex* are actually higher than those of temperate conspecifics, and hence they should have a higher P-demand (see also Daines *et al.* 2014 for a similar discussion in algae). This is of course a completely different line of reasoning, invoking evolutionary aspects rather than physiological ones. It is also argued that higher temperatures lead to an increase in protein denaturing and higher protein turn-over

<sup>1</sup>Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Biologische Anstalt Helgoland, Postfach 180, 27483 Helgoland, Germany

<sup>2</sup>University of Bremen, Bremen, Germany

<sup>3</sup>Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Bremerhaven, Germany

<sup>†</sup>Present address: Department of Marine Biology, Microbiology and Biochemistry, Cochin University of Science and Technology, Cochin 682016, India

<sup>‡</sup>Present address: Shannon Point Marine Center, Western Washington University, Anacortes, WA, USA

\*Correspondence: E-mail: maarten.boersma@awi.de

rates, thus increasing the demand for nitrogen at higher temperatures (Lemoine *et al.* 2013). In contrast, higher temperatures also imply higher metabolic rates. If there is a different temperature dependence of anabolic and catabolic processes (e.g. Karl & Fischer 2008), which has been suggested especially in relation to the wide spread phenomenon of temperature related changes in body size (Forster *et al.* 2011, 2012), and respiratory processes have a higher  $Q_{10}$  than growth (see also Larsson & Berglund 2005), then we would expect that at higher temperatures the relative demand for carbon (respiration) is higher than for N or P. Cross *et al.* (2015) make this argument, leading to a prediction of the temperature sensitivity to quality limitation which is concurrent with the observed latitudinal variability in herbivory. Lower temperatures trigger a higher necessity for nutrients relative to carbon, and thus a higher necessity for meat.

If the explanation of direct temperature-dependent differences in nutrient demands is correct, this implies that there should be a direct influence of temperature on feeding patterns and selectivity. This can be tested. First, we would expect the pattern of changing feeding modes also to be present within one species over different seasons of the year. Furthermore, in an experimental set-up, we would predict a direct dependence of the grade of herbivory on temperature, and a higher affinity for high carbon food at higher temperatures. To answer these questions, we sampled the calanoid copepod *Temora longicornis* over several seasonal cycles, and investigated the trophic level of this copepod, which can switch between herbivorous and carnivorous feeding modes (Gentsch *et al.* 2009; Boersma *et al.* 2014), using nitrogen stable isotope measurements. Furthermore, we directly investigated the selectivity of adult females of this copepod species cultured under a range of different temperatures when fed autotrophic and heterotrophic food, and investigated feeding rates of *T. longicornis* on high C : P and low C : P algae at two different temperatures.

## MATERIALS AND METHODS

### Sampling site

The samples for the present study were collected from the Helgoland Roads time series station in the German Bight of the North Sea. This sampling station is located between the island of Helgoland and the adjacent Düne island at the Kabeltonne site (54° 11.3' N, 7° 54.0' E) (Wiltshire & Manly 2004).

### Sample collection and analysis

The samples were collected over a period of 4 years from 2004 to 2008. The copepod *Temora longicornis* was sampled using a 500 µm CalCOFI net with an aperture of 100 cm and net length 400 cm (Greve *et al.* 2004). The collected *T. longicornis* were transferred to the laboratory and immediately sorted alive using a stereomicroscope. Then the copepods were individually washed in distilled water to remove particles and salt. Three groups of 10–20 individuals were transferred into tin cartridges, and dried at 60 °C for up to 24 h. Similarly,

seawater surface samples were collected and filtered, using pre-combusted (6 h, 420 °C) glass fibre filters (Whatman, GF/C, 25 mm diameter). Depending on the density of algae 200 mL–1 L was filtered. All filters were also heat dried at 60 °C for 12 h and transferred to tin capsules for further analysis. To estimate the trophic position of the *T. longicornis* females, we analysed the  $\delta^{15}\text{N}$  content of both the animals and the seston. The samples were analysed by ARGROISO-LAB (Jülich/Germany) or at the Stable Isotope Facility at University of California, Davis using an Elementar Vario EL Cube and Micro Cube elemental analyser (Elementar Analysensysteme GmbH, Hanau, Germany) which was interfaced to PDZ Europa 20-20 isotope ratio mass spectrometer (Sercon Ltd., Cheshire, UK). The difference between the  $\delta^{15}\text{N}$  value of animals and seston was used to calculate the trophic level (TL) position of the *Temora longicornis* females, using a trophic fraction constant of 3.4 per trophic level (Minagawa & Wada 1984; Vander Zanden & Rasmussen 2001). In order not to artificially inflate the number of samples, we averaged the values of the three replicates per sampling day, before entering the analysis. We then related the computed trophic level of the copepods with ambient temperatures.

### Copepod grazing experiments

For the grazing experiment on the two different prey the protocols of Löder *et al.* (2011) were followed. *T. longicornis* was collected from the field, and five adult female *T. longicornis* were sorted and transferred to 800 mL glass beakers containing 500 mL of filtered seawater. Five replicate beakers per temperature were subjected to temperatures from 10 to 28 °C (with a total of 10 different temperatures) for 24 h to precondition the animals for experiments. To determine grazing selectivity of *T. longicornis* a mixed diet of the cryptophyte *Rhodomonas salina* and the heterotrophic dinoflagellate *Oxyrrhis marina* was used. *R. salina* was cultured in F/2 culture medium, whereas *O. marina* was cultured in the same medium but pre-fed *R. salina*. Feeding conditions for the dinoflagellate before the experimental incubations were such that the food algae in the *O. marina* cultures had virtually been depleted at the time of the start of the experiment. For culture conditions and experimental protocol see Meunier *et al.* (2011). The C : N : P stoichiometry of *R. salina* and *O. marina* grown under these conditions are 320 : 38 : 1 and 175 : 25 : 1, respectively (see also Hantzsche & Boersma 2010). This demonstrates the premise of higher relative nutrient content in heterotrophs.

The experimental containers with *T. longicornis* were inoculated with *R. salina* and *O. marina* in cell densities of 8000 and 1000 cells mL<sup>-1</sup>, respectively, to provide similar concentrations of prey carbon for *T. longicornis*. The cell number of the algal cultures was determined using a CASY particle counter (Schärfe Systems, Reutlingen, Germany) (Meunier *et al.* 2013), manual counts were carried out to determine the density of *O. marina*. The experiment was conducted using a temperature table (Thomas *et al.* 1963) with a temperature gradient that ranged from 10 to 28 °C. Each of the ten temperatures had five replicates. Since the dinoflagellate *O. marina* also feeds on *R. salina*, the grazing experiment needed



appropriate controls. The controls included one that contained *R. salina* only to estimate growth rates of the alga, and another one containing a mix of *R. salina* and *O. marina* at the same densities as in the treatment with *T. longicornis*. This allows the separate estimation of growth and consumption rates of *O. marina* on *R. salina*. The experiment ran for 24 h with 15 h of light and 9 h of darkness. At the end of the experiment, the number of surviving *T. longicornis* were counted, and Lugol's iodine solution was used to arrest the growth of the algae and to preserve the *R. salina* and *O. marina* for further analysis: *O. marina* was counted using a Sedgewick Rafter Cell Counter under the microscope and *R. salina* using the CASY particle counter. Typically, in an experiment such as this one, grazing and ingestion rates are computed using the equations as presented by Frost (1972), Heinbokel (1978), and Nejstgaard *et al.* (2001) (summarised by Löder *et al.* 2011). Despite the fact that this set of equations is much used in the literature, the way these equations are set up creates an interdependence of estimates of the grazing rates of the different prey. An increased estimate of the grazing impact of *T. longicornis* on *O. marina* leads to a decrease in the calculated grazing pressure of *O. marina* on the alga and, to balance numbers, automatically also to an increase in the estimate of the grazing impact of the copepod on the alga. Hence, since we were interested in the temperature dependent patterns, rather than the exact numbers, we decided to simplify the approach, and also compared the slopes with temperature of the average numbers of *R. salina* and *O. marina* in the treatments with and without predators, using ANCOVA analysis. We used the exact temperatures measured in each separate replicate for this analysis rather than the averages of each of the rows in the temperature table. Based on the computed grazing rates, using the appropriate equations (see above), we computed the ingestion on both prey types, and from this the amount of ingested carbon and phosphorus, and the C : P ratio of the ingested material.

The final step linking herbivory to potential nutritional demands is the investigation of feeding patterns on one food source of different nutritional content at different temperatures. If a higher temperature is linked to a higher C-demand, then one would expect higher relative grazing rates on nutrient limited (C-rich) algae at higher temperatures than at lower temperatures. Thus in two additional experiments, we collected *T. longicornis* from the field, and sorted 10 adult females per treatment. These were transferred to 100 mL plastic beakers containing 100 mL of filtered seawater, and placed in one of two temperature-controlled rooms at 10 and 20 °C for 72 h to pre-condition the animals for experiments. During the first 48 h the copepods were fed seston between 30 and 150 µm and were left without food for the last 24 h to reduce any potential pre-feeding effects. To determine grazing of *T. longicornis*, two cultures of the cryptophyte *R. salina* were used, one grown in F/2 medium while the other was grown in F/2 medium without P to induce a P-limitation (Meunier *et al.* 2011, 2015). The P content of P-replete and P-depleted *R. salina* were 0.30 ( $\pm 0.07$ ) and 0.13 ( $\pm 0.01$ ) pg P cell<sup>-1</sup>, respectively, yielding C : P rates of 448 : 1 and 1100 : 1. Two experiments were carried out to investigate the robustness of the change in feeding. *T. longicornis* were caught from two

different sites at two different occasions, with about 6 weeks between them. We incubated ten *T. longicornis* in 250 mL Falcon culture flasks with either P-replete or P-depleted *R. salina* in cell densities of 1000 cells mL<sup>-1</sup>. The flasks were filled with artificial sea water at a salinity of 32 to prevent any change in *R. salina* nutritional quality. Another set of bottles were incubated without copepods to serve as control. At least three replicates per treatment were used and the flasks were placed on two plankton wheels (0.8 rpm) in temperature-controlled rooms at 10 and 20 °C. The experiment was run for 24 h and samples were taken and fixed with formalin (formaldehyde 20% buffered with hexamine). The samples were stored cool and dark before being counted with a BD ACURI C6 flow cytometer. For each sample, 500 µL of sample was processed by the flow cytometer to ensure that at least 200 *R. salina* cells were counted. Grazing rates were calculated as described above based on Frost (1972), and data were analysed with a three-way ANOVA using experiment, food type and temperature as factor and grazing rate as dependent variables.


## RESULTS

### Stable isotope analysis of natural populations

The  $\delta^{15}\text{N}$  values from the natural populations of *T. longicornis* and seston showed considerable variation with time (Fig. 1a). The difference in  $\delta^{15}\text{N}$  values between *T. longicornis* and seston was used to compute the trophic level of the copepods. A trophic level of two indicates that the only food source of the copepod is seston, and that *T. longicornis* is the primary consumer of this seston, whereas a trophic level of three indicates that there is at least one trophic level between the seston and the copepod, and hence that the copepod is in fact the secondary consumer (Fig. 1b). The trophic level position of *T. longicornis* ranged between 1.01 and 3.45, and it significantly decreased with increasing in temperature ( $r^2 = 0.32$ ;  $P < 0.0001$ ). This decrease in the trophic level position indicates that *T. longicornis* fed on a diet richer in phytoplankton at warmer temperatures, the slope of  $-0.064$  °C indicates that with every 15–16 °C of temperature increase the trophic position of *T. longicornis* decreases by one level (Fig. 1c). This relationship still holds after collating all the data into temperature bins/ groups of 4 °C (Fig. 1c, inset), to avoid the problem of autocorrelations in the data. The regression with the means of the bins against temperature is still significant, despite an n of only five points ( $r^2 = 0.85$ ;  $P = 0.03$ ). Furthermore, if we compute the change in trophic level of *T. longicornis* between two adjacent measuring points from the data series, the change in trophic level (TL) averages  $-0.14$  (SE = 0.10) with a temperature increase between two points, whereas the change in TL is positive (0.24; SE = 0.12) when the temperature decreases. This difference is significant (*t*-test;  $P = 0.01$ ).

### Temperature dependent copepod grazing on heterotrophic vs. autotrophic prey


The grazing experiment with *T. longicornis* was conducted using *R. salina* and *O. marina* as prey. All *T. longicornis* died



**Figure 1** (a)  $\delta^{15}\text{N}$  values of seston and the copepod *Temora longicornis* for samples collected from 2004 to 2008; (b) Trophic level position of *T. longicornis* (triangles) with water temperature at the moment of sampling (line); (c) Correlation of the trophic position of *T. longicornis* with temperature. The regression line (trophic level =  $2.62 - 0.064 \times \text{Temp}$ ) explains 32% of the variance in the trophic level of *T. longicornis* ( $P < 0.0001$ ;  $n = 76$ ), and indicates a loss of one trophic level with every 15–16 °C. The inset shows the average trophic level of the *T. longicornis* in 4 °C bins. Also the linear regression with only the averages ( $n = 5$ ) as dependent variables is significant ( $P = 0.03$ ).

at 28 °C, survival was very good in the other temperatures. The growth rate of *R. salina* was affected by the temperature, as indicated by the temperature dependence of the average densities in the treatment with *R. salina* alone (Fig. 2a;  $r^2 = 0.19$ ;  $P < 0.005$ ). In the presence of only *O. marina* as a grazer, we observed no temperature dependence of the *R. salina* densities ( $r^2 = 0.003$ ;  $P = 0.75$ ), whereas with two

predators present densities were again significantly correlated with temperature ( $r^2 = 0.48$ ;  $P < 0.001$ ), indicating higher consumption rates of *T. longicornis* on algae at higher temperatures. Homogeneity of slope analysis revealed that the slopes of the two significant regressions (*R. salina* alone, and *R. salina* with both predators) with temperature were not significantly different from each other (ANCOVA,  $F_{1,1} = 1.03$ ;



**Figure 2** Densities of (a) *Rhodomonas salina*, and (b) *Oxyrrhis marina* in treatments with and without predators. Regression lines indicate significant relationships between temperature and cell numbers. Dotted lines are the initial densities at the beginning of the experiment; (a) *Rhodomonas salina* alone (green triangles; Density =  $15\,080 - 244 \times \text{Temp}$ ;  $r^2 = 0.19$ ;  $P < 0.005$ ), with one predator (yellow circles), and two predators (red stars; Density =  $14\,835 - 347 \times \text{Temp}$ ;  $r^2 = 0.48$ ;  $P < 0.001$ ); (b) *Oxyrrhis marina* densities alone (green triangles), and with predator (yellow circles; Density =  $540 + 21 \times \text{Temp}$ ;  $r^2 = 0.59$ ;  $P < 0.001$ ); (c) Grazing rates of *Temora longicornis* on *R. salina* (red triangles; rate =  $-0.4 + 0.035 \times \text{Temp}$ ;  $r^2 = 0.18$ ;  $P < 0.007$ ) and *O. marina* (blue circles; rate =  $1.05 - 0.04 \times \text{Temp}$ ;  $r^2 = 0.28$ ;  $P < 0.0005$ ) indicates that the grazing on *R. salina* increases with temperature, whereas the grazing on *O. marina* decreases. All computed negative feeding rates were set to zero.

$P = 0.313$ ) but both were different from the treatment with only *Oxyrrhis* as a grazer. When comparing the average number of *O. marina* in relation to the temperature (Fig. 2b) it becomes clear that whereas the growth rate of *O. marina* is not correlated with temperature (no significant correlation,  $r^2 = 0.01$ ;  $P = 0.56$ ), the average density in the presence of *T. longicornis* is affected ( $r^2 = 0.59$ ;  $p < 0.001$ ), resulting in a significant difference in the slopes of the regression lines with temperature (ANCOVA, interaction between temperature and treatment,  $F_{1,1} = 12.90$ ;  $P < 0.001$ ). This difference remained significant even after removing the data points for the lowest temperature for *O. marina* without the copepods, thus this result was not dependent on these points. With higher temperatures the average numbers of *O. marina* cells in the experimental units increased significantly, which indicates a lower grazing pressure of *T. longicornis* on *O. marina* with increasing temperature. The data are summarised in Fig. 2c, which shows the computed grazing rates of *T. longicornis* on both potential prey (with negative grazing rates set to zero, see also Nejstgaard *et al.* (2001)). The grazing rate of *T. longicornis* on the alga *R. salina* showed a significantly positive correlation

with temperature ( $r^2 = 0.18$ ;  $P < 0.01$ ), whereas the grazing on the dinoflagellate *O. marina* was negatively correlated with temperature ( $r^2 = 0.28$ ;  $P < 0.0005$ ), thus indicating a switch in preference from 'animal' food at low temperatures to 'plant' food at higher temperatures. This result does not depend on our choice to set negative grazing rates to zero. Even with all values included (Table S1; Fig. S1a), or with negative values excluded from the analysis (Fig. S1c), the results are qualitatively the same. In fact, the slopes of the regression lines of the three different analyses are not significantly different from each other (Table S1). Based on the ingestion rates at different temperatures and the C : P ratios of both prey, we can compute the change in C : P ratio in the food taken up by the *T. longicornis* at the different temperatures. With a linear regression relating temperature to ingested C : P ( $r^2 = 0.18$ ;  $P < 0.001$ ) we estimated that at 10 °C the C : P ratio of the ingested food was 175 (essentially only taking up *O. marina*), and at 20 °C the C : P of the food taken up was 216. This means that at higher temperatures *Temora longicornis* take up ~25% more carbon relative to phosphorus.

### Temperature dependent copepod grazing on high C : P vs. low C : P algae


In the two experiments with differently grown *R. salina* we observed a significant interaction between temperature and food type (Table 1; Fig. 3). Feeding on P-limited *R. salina* increased with temperature, whereas feeding on P-replete algae decreased, consistent with the predictions and observations from the previous experiments. This shows that P-limited food is a more attractive food source at higher temperatures.

### DISCUSSION

In this study, we showed that the trophic level of the calanoid copepod *Temora longicornis* varied seasonally, leading to a significantly negative correlation between the estimated trophic level and temperature of the environment. This finding is consistent with the global patterns of the relationship between intensity of herbivory and latitude, and shows that this pattern can even be observed within one system on a seasonal scale. The experiment directly investigating the effect of temperature on dietary preference of *T. longicornis* corroborated these findings. At higher temperatures the copepod

**Table 1** ANOVA results of the analysis on the effects of experiment, temperature and food type on the grazing rate of *Temora longicornis*

Factor	Sum of squares	d.f.	Mean squares	F	P
Experiment (E)	0.116	1	0.116	16.55	< 0.001
Temperature (T)	0	1	0	0.01	0.925
Food type (F)	0.035	1	0.035	5.04	0.031
E × T	0.044	1	0.044	6.29	0.017
E × F	0.007	1	0.007	1.01	0.321
T × F	0.047	1	0.047	6.74	0.013
E × T × F	0.001	1	0.001	0.18	0.673



**Figure 3** Grazing rates (SE) of *Temora longicornis* on P-limited (C : P = 1100 : 1) and P-replete (C : P = 448 : 1) *R. salina*. A significant interaction between temperature and food quality was observed (Table 1).

showed a higher preference for algae than for heterotrophic prey. Thus the demand for nutrients is higher at lower temperatures, and/or the demand for carbon is higher at higher temperatures. These are difficult to separate, but we will argue that the latter is the more logical explanation. Furthermore the third experiment revealed a significant interaction between temperature and C : P ratio of the food on the grazing rates of the copepods, stressing the importance of the C : P ratio in determining the grazing rates.

Obviously, the correlative approach of the field survey is not without caveats. The first one is in the nature of the analysis. Correlative analyses can never infer causality, and it could well be that temperature is not the factor driving the change in  $\delta^{15}\text{N}$  signal, but other correlated factors such as the availability of nutrients (Aberle & Malzahn 2007). Unfortunately, there is no real way to resolve this. Moreover, trophic fractionation of stable nitrogen isotopes may be temperature dependent. The literature on the effect of temperature on stable isotope physiology gives conflicting information ranging from increased to decreased fractionation of stable nitrogen isotopes with temperature (Power *et al.* 2003; Barnes *et al.* 2007; Matthews & Mazumder 2008). For crustacean zooplankton we are aware of only two studies on the effects of temperature on the stable isotope physiology. Whereas Matthews & Mazumder (2008) reported no effects of temperature on the fractionation, Power *et al.* (2003) observed an increase of trophic fractionation of 0.1‰ per °C. The latter would result in a much steeper slope of the regression line in Fig. 2, and a much stronger dependency of the trophic level on the temperature. The third issue in studies like this one is the matter of finding the proper baseline. We do realise that filtering seston throughout a season does not give a clean algal (primary producer) baseline signal. Smaller heterotrophic organisms such as ciliates and flagellates are included in this fraction, but they are impossible to omit from the analysis as they are essentially the same size as the algae. To date, there is no good alternative for this, except the comparison of stable isotope signals of specific compounds, some of which are essential for herbivores, such as some amino acids, and within the herbivore tissue represent the food source signal, with those substances that undergo fractionation and represent the consumer signal (e.g. Chikaraishi *et al.* 2014). However, these techniques are still highly demanding and not suitable for routine samplings. Hence, the traditional way of filtering the seston fraction to obtain a base-line estimate is still very much in use (Hansson *et al.* 1997; Sommer & Sommer 2004; Gentsch *et al.* 2009; Guzzo *et al.* 2011; Kürten *et al.* 2013), and even if it does not resolve the absolute trophic position, the  $\delta^{15}\text{N}$  value gives a clear indication of the relative position of a consumer in the food web. We also realise that the trophic level of the *Temora* individuals reported here is sometimes low. This is caused by our choice of trophic fractionation factor for  $^{15}\text{N}$  of 3.4‰ per food level. In fact, reported values differ widely in the literature going as low as 0.1‰ per trophic level (D'Ambra *et al.* 2014). Obviously, this affects our estimates of the absolute trophic level, but using a different fraction factor would not have affected the pattern in the data at all. Of course, this trophic position does not say anything about what drives the change. If consumers were to

simply follow availability of food sources then they might be more herbivorous in summer when algae are more plentiful. In fact, we observed a significant negative correlation between total algal biomass and trophic level of *T. longicornis*, but the relationship of trophic level with temperature remained significant even after correcting for the effect of algal biomass.

Grazing experiments with more than one predator are notoriously difficult and usually show a lot of variation, potentially leading to several estimates of negative grazing rates (Löder *et al.* 2011). The corrections proposed by Nejstgaard *et al.* (2001) are useful in this respect, but they create an interdependence between the grazing rates on the different food sources, which is difficult to interpret. Moreover, they assume a constant grazing of the predator over the whole experimental period, which is not necessarily correct. Nevertheless, using these corrections we were able to show that the grazing rate of the copepod *T. longicornis* on the alga *R. salina* showed a significantly positive correlation with temperature, whereas the opposite was true for the grazing on *O. marina*, patterns that remain visible even when including all the data, or excluding all estimates of negative grazing rates (see Table S1 and Fig. S1). Especially the relationship of *T. longicornis* grazing on *O. marina* is very robust. From this we conclude that *T. longicornis* prefers 'meat' at lower temperatures and 'vegetables' at higher ones. This finding is consistent with the results from the field survey. Very few studies have experimentally investigated temperature related changes in feeding, and those that have do not yield a coherent picture. Whereas Sotka & Giddens (2009) reported an increase in feeding rates on low protein foods with temperature in at least a sub-set of the amphipods they investigated (with no change in the others), Lemoine *et al.* (2013), in their study on selective feeding in the beetle *Popillia japonica*, observed that at higher temperatures the beetles selected for plants with higher N-content. Unfortunately, in the latter study levels of allelopathy in the plants changed as well, so it is difficult to draw conclusions. Nevertheless, using the difference in nutrient content between different sources of prey, our study can provide a potential mechanistic explanation of the correlative evidence from the meta-analysis studies on latitudinal gradients of herbivory cited in the introduction.

Our results raise the question of whether the increase in feeding on lower quality plant material at higher temperatures is a luxury that generalist consumers can afford, thus broadening their food spectrum at higher temperatures, or a necessity to obtain the amount of energy necessary to pay for the increased metabolism. We suggest that is rather the latter. First, Castellani & Altunbaş (2014) reported that for the copepod *Temora longicornis* the respirational demand doubled from *c.* 5 to 10% of body carbon in the temperature range from 5 to 17.5 °C, whereas the change in egg production rate is much lower (Castellani & Altunbaş 2006). Under the assumption that respiration only requires carbon, and egg production both carbon and phosphorus this implies a higher carbon demand at higher temperatures, which can be met with the 25% higher uptake of carbon relative to phosphorus, something we indeed see in our last experiment. Moreover, Cross *et al.* (2015) argued that the threshold

elemental ratio (TER, Urabe & Watanabe 1992) should increase with temperature, even though they also state that there is a great need to measure the dependence of the TER on temperature. Traditionally, this TER indicates the ratio between carbon and a nutrient below which the nutrient N or P is no longer limiting, and potentially carbon becomes limiting (Anderson *et al.* 2005), and organisms need to excrete the nutrient. Growth should be maximal at the TER (Boersma & Elser 2006), thus animals at higher temperatures should eat more carbon relative to nutrients to maximise their growth rate.

### Implications

Our present study has three major implications. First, a methodological one: in any study on dietary preference of consumers it is very important to indicate the temperature at which the experiments were carried out, as the outcome of the experiments may be temperature dependent. Second, current changes in temperatures in many areas in the world may lead to a change in feeding preference of those organisms that have the initial dietary breadth to do so. Based on our results, we predict an increase in the overall incidence of herbivory. This may also cause a shift in species composition if some species are competitively superior herbivores while being less capable to feed on heterotrophs. Third, many water bodies in the world are undergoing re-oligotrophication, with decreasing inputs of N and P, in concert with an increase in the CO<sub>2</sub> availability. Where this might influence primary production, and increase the carbon to nutrient ratios of many primary producers, the effects of this apparent decrease in food quality might be counteracted by the increasing temperature, thus potentially leading to a higher secondary production.

### ACKNOWLEDGEMENTS

We thank Eike Gentsch and Susanna Knotz for collection and analysis of part of the samples, the captain and crew of the research vessel 'Aade' for providing us with samples, and Karen Wiltshire, Arne Malzahn and Nicole Aberle for useful discussions. Special thanks to Saskia Ohse for the technical support. This study was partly funded by Nippon Foundation (NF), through the Partnership for Observation of the Global Oceans (POGO). We extend our gratitude to Dr. F. Gerald Plumley, co-ordinator of NF-POGO Centre of Excellence Program. We gratefully acknowledge support by the German Science Foundation (DFG: AB 289/2-1) and the Helmholtz-Society (VH-NG-058). This study was further supported through the GLOBEC-Germany project (03F0320C) by the German Federal Ministry of Science and Education (BMBF).

### AUTHORSHIP

MB, BN and KLS conceived the study and collected data, KAM carried out one experiment, CLM and RMFS the others, MB wrote the first draft of the manuscript, and all authors contributed substantially to revisions.

## REFERENCES

- Aberle, N. & Malzahn, A.M. (2007). Interspecific and nutrient-dependent variations in stable isotope fractionation: experimental studies simulating pelagic multitrophic systems. *Oecologia*, 154, 291–303.
- Anderson, T.R., Hessen, D.O., Elser, J.J. & Urabe, J. (2005). Metabolic stoichiometry and the fate of excess carbon and nutrients in consumers. *Am. Nat.*, 165, 1–15.
- Barnes, C., Sweeting, C.J., Jennings, S., Barry, J.T. & Polunin, N.V.C. (2007). Effect of temperature and ration size on carbon and nitrogen stable isotope trophic fractionation. *Funct. Ecol.*, 21, 356–362.
- Behrens, M.D. & Lafferty, K.D. (2007). Temperature and diet effects on omnivorous fish performance: implications for the latitudinal diversity gradient in herbivorous fishes. *Can. J. Fish Aquat. Sci.*, 64, 867–873.
- Boersma, M. & Elser, J.J. (2006). Too much of a good thing: on stoichiometrically balanced diets and maximal growth. *Ecology*, 87, 1335–1340.
- Boersma, M., Wesche, A. & Hirche, H.-J. (2014). Predation of calanoid copepods on their own and other copepods' offspring. *Mar. Biol.*, 161, 733–743.
- Castellani, C. & Altunbaş, Y. (2006). Factors controlling the temporal dynamics of egg production in the copepod *Temora longicornis*. *Mar. Ecol. Prog. Ser.*, 308, 143–153.
- Castellani, C. & Altunbaş, Y. (2014). Seasonal change in acclimatised respiration rate of *Temora longicornis*. *Mar. Ecol. Prog. Ser.*, 500, 83–101.
- Chikaraishi, Y., Steffan, S.A., Ogawa, N.O., Ishikawa, N.F., Sasaki, Y., Tsuchiya, M. et al. (2014). High-resolution food webs based on nitrogen isotopic composition of amino acids. *Ecol. Evol.*, 4, 2423–2449.
- Clements, K.D., Raubenheimer, D. & Choat, J.H. (2009). Nutritional ecology of marine herbivorous fishes: ten years on. *Funct. Ecol.*, 23, 79–92.
- Cross, W.F., Hood, J.M., Benstead, J.P., Hury, A.D. & Nelson, D. (2015). Interactions between temperature and nutrients across levels of ecological organization. *Global Change Biol.*, 21, 1025–1040.
- Daines, S.J., Clark, J.R. & Lenton, T.M. (2014). Multiple environmental controls on phytoplankton growth strategies determine adaptive responses of the N : P ratio. *Ecol. Lett.*, 17, 414–425.
- D'Ambra, I., Carmichael, R. & Graham, W. (2014). Determination of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  and trophic fractionation in jellyfish: implications for food web ecology. *Mar. Biol.*, 161, 473–480.
- Dobberfuhl, D.R. & Elser, J.J. (2000). Elemental stoichiometry of lower food web components in arctic and temperate lakes. *J. Plankton Res.*, 22, 1341–1354.
- Elser, J.J., Dobberfuhl, D.R., Mackay, N.A. & Schampel, J.H. (1996). Organism size, life history, and N : P stoichiometry. *Bioscience*, 46, 674–684.
- Elser, J.J., O'Brien, W.J., Dobberfuhl, D.R. & Dowling, T.E. (2000). The evolution of ecosystem processes: growth rate and elemental stoichiometry of a key herbivore in temperate and arctic habitats. *J. Evol. Biol.*, 13, 845–853.
- Floeter, S.R., Behrens, M.D., Ferreira, C.E.L., Paddack, M.J. & Horn, M.H. (2005). Geographical gradients of marine herbivorous fishes: patterns and processes. *Mar. Biol.*, 147, 1435–1447.
- Forster, J., Hirst, A.G. & Woodward, G. (2011). Growth and development rates have different thermal responses. *Am. Nat.*, 178, 668–678.
- Forster, J., Hirst, A.G. & Atkinson, D. (2012). Warming-induced reductions in body size are greater in aquatic than terrestrial species. *Proc. Natl Acad. Sci. USA*, 109, 19310–19314.
- Frost, B. (1972). Effects of size and concentration of food particles on the feeding behavior of the marine planktonic copepod *Calanus pacificus*. *Limnol. Oceanogr.*, 17, 805–815.
- Gentsch, E., Kreibich, T., Hagen, W. & Niehoff, B. (2009). Dietary shifts in the copepod *Temora longicornis* during spring: evidence from stable isotope signatures, fatty acid biomarkers and feeding experiments. *J. Plankton Res.*, 31, 45–60.
- Gonzalez-Bergonzoni, I., Meerhoff, M., Davidson, T.A., Teixeira-de Mello, F., Baattrup-Pedersen, A. & Jeppesen, E. (2012). Meta-analysis shows a consistent and strong latitudinal pattern in fish omnivory across ecosystems. *Ecosystems*, 15, 492–503.
- Greve, W., Reiners, F., Nast, J. & Hoffmann, S. (2004). Helgoland Roads meso- and macrozooplankton time-series 1974 to 2004: lessons from 30 years of single spot, high frequency sampling at the only off-shore island of the North Sea. *Helgol. Mar. Res.*, 58, 274–288.
- Guzzo, M.M., Haffner, G.D., Sorge, S., Rush, S.A. & Fisk, A.T. (2011). Spatial and temporal variabilities of delta  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  within lower trophic levels of a large lake: implications for estimating trophic relationships of consumers. *Hydrobiologia*, 675, 41–53.
- Hansson, S., Hobbie, J.E., Elmgren, R., Larsson, U., Fry, B. & Johansson, S. (1997). The stable nitrogen isotope ratio as a marker of food-web interactions and fish migration. *Ecology*, 78, 2249–2257.
- Hantzsch, F.M. & Boersma, M. (2010). Dietary-induced responses in the phagotrophic flagellate *Oxyrrhis marina*. *Mar. Biol.*, 157, 1641–1651.
- Heinbokel, J.F. (1978). Studies on the functional role of tintinnids in the Southern California Bight. I. Grazing and growth rates in laboratory cultures. *Mar. Biol.*, 47, 177–189.
- Karl, I. & Fischer, K. (2008). Why get big in the cold? Towards a solution to a life-history puzzle. *Oecologia*, 155, 215–225.
- Kürten, B., Painting, S.J., Struck, U., Polunin, N.V.C. & Middelburg, J.J. (2013). Tracking seasonal changes in North Sea zooplankton trophic dynamics using stable isotopes. *Biogeochemistry*, 113, 167–187.
- Larsson, S. & Berglund, I. (2005). The effect of temperature on the energetic growth efficiency of Arctic charr (*Salvelinus alpinus* L.) from four Swedish populations. *J. Therm. Biol.*, 30, 29–36.
- Lemoine, N.P., Drews, W.A., Burkepile, D.E. & Parker, J.D. (2013). Increased temperature alters feeding behavior of a generalist herbivore. *Oikos*, 122, 1669–1678.
- Löder, M.G.J., Meunier, C., Wiltshire, K.H., Boersma, M. & Aberle, N. (2011). The role of ciliates, heterotrophic dinoflagellates and copepods in structuring spring plankton communities at Helgoland Roads, North Sea. *Mar. Biol.*, 158, 1551–1580.
- Matthews, B. & Mazumder, A. (2008). Detecting trophic-level variation in consumer assemblages. *Freshwat. Biol.*, 53, 1942–1953.
- Meunier, C.L., Hantzsch, F.M., Cunha-Dupont, A.Ö., Haafke, J., Oppermann, B., Malzahn, A.M. et al. (2011). Intraspecific selectivity, compensatory feeding and flexible homeostasis in the phagotrophic flagellate *Oxyrrhis marina*: three ways to handle food quality fluctuations. *Hydrobiologia*, 680, 53–62.
- Meunier, C.L., Schulz, K., Boersma, M. & Malzahn, A.M. (2013). Impact of swimming behaviour and nutrient limitation on predator-prey interactions in pelagic microbial food webs. *J. Exp. Mar. Biol. Ecol.*, 446, 29–35.
- Meunier, C.L., Boersma, M., Wiltshire, K.H. & Malzahn, A.M. (2015). Zooplankton eat what they need: copepod selective feeding and potential consequences for marine systems. *Oikos*, doi: 10.1111/oik.02072.
- Minagawa, M. & Wada, E. (1984). Stepwise enrichment of  $^{15}\text{N}$  along food-chains – further evidence and the relation between  $\delta^{15}\text{N}$  and animal age. *Geochim. Cosmochim. Acta*, 48, 1135–1140.
- Nejstgaard, J.C., Naustvoll, L.-J. & Sazhin, A. (2001). Correcting for underestimation of microzooplankton grazing in bottle incubation experiments with mesozooplankton. *Mar. Ecol. Prog. Ser.*, 221, 59–75.
- Persson, J., Wojewodzic, M., Hessen, D. & Andersen, T. (2011). Increased risk of phosphorus limitation at higher temperatures for *Daphnia magna*. *Oecologia*, 165, 123–129.
- Power, M., Guiguer, K. & Barton, D.R. (2003). Effects of temperature on isotopic enrichment in *Daphnia magna*: implications for aquatic food-web studies. *Rapid Commun. Mass Spectrom.*, 17, 1619–1625.
- Sommer, F. & Sommer, U. (2004).  $\delta^{15}\text{N}$  signatures of marine mesozooplankton and seston size fractions in Kiel Fjord. *Baltic Sea. J. Plankton Res.*, 26, 495–500.
- Sotka, E.E. & Giddens, H. (2009). Seawater Temperature Alters Feeding Discrimination by Cold-Temperate but not Subtropical Individuals of an Ectothermic Herbivore. *Biol. Bull.*, 216, 75–84.

- Sterner, R.W. & Elser, J.J. (2002). *Ecological Stoichiometry: The Biology of Elements from Molecules to the Biosphere*. Princeton University Press, Princeton, NJ.
- Thomas, W.H., Scotten, H.L. & Bradshaw, J.S. (1963). Thermal gradient incubators for small aquatic organisms. *Limnol. Oceanogr.*, 8, 357–360.
- Urabe, J. & Watanabe, Y. (1992). Possibility of N-limitation or P-limitation for planktonic cladocerans – An experimental test. *Limnol. Oceanogr.*, 37, 244–251.
- Vander Zanden, M.J. & Rasmussen, J.B. (2001). Variation in  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  trophic fractionation: implications for aquatic food web studies. *Limnol. Oceanogr.*, 46, 2061–2066.
- van de Waal, D.B., Verschoor, A.M., Verspagen, J.M.H., van Donk, E. & Huisman, J. (2009). Climate-driven changes in the ecological stoichiometry of aquatic ecosystems. *Front. Ecol. Environ.*, 8, 145–152.

- Wiltshire, K.H. & Manly, B.F.J. (2004). The warming trend at Helgoland Roads, North Sea: Phytoplankton response. *Helgol. Mar. Res.*, 58, 269–273.

#### SUPPORTING INFORMATION

Additional Supporting Information may be downloaded via the online version of this article at Wiley Online Library ([www.ecologyletters.com](http://www.ecologyletters.com)).

Editor, Gregor Fussmann  
Manuscript received 2 October 2015  
Manuscript accepted 5 October 2015





#### **4 SYNOPTIC DISCUSSION**

This thesis provides a characterization of the laboratory performance for two members of the zooplankton from Helgoland, females of the calanoid copepod *Temora longicornis* and meroplanktonic larvae of the polychaete *Lanice conchilega*, in relation to food quality. An extensive dataset on physiological parameters is provided for the copepods in Chapters I and II, and includes rates of ingestion, respiration, excretion, egestion, growth and reproduction. The efficiency with which these processes occur is also provided, and a budget with C and N values for the rates is constructed in order to evaluate copepod performance. In addition, the effects of food quality on carbon assimilation and turnover rates in the copepod are also evaluated (Chapter II). It was not possible to obtain such a qualitative and quantitative dataset for the polychaete larvae. Nevertheless its carbon assimilation and turnover rates while feeding on nutrient-replete diatom cultures are reported (Chapter III). The work on lipid C assimilation in the copepods and polychaete larvae, together with data available from the literature, was also used to propose the hypothesis that lipid homeostasis depends upon the total lipid content of zooplanktonic organisms (Chapter III).

The synoptic discussion is divided into three parts. Section 4.1 provides a thorough evaluation of the CSIA method used in the experiments conducted during this PhD, with a special focus on methodological problems which need to be resolved in future studies. Section 4.2 provides an evaluation of the Ecological Stoichiometry framework, which was used in this PhD to study the impacts of diet quality on the physiology of the zooplankton. It discusses what are the applications and limitations of this framework in relation to the results obtained with experiments conducted with mero- and holoplanktonic species. Section 4.3 presents a description of how communities have been impacted by climate changes and anthropogenic activities in diverse marine ecosystems across all oceans. It continues by describing how distinct geographic regions within the North Sea have been affected differently in the past decades. It further focuses on regime shifts in the area surrounding the German island of Helgoland, from where the organisms used in the experiments performed for this PhD were sampled. Experimental results are then used to infer about food quality effects on copepod performance, and on how this group might fare in future scenarios of changing food regimes.

### 4.1 ADVANTAGES AND CAVEATS OF USING COMPOUND-SPECIFIC STABLE ISOTOPE ANALYSIS

Trophic ecology has long sought to determine the relationships between prey and predator. Several techniques can be used to determine the diet of a consumer, with varying degrees of invasiveness and lethality. Visual observation was probably the first method employed, followed by analysis of blood, regurgitated dietary material, faecal matter, and of gut contents. All of these conventional methods, however, can only provide a snapshot of the diet of a consumer, the most recent items it fed upon (Michener and Kaufman, 2007; Newton, 2010). Stable isotope analysis (SIA), on the other hand, offers the possibility of diet integration over a longer period of time, while also providing information on the source of a food item and on consumer trophic level (Dalsgaard et al., 2003; Michener and Kaufman, 2007; Newton, 2010). Stable isotope ratios were first applied to studies in the field of earth sciences, but over the past 30 years ecologists have sought it to better understand interactions within species and between them and their environment (Newton, 2010). Their great usefulness for many areas arises from changes in the ratio of lighter and heavier isotopes in a compound during physical and chemical processes (Newton, 2010). Of particular use to ecological studies are the isotopes of the light elements H, C, N, O, and S. In this section I mainly discuss about the use of the  $^{13}\text{C}/^{12}\text{C}$  ratio, as this was the ratio I analyzed in my experiments, but several pertinent questions in ecology can be answered by using ratios of the other elements.

The  $^{13}\text{C}$  SIA of bulk (or aliquot) samples (BSIA) is commonly used to identify the possible food items (or foraging area) ingested by a consumer (e.g.: Chen et al., 2018; Cherel et al., 2006; Gorbatenko et al., 2014), based upon the knowledge that different photosynthetic pathways in (C3 and C4) plants result in differing  $^{13}\text{C}$  signatures (first observed by Smith and Epstein, 1971). It is also possible to estimate tissue-specific turnover rates in organisms with BSIA, though this involves multiple sampling and the use of two or more isotopically disparate food sources in laboratory experiments (Newton, 2010). In the past, the determination of stable isotope ratios was limited to BSIA (Lichtfouse, 2000). The analysis of ratios for individual compounds within complex molecules (CSIA) was only possible with the commercial production of the gas chromatograph/combustion/isotope ratio mass spectrometer (GC/C/IRMS) in the early 90's (Evershed et al., 2007; Meier-Augenstein, 1999, 2002). There are several advantages to using

CSIA over BSIA, though these depend upon the goal of an investigation. In terms of tracing the use of resources by consumers in complex marine systems, for example, BSIA cannot provide information on the isotopic signature of biochemical components, whether they are different or identical but belong to different pools (Evershed et al., 2007). As these compounds may have different functions, they may also reflect different turnovers of dietary material, i.e., provide information on temporal changes in diet. In a study on the use of food resources for reproduction in butterflies, for example, it was shown that the nonessential AAs in the eggs were derived from nectar sugar (adult diet), whereas essential AAs were obtained solely from the larval diet (O'Brien et al., 2004). In the case of a key marine copepod, *Acartia* spp., the use of CSIA of AA  $^{13}\text{C}$  allowed for the identification of a selective feeding behavior which varied seasonally (Nielsen and Winder, 2015). In general, and due to the nature of the analysis itself, CSIA will provide a greater wealth of information than BSIA (specified in Evershed et al., 2007), which may be helpful in disentangling ecological questions.

CSIA is a methodology that provides accurate and precise (up to 0.3‰ or 0.0003 atom percent excess, APE) measurements of the isotopic composition of compounds (Meier-Augenstein, 2002). It is a powerful tool for tracing the origin and fate of organic matter and to investigate biochemical processes and patterns in individuals and ecosystems (Evershed et al., 2007; Guilini et al., 2010; Meier-Augenstein, 2002; Rix et al., 2018; Wegener et al., 2012). The applications of CSIA are as broad as the fields that have used it and include, a. o., forensics, environmental science, archaeology, food science, and doping control (Benson et al., 2006; Fogel and Tuross, 2003; Lichtfouse, 2000; Ogrine et al., 2003; Schmidt et al., 2004; Simpson et al., 1999; Tobias and Brenna, 2018). There has been an increase in the use of CSIA in tracer studies, especially those aiming to quantify biochemical processes such as assimilation, turnover, and synthesis of important molecules (Boissonnot et al., 2016; Budge et al., 2008; De Troch et al., 2012; Graeve et al., 2005; Kluijver et al., 2013; Middelburg et al., 2000). This was also the case in this PhD work. Before the advent of CSIA, such metabolic studies were only possible, if at all, by using stable isotope tracers at high enrichment levels, which were analyzed with a simple GC-MS, or by using radioactive tracers (Meier-Augenstein, 1999).

Within the context of this PhD, BSIA would have certainly been advantageous time-wise, as the sample preparations prior to data acquisition runs are simpler and faster for this technique than

## SYNOPTIC DISCUSSION

---

for CSIA. The remarkably smaller data output obtained from a single run/sample analyzed with BSIA would also have made the interpretation of results a much simpler process than that required for the multiple dataset returned from a single run/sample analyzed with CSIA. This PhD project was designed with the goal to better understand the performance of zooplanktonic organisms in relation to diet nutritional quality. One of the specific goals was to obtain quantitative data on the transfer of dietary C from prey to predator, but I was also interested in investigating FA-specific assimilation, biosynthesis and bioconversion due to the importance of certain FAs in the reproductive process in copepods (Jónasdóttir et al., 2009; Thor et al., 2007). This type of information could only be achieved with CSIA.

### *Methodological constraints of CSIA in this PhD*

The use of CSIA in tracer studies is still at its early stages and variables such as the duration of exposure of target species to  $^{13}\text{C}$  and the amount of label necessary to obtain robust results are still being fine-tuned. High levels of isotopic enrichment are not desirable, as they cannot be properly detected by the GC-c-IRMS (M Graeve, pers. obs.). Low concentrations of  $\text{NaH}^{13}\text{CO}_3$  which still enable correct and reproducible estimations of C assimilation and turnover should thus be favored, as they also minimize research costs associated with the use of the bicarbonate powder. The concentration of  $\text{NaH}^{13}\text{CO}_3$  used to enrich *C. weissflogii* cultures, for example, has decreased over the years from  $200 \text{ mg L}^{-1}$  (Graeve et al., 2005) to  $15 \text{ mg L}^{-1}$  (Boissonnot et al., 2016). For the experiments conducted during this PhD, I used, for the same diatom species, a concentration of  $4 \text{ mg L}^{-1}$ . Although the atom% values that I obtained for these cultures (and for the copepods that fed on them) were low, they were above the natural concentrations (1.08 atom% or  $-30\text{‰ }^{13}\delta\text{C}$ ). This enabled me to reliably quantify the C assimilation and turnover rates in the copepods (Chapter II) and polychaetes (Chapter III) fed with the diatom diets. *Oxyrrhis marina*, the other prey item used to feed copepods and polychaetes, is a heterotrophic dinoflagellate, and had to be indirectly labelled via feeding on isotopically-enriched *R. salina*. The concentration of  $^{13}\text{C}$  I used to enrich *R. salina*,  $12 \text{ mg L}^{-1}$ , did not seem to be enough for a transfer of adequate labelling to the next trophic level in one of the *O. marina* cultures. Future studies that attempt to use this species to indirectly label non-autotrophic organisms should run pre-experiment tests with enrichment of both trophic levels in order to determine the quantity of  $^{13}\text{C}$  necessary to achieve their goals.

Labelling in the dinoflagellate cultures differed, as those fed with N-depleted (enriched) *R. salina* (Dino- cultures) showed isotopic composition above the natural concentrations and those fed with nutrient-replete *R. salina* had isotopic composition values at the edge of valid labelling. The data I gathered from the experiments cannot fully explain this result. It was not caused by poor labelling of nutrient-replete *R. salina*, as these cultures enabled the enrichment, though low, of the Dino+ culture and were also properly labelled in another experiment I conducted during the PhD (atom% =  $1.24 \pm 0.12$ , not mentioned in this thesis). Absence of feeding of *O. marina* on the nutrient-replete *R. salina* cultures was also not a problem, as suggested by a previous study (Hantzsche & Boersma, 2010) and confirmed by the flow cytometer readings (not shown in this thesis). It would be possible that Dino+ fed only partially on *R. salina* and complemented its nutrition via cannibalism (thus diluting its  $^{13}\text{C}$  enrichment), though there is evidence that only a few cells within an *O. marina* culture are cannibalistic (Martel and Flynn, 2008). The variation in  $\delta^{13}\text{C}$  and the atom% values were clearly higher in the Dino- cultures (339-1174‰ and  $> 1.47$ , respectively) than in the Dino+ (7-197‰ and  $> 1.09$ , respectively). This indicates that the difference in isotopic enrichment between the dinoflagellate cultures might have been caused by a differential labelling of *R. salina* cells when they were cultured under different nutrient conditions. Different growth and environmental conditions can trigger different pathways for C uptake by phytoplankton cells (Eek et al., 1999), which will result in varying isotopic composition of cells (Riebesell et al., 2000). Since nutrient supply can regulate phytoplankton growth under saturated light conditions, nutrient concentration should indirectly affect C isotopic fractionation. It would seem that when algae are deprived of N during growth, carbohydrates and lipids are accumulated at the expense of AAs, a shift in the flux of C which is accompanied by a lower isotopic fractionation and, thus, results in higher isotopic signal (Eek et al., 1999). The difference in labelling of *O. marina* cultures thus seems to have derived from a differential uptake of  $^{13}\text{C}$  by *R. salina* cells in relation to nutrient availability during growth. The results I recorded for isotopic composition of both *C. weissflogii* and *R. salina* cells follow this pattern, with cells grown under N-depleted conditions showing higher isotopic signal than those grown under nutrient-replete conditions. Future tracer studies using phytoplankton cultures with different nutrient concentrations should adjust the amount of labelling added in order to account for fractionation differences.

## ***SYNOPTIC DISCUSSION***

---

The low level of labelling transfer from *R. salina* to *O. marina* in the Dino+ cultures made it difficult to follow the transfer of labelled FAs from the dinoflagellates into the copepods they were fed to. The APE of copepods fed with the diatom and Dino- cultures accounted for approximately 25-35% of the atom% of their respective diets, whereas the APE of those fed with the Dino+ cultures accounted for only 7-8%. Because the labelling of the Dino+ cultures was so low to begin with, it is not possible to affirm with certainty whether the low APE and C assimilation and turnover values obtained for copepods fed with this diet were due to methodological problems, biological processes, or both. There is, however, indication that it was due to the low labelling of the Dino+ cultures. The total C assimilation values recorded from copepods feeding on labelled diets should always be equal to or higher than the increase in total lipid C. These two values were similar in copepods fed with the diatom and Dino-, but the C assimilation of copepods fed with Dino+ accounted for only 15-26% of the recorded increase in total lipid C. There is no plausible explanation for an increase in lipid C that would not be recorded as C assimilated in animals feeding on isotopically-enriched food. During this PhD I have observed that *T. longicornis* not only feeds normally on nutrient-replete *O. marina*, but it also thrives on this diet (highest ingestion, growth and egg production rates; Chapter I). This led me to conclude that, in the case of copepods fed with Dino+, the C assimilation values were highly underestimated due to methodological problems (i.e., with the isotopic enrichment of *R. salina*).

This methodological problem was the source of some academic frustration. As can be seen from Chapter I, the copepods fed on the Dino+ diet were regarded as displaying the best performance, and the FA-specific C assimilation and turnover data for this treatment could have offered many insights into why that was the case. Moreover, the inability to properly label *O. marina* also affected the experiments with the polychaete larvae. Originally, I would have been able to (a) verify whether the larvae can feed on dinoflagellates, for which no record exists, and (b) assuming it does, compare their performance on different diets. Due to the labelling problem, I was only able to investigate lipid assimilation and utilization in *L. conchilega* larvae while feeding on diatoms (Chapter III). Despite the problems, Chapters II and III are solid contributions to the field of trophic ecology, as there are few studies that have followed lipid transfer between copepods and their prey and no information in the literature regarding the feeding ecology of marine polychaete larvae. The results of this thesis also help to further

advance the establishment of guidelines for the experimental labelling of organisms in tracer studies in food webs.

**Take-home messages for Section 4.1**


- CSIA provides accurate measurements of the isotopic composition in specific compounds within molecules. Combined with the use of  $^{13}\text{C}$  enrichment in laboratory experiments, it enables the quantification of biochemical processes such as lipid C transfer and FA assimilation, turnover, and synthesis in organisms.
- Species-specific protocols for using CSIA in tracer studies need to be fine-tuned, as variables such as the duration of exposure of target species to  $^{13}\text{C}$  and the amount of label necessary to obtain robust results are still being tested.
- The uptake of C by algae and, thus, their enrichment with  $^{13}\text{C}$ , is related to photosynthetic pathways and varies according to culture conditions. As the effects of the latter on  $^{13}\text{C}$  labelling are not well described at present, it is recommended that the successful labelling of algae with  $^{13}\text{C}$  be verified prior to their use in laboratory tracer experiments.
- Tracer studies which investigate lipid transfer between three or more trophic levels should make sure that the  $^{13}\text{C}$  enrichment of the lowest trophic level is high enough to be transferred in measurable and differentiable quantities to the highest trophic level.

#### 4.2 ECOLOGICAL STOICHIOMETRY AND HOMEOSTASIS AS TOOLS TO STUDY ZOOPLANKTON PHYSIOLOGY

Zooplankton physiology, and how it may be affected directly and indirectly by climate change, can be studied from the perspective of several nutritional frameworks, such as metabolic theory, ecological stoichiometry (ES), and nutritional geometry. ES has been presented as a useful tool for investigating environmental effects on the stoichiometry of organisms and, consequently, the flow of elements through different levels of biological organization and, ultimately, the cycling of nutrients and elements at the ecosystem level (Elser et al. 1988; Sterner & Elser 2002). For the work conducted during this thesis, the physiology of two members of the zooplankton was investigated under the framework of ES.

An important concept to ES is that of homeostasis - the ability of some organisms to maintain a constant body chemical composition despite changes in the chemical composition of their environment and, more specifically, their food (Kooijman, 1995). If the assimilation of nutrients by a consumer occurs in identical proportions to their relative content in the prey, then the consumer is nonhomeostatic – the concept of “you are what you eat” (Fig. 8a). Organisms which assimilate nutrients in constant proportions to their relative dietary abundance are also nonhomeostatic (Fig. 8a), as they do not adjust their elemental composition in response to that of their prey, but they no longer are what they eat. Homeostatic regulation occurs as a feedback (in the opposite direction to that of the disruption) between the composition of the consumer and that of its prey, such that in strict homeostasis there is no variation in the former in relation to the latter (Sterner and Elser, 2002; Fig. 8b). Strict homeostasis can also occur within certain physiological pools in the organism (structure vs. energy reserves) but the relative proportions of the pools may vary with time (Kooijman 1995; Sterner and Elser, 2002). Organisms will usually fall somewhere between nonhomeostatic and strictly homeostatic, with autotrophs being usually less homeostatic than heterotrophs (Sterner and Elser, 2002). The degree of homeostasis will vary in relation to several parameters, among which the most important are likely to be the species of consumer, its life stage, the element/molecule composition of its food source, and the comprehensiveness of the dataset (Sterner and Elser, 2002).





**Figure 8:** Stoichiometric patterns which relate the stoichiometry of consumers and resources. Horizontal and vertical axes can be any type of stoichiometric measure. If the stoichiometry of a consumer always matches that of the resources it feeds upon, then their relationship is indicated by the points in the line with slope 1 and intercept 0 (“you are what you eat: no homeostasis”). The solid diagonal lines in A) represent instances when the consumer performs constant differential nutrition retention (albeit nonhomeostatic). The solid horizontal lines (slope 0, intercept  $> 0$ ) in B) characterize strictly homeostasis. Source: Sterner and Elser, 2002.

Consumers can react to the availability of prey items of differing elemental composition by changing feeding rates, food selection, production efficiency, and growth rate (Sterner and Elser, 2002). Homeostatic regulation of consumer stoichiometry can occur before and after the ingestion of nutritionally unbalanced food (i. e., prey with a different molar C:N ratio to that of the consumer) (Siuda and Dam, 2010; Sterner and Elser, 2002). Pre-ingestion regulation is related to consumer behavior, which can select higher quality prey (Jones and Flynn, 2005; Kiørboe et al., 1996) or increase ingestion of low-quality prey (compensatory feeding) (Augustin and Boersma, 2006; Plath and Boersma, 2001; Siuda and Dam, 2010). Post-ingestion regulation can be achieved through physiological processes. For individuals feeding on a nutrient-limited diet, ES predicts that consumer stoichiometry will be regulated by one or more of the following processes: (a) increase in assimilation of the limiting nutrient; (b) increase in waste production from the excess nutrient; (c) conservation of the limiting nutrient (i.e., decrease in nutrient loss) through altered biochemical pathways (Sterner and Elser, 2002). Furthermore, ES also stipulates that stoichiometric imbalance between prey and predator will negatively affect consumer growth, and that the more strictly homeostatic an organism is, the greater the effects of a nutritionally unbalanced diet on its elemental gross growth efficiency (GGE) (Sterner and Elser, 2002).

## SYNOPTIC DISCUSSION

---


Over the years several studies have employed concepts from ES to assess performance in various copepod species, some obtaining results which agree with the predictions from ES and some with contrasting findings. When studying the effects of N-limitation on the calanoid copepod *Acartia tonsa*, Jones et al. (2002) verified that the GGE of N was greater than that of C, recording a more efficient use for the limiting element. The authors also recorded higher N use efficiency (relative to C use efficiency) for copepods fed with N-limited prey than for those preying upon N-replete food, and greater C waste associated with higher food molar C:N ratio, as would be expected from ES. Checkley and Entzeroth (1985) also reported a higher efficiency in the use of N than in the use of C for tissue production in copepods. On the other hand, higher N excretion rates were reported for carnivores in comparison to herbivores (despite the more nutrient-rich diet of the former in comparison to the latter), contradicting what would be expected from ES (Vanni and McIntyre, 2016). In general, a failure in the explanation of theoretical (model) or practical (laboratory) results by the assumptions of ES has related by the available literature to the possibility that parameters other than elemental composition might be playing a part in limiting consumer growth (Anderson and Hessen, 1995; Anderson and Pond, 2000; Jones and Flynn, 2005).

### *ES framework for the study of performance in adult females of T. longicornis*

This PhD project was interested in relating the nutritional quality of prey items with the performance of *T. longicornis* by analyzing the effect of the former over each physiological process separately (Chapter I). Copepods were fed with four different diets, two of which consisting of diatoms and two of dinoflagellates, which were further divided into nutrient-replete and N-depleted growth conditions. Diet acclimation was conducted with copepods for 48h, after which they individuals were transferred to different incubation units for 24h to measure rates of ingestion, respiration, excretion, and egg and faecal pellet production. Respiration and excretion incubations were conducted without the addition of food to experimental units, whereas prey was available to copepods during grazing and EPR/FPR incubations.

The degree of homeostasis of an organism can be inferred from the plot in log scale between resource and consumer stoichiometry by calculating the inverse of the slope of the regression line. In Figure 8a, for example, it would be 1/1 for the nonhomeostatic organism. A degree of 7.7, for example, indicates strong homeostasis in a consumer (Sterner and Elser, 2002). The log

plot of resource vs. consumer stoichiometry obtained during this PhD for all *T. longicornis* samples and their respective diets (1/slope = 9.97) indicates that the adult females of this species are strictly homeostatic, at least for the range of prey molar C:N ratios investigated (Fig. 9b). This result further demands the analysis of the physiological measurements recorded during this PhD in relation to the assumptions of ES listed a few paragraphs above. This approach can indicate the appropriateness of using this framework to better understand copepod performance in relation to changing food regimes.



**Figure 9:** Resource vs. consumer stoichiometry plot obtained from samples collected during this PhD. Copepod values include adult male and female *T. longicornis*; diet values include *in situ* seston and the different diets used in the PhD experiments. A shows the molar C:N ratio as stoichiometric measure; B shows the same values under a logarithmic scale, as the slope of the regression line is used to calculate the level of homeostasis of the copepod.

Was it possible to observe an increase in the assimilation of the limiting nutrient, N, in the copepods? It is important to establish here the distinction between absorption and assimilation, which can be found in the literature with similar definitions despite their representing different processes. Absorption occurs after food ingestion and egestion by the individual and includes the energy which will be used for metabolism (in respiration and excretion) and growth (somatic and reproductive). Food assimilation only refers to the energy that is incorporated into growth. The absorption efficiency for N ( $AE_N$ , Chapter I) was fairly similar between copepods fed with different diets (60-73%). The N assimilation efficiency, which was not directly measured but can

## ***SYNOPTIC DISCUSSION***

---

be inferred from the sum of the net somatic growth ( $NSGE_N$ , Chapter I) and egg production ( $NEPE_N$ , Chapter I) efficiencies for N, was 2-3 times lower in the copepods fed with the N-limited diatoms (8-25%). These results contradict ES by reporting a decrease in the assimilation of N by copepods fed with a N-poor diet, and also suggest that most of the N absorbed by these copepods was used for maintenance and metabolism rather than for growth.

Did the waste production of C in copepods increase with the increase in the molar C:N ratio of the diets? If only the diatom diets are compared, then indeed the percentage of body C that was eliminated via FP production and via respiration was higher in copepods fed with N-limited (29% body C) than with nutrient-replete (20% body C) diatoms. The greater imbalance in molar C:N ratio between copepods and the N-limited diatoms did not result in higher waste production of C when rates are compared with those from copepods feeding on the dinoflagellate treatments (30-37% body C). It is interesting to note that, contrary to what is commonly assumed, the main pathway for eliminating excess C in the experiments conducted for this PhD was found to be FP production rather than respiration.

Were the copepods feeding on N-limited prey able to conserve their N reserves through altered biochemical pathways? It seems they were not. As a matter of fact, the lowest N excretion rates were recorded for copepods fed with nutrient-replete diets (5-8% body N), whereas the highest rate was measured for the copepods feeding on N-limited diatoms (27% body N). It has been suggested that the level of N excretion depends not only on the molar C:N ratios of prey and predator, but also on food intake, the energetic demands of respiration, and on the nature of substrates that were respired (Anderson, 1992). Given the high N excretion rates of the copepods fed with the Diat- diet, it would appear that the homeostatic regulation required to offset the nutritional imbalance of this diet had high energetic costs, and that nitrogenous substrates (amino acids and proteins) were the ones respired by copepods. The copepods fed with the Dino- diet also display high N excretion rates (15% body N) in relation to the nutrient-replete diets, despite the fairly similar molar C:N ratio between copepods and dinoflagellates in this diet. This indicates that N excretion and C respiration are not only linked to the availability of these elements in the diet, but to the biochemical molecules in which they are embedded.

And last but not least, was consumer growth negatively affected by stoichiometric imbalance between prey and predator? Growth can be divided into two components for adult female

copepods, somatic growth and reproductive growth. The results from this PhD indicate that both forms of growth were negatively affected (to various degrees) by the stoichiometric imbalance between copepods and the diet with the highest molar C:N ratio, Diat–, with the exception for the C net egg production efficiency ( $NEPE_C$ ), which was similar between all treatments. The  $NSGE_N$  seems to have been affected the most by prey-predator nutritional imbalance, and was irrelevant in copepods fed with the Diat– diet (-1% body N) and between 25-36% body N in copepods fed with the other diets.  $NEPE_N$  in copepods fed with Diat– was approximately half that of copepods fed with nutrient-replete diets (48% body N), and the C net somatic growth efficiency ( $NSGE_C$ ) in the former (20% body C) was approximately 65% of that recorded for copepods fed with the other diets.

From the ES assumptions mentioned above, only one was supported by the results from this PhD, that somatic and reproductive growth (both in terms of C and N) are negatively affected by increasing nutritional imbalance between prey and predator. The experimental results, however, mostly contradict ES assumptions, as the increase in nutritional imbalance between prey and predator did not lead to higher waste production from the excess element and resulted in a decrease in assimilation and in an increase in excretion of the limiting nutrient. Within the context of this PhD copepod performance was ultimately defined by its reproductive success. For this purpose, ES was a useful tool, as it was able to explain the differences observed between treatments in the context of varying food quality. It is evident however from the other results that ES may need a complimentary approach when other research goals are in play.

In their book about ecological stoichiometry, Sterner and Elser (2002) present the subject as a set of simple rules underlying ecological and biological complexity, capable of linking themes such as biochemical composition of subcellular structures, food web dynamics, and gene organization. The intention of being a unifying concept linking several scales of ecological patterns and processes has not been fully achieved by ES (Welti et al., 2017). Although it is necessary to understand prey-predator interactions in terms of the chemical nature of life, the authors themselves (and others) acknowledge the somewhat reductionist approach (Mitra and Flynn, 2005; Sperfeld et al., 2017; Sterner and Elser, 2002; Tang and Dam, 1999), as food substrates exist as compounds rather than as individual elements. It seems the ES approach would be better taken advantage of if used jointly with other approaches to nutritional ecology. Ecological

## SYNOPTIC DISCUSSION


---

stoichiometry should also consider, for example, the importance of biochemical compounds, or integrate concepts from theories such as trait-based ecology and metabolic theory of ecology (as also suggested by Anderson and Pond, 2000; Anderson et al., 2004; Meunier et al., 2017; Moe et al., 2005; Vanni and McIntyre, 2016; Ventura, 2006; a.o.).

### *Biochemical (lipid) homeostasis patterns in zooplankton*

Homeostasis has been said to be an ecological and evolutionary force (Sterner and Elser, 2002). From early to later prokaryotes, which were followed by unicellular and multicellular eukaryotes, respectively, homeostatic ability increased with the evolutionary development of ion pumps and the increase in cellular compartmentalization and biochemical complexity (Williams and Fraústo da Silva, 1996). But homeostatic regulation has an associated energetic cost and, as all activities which use up energy, will affect individual fitness (Sterner and Elser, 2002). What are the benefits of homeostatic regulation that offset this extra energy expenditure?

Chapter III provides one potential answer to this question. While studying the assimilation of dietary C into FAs of the meroplanktonic larvae of *L. conchilega* it was possible to observe that the FA profile of the individuals did not resemble that of the food items, as is common for organisms with intermediate total lipid contents, such as small copepods. On the contrary, *L. conchilega* larvae were able to selectively accumulate and/or metabolize certain FAs, thus regulating their lipid composition. FA homeostasis may be energetically simple when an organism feeds upon a prey of similar lipid composition. Just like elemental homeostasis, however, it will require that energy be allocated to bioconversion if prey items have a different biochemical make-up than that of the consumer. *Lanice conchilega* larvae, as most lecithotrophic and planktotrophic meroplanktonic larvae, are lipid-poor, as they spend most energy ingested to sustain metabolism and to grow into juveniles. These lipid-poor organisms likely have specific growth requirements (in terms of FAs) to sustain body function, so FA homeostasis is their solution to obtaining the nutrition they need regardless of the FA availability in the diet. One of the FAs *L. conchilega* was able to bioconvert, for example, was 22:5(n-3), an essential FA for consumers due to its role in membrane formation. This FA was not present in the diet with which the polychaete was fed, and had it not been able to bioconvert it from other precursor FAs its development and growth might have been physiologically compromised.



**Figure 10:** Lipid homeostasis patterns and associated total lipid content characteristics in zooplanktonic organisms. Sources of images: Jónasdóttir, 2017; Pérez-Losada et al., 2009; SERTC, 2018; Stübing, 2004; Vidal et al., 2010; WoRMS, 2018.

Based on these PhD results and on data available in the literature (Boissonnot et al., 2016; Graeve et al., 2005; Lee et al., 2006 and references therein; other references shown in Table 3 from Chapter III) it was possible to establish a hypothesis regarding patterns of lipid homeostasis in three different groups within the zooplankton (Fig. 10). Lipid-poor meroplanktonic larvae (with the exception of those from hermatypic and soft coral) have specific FA requirements to sustain body functions, and will thus display lipid homeostasis. *Lanice conchilega* larvae are an example of an organism with this strategy, as described above. Holoplanktonic organisms which have intermediate total lipid contents, such as small calanoid copepods like *T. longicornis*, have a modest ability to form lipid reserves in order to maximize reproductive output. These organisms are likely nonhomeostatic and assimilate dietary FAs in an unmodified manner. Lipid-rich holoplanktonic organisms, such as large polar calanoid copepods, need to store energy in the

## ***SYNOPTIC DISCUSSION***

---

most dense and effective way for overwintering and buoyancy purposes. These organisms are semi-homeostatic, as they will bioconvert dietary FAs into long chain MUFAs / wax esters.

It is also important to highlight that the majority of information on zooplankton feeding ecology in the North Sea (and in other areas) is available for copepods. Although they dominate the zooplankton in the region, countless other groups of organisms are being or will likely also be affected by environmental changes. In the North Sea, despite the decreasing abundances of holoplankton, data collected between 1958 and 1995 showed an increase in abundance of meroplanktonic organisms (Lindley and Batten, 2002). The meroplanktonic larvae of benthic echinoderm species, for example, now dominate numerically the summer plankton community in the North Sea (Kirby et al., 2007), and their peak occurrence is, on average, 47 days earlier in the season (Edwards and Richardson, 2004). If the entire ecosystem is to be better understood, especially under climate change scenarios, then other members of the zooplankton and their reaction to changing food regimes must also be characterized. In particular, the effects of regime shifts on the meroplankton should receive greater attention, as they will eventually propagate to the benthic populations. Furthermore, some species which have meroplanktonic larvae play a key role in their environment – *L. conchilega*, for example, is an ecosystem engineer whose presence is associated with higher benthic species richness and composition and faunal abundance (Callaway, 2006; Rabaut et al., 2007; Zühlke et al., 1998; Zühlke, 2001). It is important to better understand how these key species are affected by environmental changes, as any decrease in their populations may result in further shifts in ecosystem structure and reinforce regime shifts.



**Take-home messages for Section 4.2**

- ES explains the balance of energy and materials between prey and predator in terms of their elemental composition, and can be a useful tool for investigating environmental effects on the stoichiometry and, thus, physiology of organisms.
- Consumers can be characterized by their ability to handle nutrient-limited prey. ES assumes that the ingestion of nutritionally unbalanced diets by a consumer can lead to increasing assimilation of the limiting nutrient and waste production from the excess nutrient, decreasing (limiting) nutrient loss, and negative effects on consumer growth.
- ES was a useful tool to explain differences in copepod reproductive success in relation to diet nutritional quality. It failed, however, to explain other physiological results. This indicates a need for complimentary and combined approaches in nutritional ecology studies that also consider the biochemical composition of prey and predator.
- Patterns of lipid homeostasis in zooplankton can be related to the total lipid content of organisms. Lipid-poor meroplanktonic larvae, like those of *L. conchilega*, are quasi-homeostatic and can bioconvert dietary FAs to fulfill physiological/metabolic requirements. Holoplankton with intermediate lipid content, such as *T. longicornis* and other small temperate calanoids, are nonhomeostatic and assimilate dietary FAs in an unmodified manner, accumulating small lipid reserves for investment in reproduction. Lipid-rich holoplankton, such as large polar calanoid copepods, are semi-homeostatic and bioconvert dietary FAs into wax esters, forming large energy reserves for overwintering purposes.
- Studies on zooplankton feeding ecology must also address meroplanktonic organisms, which are understudied despite their ecological and economic importance.

### 4.3 CLIMATE CHANGE AND MARINE REGIME SHIFTS

Climate change in marine environments can be inferred from alterations in planktonic community structure, abundance, phenology and distribution (Hays et al., 2005). The plankton can, at times, be a more sensitive indicator of climate change than environmental variables themselves (Taylor et al., 2002), as biological patterns and ecological processes are profoundly influenced by variations in climate (Stenseth et al., 2004). On an individual level, changes such as increasing temperatures and decreasing nutrient availabilities can directly influence the physiology of an organism by increasing metabolic rates and through regulatory processes associated with altered food quality. Changes in temperature (i.e., global warming), both those which have already been observed and those which are still predicted for this century, are comparable in magnitude to the major global changes that the Earth experienced in the previous 65 million years (Diffenbaugh and Field, 2013). Combined land and ocean surface temperatures, for example, have risen globally by an average of  $0.85 \pm 0.2^{\circ}\text{C}$  in the past 140 years (IPCC, 2014). Sea surface temperatures (SST) have increased by approximately  $1^{\circ}\text{C}$  on the coast of Western Australia in the second half of the 20<sup>th</sup> century (Hobday and Pecl, 2014) and by 0.2 to  $0.6^{\circ}\text{C decade}^{-1}$  on the North Atlantic since the late 1970's (Dulvy et al., 2008). As the climate changes, species which cannot tolerate the environmental alterations will have to either adapt, relocate, or face possible extinction. Responses to climate change are species-specific, and current interaction patterns between organisms can be disrupted and new ones can be formed (Alheit et al., 2005; Pecl et al., 2017).

#### *Regime shifts*

Regime shifts can be defined as abrupt, substantial, and persistent changes to the state (structure, dynamics, and feedback mechanisms) of a natural system (Reid et al., 2016), which will lead to its reorganization into a different set of mutually reinforcing structures and processes (Rocha et al., 2014). They usually encompass multiple and interacting variables and affect several trophic levels and key structural species (Lees et al., 2006; Möllmann et al., 2014). Regime shifts have been recorded for marine environments such as salt marshes, sea ice, coral reefs, mangroves, and seagrass beds across all oceans (Alheit et al., 2005; Anderson and Piatt, 1999; Auber et al., 2015; Hughes, 1994; Lees et al., 2006, and references therein; Llope et al., 2012; Rocha et al., 2014). An increase in SST and stronger westerly winds caused a shift to a warm regime in the Gulf of

Alaska in 1977, which resulted in an increase in groundfish recruitment and salmon catches, a decrease in forage fish populations, which negatively affected birds and mammals, and overall community reorganization over the next 2-20 years (Anderson and Piatt, 1999). According to the authors, the community in the Gulf of Alaska was expected to bounce back to a cold regime in the early 2000's, in accordance with previous ecosystem changes, but this was not observed in later studies, as temperatures continued rising between 2001-2005 (Litzow, 2006). In the eastern English Channel, the Atlantic Multidecadal Oscillation caused a shift from a cold to a warm water regime in the mid 1990's and resulted in an increase in large-sized, higher trophic level fish and a decrease of small-bodied, low trophic level fish, which migrated northward to escape the warmer waters (Auber et al., 2015). In the eastern Pacific, alternations between cold and warm water regimes, characterized by high abundances of anchovy and sardine, respectively, have been reported to occur every ~ 25 years since the beginning of the 20<sup>th</sup> century (Alheit and Niquen, 2004; Chavez et al., 2003), but future global warming might change this stable state shifting pattern to a more permanent regime shift.

Dramatic and possibly catastrophic ecological, economic and social consequences can arise from ecological regime shifts (Alheit and Hagen, 2002; Alheit and Niquen, 2004; Hughes, 1994; Scheffer et al., 2001), as they often result in collapsing ecosystems and loss of services and goods. This is the case, for example, when pelagic fish such as sardine, anchovy, and herring are involved, as they comprise about one quarter of the total annual world fisheries catch (Hunter and Alheit, 1995) and are an important food source for larger fish, birds, and mammals (Crawford et al., 1991; Trillmich and Ono, 1991). Furthermore, regime shifts may also affect climate conditions and influence biogeochemical cycles of several elements, including the biological C pump, via feedback mechanisms (Alheit, 2009; Boyd and Doney, 2003). It is, thus, vital that the drivers and mechanisms causing regime shifts be better understood (Fisher et al., 2015). In past studies, marine regime shifts were often associated to a specific cause (Conversi et al., 2015), but a recent review identified 54 drivers which were, individually or jointly, responsible for 13 different types of global marine regime shifts and resulted in impacts on ecosystem biodiversity, fisheries provision, and primary production, a.o. (Rocha et al., 2014). Global warming, nutrient inputs, and fishing have been suggested to be major drivers in marine regime shifts (Ichii et al., 2017; Jiao, 2009 and references therein; Lees et al., 2006 and

## ***SYNOPTIC DISCUSSION***

---

references therein; Rocha et al., 2014), with the former causing changes in key physiological, demographic, and community-scale processes (Bennett et al., 2015; Wernberg et al., 2010).

The dynamics of regime shifts can only be unraveled by better understanding predator-prey interactions (endogenous mechanisms) and spatial dynamics of climate influences (exogenous drivers) (Conversi et al., 2015; Fisher et al., 2015). The zooplankton is commonly regarded as a key group in marine ecosystems, given that it is the bridge between primary production and higher trophic levels. Any alteration in the predator-prey interaction at the base of the food web would, thus, impact energy transfer in the system as a whole, potentially affecting even commercial fisheries. A better understanding of the stability of the trophic interactions in the plankton could help improve the assessment of the effects of environmental changes on marine systems (Llope et al., 2012). It is also important to consider the spatial and temporal variability of the drivers affecting these trophic interactions, as they may affect contiguous ecosystems in different ways (Llope et al., 2012). The North Sea is a good example of how changes in climate will affect trophic interactions in different ways depending on the spatial scale considered.

### *The regime shifts in the southern North Sea*

The establishment of monitoring programs such as the Continuous Plankton Recorder (CPR) and the Helgoland Roads time series likely resulted on the North Sea being one of the most intensively sampled marine systems in the world (~ 500,000 samples from CPR alone, McQuatters-Gollop et al., 2015). The analysis of data gathered by such time series has revealed the high variability characteristic of the North Sea (Llope et al., 2012). As previously mentioned in this thesis, at least 3 regime shifts have already been described for the North Sea, in the late 70's and 80's and early 2000's (Alheit, 2009; Alvarez-Fernandez et al., 2012; Beaugrand, 2004; Beaugrand et al., 2003; Becker and Pauly, 1996; Edwards et al., 2002; Kirby et al., 2008; Llope et al., 2012; Payne et al., 2009; Reid et al., 2003) with changes in all trophic levels, from phytoplankton to zooplankton, benthos, and nekton (Alheit et al., 2005). When this wealth of data is analyzed in relation to the different regions in the North Sea, it seems the effects of climate change were similar between the regions from 1958 and 1972, but varied according to spatial scale between 1972 and 2004 (Llope et al., 2012). In the second period top-down control mechanisms seem to prevail between phyto- and zooplankton on most areas in the North Sea,

especially off the Norwegian coast and the Skagerrak, but in the southern reach the control mechanisms are identified as bottom-up (Llope et al., 2012).

The work conducted during this PhD was performed in the area surrounding the small German island of Helgoland, in the southern North Sea, an ecosystem which has seen a considerable amount of changes in the past decades. One of the components of this thesis, the investigation into copepod performance in a changing ocean, was motivated by these major changes in environmental and biotic parameters. Between 1979 and 2011, water temperatures increased by 1.2°C and loading of N and P decreased from 25.32 to 13.06 mol N L<sup>-1</sup> and from 0.83 to 0.56 mol P L<sup>-1</sup>, with molar N:P ratio going from 30.5 to 39.4 and down to 23.3 (Boersma et al., 2015). Light penetration and wind speeds have also been reported to increase in that period of time (Wiltshire and Boersma, 2016). In the phytoplankton compartment, the abundances of two important food items for zooplankton predators have also changed. Dinoflagellate abundances declined from 3.82 to 2.14 \*10<sup>9</sup> µm<sup>3</sup> L<sup>-1</sup> (Boersma et al., 2015), whereas diatom abundances increased, especially that of large species (Wiltshire et al., 2010) which are difficult to ingest for organisms such as copepods (Roy et al., 1989). Members of the zooplankton were also affected, with temperature-related changes in phenology reported for copepods and gelatinous zooplankton (Boersma et al., 2015; Schlüter et al., 2010; Wiltshire and Boersma, 2016). Local calanoid copepod densities also drastically decreased (by approx. eightfold) since 1985, with the abundances of *T. longicornis* plummeting from 2000 ind m<sup>-3</sup> then to < 250 ind m<sup>-3</sup> in 2010 (Boersma et al., 2015). In order to better understand and predict how the Helgoland system will react in the present and near future to such changes, I designed experiments that attempted to reproduce environmental scenarios familiar to the region. This allowed me to look into the interaction between an important member of the zooplankton, the calanoid copepod *Temora longicornis*, and two of its important prey items, the diatom *Conticribra weissflogii* and the dinoflagellate *Oxyrrhis marina*, under two different nutrient availability scenarios.

#### *Investigating zooplankton performance in relation to diet nutritional quality*

Copepod performance was evaluated in two steps. First, their physiological responses to diet nutritional quality were quantified in terms of ingestion, respiration, excretion, egestion, growth, and reproduction rates. The efficiency with which these processes occurred, in terms of C and N, was also evaluated, and a budget was constructed with all of the rates in order to help understand

## ***SYNOPTIC DISCUSSION***

---

copepod responses to diet treatments. Ultimately, copepod performance was defined in terms of reproductive capacity, as this parameter defines the success and continuation of a population. The addition of  $^{13}\text{C}$ -labelled sodium bicarbonate ( $\text{NaH}^{13}\text{CO}_3$ ) to the diet cultures enabled the quantification of C transfer between prey and predator, such that it was possible to estimate C assimilation in relation to C ingestion rates. The analysis of  $^{13}\text{C}$ -enriched copepod samples via CSIA also provided FA-specific information that helped me better explain and understand the physiological data gathered during the experiments. The next pages contain a summary of my major results, whose meaning I discuss on a population scale, in terms of copepod performance, and on an ecosystem scale, within the context of the changes recorded for the Helgoland region.

The quantity of energy ingested by copepods did not match, at times, the energy used by the individuals. Although the rate measurements were recorded in the same fashion for all replicates fed with all diet, some treatments had a nearly balanced budget (e.g. Fig. 3C, Chapter I), in which energy consumption matched its utilization, while others revealed unbalanced budgets (e.g. Fig. 3F, Chapter I). I hypothesized that this reflected an underestimation of ingestion rates due to copepods feeding on eggs and FPs, as these were not physically separated in the glass bottles used in the grazing incubation. These alternate food sources were unaccounted for when determining ingestion rates, and were likely ingested in quantities inversely proportional to the quality of the diet treatments provided to copepods. It would seem that, by feeding on their own eggs and FPs, copepods could have complemented nutritionally inadequate diets, though this response mechanism is probably not relevant for the species in its natural environment.

The interesting result regarding ingestion rates is obtained by combining them with the C assimilation rates. When C assimilation is calculated as the percentage of dietary C that was ingested by copepods, no significant difference is found between the different treatments. This implied that food quality did not affect the assimilability of food, only the rate at which it was ingested. Methodological problems unfortunately prevented the verification of this pattern in the Dino+ treatment, which returned the highest EPR and was, thus, considered to provide the highest nutritional quality among the diets tested. Nevertheless the finding about diet quality affecting ingestion but not C assimilation rates is still relevant to the evaluation of copepod performance while feeding on the other treatments. For immature copepods, this result would have meant different time and energy investments for a similar C assimilation in relation to the

food available in the environment. As these copepods were females, we can explain the ingestion of more or less C with investment in reproductive output. The superiority of dinoflagellates as quality food for reproductive purposes in copepods, even when the former are nutritionally unbalanced, was further evidenced by the second highest EPR values, recorded for individuals who had the lowest ingestion rates while feeding on N-limited *O. marina*.

The caveat of the EPR results is that egg production, albeit the first step, is only one part of the process of producing viable offspring which survive into adulthood. The lack of information on how egg viability and juvenile survival are affected by diet quality limits my ability to claim with full certainty that *T. longicornis* fares better when feeding on dinoflagellates than on diatoms. I was able to record, however, different levels of investment of body N into reproduction between the treatments, which resulted in eggs with differing molar C:N ratios. This result indicates that copepods feeding on N-limited diatoms will likely produce eggs with a low N:C content. Studies on the hatching success of copepod eggs and nauplii survival in relation to egg molar C:N ratio are not available for *T. longicornis*, but can be found for the calanoid, *Calanus helgolandicus*, albeit with contradicting results. Guisande and Harris (1995) found that increasing egg sizes, which are controlled by food quantity and positively correlated to egg protein, carbohydrate, and lipid contents, lead to increases in egg hatching success and in nauplii survival. Pond et al. (1996), on the other hand, reported that the C and N contents of eggs are inversely correlated with egg viability, and not at all correlated with nauplii survival. The same trend is visible if one converts the C and N content data from the latter into molar C:N ratio. It remains to be seen whether and how egg molar C:N ratio will affect the success of the reproductive process in *T. longicornis*. In light of the aspects discussed, I believe the results obtained during this PhD are a very good indication that *T. longicornis* females perform better when feeding on dinoflagellates than on diatoms, even if N levels in the environment are limiting.

The availability of nutrients for algal growth in the North Sea is expected to become even more dependent on the influx from the North Atlantic in the future (Boersma et al., 2015). Dinoflagellates are poor competitors for nitrate whereas large diatoms, which possess a N “storage-adapted” strategy, have the advantage in the competition for this nutrient (Litchman et al., 2007). The availability of silicate and nitrate in the water is, thus, a major determinant of the ratio between diatoms and flagellates (Black et al., 2016). Changes in diatom and dinoflagellate

## ***SYNOPTIC DISCUSSION***

---

abundances and nutritional quality have the potential to greatly impact their zooplanktonic predators, as indicated by the work conducted during this PhD with *T. longicornis* females, and to affect the entire system in the near future (Wiltshire and Manly, 2004; Wiltshire et al., 2008). As a matter of fact, the decrease in calanoid copepod abundance in Helgoland in the past decades, although correlated with increases in seawater temperature, was indirectly related to the decrease in N and P input in the region (Boersma et al., 2015; Wiltshire and Boersma, 2016), and a similar trend has been suggested for other regions in the North Sea (Llope et al., 2012).

If the trophic interactions between calanoid copepods and their food items are only analyzed in relation to decreasing nutrient input, the logical conclusion is that their populations will continue to be negatively affected around Helgoland, as the availability of the nutritionally best prey (dinoflagellates) decreases and that of the worst quality (N-limited diatoms) increases. It has been suggested, however, that temperature is the key physical variable impacting phytoplankton, zooplankton, and fish species in the North Sea (Alheit et al., 2005). Increasing temperatures in the region (Wiltshire et al., 2010) will likely result in higher metabolic (respiration) costs for copepods (Castellani and Altunbas, 2014). These can only be met by increasing the demand (ingestion) for C. The effects of feeding on prey with higher C:nutrient ratios (due to the decrease in N and P loading) would be dampened by the ingestion of a higher relative C content, which would offset increased metabolic demand (Boersma et al., 2016; Malzahn et al., 2016). In such a scenario, the suitability of N-limited diatoms as “quality” food for copepods changes, but the end result for copepod performance and population success remains grim. And that is if further changes in copepod and diatom phenologies, which have shifted (by different processes) to a month earlier in the season during the last 50 years (Wiltshire and Boersma, 2016), do not lead to a mismatch between prey and predator, which it might in future (Wiltshire and Boersma, 2016). Alternatively, copepods feeding on high C:N diets in a warming scenario will offset the higher demand for C with increased ingestion of food and from the excess C in it, remaining limited by N and favoring the selection of food with lower C:N ratios (Anderson et al., 2017).

The full scale of the effects and feedback mechanisms of climate change in Helgoland are yet to be clarified and merit further investigation. Concomitant changes in all of these variables make it difficult to evaluate how copepods will react to future prey composition changes in the area. Although the work for this thesis refers to organisms from a very specific location, a small island



in the German Bight, the ubiquitousness of the species and of the environmental variable (N availability) studied enable the comparison of my results with those from several marine ecosystems. I believe my work represents a robust contribution towards a better understanding of marine systems and how they are affected by regime shifts.

**Take-home messages for Section 4.3**

- Regime shifts are abrupt, substantial, and persistent changes to the state of an ecosystem. They encompass multiple and interacting variables and affect several trophic levels and key species. It is thus necessary to understand how key organisms, such as copepods, will react to changing food regimes in their ecosystems.
- Major ecosystem changes have been recorded around Helgoland in the past 5 decades: increasing water temperatures, light penetration, wind speeds, and diatom abundance; decreasing nutrient loading, dinoflagellate abundance, and calanoid copepod densities; and temperature-related changes in copepod and zooplankton phenologies.
- Copepod performance in relation to food nutritional quality was ultimately evaluated by their egg production rates. Dinoflagellates were a food source of superior nutritional quality than diatoms, but further studies need to assess the effect of these diets on egg viability and juvenile survival. Concomitant changes in more than one environmental variable make it difficult to fully evaluate and predict how copepods will react to future prey composition changes in Helgoland.
- The ubiquitousness of the species and environmental variable studied enable the comparison of these results with those from other calanoids and marine ecosystems.



## **5 PERSPECTIVES**

Climate change and anthropogenic activities can affect ecosystems at all different levels of biological organization. In order for society to better understand the full scale of these (often synergistic) effects, a combined scientific effort must be adopted and make full use of approaches involving monitoring, modelling, and laboratory and field experiments. The long-term biological data sets obtained from time series such as Helgoland Roads provide an important temporal (and sometimes spatial) record of the patterns in ecosystem change (Alheit and Niquen, 2004; Pecl et al., 2017). Although laboratory and field experiments are unable to address all parameters at play in the investigation of ecosystem change, they can provide valuable insights into the feedback mechanisms generated from alterations. It is necessary to join monitoring data and experimental results in ecosystem modelling exercises in order to empirically test various mechanisms of regime shifts, and proposed courses of actions (Fisher et al., 2015). Only then will we be able to predict, prevent and/or respond to regime shifts.

I would recommend that more comprehensive experimental approaches be chosen for future scientific investigations into the effects of regime shifts on the zooplankton. Regarding the Helgoland region, I think it is important to investigate the combined effects of temperature and nutrient input on the planktonic community, though other parameters should also receive scientific attention. Temperature affects the ingestion of food, metabolism, and growth of individuals, and has a complex and poorly-understood impact on their nutritional requirements (Anderson et al., 2017). Given the fact that many zooplanktonic invertebrates are limited in nutrients (Hessen, 1992), it becomes clear the importance of investigating the synergistic effect of temperature and nutrient input on the zooplankton. These should be tested in terms of organism performance through all life stages, both in controlled conditions in the laboratory and in field mesocosm experiments, with results between both approaches compared. Nutrient limitation triggers changes at the molecular level which affect organism homeostasis (Sperfeld et al., 2017), and it would be interesting that the data gathered from experiments be analyzed under the framework of ES and another approach to nutritional ecology. I suggest that this should be the dynamic energy budget theory, which models energy uptake and usage for all vital physiological rates (as investigated during this thesis) at all life stages (Kooijman, 1995).

## ***PERSPECTIVES***

---

Last but not least, I also strongly encourage the procedural refinement for using labelled bicarbonate in tracing studies with organisms. In order to obtain accurate and pertinent results, it is important to understand how the phytoplankton incorporates  $^{13}\text{C}$  during photosynthesis and how this process is affected by culture conditions, such as light intensity, photoperiod, nutrient availability, and so on. This is a methodology with great potential to resolve unanswered biochemical questions and it should be used accordingly. In order to answer these questions of environmental and societal relevance, we must first perform the repetitive studies required to define the quantities of  $\text{NaH}^{13}\text{CO}_3$  that should be used for labelling organisms so that energy flow can be traced between trophic levels.

**REFERENCES**

- Ackman, R. G., Tocher, C. S., and McLachlan, J. (1968) Marine phytoplankter fatty acids. *J. Fish. Res. Board. Can.*, **25**, 1603–1620.
- Alheit, J., and Niquen, M. (2004) Regime shifts in the Humboldt Current ecosystem. *Prog. Oceanogr.*, **60**, 201–222.
- Alheit, J., and Hagen, E. (2002) Climate variability and historical NW European fisheries. In: Wefer, G., Berger, W. H., Behre, K.-E., and Jansen E. (eds.), *Climate development and history of the North Atlantic realm*. Springer-Verlag, Berlin, pp. 435–445.
- Alheit, J., Möllmann, C., Dutz, J., Kornilovs, G., Loewe, P., Mohrholz, V., and Wasmund, N. (2005) Synchronous ecological regime shifts in the central Baltic and the North Sea in the late 1980s. *ICES J. Mar. Sci.*, **62**, 1205–1215.
- Alheit, J. (2009) Consequences of regime shifts for marine food webs. *Int. J. Earth Sci.*, **98**, 261–268.
- Almeda, R., Alcaraz, M., Calbet, A., and Saiz, E. (2011) Metabolic rates and carbon budget of early developmental stages of the marine cyclopoid copepod *Oithona davisae*. *Limnol. Oceanogr.*, **56**, 403–414.
- Almeda, R., Pedersen, T. M., Jakobsen, H. H., Alcaraz, M., Calbet, A., and Hansen, B. W. (2009) Feeding and growth kinetics of the planktotrophic larvae of the spionid polychaete *Polydora ciliata* (Johnston). *J. Exp. Mar. Biol. Ecol.*, **382**, 61–68.
- Alvarez-Fernandez, S., Lindeboom, H., and Meesters, E. (2012) Temporal changes in plankton of the North Sea: Community shifts and environmental drivers. *Mar. Ecol. Prog. Ser.*, **462**, 21–38.
- Alves, R. M. S., van Colen, C., Vincx, M., Vanaverbeke, J., de Smet, B., Guarini, J.-M., Rabaut, M., and Bouma, T. J. (2017) A case study on the growth of *Lanice conchilega* (Pallas, 1766) aggregations and their ecosystem engineering impact on sedimentary processes. *J. Exp. Mar. Biol. Ecol.*, **489**, 15–23.
- Ambler, J. W. (1986) Effect of food quantity and quality on egg production of *Acartia tonsa* Dana from East Lagoon, Galveston, Texas. *Estuarine, Coastal Shelf Sci.*, **23**, 183–196.
- Anderson, M. J., Gorley, R. N., and Clarke, K. R. (2008) PERMANOVA+ for PRIMER: guide to software and statistical methods. PRIMER-E Ltd.
- Anderson, P. J., and Piatt, J. F. (1999) Community reorganization in the Gulf of Alaska following ocean climate regime shift. *Mar. Ecol. Prog. Ser.*, **189**, 117–123.
- Anderson, T. R. (1992) Modelling the influence of food C:N ratio, and respiration on growth and nitrogen excretion in marine zooplankton and bacteria. *J. Plankton Res.*, **14**, 1645–1671.

## ***REFERENCES***

---

- Anderson, T. R., Boersma, M., Raubenheimer, D. (2004) Stoichiometry: Linking elements to biochemicals. *Ecology*, **85**, 1193–1202.
- Anderson, T. R., Hessen, D. O. (1995) Carbon or nitrogen limitation in marine copepods? *J. Plankton Res.*, **17**, 317–331.
- Anderson, T. R., Hessen, D. O., Boersma, M., Urabe, J., and Mayor, D. J. (2017) Will invertebrates require increasingly carbon-rich food in a warming world? *Am. Nat.* **190**, 725–742.
- Anderson, T. R., Hessen, D. O., Elser, J. J., and Urabe, J. (2005) Metabolic stoichiometry and the fate of excess carbon and nutrients in consumers. *Am. Nat.* **165**, 1–15.
- Anderson, T. R., and Pond, D. W. (2000) Stoichiometric theory extended to micronutrients: Comparison of the roles of essential fatty acids, carbon, and nitrogen in the nutrition of marine copepods. *Limnol. Oceanogr.*, **45**, 1162–1167.
- Arendt, K. E., Jónasdóttir, S. H., Hansen, P. J., and Gärtner, S. (2005) Effects of dietary fatty acids on the reproductive success of the calanoid copepod *Temora longicornis*. *Mar. Biol.*, **146**, 513–530.
- Auber, A., Travers-Trolet, M., Villanueva, M. C., and Ernande, B. (2015) Regime shift in an exploited fish community related to natural climate oscillations. *PLoS One*, **10**, e0129883.
- Augustin, C. B., and Boersma, M. (2006) Effects of nitrogen stressed algae on different *Acartia* species. *J. Plankton Res.*, **28**, 429–436.
- Bakker, C., and van Rijswijk, P. (1987) Development time and growth rate of the marine calanoid copepod *Temora longicornis* as related to food conditions in the Oosterchelde estuary (Southern North Sea). *Neth. J. Sea Res.*, **21**, 125–141.
- Båmstedt, U. (1986) The biological chemistry of marine copepods. In: Corner, E. D. S., and O'Hara, S. C. M. (eds.), *The biological chemistry of marine copepods*. Clarendon Press, Oxford, pp. 1–58.
- Beaugrand, G. (2004) The North Sea regime shift: evidence, causes, mechanisms and consequences. *Progr. Oceanogr.*, **60**, 245–262.
- Beaugrand, G., Brander, K. M., Lindley, J. A., Souissi, S., and Reid, P. C. (2003) Plankton effect on cod recruitment in the North Sea. *Nature*, **426**, 661–664.
- Becker, G. A., and Pauly, M. (1996) Sea surface temperature changes in the North Sea and their causes. *ICES J. Mar. Sci.*, **53**, 887–898.
- Bennett, S., Wernberg, T., Harvey, E. S., Santana-Garcon, J., and Saunders, B. J. (2015) Tropical herbivores provide resilience to a climate-mediated phase shift on temperate reefs. *Ecol. Lett.* **18**, 714–723.
- Benson, S., Lennard, C., Maynard, P., and Roux, C. (2006) Forensic applications of isotope ratio mass spectrometry—A review. *Forensic Sci. Int.*, **157**, 1–22.

- Bhaud, M. (1988) The two planktonic larval periods of *Lanice conchilega* (Pallas, 1766) annelida polychaeta, a peculiar example of the irreversibility of evolution. *Ophelia*, **29**, 141–152.
- Bhaud, M., and Cazaux, C. P. (1990) Buoyancy characteristics of *Lanice conchilega* (Pallas) larvae (Terebellidae). Implications for settlement. *J. Exp. Mar. Biol. Ecol.*, **141**, 31–45.
- Black, K. P., Longmore, A. R., Hamer, P. A., Lee, R., Swearer, S. E., and Jenkins, G. P. (2016) Linking nutrient inputs, phytoplankton composition, zooplankton dynamics and the recruitment of pink snapper, *Chrysophrys auratus*, in a temperate bay. *Estuarine, Coastal Shelf Sci.*, **183**, 150–162.
- Boersma, M., Wiltshire, K. H., Kong, S. M., Greve, W., and Renz, J. (2015) Long-term change in the copepod community in the southern German Bight. *J. Sea Res.*, **101**, 41–50.
- Boersma, M., Mathew, K. A., Niehoff, B., Schoo, K. L., Franco-Santos, R. M., and Meunier, C. L. (2016) Temperature driven changes in the diet preference of omnivorous copepods: no more meat when it's hot? *Ecol. Lett.* **19**, 45–53.
- Boissonnot, L., Niehoff, B., Hagen, W., Søreide, J. E., and Graeve, M. (2016) Lipid turnover reflects life-cycle strategies of small-sized Arctic copepods. *J. Plankton Res.*, **38**, 1420–1432.
- Boyd, P. W., and Doney, S. C. (2003) The impact of climate change and feedback processes on the ocean carbon cycle. In: Fasham, M. J. R. (ed.), *Ocean Biogeochemistry. Global Change — The IGBP Series* (closed). Springer, Berlin, Heidelberg.
- Braber, L., and de Groot, S. J. (1973) The food of five flatfish species (Pleuronectiformes) in the southern North Sea. *Neth. J. Sea Res.*, **6**, 163–172.
- Braeckman, U., Provoost, P., Sabbe, K., Soetaert, K., Middelburg, J. J., Vincx, M., and Vanaverbeke, J. (2012) Temporal dynamics in the diet of two marine polychaetes as inferred from fatty acid biomarkers. *J. Sea Res.*, **68**, 6–19.
- Bromke, M. A., Sabir, J. S., Alfassi, F. A., Hajarrah, N. H., Kabli, S. A., Al-Malki, A. L., Ashworth, M. P., Méret, M., Jansen, R. K., and Willmitzer, L. (2015). Metabolomic profiling of 13 diatom cultures and their adaptation to nitrate-limited growth conditions. *PLoS One*, 10: e0138965.
- Budge, S. M., Iverson, S. J., and Koopman, H. N. (2006) Studying trophic ecology in marine ecosystems using fatty acids: a primer on analysis and interpretation. *J. Exp. Biol.*, **214**, 3790–3800.
- Budge, S. M., Wooller, M. J., Springer, A. M., Iverson, S. J., McRoy, C. P., and Divoky, G. J. (2008) Tracing carbon flow in an arctic marine food web using fatty acid-stable isotope analysis. *Oecologia*, **157**, 117–129.
- Bühr, K.-J. (1976) Suspension-feeding and assimilation efficiency in *Lanice conchilega* (Polychaeta). *Mar. Biol.*, **38**, 373–383.
- Bühr, K.-J., and Winter, J. E. (1977) Distribution and maintenance of a *Lanice conchilega* association in the Weser Estuary (FRG), with special reference to the suspension-feeding behaviour of *Lanice*

## **REFERENCES**

---

- conchilega*. In: Keegan, B. F., Ceidigh, P. O., and Boaden, P. J. S. (eds.) Biology of benthic organisms. Pergamon, pp. 101–113.
- Callaway, R. (2003) Juveniles stick to adults: recruitment of the tube-dwelling polychaete *Lanice conchilega* (Pallas, 1766). *Hydrobiologia*, **503**, 121–130.
- Callaway, R. (2006) Tube worms promote community change. *Mar. Ecol. Prog. Ser.*, **308**, 49–60.
- Castellani, C., and Altunbas, Y. (2014) Seasonal change in acclimatised respiration rate of *Temora longicornis*. *Mar. Ecol. Prog. Ser.*, **500**, 83–101.
- Chavez, F. P., Ryan, J., Lluch-Cota, S. E., and Ñiquen, M. (2003) From anchovies to sardines and back: Multidecadal change in the Pacific Ocean. *Science*, **229**, 217–221.
- Checkley, D. M., and Entzeroth, L. C. (1985) Elemental and isotopic fractionation of carbon and nitrogen by marine, planktonic copepods and implications to the marine nitrogen cycle. *J. Plankton Res.*, **7**, 553–568.
- Chen, M., Kim, D., Liu, H., and Kang, C.-K. (2018) Variability in copepod trophic levels and feeding selectivity based on stable isotope analysis in Gwangyang Bay of the southern coast of the Korean Peninsula. *Biogeosciences*, **15**, 2055–2073.
- Cherel, Y., Phillips, R. A., Hobson, K. A., and McGill, R. (2006) Stable isotope evidence of diverse species-specific and individual wintering strategies in seabirds. *Biol. Lett.*, **2**, 301–303.
- Clarke, K. R., and Gorley, R. N. (2015) PRIMER v7: User Manual/Tutorial. PRIMER-E, Plymouth.
- Conover, R. J. (1978). Transformation of organic matter. In: Kinne, O. (ed.) Marine Ecology. Vol IV Dynamics. Wiley, New York, pp. 221–499.
- Conversi, A., Dakos, V., Gårdmark, A., Ling, S., Folke, C., Mumby, P. J., Greene, C., et al. (2015) A holistic view of marine regime shifts. *Phil. Trans. R. Soc. B*, **370**, 20130279.
- Cotonnec, G., Brunet, C., Sautour, B. and Thoumelin, G. (2001). Nutritive value and selection of food particles by copepods during a spring bloom of *Phaeocystis* sp. in the English Channel, as determined by pigment and fatty acid analyses. *J. Plankton Res.*, **23**, 693–703.
- Crawford, R. J. M., Ryan, P. G., and Williams, A. J. (1991) Seabird consumption and production in the Benguela and Western Agulhas ecosystems. *S. Afr. J. Marine Sci.*, **11**, 357–375.
- Dalsgaard, J., John, M. S., Kattner, G., Müller-Navarra, D., and Hagen, W. (2003) Fatty acid trophic markers in the pelagic marine environment. *Adv. Mar. Biol.*, **46**, 225–340.
- De Smet, B., Braeckman, U., Soetaert, K., Vincx, M., and Vanaverbeke, J. (2016) Predator effects on the feeding and bioirrigation activity of ecosystem-engineered *Lanice conchilega* reefs. *J. Exp. Mar. Biol. Ecol.*, **475**, 31–37.



- De Smet, B., Godet, L., Fournier, J., Desroy, N., Jaffré, M., Vincx, M., and Rabaut, M. (2013) Feeding grounds for waders in the Bay of the Mont Saint-Michel (France): the *Lanice conchilega* reef serves as an oasis in the tidal flats. *Mar. Biol.*, **160**, 751–761.
- De Troch, M., Boeckx, P., Cnudde, C., Van Gansbeke, D., Vanreusel, A., Vincx, M., and Caramujo, M. J. (2012) Bioconversion of fatty acids at the basis of marine food webs: insights from a compound-specific stable isotope analysis. *Mar. Ecol. Prog. Ser.*, **465**, 53–67.
- Debs, C. A. (1984) Carbon and nitrogen budget of the calanoid copepod *Temora stylifera*: effect of concentration and composition of food. *Mar. Ecol. Prog. Ser.*, **15**, 213–223.
- DeMott, W. R. (1988) Discrimination between algae and artificial particles by freshwater and marine copepods. *Limnol. Oceanogr.*, **33**, 397–408.
- Diffenbaugh, N. S., and Field, C. B. (2013) Changes in ecologically critical terrestrial climate conditions. *Science*, **341**, 486–492.
- Dubois, S., Orvain, F., Marin-Leal, J. C., Ropert, M., and Lefebvre, S. (2007). Small-scale spatial variability of food partitioning between cultivated oysters and associated suspension-feeding species, as revealed by stable isotopes. *Mar. Ecol. Prog. Ser.*, **336**, 151–160.
- Ducrotoy, J. P., Elliot, M., and De Jonge, V. N. (2000) The North Sea. *Mar. Pollut. Bull.*, **41**, 5–23.
- Dulvy, N. K., Rogers, S. I., Jennings, S., Stelzenmüller, V., Dye, S. R., and Skjoldal, H. R. (2008) Climate change and deepening of the North Sea fish assemblage: a biotic indicator of warming seas. *J. Appl. Ecol.*, **45**, 1029–1039.
- Edwards, M., and Richardson, A. J. (2004) Impact of climate change on marine pelagic phenology and trophic mismatch. *Nature*, **430**, 881–884.
- Edwards, M., Beaugrand, G., Reid, P. C., Rowden, A. A., and Jones, M. B. (2002) Ocean climate anomalies and the ecology of the North Sea. *Mar. Ecol. Prog. Ser.*, **239**, 1–10.
- Eek, M. K., Whiticar, M. J., Bishop, J. K. B., and Wong, C. S. (1999) Influence of nutrients on carbon isotope fractionation by natural populations of Prymnesiophyte algae in NE Pacific. *Deep-Sea Res. Pt. II*, **46**, 2863–2876.
- Evans, F. (1981) An investigation into the relationship of sea temperature and food supply to the size of the planktonic copepod *Temora longicornis* Müller in the North Sea. *Estuarine, Coastal Shelf Sci.*, **13**, 145–158.
- Evershed, R. P., Bull, I. D., Corr, L. T., Crossman, Z. M., van Dongen, B. E., Evans, C. J., Jim, S., Mottram, H. R., Mukherjee, A. J., and Pancost, R. D. (2007) Compound specific stable isotope analysis in ecology and paleoecology. In: Michener, R. H., and Lajtha, K. (eds.), *Stable isotopes in ecology and environmental science*. Blackwell Publishing, UK, pp. 480–540.

## **REFERENCES**

---

- Evjemo, J. O., and Olsen, Y. (1997) Lipid and fatty acid content in cultivated live feed organisms compared to marine copepods. *Hydrobiologia*, **358**, 159–162.
- Falk-Petersen, S., Sargent, J. R., Henderson, J., Hegseth, E. N., Hop, H., and Okolodkov, Y. B. (1998) Lipids and fatty acids in ice algae and phytoplankton from the Marginal Ice Zone in the Barents Sea. *Polar Biol.*, **20**, 41–47.
- Fischer, A. (2013) The Helgoland manual of animal development. Notes and laboratory protocols on marine invertebrates. Pfeil, München. 192pp.
- Fisher, J. A. D., Casini, M., Frank, K. T., Möllmann, C., Leggett, W. C., and Daskalov, G. (2015) The importance of within-system spatial variation in drivers of marine ecosystem regime shifts. *Phil. Trans. R. Soc. B*, **370**, 20130271.
- Fogel, M. L., and Tuross, N. (2003) Extending the limits of paleodietary studies of humans with compound specific carbon isotope analysis of amino acids. *J. Archaeol. Sci.*, **30**, 535–545.
- Franco-Santos, R. M., Auel, H., Boersma, M., De Troch, M., Meunier, C. L., and Niehoff, B. (2018). Bioenergetics of the copepod *Temora longicornis* under different nutrient regimes. *J. Plankton Res.*, **40**, 420–435. doi:10.1093/plankt/fby016.
- Franz, H. G., Colebrook, J. M., Gamble, J. C., and Krause, M. (1991) The zooplankton of the North Sea. *Neth. J. Sea Res.*, **28**, 1–52.
- Fraser, A. J., Sargent, J. R., and Gamble, J. C. (1989) Lipid class and fatty acid composition of *Calanus finmarchicus* (Gunnerus), *Pseudocalanus* sp. and *Temora longicornis* Muller from a nutrient-enriched seawater enclosure. *J. Exp. Mar. Biol. Ecol.*, **130**, 81–92.
- Frost, B. W. (1972) Effects of size and concentration of food particles on the feeding behavior of the marine planktonic copepod *Calanus pacificus*. *Limnol. Oceanogr.*, **17**, 805–815.
- Gentsch, E., Kreibich, T., Hagen, W., Niehoff, B. (2009) Dietary shifts in the copepod *Temora longicornis* during spring: evidence from stable isotope signatures, fatty acid biomarkers and feeding experiments. *J. Plankton Res.*, **31**, 45–60.
- Godet, L., Toupoint, N., Olivier, F., Fournier, J., and Retière, C. (2008) Considering the functional value of common marine species as a conservation stake: The case of Sandmason Worm *Lanice conchilega* (Pallas 1766) (Annelida, Polychaeta) beds. *Ambio*, **37**, 347–355.
- Gorbatenko, K. M., Lazhentsev, A. E., and Kiyashko, S. I. (2014) Seasonal dynamics of the trophic status of zooplankton in the Sea of Okhotsk (based on data from stable carbon and nitrogen isotope analysis). *Russ. J. Mar. Biol.*, **40**, 519–531.
- Graeve, M., Albers, C., and Kattner, G. (2005) Assimilation and biosynthesis of lipids in Arctic *Calanus* species based on feeding experiments with a <sup>13</sup>C labelled diatom. *J. Exp. Mar. Biol. Ecol.*, **317**, 109–125.

- Graeve, M., Kattner, G., and Hagen, W. (1994) Diet-induced changes in the fatty acid composition of Arctic herbivorous copepods: Experimental evidence of trophic markers. *J. Exp. Mar. Biol. Ecol.*, **182**, 97–110.
- Guilini, K., van Oevelen, D., Soetaert, K., Middelburg, J. J., and Vanreusel, A. (2010) Nutritional importance of benthic bacteria for deep-sea nematodes from the Arctic ice margin: Results of an isotope tracer experiment. *Limnol. Oceanogr.*, **55**, 1977–1989.
- Guillard, R. R. L. (1975) Culture of phytoplankton for feeding marine invertebrates. In: Smith, W. L., and Chanley, M. H. (eds.), *Culture of marine invertebrate animals*. Springer, US, pp. 29–60.
- Guisande, C., and Harris, R. (1995) Effect of total organic content of eggs on hatching success and naupliar survival in the copepod *Calanus helgolandicus*. *Limnol. Oceanogr.*, **40**, 476–482.
- Guisande, C., Riveiro, I., and Maneiro, I. (2000) Comparisons among the amino acid composition of females, eggs and food to determine the relative importance of food quantity and food quality to copepod reproduction. *Mar. Ecol. Prog. Ser.*, **202**, 135–142.
- Halsband, C., and Hirche, H. J. (2001) Reproductive cycles of dominant calanoid copepods in the North Sea. *Mar. Ecol. Prog. Ser.*, **209**, 219–229.
- Halsband-Lenk, C., Carlotti, F., and Greve, W. (2004) Life-history strategies of calanoid congeners under two different climate regimes: a comparison. *ICES J. Mar. Sci.*, **61**, 709–720.
- Hansen, B. W. (1999) Cohort growth of planktotrophic polychaete larvae - are they food limited? *Mar. Ecol. Prog. Ser.*, **178**, 109–119.
- Hantzsche, F. M., and Boersma, M. (2010) Dietary-induced responses in the phagotrophic flagellate *Oxyrrhis marina*. *Mar. Biol.*, **157**, 1641–1651.
- Harris, R., Wiebe, P., Lenz, J., Skjoldal, H. R., and Huntley, M. (eds.) (2000) *ICES Zooplankton Methodology Manual*. Academic Press.
- Harrison, P. J., Thompson, P. A., and Calderwood, G. S. (1990) Effects of nutrient and light limitation on the biochemical composition of phytoplankton. *J. Appl. Phycol.*, **2**, 45–56.
- Hays, G. C., Richardson, A. J., and Robinson, C. Climate change and marine plankton. *TRENDS Ecol. Evol.*, **20**, 337–344.
- Helland, S., Terjesen, B. F., and Berg, L. (2003) Free amino acid and protein content in the planktonic copepod *Temora longicornis* compared to *Artemia franciscana*. *Aquaculture*, **215**, 213–228.
- Henderson, R. J., Hegseth, E. N., and Park, M. T. (1998) Seasonal variation in lipid and fatty acid composition of ice algae from the Barents Sea. *Polar Biol.*, **20**, 48–55.
- Hessen, D. O. (1992) Nutrient element limitation of zooplankton production. *Am Nat*, **140**, 799–814.
- Hessen, D. O., and Anderson, T. R. (2008) Excess carbon in aquatic organisms and ecosystems: physiological, ecological, and evolutionary implications. *Limnol. Oceanogr.*, **53**, 1685–96.

## **REFERENCES**

---

- Hickel, W. (1975) The mesozooplankton in the Wadden Sea of Sylt (North Sea). *Helgoländer Wissenschaftliche Meeresunterssungen*, **27**, 254–262.
- Hickel, W., Mangelsdorf, P., and Berg, J. (1993). The human impact in the German Bight- Eutrophication during three decades (1962-1991). *Helgoländer Meeresunters.*, **47**, 243–263.
- Hinder, S. L., Hays, G. C., Edwards, M., Roberts, E. C., Walne, A. W., and Gravenor, M. B. (2012) Changes in marine dinoflagellate and diatom abundance under climate change. *Nat. Clim. Change*, **2**, 271–275.
- Hirst, A. G., and Bunker, A. J. (2003) Growth of marine planktonic copepods: Global rates and patterns in relation to chlorophyll *a*, temperature, and body weight. *Limnol. Oceanogr.*, **48**, 1988–2010.
- Hobday, A. J., and Pecl, G. T. (2014) Identification of global marine hotspots: sentinels for change and vanguards for adaptation action. *Rev. Fish Biol. Fisher.*, **24**, 415–425.
- Holthe, T. (1986) Evolution, systematics, and distribution of the Polychaeta Terebellomorpha, with a catalogue of the taxa and a bibliography. *Gunneria*, **55**, 1–236.
- Hughes, T. P. (1994) Catastrophes, phase shifts, and large-scale degradation of a Caribbean coral reef. *Science*, **265**, 1547–1551.
- Hunter, J. R., and Alheit, J. (1995) International GLOBEC small pelagic fishes and climate change program. GLOBEC Report No. 8. 72 pp.
- Huntley, M., and Boyd, C. (1984) Food limited growth of marine zooplankton. *Am. Nat.*, **124**, 455–478.
- Ianora, A. (1998) Copepod life history traits in subtemperate regions. *J. Mar. Syst.*, **15**, 337–349.
- ICES (2016) ICES Ecosystem Overview – Greater North Sea Ecoregion. Available at: [http://www.ices.dk/sites/pub/Publication%20Reports/Advice/2016/2016/Greater\\_North\\_Sea\\_Eco\\_region-Ecosystem\\_overview.pdf](http://www.ices.dk/sites/pub/Publication%20Reports/Advice/2016/2016/Greater_North_Sea_Eco_region-Ecosystem_overview.pdf). Accessed on September 10, 2018.
- Ichii, T., Nishikawa, H., Igarashi, H., Okamura, H., Mahapatra, K., Sakai, M., Wakabayashi, T., Inagaki, D., and Okada, Y. (2017) Impacts of extensive driftnet fishery and late 1990s climate regime shift on dominant epipelagic nekton in the Transition Region and Subtropical Frontal Zone: Implications for fishery management. *Prog. Oceanogr.*, **150**, 35–47.
- Ihaka, R., and Gentleman, R. (1993) R (Version 3.4.4) [software]. Retrieved from <https://cran.r-project.org/bin/windows/base/old/3.2.4/>.
- Ikeda, T. (1976) The effect of laboratory conditions on the extrapolation of experimental measurements to the ecology of marine zooplankton I. Effect of feeding condition on the respiration rate. *Bull. Plankton Soc. Jpn.*, **23**, 51–60.
- Ikeda, T., Kanno, Y., Ozaki, K., and Shinada, A. (2001) Metabolic rates of epipelagic marine copepods as a function of body mass and temperature. *Mar. Biol.*, **139**, 587–596.

- IMM (Inter Ministerial Meeting) (1997) Assessment Report on Fisheries and Related Species and Habitat Issues. Danish Environmental Protection Agency, Copenhagen.
- IPCC - Intergovernmental Panel on Climate Change (2014) Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Available at: [www.ipcc.ch/report/ar5/syr/](http://www.ipcc.ch/report/ar5/syr/). Accessed on September 17, 2018.
- Jiao, Y. (2009) Regime shift in marine ecosystems and implications for fisheries management, a review. *Rev. Fish Biol. Fisher.*, **19**, 177–191.
- Jónasdóttir, S. H., Fields, D., and Pantoja, S. (1995) Copepod egg production in Long Island Sound, USA, as a function of the chemical composition of seston. *Mar. Ecol. Prog. Ser.*, **119**, 87–98.
- Jónasdóttir, S. H. (2017) The role of vertically migrating zooplankton in biogeochemical flux. Presentation for the Ocean Carbon and Biogeochemistry (OCB) meeting in Woods Hole, USA, on 28 June 2017.
- Jónasdóttir, S. H., Kiørboe, T., Tang, K. W., St John, M., Visser, A. W., Saiz, E., Dam, H. G. (1998) Role of diatoms in copepod production: good, harmless or toxic? *Mar. Ecol. Prog. Ser.*, **172**, 305–308.
- Jónasdóttir, S. H., Visser, A. W., and Jespersen, C. (2009) Assessing the role of food quality in the production and hatching of *Temora longicornis* eggs. *Mar. Ecol. Prog. Ser.*, **382**, 139–150.
- Jones, R. H., and Flynn, K. J. (2005) Nutritional status and diet composition affect the value of diatoms as copepod prey. *Science*, **307**, 1457–1459.
- Jones, R. H., Flynn, K. J., and Anderson, T. R. (2002) Effect of food quality on carbon and nitrogen growth efficiency in the copepod *Acartia tonsa*. *Mar. Ecol. Prog. Ser.*, **235**, 147–156.
- Juul-Pedersen, T., Nielsen, T. G., Michel, C., Møller, E. F., Tiselius, P., Thor, P., Olesen, M., Selander, E., et al. (2006) Sedimentation following the spring bloom in Disko Bay, West Greenland, with special emphasis on the role of copepods. *Mar. Ecol. Prog. Ser.*, **314**, 239–255.
- Kattner, G., Gercken, G., and Eberlein, K. (1983) Development of lipids during a spring plankton bloom in the Northern North Sea. I. Particulate fatty acids. *Mar. Chem.*, **14**, 149–162.
- Kiørboe, T., and Nielsen, T. G. (1994) Regulation of zooplankton biomass and production in a temperate, coastal ecosystem. I. Copepods. *Limnol. Oceanogr.*, **39**, 493–507.
- Kiørboe, T., Møhlenberg, F., and Hamburger, K. (1985) Bioenergetics of the planktonic copepod *Acartia tonsa*: relation between feeding, egg production and respiration, and composition of specific dynamic action. *Mar. Ecol. Prog. Ser.*, **26**, 85–97.
- Kiørboe, T., Saiz, E., and Viitasalo, M. (1996) Prey switching behaviour in the planktonic copepod *Acartia tonsa*. *Mar. Ecol. Prog. Ser.*, **143**, 65–75.

## ***REFERENCES***

---

- Kirby, R. R., Beaugrand, G., and Lindley, J. A. (2008) Climate-induced effects on the meroplankton and the benthic-pelagic ecology of the North Sea. *Limnol. Oceanogr.*, **53**, 1805–1815.
- Kirby, R. R., Beaugrand, G., Lindley, J. A., Richardson, A. J., Edwards, M., and Reid, P. C. (2007) Climate effects and benthic-pelagic coupling in the North Sea. *Mar. Ecol. Prog. Ser.*, **330**, 31–38.
- Klein Breteler, W. C. M., Schogt, N., and Rampen, S. (2005) Effect of diatom nutrient limitation on copepod development: role of essential lipids. *Mar. Ecol. Prog. Ser.*, **291**, 125–133.
- Kluijver, A., Soetaert, K., Czerny, J., Schulz, K. G., Boxhammer, T., Riebesell, U., and Middelburg, J. J. (2013) A <sup>13</sup>C labelling study on carbon fluxes in Arctic plankton communities under elevated CO<sub>2</sub> levels. *Biogeosciences*, **10**, 1425–1440.
- Kooijman, S. A. L. M. (1995) The stoichiometry of animal energetics. *J. Theor. Biol.*, **177**, 139–149.
- Kreibich, T., Saborowski, R., Hagen, W., and Niehoff, B. (2008) Short-term variation of nutritive and metabolic parameters in *Temora longicornis* females (Crustacea, Copepoda) as a response to diet shift and starvation. *Helgol. Mar. Res.*, **62**, 241–249.
- Kreibich, T., Saborowski, R., Hagen, W., and Niehoff, B. (2011) Influence of short-term nutritional variations on digestive enzyme and fatty acid patterns of the calanoid copepod *Temora longicornis*. *J. Exp. Mar. Biol. Ecol.*, **407**, 182–189.
- Kuijper, L. D. J., Anderson, T. R., and Kooijman, S. A. L. M. (2004) C and N gross growth efficiencies of copepod egg production studied using a Dynamic Energy Budget model. *J. Plankton Res.*, **26**, 213–226.
- Larink, O., and Westheide, W. (2011) Coastal plankton. Photo guide for European seas. Pfeil, München. 204pp.
- Lee, R. F., Nevenzel, J. C., and Paffenhöfer, G. A. (1971) Importance of wax esters and other lipids in the marine food chain: phytoplankton and copepods. *Mar. Biol.*, **9**, 99–108.
- Lee, R. F., Hagen, W., and Kattner, G. (2006) Lipid storage in marine zooplankton. *Mar. Ecol. Prog. Ser.*, **307**, 273–306.
- Lee, W. Y., and Yan, H. Y. (1994) Energy budget of the female harpacticoid *Tisbe carolinensis* (Copepoda, Crustacea). *Comp. Biochem. Physiol.*, **109**, 817–820.
- Lees, K., Pitois, S., Scott, C., Frid, C., and Mackinson, S. (2006) Characterizing regime shifts in the marine environment. *Fish Fish.*, **7**, 104–127.
- Lichtfouse, E. (2000) Compound-specific isotope analysis. Application to archaeology, biomedical sciences, biosynthesis, environment, extraterrestrial chemistry, food science, forensic science, humic substances, microbiology, organic geochemistry, soil science and sport. *Rapid Commun. Mass Spectrom.*, **14**, 1337–1344.
- Lindeman, R. L. (1942) The trophic-dynamic aspect of ecology. *Ecology*, **23**, 399–417.

- Lindley, J. A., Batten, S. D. (2002) Long-term variability in the diversity of North Sea zooplankton. *J. Mar. Biol. Assoc. UK*, **82**, 31–40.
- Litchman, E., Klausmeier, C. A., Schofield, O. M., and Falkowski, P. G. (2007) The role of functional traits and trade-offs in structuring phytoplankton communities: scaling from cellular to ecosystem level. *Ecol. Lett.* **10**, 1170–1181.
- Litzow, M. A. (2006) Climate regime shifts and community reorganization in the Gulf of Alaska: how do recent shifts compare with 1976/1977? *ICES J. Mar. Sci.*, **63**, 1386–1396.
- Llope, M., Licandro, P., Chan, K.-S., and Stenseth, N. C. (2012) Spatial variability of the plankton trophic interaction in the North Sea: a new feature after the early 1970s. *Glob. Change Biol.*, **18**, 106–117.
- Löder, M. G. J., Meunier, C., Wiltshire, K. H., Boersma, M., and Aberle, N. (2011) The role of ciliates, heterotrophic dinoflagellates and copepods in structuring spring plankton communities at Helgoland Roads, North Sea. *Mar. Biol.*, **158**, 1551–1580.
- Maar, M., Nielsen, T. G., Gooding, S., Tønnesson, K., Tiselius, P., Zervoudaki, S., Christou, E., Sell, A., et al. (2004) Trophodynamic function of copepods, appendicularians and protozooplankton in the late summer zooplankton community in the Skagerrak. *Mar. Biol.*, **144**, 917–933.
- Malzahn, A. M., and Boersma, M. (2012) Effects of poor food quality on copepod growth are dose dependent and non-reversible. *Oikos*, **121**, 1408–1416.
- Malzahn, A. M., Aberle, N., Clemmesen, C., and Boersma, M. (2007) Nutrient limitation of primary producers affects planktivorous fish condition. *Limnol. Oceanogr.*, **52**, 2062–2071.
- Malzahn, A. M., Doerfler, D., and Boersma, M. (2016) Junk food gets healthier when it's warm. *Limnol. Oceanogr.*, **61**, 1677–1685.
- Malzahn, A. M., Hantzsche, F., Schoo, K. L., Boersma, M., and Aberle, N. (2010). Differential effects of nutrient-limited primary production on primary, secondary or tertiary consumers. *Oecologia*, **162**, 35–48.
- Marshall, S. M. (1973) Respiration and feeding in copepods. *Adv. Mar. Biol.*, **11**, 57–120.
- Martel, C. M., and Flynn, K. J. (2008) Morphological controls on cannibalism in a planktonic marine phagotroph. *Protist*, **159**, 41–51.
- Mayzaud, P. (1976) Respiration and nitrogen excretion of zooplankton. IV. The influence of starvation on the metabolism and the biochemical composition of some species. *Mar. Biol.*, **37**, 47–58.
- McConnaughey, T. (1978) Ecosystems naturally labelled with Carbon-13: Applications to the study of consumer food-webs. Thesis, University of Alaska in Fairbanks, Alaska.
- McHugh, D. (1993) A comparative study of reproduction and development in the polychaete family Terebellidae. *Biol. Bull.*, **185**, 153–167.

## **REFERENCES**

---

- McQuatters-Gollop, A., Edwards, M., Helaouët, P., Johns, D. G., Owens, N. J. P., Raitsos, D. E., Schroeder, D., Skinner, J., and Stern, R. F. (2015) The Continuous Plankton Recorder survey: How can long-term phytoplankton datasets contribute to the assessment of Good Environmental Status? *Estuar. Coast. Shelf. S.*, **162**, 88–97.
- Meier-Augenstein, W. (1999) Applied gas chromatography coupled to isotope ratio mass spectrometry. *J. Chromatogr. A*, **842**, 351–371.
- Meier-Augenstein, W. (2002) Stable isotope analysis of fatty acids by gas chromatography–isotope ratio mass spectrometry. *Anal. Chim. Acta*, **465**, 63–79.
- Meunier, C. L., Haafke, J., Oppermann, B., Boersma, M., and Malzahn, A. M. (2012) Dynamic stoichiometric response to food quality fluctuations in the heterotrophic dinoflagellate *Oxyrrhis marina*. *Mar. Biol.*, **159**, 2241–2248.
- Meunier, C. L., Malzahn, A. M. and Boersma, M. (2014) A new approach to homeostatic regulation: towards a unified view of physiological and ecological concepts. *PLoS ONE*, **9**, e107737.
- Michener, R. H., and Kaufman, L. (2007) Stable isotope ratios as tracers in marine food webs: An update. In: Michener, R. H., and Lajtha, K. (eds.), *Stable isotopes in ecology and environmental science*. Blackwell Publishing, UK, pp. 238–282.
- Middelburg, J. J., Barranguet, C., Boschker, H. T. S., Herman, P. M. J., Moens, T., and Heip, C. H. R. (2000) The fate of intertidal microphytobenthos carbon: An in situ <sup>13</sup>C-labeling study. *Limnol. Oceanogr.*, **45**, 1224–1234.
- Miller, C. A., and Glibert, P. M. (1998) Nitrogen excretion by the calanoid copepod *Acartia tonsa*: results of mesocosm experiments. *J. Plankton Res.*, **20**, 1767–1780.
- Miralto, A., Barone, G., Romano, G., Poulet, S. A., Ianora, A., Russo, G. L., Buttino, I., Mazzarella, G., et al. (1999) The insidious effect of diatoms on copepod reproduction. *Nature*, **402**, 173–176.
- Mitra, A., Flynn, K. J. (2005) Predator–prey interactions: is ‘ecological stoichiometry’ sufficient when good food goes bad? *J. Plankton Res.*, **27**, 393–399.
- Moe, S. J., Stelzer, R. S., Forman, M. R., Harpole, W. S., Daufresne, T., and Yoshida, T. (2005) Recent advances in ecological stoichiometry: insights for population and community ecology. *Oikos*, **109**, 29–39.
- Møller, E. F. (2007) Production of dissolved organic carbon by sloppy feeding in the copepods *Acartia tonsa*, *Centropages typicus*, and *Temora longicornis*. *Limnol. Oceanogr.*, **52**, 79–84.
- Møller, E. F., Thor, P., and Nielsen, T. G. (2003) Production of DOC by *Calanus finmarchicus*, *C. glacialis* and *C. hyperboreus* through sloppy feeding and leakage from fecal pellets. *Mar. Ecol. Prog. Ser.*, **262**, 185–191.



- Möllmann, C., Folke, C., Edwards, M., and Conversi, A. (2014) Marine regime shifts around the globe: theory, drivers and impacts. *Phil. Trans. R. Soc. B*, **370**, 20130260.
- Montagnes, D. J. S., Lowe, C. D., Roberts, E. C., Breckels, M. N., Boakes, D. E., Davidson, K., Keeling, P. J., Slamovits, C. H., Steinke, M., Yang, Z., and Watts, P. C. (2011) An introduction to the special issue: *Oxyrrhis marina*, a model organism? *J. Plankton Res.*, **33**, 549–554.
- Nejstgaard, J. C., Hygum, B. H., Naustvoll, L.-J., and Båmstedt, U. (2001) Zooplankton growth, diet and reproductive success compared in simultaneous diatom- and flagellate-microzooplankton-dominated plankton blooms. *Mar. Ecol. Prog. Ser.*, **221**, 77–91.
- Newton, J. (2010) Stable isotope ecology. In: Encyclopedia of Life Sciences. John Wiley, Chichester.
- Niehoff, B., Kreibich, T., Saborowski, R., and Hagen, W. (2015) Feeding history can influence physiological responses of copepods: an experimental study comparing different cohorts of *Temora longicornis* from the Southern North Sea. *J. Exp. Mar. Biol. Ecol.*, **469**, 143–149.
- Nielsen, J. M., and Winder, M. (2015) Seasonal dynamics of zooplankton resource use revealed by carbon amino acid stable isotope values. *Mar. Ecol. Prog. Ser.*, **531**, 143–154.
- Nobili, R., Robinson, C., Buitenhuis, E., and Castellani, C. (2013) Food quality regulates the metabolism and reproduction of *Temora longicornis*. *Biogeosci. Discuss.*, **10**, 3203–3239.
- O’Brien, D. M., Boggs, C. L., and Fogel, M. L. (2004) The amino acids used in reproduction by butterflies: A comparative study of dietary sources using compound-specific stable isotope analysis. *Physiol. Biochem. Zool.*, **78**, 819–827.
- Ogrine, N., Košir, I. J., Spangenberg, J. E., and Kidrič, J. (2003) The application of NMR and MS methods for detection of adulteration of wine, fruit juices, and olive oil. A review. *Anal. Bioanal. Chem.*, **376**, 424–430.
- Paffenhöfer, G. A., Ianora, A., Miralto, A., Turner, J. T., Kleppel, G. S., Ribeira d’Alcalá, M., Casotti, R., Caldwell, G. S., et al. (2005) Colloquium on diatom–copepod interactions. *Mar. Ecol. Prog. Ser.*, **286**, 293–305.
- Parrish, C. C., and Wangersky, P. J. (1990) Growth and lipid class composition of the marine diatom, *Chaetoceros gracilis*, in laboratory and mass culture turbidostats. *J. Plankton Res.*, **12**, 1011–1021.
- Payne, M. R., Hatfield, E. M. C., Dickey-Collas, M., Falkenhaug, T., Gallego, A., Gröger, J., Licandro, P., Llope, M., et al. (2009) Recruitment in a changing environment: the 2000s North Sea herring recruitment failure. *ICES J. Mar. Sci.*, **66**, 272–277.
- Pecl, G. T., Araújo, M. B., Bell, J. D., Blanchard, J., Bonebrake, T. C., Chen, I., Clark, T. D. et al. (2017) Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science*, **355**, eaai9214.

## **REFERENCES**

---

- Pedersen, T. M., Almeda, R., Fotel, F. L., Jakobsen, H. H., Mariani, P., and Hansen, B. W. (2010) Larval growth in the dominant polychaete *Polydora ciliata* is food-limited in a eutrophic Danish estuary (Isefjord). *Mar. Ecol. Prog. Ser.*, **407**, 99–110.
- Pérez-Losada, M., Høeg, J. T., and Crandall, K. A. (2009) Remarkable convergent evolution in specialized parasitic Thecostraca (Crustacea). *BMC Biol.*, **7**, [doi:10.1186/1741-7007-7-15](https://doi.org/10.1186/1741-7007-7-15).
- Petersen, B., and Exo, K.-M. (1999) Predation of waders and gulls on *Lanice conchilega* tidal flats in the Wadden Sea. *Mar. Ecol. Prog. Ser.*, **178**, 229–240.
- Peterson, W. T., and Kimmerer, W. J. (1994) Processes controlling recruitment of the marine calanoid copepod *Temora longicornis* in Long Island Sound: Egg production, egg mortality, and cohort survival rates. *Limnol. Oceanogr.*, **39**, 1594–1605.
- Pizzetti, I., Fuchs, B. M., Gerdts, G., Wichels, A., Wiltshire, K. H., and Amann, R. (2011) Temporal variability of coastal *Planctomycetes* Clades at Kabeltonne Station, North Sea. *Appl. Environ. Microb.*, **77**, 5009-5017.
- Plath, K., and Boersma, M. (2001) Mineral limitation of zooplankton: stoichiometric constraints and optimal foraging. *Ecology*, **82**, 1260–1269.
- Pond, D., Harris, R., Head, R., and Harbour, D. (1996) Environmental and nutritional factors determining seasonal variability in the fecundity and egg viability of *Calanus helgolandicus* in coastal waters off Plymouth, UK. *Mar. Ecol. Prog. Ser.*, **143**, 45–63.
- Rabaut, M. (2009) *Lanice conchilega*, fisheries and marine conservation – Towards an ecosystem approach to marine management. Thesis, Ghent University, Belgium.
- Rabaut, M., Guilini, K., Van Hoey, G., Vincx, M., and Degraer, S. (2007) A bio-engineered soft-bottom environment: The impact of *Lanice conchilega* on the benthic species-specific densities and community structure. *Estuar. Coast. Shelf. S.*, **75**, 525–536.
- Reid, P. C., Borges, M. D., Svendsen, E. (2001) A regime shift in the North Sea circa 1988 linked to changes in the North Sea horse mackerel fishery. *Fish. Res.*, **50**, 163–171.
- Reid, P. C., Edwards, M., Beaugrand, G., Skogen, M., and Stevens, D. (2003) Periodic changes in the zooplankton of the North Sea during the twentieth century linked to oceanic inflow. *Fish. Oceanogr.*, **12**, 260–269.
- Reid, P. C., Hari, R. E., Beaugrand, G., Livingstone, D. M., Marty, C., Straile, D., Barichivich, J., et al. (2016). Global impacts of the 1980s regime shift. *Glob. Change Biol.*, **22**, 682–703.
- Reitan, K. I., Rainuzzo, J. R., and Olsen, Y. (1994) Effect of nutrient limitation on fatty acid and lipid content of marine microalgae. *J. Phycol.*, **30**, 972–979.

- Riebesell, U., Revill, A. T., Holdsworth, D. G., and Volkman, J. K. (2000) The effects of varying CO<sub>2</sub> concentration on lipid composition and carbon isotope fractionation in *Emiliana huxleyi*. *Geochim. Cosmochim. Acta*, **64**, 4179–4192.
- Rix, L., de Goeij, J. M., van Oevelen, D., Struck, U., Al-Horani, F. A., Wild, C., and Naumann, M. S. (2018) Reef sponges facilitate the transfer of coral-derived organic matter to their associated fauna via the sponge loop. *Mar. Ecol. Prog. Ser.*, **589**, 85–96.
- Roberts EC, Wootton EC, Davidson K, Jeong HJ, Lowe CD, Montagnes DJS (2011) Feeding in the dinoflagellate *Oxyrrhis marina*: linking behaviour with mechanisms. *Journal of Plankton Research* 33, 603-614.
- Rocha, J., Yletyinen, J., Biggs, R., Blenckner, T., and Peterson, G. (2014) Marine regime shifts: drivers and impacts on ecosystems services. *Phil. Trans. R. Soc. B*, **370**, 20130273.
- Roessler, P. G. (1990) Environmental control of glycerolipid metabolism in microalgae: Commercial implications and future research directions. *J. Phycol.*, **26**, 393–399.
- Ropert, M., and Dauvin, J.-C. (2000) Renewal and accumulation of a *Lanice conchilega* (Pallas) population in the baie des Veys, western Bay of Seine. *Oceanol. Acta*, **23**, 529–546.
- Ropert, M., and Gouletquer, P. (2000) Comparative physiological energetics of two suspension feeders: polychaete annelid *Lanice conchilega* (Pallas 1766) and Pacific cupped oyster *Crassostrea gigas* (Thunberg 1795). *Aquaculture*, **181**, 171–189.
- Roy, S., Harris, R. P., and Poulet, S. A. (1989) Inefficient feeding by *Calanus helgolandicus* and *Temora longicornis* on *Coscinodiscus wailesii*: quantitative estimation using chlorophyll-type pigments and effects on dissolved free amino acids. *Mar. Ecol. Prog. Ser.*, **52**, 145–153.
- Saiz, E., and Calbet, A. (2011) Copepod feeding in the ocean: scaling patterns, composition of their diet and the bias of estimates due to microzooplankton grazing during incubations. *Hydrobiologia*, **666**, 181–196.
- Scheffer, M., Carpenter, S., Foley, J. A., Folke, C., and Walker, B. (2001) Catastrophic shifts in ecosystems. *Nature*, **413**, 591–596.
- Schlüter, M. H., Merico, A., Reginatto, M., Boersma, M., Wiltshire, K. H., and Greve, W. (2010) Phenological shifts of three interacting zooplankton groups in relation to climate change. *Global Change Biol.*, **16**, 3144–3153.
- Schmidt, T. C., Zwank, L., Elsner, M., Berg, M., Meckenstock, R. U., and Haderlein, S. B. (2004) Compound-specific stable isotope analysis of organic contaminants in natural environments: a critical review of the state of the art, prospects, and future challenges. *Anal. Bioanal. Chem.*, **378**, 283–300.

## **REFERENCES**

---

- SERTC - Southeastern Regional Taxonomic Center (2018) Interactive Key to the Species of Chaetognatha from the South Atlantic Bight and northern Gulf of Mexico. Available at: <http://www.dnr.sc.gov/marine/sertc/Chaetognath%20key/Chaetognath%20key.htm>. Accessed on October 29, 2018.
- Simpson, I. A., Bol, R., Bull, I. D., Evershed, R. P., Petzke, K.-J., and Dockrill, S. J. (1999) Interpreting early land management through compound specific stable isotope analyses of archaeological soils. *Rapid Commun. Mass Spectrom.*, **13**, 1315–1319.
- Siuda, A. N. S., and Dam, H. G. (2010) Effects of omnivory and predator–prey elemental stoichiometry on planktonic trophic interactions. *Limnol. Oceanogr.*, **55**, 2107–2116.
- Smayda, T. J. (1997) Harmful algal blooms: Their ecophysiology and general relevance to phytoplankton blooms in the sea. *Limnol. Oceanogr.*, **42**, 1137–1153.
- Smetacek, V. S. (1999) Diatoms and the ocean carbon cycle. *Protist*, **150**, 25–32.
- Smith, B. N., and Epstein, S. (1971) Two categories of  $^{13}\text{C}/^{12}\text{C}$  ratios for higher plants. *Plant Physiol.*, **47**, 380–384.
- Sperfeld, E., Wagner, N. D., Halvorson, H. M., Malishev, M., and Raubenheimer, D. (2017) Bridging Ecological Stoichiometry and Nutritional Geometry with homeostasis concepts and integrative models of organism nutrition. *Funct. Ecol.*, **31**, 286–296.
- Stenseth, N. C., Ottersen, G., Hurrell, J. W., and Belgrano, A. (2004) Marine ecosystems and climate variation. The North Atlantic – A comparative perspective. Oxford University Press, Oxford. 252 pp.
- Sterner, R. W., and Elser, J. J. (2002) Ecological stoichiometry: the biology of elements from molecules to the biosphere. Princeton University Press.
- Sterner, R. W., and Schulz, K. L. (1998) Zooplankton nutrition: recent progress and a reality check. *Aquat. Ecol.*, **32**, 261–279.
- Strasser, M., and Pieloth, U. (2001) Recolonization pattern of the polychaete *Lanice conchilega* on an intertidal sand flat following the severe winter of 1995/96. *Helgol. Mar. Res.*, **55**, 176–181.
- Stübing, D. (2004) Lipid biochemistry of Antarctic euphausiids – energetic adaptations and a critical appraisal of trophic biomarkers. Thesis, Universität Bremen, Germany.
- Tang, K. W., and Dam, H. G. (1999) Limitation of zooplankton production: beyond stoichiometry. *Oikos*, **84**, 537–542.
- Taylor, A. H., Allen, J. I., and Clark, P. A. (2002) Extraction of a weak climatic signal by an ecosystem. *Nature*, **416**, 629–632.
- Thor, P. (2003) Elevated respiration rates of the neritic copepod *Acartia tonsa* during recovery from starvation. *J. Exp. Mar. Biol. Ecol.*, **283**, 133–143.

- Thor, P., and Wendt, I. (2010) Functional response of carbon absorption efficiency in the pelagic calanoid copepod *Acartia tonsa* Dana. *Limnol. Oceanogr.*, **55**, 1779–1789
- Thor, P., Koski, M., Tang, K. W., and Jónasdóttir, S. H. (2007) Supplemental effects of diet mixing on absorption of ingested organic carbon in the marine copepod *Acartia tonsa*. *Mar. Ecol. Prog. Ser.*, **331**, 131–138.
- Tobias, H. J., and Brenna, J. T. (2018) Cryofocus fast gas chromatography combustion isotope ratio mass spectrometry for rapid detection of synthetic steroid use in sport doping. *Analyst*, **143**, 1124–1132.
- Trillmich, F., and Ono, K. A. (1991). Pinnipeds and El Niño – Responses to environmental stress. Vol. 88. Springer Verlag, Berlin. 293 pp.
- van Donk, E., and Hessen, D. O. (1993) Grazing resistance in nutrient-stressed phytoplankton. *Oecologia*, **93**, 508–511.
- van Donk, E., and Hessen, D. O. (1995) Reduced digestibility of UV-B stressed and nutrient-limited algae by *Daphnia magna*. *Hydrobiologia*, **307**, 147–151.
- Van Hoey, G. (2006) Spatio-temporal variability within the macrobenthic *Abra alba* community, with emphasis on the structuring role of *Lanice conchilega*. Thesis, Ghent University, Belgium.
- Van Hoey, G., Guilini, K., Rabaut, M., Vincx, M., and Degraer, S. (2008) *Mar. Biol.*, **154**, 1009–1019.
- van Someren Gréve, H., Almeda, R., Lindegren, M., and Kiørboe, T. (2017) Gender-specific feeding rates in planktonic copepods with different feeding behavior. *Jo. Plankton Res.*, **39**, 631–644.
- Vanni, M. J., and McIntyre, P. B. (2016) Predicting nutrient excretion of aquatic animals with metabolic ecology and ecological stoichiometry: a global synthesis. *Ecology*, **97**, 3460–3471.
- Ventura, M. (2006) Linking biochemical and elemental composition in freshwater and marine crustacean zooplankton. *Mar. Ecol. Prog. Ser.*, **327**, 233–246.
- Vidal, E. A. G., Fuentes, L., and Silva, L. B. (2010) Defining *Octopus vulgaris* populations: A comparative study of the morphology and chromatophore pattern of paralarvae from Northeastern and Southwestern Atlantic. *Fish. Res.*, **106**, 199–208.
- Watts, P. C., Martin, L. E., Kimmance, S. A., Montagnes, D. J. S., and Lowe, C. D. (2011) The distribution of *Oxyrrhis marina*: a global disperser or poorly characterized endemic? *J. Plankton Res.*, **33**, 579–589.
- Weers, P. M. M., and Gulati, R. D. (1997) Growth and reproduction of *Daphnia galeata* in response to changes in fatty acids, phosphorus, and nitrogen in *Chlamydomonas reinhardtii*. *Limnol. Oceanogr.*, **42**, 1584–1589.

## **REFERENCES**

---

- Wegener, G., Bausch, M., Holler, T., Thang, N. M., Mollar, X. P., Kellermann, M. Y., Hinrichs, K-U., and Boetius, A. (2012) Assessing sub-seafloor microbial activity by combined stable isotope probing with deuterated water and <sup>13</sup>C-bicarbonate. *Env. Microbiol.*, **14**, 1517–1527.
- Welti, N., Striebel, M., Ulseth, A. J., Cross, W. F., DeVilbiss, S., Glibert, P. M., Guo, L., et al. (2017) Bridging food webs, ecosystem metabolism, and biogeochemistry using ecological stoichiometry theory. *Front. Microbiol.*, **8**, 1298.
- Wendt, I., and Thor, P. (2015) Influence of prey species and concentration on egg production efficiency and hatching success in *Acartia tonsa* Dana (Copepoda, Calanoida). *Crustaceana*, **88**, 675–687.
- Wernberg, T., Thomsen, M. S., Tuya, F., Kendrick, G. A., Staehr, P. A., and Toohey, B. D. (2010) Decreasing resilience of kelp beds along a latitudinal temperature gradient- potential implications for a warmer future. *Ecol. Lett.* **13**, 685–694.
- Wesche, A., Wiltshire, K. H., & Hirche, H. J. (2007). Overwintering strategies of dominant calanoid copepods in the German Bight, southern North Sea. *Marine Biology*, 151, 1309–1320. doi: 10.1007/s00227-006-0560-5
- Williams, R. J. P., and Fraústo da Silva, J. J. R. (1996) The natural selection of the chemical elements. The environment and life's chemistry. Clarendon Press, Oxford. 646 pp.
- Wiltshire, K. H., and Boersma, M. (2016) Meeting in the middle: on the interactions between microalgae and their predators or zooplankton and their food. In: Glibert, P. M., and Kana, T. M. (eds.), *Aquatic Microbial Ecology and Biogeochemistry: A dual perspective*. Springer, Switzerland.
- Wiltshire, K. H., and Manly, B. F. J. (2004) The warming trend at Helgoland Roads, North Sea: phytoplankton response. *Helgol. Mar. Res.*, **58**, 269–273.
- Wiltshire, K. H., Boersma, M., Carstens, K., Kraberg, A. C., Peters, S., and Scharfe, M. (2015) Control of phytoplankton in a shelf sea: Determination of the main drivers based on the Helgoland Roads Time Series. *J Sea Res.*, **105**, 42–52.
- Wiltshire, K. H., Kraberg, A., Bartsch, I., Boersma, M., Franke, H-D., Freund, J., Gebühr, C., Gerdtts, G., et al. (2010) Helgoland Roads, North Sea: 45 Years of Change. *Estuaries Coasts*, **33**, 295–310.
- Wiltshire, K. H., Malzahn, A. M., Wirtz, K., Greve, W., Janisch, S., Mangelsdorf, P., Manly, B. F. J., Boersma, M. (2008) Resilience of North Sea phytoplankton spring bloom dynamics: An analysis of long-term data at Helgoland Roads. *Limnol. Oceanogr.*, **53**, 1294–1302.
- WoRMS - World Record of Marine Species (2018) *Siriella clausii* (G.O. Sars, 1877) – Tab. 31. Available at: <http://www.marinespecies.org/aphia.php?p=image&tid=120210&pic=85617> Accessed on October 29, 2018.
- Zühlke, R. (2001) Polychaete tubes create ephemeral community patterns: *Lanice conchilega* (Pallas, 1766) associations studied over six years. *J. Sea Res.*, **46**, 261–272.

---

## ***REFERENCES***

Zühlke, R., Blome, D., Heinz van Bernem, K., and Dittmann, S. (1998). Effects of the tube-building polychaete *Lanice conchilega* (Pallas) on benthic macrofauna and nematodes in an intertidal sandflat. *Senck. Marit.*, **29**, 131–138.





Name: Rita Melo Franco Santos

Ort, Datum: Bremen, 01 November 2018

Anschrift: U1 6 Capri Drive, Sandy Bay, Hobart, 7005, Australia

## ERKLÄRUNG

Hiermit erkläre ich, dass ich die Doktorarbeit mit dem Titel:

Zooplankton performance in a changing ocean: Adaptive capacities to a shifting food regime in the  
North Sea

selbstständig verfasst und geschrieben habe und außer den angegebenen Quellen keine weiteren Hilfsmittel verwendet habe.

Ebenfalls erkläre ich hiermit, dass es sich bei den von mir abgegebenen Arbeiten um drei identische Exemplare handelt.

Rita Melo Franco Santos  
(Unterschrift)

