

A decomposition theorem for fuzzy set-valued random variables

Giacomo Aletti^{*1,2} and Enea G. Bongiorno^{†1}

¹Dipartimento di Matematica, Università degli Studi di Milano

²ADAMSS CENTRE (ADvanced Applied Mathematical and Statistical Sciences)

December 24, 2012

Abstract

In this paper, a decomposition theorem for a (square integrable) fuzzy random variable FRV is proposed. The paper is mainly divided in two parts. In the first part, for any FRV X , we define the Hukuhara set as the family of (deterministic) fuzzy sets C for which the Hukuhara difference $X \ominus_H C$ exists almost surely; in particular, we prove that such a family is a closed (with respect to different well known metrics) convex subset of the family of all fuzzy sets. In the second part, we prove that any square integrable FRV can be decomposed, up to a random translation, as the sum of a FRV Y and an element C' chosen uniquely (thanks to a minimization argument) in the Hukuhara set. This decomposition allows us to characterize fuzzy random translations; in particular, a FRV is a fuzzy random translation if and only if its Aumann expectation equals C' (given by the above decomposition) up to a deterministic translation. Examples and open problems are also presented.

Keywords: Fuzzy random variable; Hukuhara difference; decomposition theorem; randomness defuzzification;

Revised Personal Version of the text of the final journal article:

G. Aletti, E. G. Bongiorno. *A decomposition theorem for fuzzy set-valued random variables*. Fuzzy Sets and Systems. (doi: 10.1016/j.fss.2012.11.005).
<http://dx.doi.org/10.1016/j.fss.2012.11.005>

It reflects changes made in the peer review process.

1 Introduction

It is widely known (e.g. [6, Theorem 6.1.7]) that a Gaussian fuzzy random variable (FRV) may be decomposed as

$$X = \mathbb{E}X \oplus \mathbb{I}_\xi, \quad (1)$$

where $\mathbb{E}X$ is the expectation of X in the Aumann sense, ξ is a Gaussian random element in \mathbb{R}^d with $\mathbb{E}\xi = 0$ and $\mathbb{I}_A : \mathbb{R}^d \rightarrow \{0, 1\}$ denotes the indicator function of any $A \subseteq \mathbb{R}^d$

$$\mathbb{I}_A(x) = \begin{cases} 1, & \text{if } x \in A, \\ 0, & \text{otherwise.} \end{cases}$$

We write \mathbb{I}_a instead of $\mathbb{I}_{\{a\}}$ whenever $A = \{a\}$ is a singleton. Roughly speaking, a Gaussian FRV X is just a deterministic fuzzy set (its expected value $\mathbb{E}X$) up to a Gaussian translation ξ which carries out all the randomness of X . In this view, Equation (1) entails a *randomness defuzzification* for the Gaussian FRV X according to which the underlying probability structure can be defined just only on \mathbb{R}^d and no

*giacomo.aletti@unimi.it

†enea.bongiorno@unimi.it

longer on \mathbb{F} , the space of normal fuzzy sets with compact convex level sets. Even the defuzzification is complete both in “shape” and in randomness when one tries to define a fuzzy Brownian motion: if X is a fuzzy Gaussian process with null expectation (in some sense), then $X = \mathbb{I}_{b_t}$ with b_t a brownian motion in \mathbb{R}^d , see [2] for details. Clearly, such randomness defuzzification occurs also in the case of a random translation ξ (not necessarily Gaussian) of a deterministic fuzzy set ν ; and Equation (1) still holds with $\mathbb{E}X$ being equal ν up to a translation. From the modeling point of view, the (partial or complete) defuzzification occurrence results in a simplification of the framework chosen to describe the considered phenomenon, so that X can be decomposed as in (1). In other words, defuzzification may tend to mislead the chosen setting: one needs less than a fuzzy framework.

In this paper, we propose a generalization of Equation (1): any FRV $X \in L^2[\mathbb{F}]$ can be decomposed by means of

$$X(\omega) = C \oplus Y(\omega) \oplus \mathbb{I}_{\text{Ste}(X(\omega))}, \quad (2)$$

for \mathbb{P} -almost every $\omega \in \Omega$. In particular, we start to consider a \mathbb{P} -almost surely centered FRV $X \in L^2[\mathbb{F}]$ and the family Θ_X of all centered deterministic $\eta \in \mathbb{F}$ for which the Hukuhara difference $X \ominus_H \eta$ exists almost surely. Then, we prove that Θ_X is not empty, convex and closed in \mathbb{F} with respect to ρ , where ρ can be chosen between d_2 (that corresponds to the L^2 metric in the space of support functions) and D_θ (the mid-spread metric; see [5, 11]). Further,

$$C = \arg \min_{\eta \in \Theta_X} \mathbb{E}[\rho(X, \eta)^2]$$

is unique and there exists a (\mathbb{P} -almost surely centered) FRV Y such that $X(\omega) = C \oplus Y(\omega)$. Whenever X is not centered, the decomposition is given by (2). Hence, C and $Y \oplus \mathbb{I}_{\text{Ste}(X)}$ may be considered as the deterministic part (with respect to \oplus) and the random part of X respectively.

In view of such a decomposition, we are able to handle all kind of defuzzifications: they occur whenever Y or both C and Y are not fuzzy set-valued. In particular, we have a defuzzification in randomness or a complete defuzzification respectively.

In the first case (i.e. Y is \mathbb{P} -almost surely \mathbb{I}_0), X is a random fuzzy translation of C . Further, because the Aumann expectation $\mathbb{E}X$ is the (unique) Frèchet expectation with respect to d_2 , i.e.

$$\mathbb{E}X = \arg \min_{\eta \in \mathbb{F}} \mathbb{E}[d_2(X, \eta)^2],$$

we obtain immediately that a FRV X is a random translation of C if and only if $\mathbb{E}X$ is equal to C up to a translation. (Clearly a fuzzy random translation may be even characterize by its variance: in fact, let $\text{Var}(X) = \mathbb{E}[d_2(X, \mathbb{E}X)^2]$ denote the Frèchet variance of the FRV X , then X is a fuzzy random translation if and only if $\text{Var}(X - \text{Ste}(X)) = 0$.)

The existence and uniqueness of decomposition (2) seem to depend on the chosen metric ρ . In fact, problems may arise if different metrics are considered, in this paper, besides d_2 and D_θ , we discuss the case of ρ being the Hausdorff metric δ_H^∞ .

The paper is organized as following. Section 2 introduces necessary notations and literature results. Section 3 studies properties of the Hukuhara set Θ_X whilst Section 4 presents the decomposition theorem of FRV, provides some examples and open problems.

2 Preliminaries

Denote by \mathbb{K} the class of non-empty compact convex subsets of \mathbb{R}^d , endowed with the Hausdorff metric

$$\delta_H(A, B) = \max\{\sup_{a \in A} \inf_{b \in B} \|a - b\|, \sup_{b \in B} \inf_{a \in A} \|a - b\|\},$$

and the operations

$$A + B = \{a + b : a \in A, b \in B\}, \quad \lambda \cdot A = \lambda A = \{\lambda a : a \in A\} \quad \text{with } \lambda > 0.$$

For a non-empty closed convex set $A \subset \mathbb{R}^d$ the *support function* $s_A : S^{d-1} \rightarrow \mathbb{R}$ is defined by

$$s_A(x) = \sup\{\langle x, a \rangle : a \in A\}, \quad \text{for } x \in S^{d-1},$$

where $\langle \cdot, \cdot \rangle$ is the scalar product in \mathbb{R}^d and $S^{d-1} = \{x \in \mathbb{R}^d : \|x\| = 1\}$ is the unit sphere in \mathbb{R}^d . The *Steiner point* of $A \in \mathbb{K}$ is defined by

$$\mathbf{ste}(A) = \frac{1}{v_d} \int_{S^{d-1}} x s_A(x) \, dx$$

where $x \in S^{d-1}$ varies over the unit vectors of \mathbb{R}^d , dx is the Lebesgue measure on S^{d-1} , and v_d is the volume of the unit ball of \mathbb{R}^d .

Fuzzy Sets. A *fuzzy set* is a map $\nu : \mathbb{R}^d \rightarrow [0, 1]$. Let \mathbb{F} denote the family of all fuzzy sets ν , which satisfy the following conditions.

1. ν is an upper semicontinuous function, i.e. for each $\alpha \in (0, 1]$, the *cut set* or the α -*level set* $\nu_\alpha = \{x \in \mathbb{R}^d : \nu(x) \geq \alpha\}$ is a closed subset of \mathbb{R}^d .
2. ν is normal; i.e. $\nu_1 = \{x \in \mathbb{R}^d : \nu(x) = 1\} \neq \emptyset$.
3. The support set $\nu_0 = \overline{\{x \in \mathbb{R}^d : \nu(x) > 0\}}$ of ν is compact; hence every ν_α is compact for $\alpha \in (0, 1]$.
4. For any $\alpha \in [0, 1]$, ν_α is a convex subset of \mathbb{R}^d .

For any $\nu \in \mathbb{F}$ define the *support function* of ν as follows:

$$s_\nu(x, \alpha) = s_{\nu_\alpha}(x), \quad (3)$$

for $(x, \alpha) \in S^{d-1} \times [0, 1]$. Let us endow \mathbb{F} with the operations

$$(\nu \oplus \eta)_\alpha = \nu_\alpha + \eta_\alpha, \quad (\lambda \nu)_\alpha = (\lambda \odot \nu)_\alpha = \lambda \cdot \nu_\alpha, \quad \text{with } \nu, \eta \in \mathbb{F}, \lambda > 0 \text{ and } \alpha \in [0, 1]$$

(so that $(\mathbb{F}, \oplus, \odot)$ is a convex cone), and with the metrics

$$\delta_H^\infty(\nu, \eta) = \sup\{\alpha \in [0, 1] : \delta_H(\nu_\alpha, \eta_\alpha)\},$$

$$d_2(\nu, \eta) = \left(\int_0^1 \int_{S^{d-1}} |s_\nu(\alpha, u) - s_\eta(\alpha, u)|^2 \, du \, d\alpha \right)^{\frac{1}{2}},$$

where $d\alpha$ and du denote the Lebesgue measure on $[0, 1]$ and the normalized Lebesgue measure on S^{d-1} respectively. It is known that $(\mathbb{F}, \delta_H^\infty)$ is a complete metric space while (\mathbb{F}, d_2) is not (cf. [4, Chapter 7]); in the following we even use the mid-spread metric D_θ that we introduce later in this section.

Let $\nu, \eta \in \mathbb{F}$, the Hukuhara difference $\nu \ominus_H \eta$ is defined by

$$\nu \ominus_H \eta = \tau \quad \Leftrightarrow \quad \tau \oplus \eta = \nu,$$

whenever τ exists (in \mathbb{F}), it is unique.

The *generalized Steiner point* of $\nu \in \mathbb{F}$ is defined by

$$\mathbf{Ste}(\nu) = \int_{[0,1]} \mathbf{ste}(\nu_\alpha) \, d\alpha,$$

where $d\alpha$ is the Lebesgue measure on $[0, 1]$. In other words, $\mathbf{Ste}(\nu)$ may be seen as a weighted average of steiner points of the level sets of ν . The following properties are satisfied (cf. [12]).

1. For any $\nu \in \mathbb{F}$, $\mathbf{Ste}(\nu) \in \nu_0$.
2. For any $\nu, \eta \in \mathbb{F}$, $\mathbf{Ste}(\nu \oplus \eta) = \mathbf{Ste}(\nu) + \mathbf{Ste}(\eta)$.
3. $\mathbf{Ste} : \mathbb{F} \rightarrow \mathbb{R}^d$ is continuous with respect to δ_H^∞ (and hence with respect to d_2) and to the euclidean distance on \mathbb{R}^d .

On the support functions of fuzzy sets; Bobylev's characterization of a fuzzy set.

The support function for a fuzzy set $\nu \in \mathbb{F}$ can be defined equivalently on the closed unit ball $B(0, 1) = \{x \in \mathbb{R}^d : \|x\| \leq 1\} \subset \mathbb{R}^d$ instead of the unit sphere S^{d-1} by

$$s_\nu^* : \begin{array}{l} B(0, 1) \rightarrow \mathbb{R} \\ x \mapsto s_\nu^*(x) = \max\{\langle x, y \rangle : y \in \mathbb{R}^d, \nu(y) \geq \|x\|\}. \end{array} \quad (4)$$

Note that the maximum is actually attained since it is being applied to a linear function over the compact set $\nu_{\|x\|}$. The first step is to analyze the relationship between both definitions of support function. The following results gather this relationship.

Proposition 1 *Let $\nu \in \mathbb{F}$. Let s_ν and s_ν^* be defined by Equation (3) and Equation (4) respectively.*

- If we define $f : S^{d-1} \times [0, 1] \rightarrow \mathbb{R}$ as

$$f(x, \alpha) = \begin{cases} \frac{1}{\alpha} s_\nu^*(\alpha x), & \text{if } \alpha \neq 0, \\ \sup_{y \in \nu_0} \langle y, x \rangle, & \text{if } \alpha = 0, \end{cases}$$

then $f = s_\nu$.

- If we define $f^* : B(0, 1) \rightarrow \mathbb{R}$ as

$$f^*(x) = \begin{cases} \|x\| s_\nu \left(\frac{x}{\|x\|}, \|x\| \right), & \text{if } x \neq 0, \\ 0, & \text{if } x = 0, \end{cases}$$

then $f^* = s_\nu^*$.

Proof. First case. For all $\alpha \in (0, 1]$, we have

$$\begin{aligned} f(x, \alpha) &= \frac{1}{\alpha} s_\nu^*(\alpha x) = \frac{1}{\alpha} \max\{\langle \alpha x, y \rangle : y \in \mathbb{R}^d, \nu(y) \geq \|\alpha x\| = \alpha\} \\ &= \max\{\langle x, y \rangle : y \in \mathbb{R}^d, \nu(y) \geq \alpha\} = \sup\{\langle x, y \rangle : y \in \nu_\alpha\} = s_\nu(x, \alpha). \end{aligned}$$

Finally, by definition, $f(x, 0) = s_\nu(x, 0)$ for each $x \in S^{d-1}$.

Second case. Let $x \in B(0, 1) \setminus \{0\}$, thus we have

$$\begin{aligned} f^*(x) &= \|x\| s_\nu \left(\frac{x}{\|x\|}, \|x\| \right) = \|x\| \sup \left\{ \left\langle \frac{x}{\|x\|}, y \right\rangle : y \in \nu_{\|x\|} \right\} \\ &= \max\{\langle x, y \rangle : y \in \nu_{\|x\|}\} = \max\{\langle x, y \rangle : y \in \mathbb{R}^d, \nu(y) \geq \|x\|\} = s_\nu^*(x). \end{aligned}$$

Finally, $f^*(0) = 0 = s_\nu^*(0)$. ■

Since s_ν and s_ν^* are equivalent ways to identify a fuzzy sets, they can be used interchangeably. In the following we will use mainly s_ν . We will use s_ν^* only in the first part of the proof of Proposition 5 in combination with the Bobylev's characterization for fuzzy sets (Proposition 2).

Proposition 2 (Bobylev's characterization, [1, 4]) *Let $f^* : B(0, 1) \rightarrow \mathbb{R}$. It is a support function (as defined by Equation (4)) of some fuzzy set $\nu \in \mathbb{F}$ if and only if f^* satisfies the following properties*

(P-1) f^* is upper semicontinuous, i.e.,

$$f^*(x) = \limsup_{y \rightarrow x} f^*(y), \quad \forall x \in B(0, 1).$$

(P-2) f^* is positively semihomogeneous, i.e.,

$$\lambda f^*(x) \leq f^*(\lambda x), \quad \forall \lambda \in (0, 1], \forall x \in B(0, 1).$$

(P-3) f^* is quasiadditive, i.e.,

$$\|x\| f^* \left(\lambda \frac{x}{\|x\|} \right) \leq \|x_1\| f^* \left(\lambda \frac{x_1}{\|x_1\|} \right) + \|x_2\| f^* \left(\lambda \frac{x_2}{\|x_2\|} \right),$$

for every $\lambda \in (0, 1]$, and $x, x_1, x_2 \in \mathbb{R}^d \setminus \{0\}$, with $x = x_1 + x_2$.

(P-4) f^* is normal, i.e.,

$$f^*(x) + f^*(-x) \geq 0, \quad \forall x \in B(0, 1).$$

(P-5) $f^*(\cdot)/\|\cdot\|$ is bounded, i.e.,

$$\sup \{|f^*(x)|/\|x\| : x \in B(0, 1) \setminus \{0\}\} < +\infty.$$

(P-6) $f^*(0) = 0$.

In particular, $\nu \in \mathbb{F}$ is such that,

- for $x \in \mathbb{R}^d$, $\nu(x) = \max\{\alpha \in [0, 1] : \langle x, y \rangle \leq f^*(y), \forall y \in B(0, 1), |y| = \alpha\}$,
- for $\alpha \in (0, 1]$, $\nu_\alpha = \{x \in \mathbb{R}^d : \langle x, y \rangle \leq f^*(y), \forall y \in B(0, 1), |y| = \alpha\}$,
- $\nu_0 = \bigcup_{\alpha \in (0, 1]} \nu_\alpha$.

Embeddings and metrics. Let $C(S^{d-1})$ denote the Banach space of all continuous functions v on S^{d-1} with respect to the norm $\|v\|_C = \sup_{x \in S^{d-1}} |v(x)|$. Let $\overline{\mathcal{C}} := \overline{\mathcal{C}}([0, 1], C(S^{d-1}))$ be the set of all functions $f : [0, 1] \rightarrow C(S^{d-1})$ such that f is bounded, left continuous with respect to $\alpha \in (0, 1]$, right continuous at 0, and f has right limit for any $\alpha \in (0, 1)$. Then we have that $\overline{\mathcal{C}}$ is a Banach space with the norm $\|f\|_{\overline{\mathcal{C}}} = \sup_{\alpha \in [0, 1]} \|f(\alpha)\|_C$.

Let $\mathcal{L} := L^2([0, 1] \times S^{d-1}; \mathbb{R})$ be the Hilbert space of square integrable real-valued functions defined on $[0, 1] \times S^{d-1}$ and let us denote by $\langle \cdot, \cdot \rangle_{\mathcal{L}}$ and $\|\cdot\|_{\mathcal{L}}$ its usual inner product and norm respectively.

It is known, cf. [6, 9, 10], that the injection j defined by

$$\begin{aligned} j : \mathbb{F} &\rightarrow \overline{\mathcal{C}} \cap \mathcal{L} \\ \nu &\mapsto j(\nu) = s_\nu, \end{aligned} \tag{5}$$

satisfies the following properties:

1. $j(\lambda_1 \nu \oplus \lambda_2 \eta) = \lambda_1 j(\nu) + \lambda_2 j(\eta)$, $\nu, \eta \in \mathbb{F}$ and $\lambda_1, \lambda_2 \geq 0$.
2. j is an isometric mapping, i.e. for every $\nu, \eta \in \mathbb{F}$,

$$\delta_H^\infty(\nu, \eta) = \|j(\nu) - j(\eta)\|_{\overline{\mathcal{C}}}, \quad \text{and} \quad d_2(\nu, \eta) = \|j(\nu) - j(\eta)\|_{\mathcal{L}}.$$

The *mid-spread metric* D_θ^φ , introduced by [11] and extended in [5], is defined as follow. Let $f \in \mathcal{L}$, it can be decomposed as $f = \text{mid } f + \text{spr } f$ where

$$\text{mid } f(u, \alpha) = \frac{f(u, \alpha) - f(-u, \alpha)}{2}, \quad \text{spr } f(u, \alpha) = \frac{f(u, \alpha) + f(-u, \alpha)}{2},$$

with $(u, \alpha) \in S^{d-1} \times [0, 1]$. Let $\theta \in (0, \infty)$. For $f, g \in \mathcal{L}$ consider the value

$$\langle f, g \rangle_\theta^\varphi = [\text{mid } f, \text{mid } g]^\varphi + \theta [\text{spr } f, \text{spr } g]^\varphi,$$

with

$$[f, g]^\varphi = \int_0^1 \int_{S^{d-1}} f(u, \alpha) g(u, \alpha) \, du \, d\varphi(\alpha),$$

where φ is a weighting probability measure, absolutely continuous (with respect to the Lebesgue measure $d\alpha$) probability measure on $([0, 1], \mathcal{B}_{[0, 1]})$ with positive mass function in $(0, 1)$. $(\mathcal{L}, \langle \cdot, \cdot \rangle_\theta^\varphi)$ is a separable Hilbert space (whose norm is denoted by $\|\cdot\|_\theta^\varphi$). Then, embedding $j : \mathbb{F} \rightarrow \mathcal{L}$ leads to the following metric definition on \mathbb{F}

$$D_\theta^\varphi(\nu, \eta) = \langle s_\nu - s_\eta, s_\nu - s_\eta \rangle_\theta^\varphi = (\|s_\nu - s_\eta\|_\theta^\varphi)^2,$$

for any $\nu, \eta \in \mathbb{F}$. We omit φ and we write $\langle \cdot, \cdot \rangle_\theta$, $\|\cdot\|_\theta$ and D_θ whenever φ is the Lebesgue measure on $[0, 1]$. Note that, the topology induced by D_θ on \mathbb{F} is equivalent to the one induced by d_2 :

$$\begin{aligned} (\|f\|_{\mathcal{L}})^2 &\leq (\|f\|_\theta)^2 \leq \theta (\|f\|_{\mathcal{L}})^2, & \text{if } \theta > 1, \\ \theta (\|f\|_{\mathcal{L}})^2 &\leq (\|f\|_\theta)^2 \leq (\|f\|_{\mathcal{L}})^2, & \text{if } 0 < \theta \leq 1. \end{aligned}$$

Note that $\|\cdot\|_\theta^\varphi$ remains equivalent to $\|\cdot\|_{\mathcal{L}}$ in a lot of likely situations (as example, whenever the Radon–Nikodym derivative satisfies $0 < h \leq \frac{d\varphi}{d\alpha} \leq H$ for some $h, H \in \mathbb{R}$).

Fuzzy random variables. Let $(\Omega, \mathfrak{F}, \mathbb{P})$ be a complete probability space. A *fuzzy set-valued random variable* (FRV) is a function $X : \Omega \rightarrow \mathbb{F}$, such that $X_\alpha : \omega \mapsto X(\omega)_\alpha$ are random compact convex sets for every $\alpha \in (0, 1]$ (i.e. X_α is a \mathbb{K} -valued function measurable w.r.t. $\mathcal{B}_{\mathbb{K}}$, the Borel σ -algebra on \mathbb{K} generated by the metric δ_H). It has been proven in [3] that this measurability definition is equivalent to the $\mathcal{B}(\mathbb{F}, d_2)$ -measurability, hence to the $\mathcal{B}(\mathbb{F}, D_\theta)$ -measurability, and, it is necessary (but not sufficient) for the $\mathcal{B}(\mathbb{F}, \delta_H^\infty)$ -measurability, where $\mathcal{B}(\mathbb{F}, D)$ is the Borel σ -algebra defined on \mathbb{F} w.r.t. the metric D . As a consequence of continuity of $\mathbf{Ste}(\cdot)$, if X is a FRV, then $\mathbf{Ste}(X)$ is a random element in \mathbb{R}^d . A FRV X is *integrably bounded* and we write $X \in L^1[\Omega; \mathbb{F}]$, if $\mathbb{E}[\sup_{x \in X_0} \|x\|] < +\infty$. The (Aumann) *expected value* of $X \in L^1[\Omega; \mathbb{F}]$, denoted by $\mathbb{E}[X]$, is a fuzzy set such that, for every $\alpha \in [0, 1]$,

$$(\mathbb{E}[X])_\alpha = \int_{\Omega} X_\alpha \, d\mathbb{P} = \{\mathbb{E}(f) : f \in L^1[\Omega; \mathbb{R}^d], f \in X_\alpha \, \mathbb{P} - \text{a.e.}\}.$$

It should be pointed out that, whenever $\mathbb{E}[(\sup_{x \in X_0} \|x\|)^2] < +\infty$ (we write $X \in L^2[\Omega; \mathbb{F}]$), the expected value in the Aumann's sense is even the Fréchet expectation with respect to d_2 (e.g. [8]), i.e.

$$\mathbb{E}X = \arg \min_{\eta \in \mathbb{F}} \mathbb{E}[d_2(X, \eta)^2].$$

In view of above measurability consideration and from embedding (5) it follows that every FRV X can be regarded as a random element in \mathcal{L} , where $s_X(\cdot, \cdot)(\omega) = s_{X(\omega)}(\cdot, \cdot)$. Moreover, if $X \in L^1[\Omega; \mathbb{F}]$, for any $(x, \alpha) \in \mathbb{R}^d \times [0, 1]$, $s_{X(\cdot)}(x, \alpha) \in L^1[\Omega; \mathbb{R}]$ and

$$\mathbb{E}[s_X(x, \alpha)] = s_{\mathbb{E}X}(x, \alpha). \quad (6)$$

Finally, let $L^2[\Omega; \mathcal{L}] := \{f : \Omega \rightarrow \mathcal{L} \text{ s.t. } (\mathbb{E}[\|f\|_{\mathcal{L}}^2])^{1/2} < +\infty\}$. It is a Hilbert space with inner product $\mathbb{E}\langle f, g \rangle_{\mathcal{L}}$. Then, the map

$$\begin{aligned} J : L^2[\Omega; \mathbb{F}] &\rightarrow L^2[\Omega; \mathcal{L}] \\ X &\mapsto J(X) = j(X(\cdot)) = s_{X(\cdot)}, \end{aligned}$$

is well-defined and induces an isometry on $L^2[\Omega; \mathcal{L}]$; in particular, for every $X, Y \in L^2[\Omega; \mathbb{F}]$,

$$(\Delta_2(X, Y))^2 := \mathbb{E}[d_2(X, Y)^2] = \mathbb{E}[\langle J(X) - J(Y), J(X) - J(Y) \rangle_{\mathcal{L}}].$$

Clearly, $L^2[\Omega; \mathbb{F}]$ is, by means of J , even isometric to (the Hilbert space) $(L^2[\Omega; \mathcal{L}], \mathbb{E}\langle \cdot, \cdot \rangle_{\theta})$, with

$$(\Delta_{2, \theta}(X, Y))^2 := \mathbb{E}[D_\theta(X, Y)^2] = \mathbb{E}[\langle J(X) - J(Y), J(X) - J(Y) \rangle_{\theta}].$$

3 Hukuhara set

In this section we define the *Hukuhara set* associated to a FRV X , namely Θ_X . We provide some properties of Θ_X most of which turn out to be useful in the next section where a decomposition theorem for fuzzy random variables is set.

Let ν be in \mathbb{F} such that $\mathbf{Ste}(\nu) = 0$ and consider

$$\begin{aligned} \theta_\nu &= \{\eta \in \mathbb{F} : \mathbf{Ste}(\eta) = 0 \text{ and } \exists (\nu \ominus_H \eta)\} \\ &= \{\eta \in \mathbb{F} : \mathbf{Ste}(\eta) = 0 \text{ and } \exists \tau \in \mathbb{F} \text{ s.t. } \eta \oplus \tau = \nu\}; \end{aligned}$$

i.e. the family of those centered convex compact fuzzy sets η for which the Hukuhara difference $\nu \ominus_H \eta$ does exist. Note that θ_ν is not empty, since $\mathbb{I}_0, \nu \in \theta_\nu$ and $\{\lambda \odot \nu\}_{\lambda \in [0, 1]} \subseteq \theta_\nu$. Clearly, if $\eta \in \theta_\nu$ and τ is the Hukuhara difference between ν and η , then $\tau \in \theta_\nu$.

Proposition 3 θ_ν is a closed subset in $(\mathbb{F}, \delta_H^\infty)$.

Proof. Let $\{\eta_n\} \subset \theta_\nu$ be a convergent sequence with limit $\eta \in \mathbb{F}$ with respect to δ_H^∞ , we have to prove that $\eta \in \theta_\nu$. Equivalently, we have to prove that there exists $\tau \in \mathbb{F}$ such that $\eta \oplus \tau = \nu$. For each

$n = 1, 2, \dots$ there exist $\tau_n \in \mathbb{F}$ such that $\eta_n \oplus \tau_n = \nu$. Thus, the idea is to prove that $\{\tau_n\}_{n=1}^\infty$ converges, w.r.t. δ_H^∞ , to some $\tau \in \mathbb{F}$ such that $\eta \oplus \tau = \nu$. Note that

$$\begin{aligned} \delta_H^\infty(\tau_m, \tau_n) &= \|s_{\tau_m} - s_{\tau_n}\|_{\overline{C}} \\ &= \|(s_{\tau_m} + s_{\eta_m}) - (s_{\tau_n} + s_{\eta_n}) + s_{\eta_n} - s_{\eta_m}\|_{\overline{C}} \\ &= \|s_\nu - s_\nu + s_{\eta_n} - s_{\eta_m}\|_{\overline{C}} \\ &= \|s_{\eta_n} - s_{\eta_m}\|_{\overline{C}} = \delta_H^\infty(\eta_n, \eta_m) \rightarrow 0, \quad \text{for } n, m \rightarrow \infty \end{aligned}$$

where we use the isometry $\tau \mapsto s_\tau$ (first and last equalities) and the fact that η_n, η_m belong to θ_ν (third equality). Thus $\{\tau_n\}_{n \in \mathbb{N}}$ is a Cauchy sequence in $(\mathbb{F}, \delta_H^\infty)$ that is a complete metric space (e.g. [6, Theorem 5.1.6]), and hence there exists τ in \mathbb{F} such that $\tau_n \rightarrow \tau$. As a consequence, $\eta_n \oplus \tau_n \rightarrow \eta \oplus \tau$ for $n \rightarrow \infty$ combined with

$$0 = \delta_H^\infty(\eta_n \oplus \tau_n, \nu),$$

guarantees that $\eta \oplus \tau = \nu$ and hence $\eta \in \theta_\nu$; that is the thesis. \blacksquare

In what follows we need the next lemma according to which a fuzzy set can be defined starting from its α -cuts.

Lemma 4 (See [4, Proposition 6.1.7, p.39]) *If $\{C_\alpha\}_{\alpha \in [0,1]}$ satisfies*

- (a) C_α is a non empty, compact, convex subset of \mathbb{R}^d , for every $\alpha \in [0, 1]$;
- (b) $C_\beta \subseteq C_\alpha$ for $0 \leq \alpha \leq \beta \leq 1$;
- (c) $C_\alpha = \bigcap_{i=1}^\infty C_{\alpha_i}$ for all sequences $\{\alpha_i\}_{i \in \mathbb{N}}$ in $[0, 1]$ converging from below to α , i.e. $\alpha_i \uparrow \alpha$ in $[0, 1]$;

then the function

$$\nu(x) = \begin{cases} 0, & \text{if } x \notin C_0, \\ \sup\{\alpha \in [0, 1] : x \in C_\alpha\}, & \text{if } x \in C_0, \end{cases}$$

is an element of \mathbb{F} with $\nu_\alpha = C_\alpha$ for any $\alpha \in (0, 1]$ and

$$\nu_0 = \overline{\bigcup_{\alpha \in (0,1]} C_\alpha} \subseteq C_0.$$

Let X be a FRV. For the sake of simplicity and without loss of generality, let us suppose that $\mathbf{Ste}(X) = 0$; otherwise one can always consider its associated centered FRV $\tilde{X} = X - \mathbb{1}_{\mathbf{Ste}(X)}$. The next theorem defines the *Hukuhara set* Θ_X associated to X , and provides some properties of Θ_X .

Proposition 5 *Let X be a FRV with $\mathbf{Ste}(X) = 0$. If $\eta \in \mathbb{F}$, then $E = \{\eta \in \theta_X\} := \{\omega \in \Omega : \eta \in \theta_{X(\omega)}\}$ is measurable in (Ω, \mathfrak{F}) . Moreover, if $\Theta_X = \{\eta \in \mathbb{F} : \mathbb{P}(\eta \in \theta_X) = 1\}$, then the following statements hold.*

- (i) Θ_X is non-empty: $\mathbb{1}_0 \in \Theta_X$.
- (ii) $\eta \in \Theta_X$ if and only if there exist a FRV Y such that $\eta \oplus Y = X$, \mathbb{P} -a.s.. If $X \in L^2[\Omega; \mathbb{F}]$, then Y is in $L^2[\Omega; \mathbb{F}]$ too.
- (iii) Θ_X is a convex subset in (\mathbb{F}, \oplus) . As a consequence, if $\eta \in \Theta_X$, then $\{\lambda\eta\}_{\lambda \in [0,1]} \subseteq \Theta_X$ with $0 \cdot \eta = \mathbb{1}_0 \in \Theta_X$.
- (iv) Θ_X is a closed subset of $(\mathbb{F}, \delta_H^\infty)$.
- (v) Θ_X is a closed subset of (\mathbb{F}, d_2) .
- (vi) Θ_X is a closed subset of (\mathbb{F}, D_θ) and, more in general, of $(\mathbb{F}, D_\theta^\varphi)$.

Proof. By definition of $\theta_{X(\omega)}$, we have

$$\begin{aligned} E &= \{\omega \in \Omega : \mathbf{Ste}(\eta) = 0 \text{ and } \exists (X(\omega) \ominus_H \eta)\} \\ &= \{\omega \in \Omega : \mathbf{Ste}(\eta) = 0\} \cap \{\omega \in \Omega : \exists Y_\omega \in \mathbb{F}, \text{ s.t. } s_\eta + s_{Y_\omega} = s_{X(\omega)}\}, \end{aligned}$$

where $s_{X(\omega)}$ and s_η are defined as in (3). Let f_ω be the real-valued map defined on $S^{d-1} \times [0, 1]$ by $f_\omega = s_{X(\omega)} - s_\eta$, and define $f_\omega^* : B(0, 1) \rightarrow \mathbb{R}$ as

$$f_\omega^*(x) = \begin{cases} \|x\| f_\omega\left(\frac{x}{\|x\|}, \|x\|\right), & \text{if } x \neq 0, \\ 0, & \text{if } x = 0, \end{cases}$$

then, thanks to Bobylev's characterization (Proposition 2),

$$\begin{aligned} E &= \{\omega \in \Omega : \mathbf{Ste}(\eta) = 0\} \cap \{\omega \in \Omega : f_\omega^* \text{ satisfies (P-1), } \dots, \text{ (P-6)}\} \\ &= E_0 \cap E_1 \cap \dots \cap E_6, \end{aligned}$$

where $E_0 = \{\omega \in \Omega : \mathbf{Ste}(\eta) = 0\}$ and $E_i = \{\omega \in \Omega : f_\omega^* \text{ satisfies (P-}i\text{)}\}$ for $i = 1, \dots, 6$. Clearly, E is measurable if E_0, \dots, E_6 are so. Since η is a deterministic fuzzy set, E_0 is either the empty set or the whole Ω ; hence E_0 is measurable. Further, note that each E_i ($i = 1, \dots, 4, 6$) can be written as $E_i = \{\omega : g_i(\omega) \leq 0\}$ and $E_5 = \{\omega : g_5(\omega) < +\infty\}$ where

$$\begin{aligned} g_1 &= \sup\{|\limsup_{y \rightarrow x} f_\omega^*(y) - f_\omega^*(x)| : x \in B(0, 1)\}, \\ g_2 &= \sup\{\lambda f_\omega^*(x) - f_\omega^*(\lambda x) : \lambda \in (0, 1], x \in B(0, 1)\}, \\ g_3 &= \sup\left\{\|x\| f_\omega^*\left(\lambda \frac{x}{\|x\|}\right) - \|x_1\| f_\omega^*\left(\lambda \frac{x_1}{\|x_1\|}\right) - \|x_2\| f_\omega^*\left(\lambda \frac{x_2}{\|x_2\|}\right) \right. \\ &\quad \left. : \lambda \in (0, 1], x, x_1, x_2 \in \mathbb{R}^d \setminus \{0\}, \text{ with } x = x_1 + x_2\right\}, \\ g_4 &= -\inf\{f_\omega^*(x) + f_\omega^*(-x) : x \in B(0, 1)\}, \\ g_5 &= \sup\left\{\frac{|f_\omega^*(x)|}{\|x\|} : x \in B(0, 1) \setminus \{0\}\right\}, \\ g_6 &= |f_\omega^*(0)|. \end{aligned}$$

Clearly $\omega \mapsto g_i(\omega)$ are measurable maps and hence E is a measurable event in the σ -algebra \mathfrak{F} .

ITEM (i). Since \mathbb{I}_0 belongs to Θ_X , Θ_X is not empty.

ITEM (ii). The sufficiency is trivial, let us prove the necessity. Let $E^c = \Omega \setminus E = \{\omega \in \Omega : \eta \notin \theta_{X(\omega)}\}$, by hypothesis $\mathbb{P}(E) = 1$ and $\mathbb{P}(E^c) = 0$. For every $\omega \in \Omega \cap E$, there exists $Y_\omega \in \mathbb{F}$ such that $\eta \oplus Y_\omega = X(\omega)$. Let us consider the map

$$\begin{aligned} Y : \Omega &\rightarrow \mathbb{F} \\ \omega &\mapsto Y(\omega) = \begin{cases} Y_\omega, & \omega \in \Omega \cap E, \\ \mathbb{I}_0, & \omega \in E^c. \end{cases} \end{aligned} \quad (7)$$

Since $s_Y = s_X - s_\eta$ \mathbb{P} -almost surely, s_Y is measurable. Hence, the map Y defined above, is the FRV we are looking for.

Moreover, let $X \in L^2[\Omega; \mathbb{F}]$, then s_X and hence $s_Y = s_X - s_\eta$ belong to $L^2[\Omega; \mathcal{L}]$.

ITEM (iii). Consider $\eta^1, \eta^2 \in \Theta_X$. By ITEM (ii) we know that there exist two FRV Y^1, Y^2 with values in \mathbb{F} such that \mathbb{P} -a.s. $\eta^1 \oplus Y^1 = X$ and $\eta^2 \oplus Y^2 = X$. For any $\lambda \in [0, 1]$, the following hold

$$\lambda(\eta^1 \oplus Y^1) = \lambda X, \quad (1 - \lambda)(\eta^2 \oplus Y^2) = (1 - \lambda)X, \quad \mathbb{P} - a.s.$$

so that

$$[\lambda\eta^1 \oplus (1 - \lambda)\eta^2] \oplus Y = X, \quad \mathbb{P} - a.s.$$

with $Y = \lambda Y^1 \oplus (1 - \lambda)Y^2$ \mathbb{P} -a.s.. Hence $\lambda\eta^1 \oplus (1 - \lambda)\eta^2 \in \Theta_X$.

To prove the last part consider $\eta \in \Theta_X$, then $\lambda\eta = \lambda\eta \oplus (1 - \lambda)\mathbb{I}_0 \in \Theta_X$ for any $\lambda \in [0, 1]$.

ITEM (iv). Consider a sequence $\{\eta_n\}_{n=1}^\infty \subset \Theta_X$ converging to $\eta \in \mathbb{F}$ in $(\mathbb{F}, \delta_H^\infty)$, i.e.

$$\delta_H^\infty(\eta, \eta_n) \rightarrow 0, \text{ as } n \rightarrow \infty.$$

We have to prove that $\eta \in \Theta_X$.

For any $n \in \mathbb{N}$, let $E_n = \{\omega \in \Omega : \eta_n \in \theta_{X(\omega)}\}$ and Y_n a FRV as in (ii), then, for any $\omega \in \Omega \cap E_n$, $\eta_n \oplus Y_n(\omega) = X(\omega)$. Further, for any $N \in \mathbb{N}$, there exist $n, m \geq N$ such that, for any $\omega \in \Omega \cap E_n \cap E_m$,

$$\delta_H^\infty(Y_m(\omega), Y_n(\omega)) = \delta_H^\infty(\eta_m, \eta_n) < \frac{1}{N}.$$

Thus, the completeness of $(\mathbb{F}, \delta_H^\infty)$ guarantees that, for every $\omega \in \Omega \cap \bigcap_n E_n$, $\{Y_n(\omega)\}_{n \in \mathbb{N}}$ converges w.r.t. δ_H^∞ to some $Y_\omega \in \mathbb{F}$. Further, for every $\omega \in \Omega \cap \bigcap_n E_n$ and $n \in \mathbb{N}$ the following inequalities hold

$$\begin{aligned} 0 \leq \delta_H^\infty(X(\omega), \eta \oplus Y_\omega) &\leq \delta_H^\infty(X(\omega), \eta_n \oplus Y_n(\omega)) + \delta_H^\infty(\eta_n \oplus Y_n(\omega), \eta \oplus Y_\omega) \\ &\leq 0 + \delta_H^\infty(\eta_n, \eta) + \delta_H^\infty(Y_n(\omega), Y_\omega) \rightarrow 0 \end{aligned}$$

where we use the triangular inequality, the identity $X(\omega) = \eta_n \oplus Y_n(\omega)$, and the inequality (cf. [4, p.53])

$$\delta_H^\infty(\nu^1 \oplus \nu^2, \eta^1 \oplus \eta^2) \leq \delta_H^\infty(\nu^1, \eta^1) + \delta_H^\infty(\nu^2, \eta^2), \quad \nu^i, \eta^i \in \mathbb{F}, \quad i = 1, 2.$$

Then $X = \eta \oplus Y$ \mathbb{P} -a.s., and Y is the FRV defined by Equation (7). Thus we have the thesis; the limit of the convergent sequence $\{\eta_n\} \subseteq \Theta_X$ belongs to Θ_X too.

ITEM (v). Let us consider a sequence $\{\eta_n\}_{n=1}^\infty \subset \Theta_X$ converging to $\eta \in \mathbb{F}$ in (\mathbb{F}, d_2) , i.e.

$$d_2(\eta, \eta_n) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

We have to prove that $\eta \in \Theta_X$. In this case, (\mathbb{F}, d_2) is not complete and, hence, we can not repeat exactly all arguments in (iv). In particular, for any $n \in \mathbb{N}$ and for every $\omega \in \Omega \cap E_n = \{\omega \in \Omega : \eta_n \in \theta_{X(\omega)}\}$, there exist $Y_n(\omega)$ such that $\eta_n \oplus Y_n(\omega) = X(\omega)$ and, using analogous arguments of those in Proposition 3,

$$d_2(Y_m(\omega), Y_n(\omega)) = \left(\int_0^1 \int_{S^{d-1}} |s_{Y_m(\omega)}(\alpha, u) - s_{Y_n(\omega)}(\alpha, u)|^2 \, du \, d\alpha \right)^{\frac{1}{2}} = d_2(\eta_m, \eta_n) \rightarrow 0,$$

as $n \rightarrow \infty$ and where $d\alpha$ and du denote the Lebesgue measure on $[0, 1]$ and the normalized Lebesgue measure on S^{d-1} respectively. Thus, for every $\omega \in \Omega \cap \bigcap_n E_n$, $\{s_{Y_n(\omega)}\}_{n \in \mathbb{N}}$ is a Cauchy sequence in the Hilbert space $\mathcal{L} (= L^2([0, 1] \times S^{d-1}; \mathbb{R}))$ and it admits limit in \mathcal{L} , namely f_ω . Since

$$\|s_{Y_n(\omega)} - (s_{X(\omega)} - s_\eta)\|_{\mathcal{L}} = \|(s_{Y_n(\omega)} - s_{X(\omega)}) + s_\eta\|_{\mathcal{L}} = \|s_\eta - s_{\eta_n}\|_{\mathcal{L}} \rightarrow 0,$$

necessarily we have

$$s_{Y_n(\omega)} \xrightarrow{L^2} f_\omega = s_{X(\omega)} - s_\eta, \quad \forall \omega \in \Omega \cap \bigcap_n E_n.$$

Note that, f_ω is not necessarily the support function of some element in \mathbb{F} . In other words, for every $\omega \in \Omega \cap \bigcap_n E_n$, $\{Y_n(\omega)\}_{n \in \mathbb{N}}$ is a Cauchy sequence in the non-complete space (\mathbb{F}, d_2) , but under the embedding j , Equation (5), we have that the sequence $\{j(Y_n(\omega))\}_{n \in \mathbb{N}} = \{s_{Y_n(\omega)}\}_{n \in \mathbb{N}}$ is a Cauchy sequence that admits limit in the Hilbert space \mathcal{L} . But, in general, this limit is not the image under j of some element of \mathbb{F} . We claim that, for every $\omega \in \Omega \cap \bigcap_n E_n$, there exists $Y_\omega \in \mathbb{F}$ such that $s_{Y_\omega} = f_\omega = s_{X(\omega)} - s_\eta$. This allows us to deduce the thesis because, defining the FRV Y as in Equation (7), we have that $\eta \oplus Y = X$ holds \mathbb{P} -a.s..

In fact, let us consider the family $\{C_\alpha\}_{\alpha \in [0, 1]}$ of subsets of \mathbb{R}^d defined by

$$C_\alpha = \{y \in \mathbb{R}^d : \langle y, u \rangle \leq f_\omega(\alpha, u), \forall u \in S^{d-1}\}, \quad \alpha \in [0, 1].$$

In what follows, let $\omega \in \Omega \cap \bigcap_n E_n$, we prove that the family $\{C_\alpha\}_{\alpha \in [0, 1]}$ satisfies (a), (b), (c) from Lemma 4, and it defines uniquely a fuzzy set ν whose support function is, clearly, f_ω . Thus the fuzzy set ν defined in Lemma 4 is just the $Y(\omega)$ in \mathbb{F} we are looking for.

(a). Let $\alpha \in [0, 1]$.

C_α is non-empty: since $\eta_\alpha \subseteq (X(\omega))_\alpha$, then for every $u \in S^{d-1}$

$$f_\omega(\alpha, u) = s_{X(\omega)}(\alpha, u) - s_\eta(\alpha, u) \geq 0 = \langle 0, u \rangle, \quad (8)$$

i.e. $0 \in C_\alpha$.

C_α is convex: let $\lambda \in [0, 1]$ and $y_1, y_2 \in C_\alpha$, for every $u \in S^{d-1}$

$$\langle \lambda y_1 + (1 - \lambda)y_2, u \rangle \leq \lambda f_\omega(\alpha, u) + (1 - \lambda)f_\omega(\alpha, u) = f_\omega(\alpha, u)$$

i.e. $\lambda y_1 + (1 - \lambda)y_2 \in C_\alpha$.

C_α is compact: we have to prove that it is a bounded closed subset of \mathbb{R}^d . Note that $\{0\} \subseteq \eta_\alpha \subseteq (X(\omega))_\alpha$, then $s_{X(\omega)}(\alpha, u) \geq s_\eta(\alpha, u) \geq 0$ for each $u \in S^{d-1}$ and $s_{X(\omega)}(\alpha, u) \geq s_{X(\omega)}(\alpha, u) - s_\eta(\alpha, u) = f_\omega(\alpha, u)$. This implies that $\langle y, u \rangle$ is bounded for every $u \in S^{d-1}$ and hence that $C_\alpha \subseteq \mathbb{R}^d$ is bounded. On the other hand, let $\{y_n\} \subset C_\alpha$ be convergent to $y \in \mathbb{R}^d$, then, for every $n \in \mathbb{N}$ and $u \in S^{d-1}$,

$$\langle y_n, u \rangle \leq f_\omega(\alpha, u),$$

and passing to the limit we obtain the same inequality for y and for every $u \in S^{d-1}$; i.e. $y \in C_\alpha$. This fact allows us to conclude that C_α is closed and hence compact.

(b). Let $0 \leq \alpha \leq \beta \leq 1$. Note that, for every $n \in \mathbb{N}$ and $u \in S^{d-1}$, $s_{Y_n(\omega)}(\beta, u) \leq s_{Y_n(\omega)}(\alpha, u)$. Let $n \rightarrow \infty$, then $f_\omega(\beta, u) \leq f_\omega(\alpha, u)$ for every $u \in S^{d-1}$; i.e., for every $u \in S^{d-1}$ and $n \in \mathbb{N}$, $s_{Y_n(\omega)}$ and f_ω are non-increasing functions with respect to α . Now, let us consider $y \in C_\beta$, then for every $u \in S^{d-1}$, $\langle y, u \rangle \leq f_\omega(\beta, u) \leq f_\omega(\alpha, u)$; i.e. $y \in C_\alpha$ and $C_\beta \subseteq C_\alpha$.

(c). Let $\{\alpha_i\}_{i \in \mathbb{N}} \subset [0, 1]$ such that $\alpha_i \uparrow \alpha$ as i tends to infinity, that is $\alpha_i \leq \alpha_{i+1}$ and $\alpha_i \rightarrow \alpha$ as $i \rightarrow \infty$. Because of $\alpha_i \leq \alpha$ and (b), we have $C_\alpha \subseteq C_{\alpha_i}$ and $C_\alpha \subseteq \bigcap_{i \in \mathbb{N}} C_{\alpha_i}$. It remains to show the opposite inclusion. To do this let $y \in \bigcap_{i \in \mathbb{N}} C_{\alpha_i}$, i.e. $y \in C_{\alpha_i}$ for all $i \in \mathbb{N}$ or, equivalently,

$$\langle y, u \rangle \leq f_\omega(\alpha_i, u), \quad \text{for every } i \in \mathbb{N}, u \in S^{d-1}. \quad (9)$$

Note that, for every $u \in S^{d-1}$, $f_\omega(\cdot, u)$ is left-continuous with respect to α because it is the difference of two left-continuous functions (cf. Equation (8)). Hence, for the arbitrariness of i in (9), as i tends to infinity we get $\langle y, u \rangle \leq f_\omega(\alpha, u)$; i.e. $y \in C_\alpha$.

ITEM (vi). The D_θ case is a consequence of ITEM (v) and of the equivalence of d_2 and D_θ . The proof in the D_θ^φ case is exactly the same of ITEM (v) with $(\mathcal{L}, \langle \cdot, \cdot \rangle_\theta^\varphi)$ and D_θ^φ substituting $(\mathcal{L}, \langle \cdot, \cdot \rangle_\mathcal{L})$ and d_2 respectively. \blacksquare

Remark 6 Using the definition of fuzzy sets given in [5], the 0-level of fuzzy sets can be unbounded generalizing our framework. In this case, the space of fuzzy sets is not only separable but also complete since the D_θ^φ metric isometrically embeds the considered space onto a closed convex cone of a Hilbert space. Under this extended framework, properties of the Hukuhara set (Proposition 5) may be easily obtained. On the other hand, the Steiner point is not always defined and, hence, a characterization point (with similar properties: additivity and continuity) for this extended framework should be found for completeness. \square

4 Hukuhara decomposition

Now, we are ready to prove the decomposition (2) presented in the Introduction that is the main result of this paper. Roughly speaking, (2) allows to decompose uniquely (thanks to a minimization argument) any $X \in L^2[\Omega; \mathbb{F}]$ as the sum of a deterministic convex fuzzy set $H_X^\perp \in \Theta_X$, a FRV Y (that carries the whole “fuzzy randomness” of X) and a random translation in \mathbb{R}^d (that provides the “spatial” location of X).

We present the main result in Corollary 9 as a consequence of the centered case proved in Theorem 7.

Theorem 7 *Let $X \in L^2[\Omega; \mathbb{F}]$ with $\mathbf{Ste}(X) = 0$. Thus there exists $H_X^\perp \in \mathbb{F}$ with $\mathbf{Ste}(H_X^\perp) = 0$ and $Y \in L^2[\Omega; \mathbb{F}]$ with $\mathbf{Ste}(Y) = 0$ \mathbb{P} -a.s. such that X decomposes according to*

$$X(\omega) = H_X^\perp \oplus Y(\omega), \quad (10)$$

for \mathbb{P} -almost all $\omega \in \Omega$. In particular, H_X^\perp is the unique element in \mathbb{F} that satisfies (10) and minimizes $\Delta_2(X, \eta)$ (with η varying in Θ_X); i.e., there exists a unique $H_X^\perp \in \Theta_X$ such that

$$H_X^\perp := \arg \min_{\eta \in \Theta_X} \Delta_2(X, \eta) = \arg \min_{\eta \in \Theta_X} \mathbb{E}[(d_2(X, \eta))^2].$$

Hence Y is the unique (except on a \mathbb{P} -negligible set) FRV such that its support function is given by $s_Y = s_X - s_{H_X^\perp}$. Moreover, H_X^\perp is a maximal element in Θ_X with respect to the level-wise set inclusion; that is, if $C \in \Theta_X$ with $(H_X^\perp)_\alpha \subseteq C_\alpha$ for any $\alpha \in [0, 1]$, then $H_X^\perp = C$.

Corollary 8 *Let $X \in L^2[\Omega; \mathbb{F}]$, $D \in \mathbb{F}$ with $\mathbf{Ste}(X) = \mathbf{Ste}(D) = 0$ and $X' = X \oplus D$. Then $H_{X'}^\perp = H_X^\perp \oplus D$.*

Corollary 9 *Let $X \in L^2[\Omega; \mathbb{F}]$. Thus there exists $H_X^\perp \in \mathbb{F}$ with $\mathbf{Ste}(H_X^\perp) = 0$ and $Y \in L^2[\Omega; \mathbb{F}]$ with $\mathbf{Ste}(Y) = 0$ \mathbb{P} -a.s. such that X decomposes uniquely according to Equation (2)*

$$X(\omega) = H_X^\perp \oplus Y(\omega) \oplus \mathbb{I}_{\mathbf{Ste}(X(\omega))}, \quad (2)$$

for \mathbb{P} -almost all $\omega \in \Omega$. In particular, it holds

$$H_X^\perp = H_{\tilde{X}}^\perp, \quad Y = \tilde{Y},$$

with $H_{\tilde{X}}^\perp$ and \tilde{Y} given by Theorem 7 applied to the FRV $\tilde{X} = X \oplus \mathbb{I}_{-\text{Ste}(X)}$.
 Moreover, if $D \in \mathbb{F}$ and $X' = X \oplus D$ then

$$X' = H_{X'}^\perp \oplus Y' \oplus \mathbb{I}_{\text{Ste}(X')} = (H_X^\perp \oplus D \oplus \mathbb{I}_{-\text{Ste}(D)}) \oplus \tilde{Y} \oplus \mathbb{I}_{\text{Ste}(X) + \text{Ste}(D)}, \quad \mathbb{P} - \text{a.s.}$$

Proof of Theorem 7. Since Θ_X collects all the elements of \mathbb{F} for which (10) holds, we have to prove that there exists a unique element in Θ_X that minimizes the map $\eta \in \Theta_X \rightarrow \mathbb{E}[(d_2(X, \eta))^2]$.

At first note that Θ_X can be seen as a subset of $L^2[\Omega; \mathbb{F}]$; in fact, for each $\eta \in \Theta_X$ the constant map $\omega \mapsto \eta$ is an element of $L^2[\Omega; \mathbb{F}]$ since

$$\mathbb{E}[(\sup_{x \in \eta_0} \|x\|)^2] = (\sup_{x \in \eta_0} \|x\|)^2 < +\infty.$$

Moreover, Θ_X is closed in $(L^2[\Omega; \mathbb{F}], \Delta_2)$ as a consequence of

$$\Delta_2(\nu, \eta) = \{\mathbb{E}[(d_2(\nu, \eta))^2]\}^{1/2} = d_2(\nu, \eta),$$

for any couples $\nu, \eta \in \mathbb{F}$, and thanks to the fact that Θ_X is closed in (\mathbb{F}, d_2) , see Proposition 5.

Thus the minimization problem is equivalent to prove that there exists a unique projection of X onto Θ_X that is a closed convex subset of $L^2[\Omega; \mathbb{F}]$ endowed with the metric Δ_2 . Since $(L^2[\Omega; \mathbb{F}], \Delta_2)$ embeds isometrically in the Hilbert space $(L^2[\Omega; \mathcal{L}], \mathbb{E}[\langle \cdot, \cdot \rangle_{\mathcal{L}}])$ through map J (see Section 2), there exists a unique element $H_{\tilde{X}}^\perp \in \Theta_X$ that realizes the required minimum

$$H_{\tilde{X}}^\perp = \underset{\eta \in \Theta_X}{\text{arg min}} \mathbb{E}[(d_2(X, \eta))^2].$$

As a consequence of $H_{\tilde{X}}^\perp \in \Theta_X$ and of (ii) in Proposition 5, the FRV Y is defined through its support function $s_Y = s_X - s_{H_{\tilde{X}}^\perp}$.

Finally, let C be as in the thesis; thus inclusions $(H_{\tilde{X}}^\perp)_\alpha \subseteq C_\alpha \subseteq X_\alpha$ imply $s_X - s_C \leq s_X - s_{H_{\tilde{X}}^\perp}$. Then, by definitions of $H_{\tilde{X}}^\perp$ and d_2 , necessarily $C = H_{\tilde{X}}^\perp$ holds. \blacksquare

The chosen notation wants to recall the line of the proof; $H_{\tilde{X}}^\perp$ is obtained as the (necessarily unique) projection of the FRV X on its Hukuhara set Θ_X . Further, we want to stress out that the suffix X does not mean that $H_{\tilde{X}}^\perp$ is random; in fact, it does not depend on ω but rather it is a deterministic element of \mathbb{F} (that is a constant element in $L^2[\Omega; \mathbb{F}]$) that depends on the whole map $\omega \mapsto X(\omega)$. A qualitative graphical interpretation of Theorem 7 is given in Figure 1.

About different metrics and open problems: Although in this paper different metric for fuzzy sets are used, the decomposition in Theorem 7 is just proved for the d_2 metric. Thus, the following question arises spontaneously: *what can we say if d_2 is replaced with another metric?* Note that, two facts are mainly involved in the proof of Theorem 7:

1. Θ_X is a non-empty, closed convex subset of $L^2[\mathbb{F}]$ with respect to d_2 .
2. $(L^2[\mathbb{F}], d_2)$ embeds isometrically in a Hilbert space.

These facts allow to project X onto Θ_X in a unique way. As a consequence, arguments in Theorem 7 still hold for every metric that satisfies the preceding properties. This is the case of D_θ but not of δ_H^∞ (because a Hilbert space is not available). Anyway, changing metric several questions may arise.

Let us consider at first the case of D_θ . Although d_2 and D_θ induce the same topology on \mathbb{F} , the projection $\widetilde{H_{\tilde{X}}^\perp}$ obtained w.r.t. D_θ , in general, differs from $H_{\tilde{X}}^\perp$. It is a conjecture of the authors that $\widetilde{H_{\tilde{X}}^\perp} = H_{\tilde{X}}^\perp$.

In the case of δ_H^∞ , $L^2[\mathbb{F}]$ can not be embedded in a Hilbert space and, hence, a projection theorem can not be used. Anyway, the map $\delta_H^\infty(X, \cdot)$ is continuous and convex and, hence, admits at least a minimum on Θ_X . Nevertheless, the minimum is not necessarily achieved in a unique way and, even if it was unique, it should be compared with $H_{\tilde{X}}^\perp$ and $\widetilde{H_{\tilde{X}}^\perp}$. Author's conjecture is that both $H_{\tilde{X}}^\perp$ and $\widetilde{H_{\tilde{X}}^\perp}$ minimize $\mathbb{E}[\delta_H^\infty(X, \cdot)^2]$ on Θ_X .




Figure 1: A qualitative graphical interpretation of some results of Section 3 and Section 4. In particular, Θ_X is represented as a closed convex subset of \mathbb{F} containing the origin and such that, for any $\eta \in \Theta_X$ and $\lambda \in [0, 1]$, $\lambda\eta \in \Theta_X$. Hence, H_X^\perp is the projection of X on Θ_X , as a subset of $L^2[\Omega; \mathbb{F}]$, with respect to the metric $\mathbb{E}[d_2(\cdot, \cdot)^2]$, this also guarantees the uniqueness of H_X^\perp since the cone $L^2[\Omega; \mathbb{F}]$ is embeddable in the Hilbert space $L^2[\Omega; \mathcal{L}]$ through the isometry $X \mapsto j(X)$. Finally the following inclusions or embeddings are qualitatively represented: $\Theta_X \subseteq \mathbb{F} \hookrightarrow L^2[\Omega; \mathbb{F}] \hookrightarrow L^2[\Omega; \mathcal{L}]$.

Fuzzy random translations. Theorem 7 allows us to characterize fuzzy random translation by means of the Aumann expectation.

Definition 10 A FRV X is a *translation* if there exists $\nu \in \mathbb{F}$ with $\mathbf{Ste}(\nu) = 0$ such that

$$X(\omega) = \nu \oplus \mathbb{I}_{\mathbf{Ste}(X)}.$$

Clearly, every Gaussian FRV X is a FRV translation with $\nu \oplus \mathbb{I}_{\mathbb{E}[\mathbf{Ste}(X)]} = \mathbb{E}X$. Another sufficient condition for X to be a FRV translation is given by Proposition 11, while a necessary and sufficient condition is stated in Proposition 12.

Proposition 11 *Let X be a FRV such that $\mathbb{E}X = \mathbb{I}_c$ where $c \in \mathbb{R}^d$. Then $X = \mathbb{I}_\xi$ \mathbb{P} -a.s. for some random element ξ in \mathbb{R}^d . (Clearly X is a FRV translation.)*

Proof. Thesis may be obtained as in [2, Theorem 6], or, whenever $X \in L^2[\Omega; \mathbb{F}]$, as corollary of the Theorem 7 and Proposition 12. \blacksquare

Let X be a FRV translation, and $\tilde{X} = X \oplus \mathbb{I}_{-\mathbf{Ste}(X)}$. Then it easy to check that

$$X = H_X^\perp \oplus \mathbb{I}_{\mathbf{Ste}(X)}, \quad \mathbb{P} - \text{a.s.} \quad (11)$$

Proposition 12 *Let $X \in L^2[\Omega; \mathbb{F}]$. X is a FRV translation if and only if H_X^\perp satisfies*

$$\mathbb{E}X = H_X^\perp \oplus \mathbb{I}_{\mathbb{E}[\mathbf{Ste}(X)]} \quad (12)$$

with $\mathbb{E}X$ being the Aumann expectation; in other words, H_X^\perp is equal to $\mathbb{E}X$ up to a translation.

Proof. For the “only if” part, in order to obtain Equation (12), it is sufficient to compute the expectation in Equation (11).

“If” part. For the sake of simplicity, let us assume that $\mathbf{Ste}(X) = 0$, a straightforward argument extends the result in the more general case of a FRV with non-null $\mathbf{Ste}(X)$. Then, in term of support functions, Equation (10) becomes

$$s_X = s_{H_X^\perp} + s_Y = s_{\mathbb{E}X} + s_Y, \quad \mathbb{P} - \text{a.s.}$$

where we use the fact that $H_X^\perp = \mathbb{E}X$. Computing expectation of both sides and using (6), we get $s_{\mathbb{E}Y} = 0$. Hence $Y = \mathbb{I}_\xi$ a.s. for some random element ξ in \mathbb{R}^d (cf. [2]). \blacksquare

Remark 13 Whenever $X \in L^2[\Omega; \mathbb{F}]$, in view of Theorem 7 and Proposition 12, we get a proof of Proposition 11. In fact, suppose that $\mathbb{E}X = \mathbb{I}_c$ for some $c \in \mathbb{R}^d$, and compute expectation of both sides in Equation (10)

$$\mathbb{I}_c = \mathbb{E}X = H_X^\perp \oplus \mathbb{E}Y.$$

Hence, for any $\alpha \in [0, 1]$, $(H_X^\perp)_\alpha$ is a subset of $\{c\}$ up to a translation, that is $(H_X^\perp)_\alpha$ is a singleton as well as $(\mathbb{E}Y)_\alpha$. Then $H_X^\perp = \mathbb{I}_{c'}$ for some $c' \in \mathbb{R}^d$, i.e. H_X^\perp equals $\mathbb{E}X$ up to a translation and, by Proposition 12, X is a FRV translation that implies $Y = \mathbb{I}_\xi$ for some random element in \mathbb{R}^d . Finally, Equation (10) becomes

$$X = H_X^\perp \oplus Y = \mathbb{I}_{c'} \oplus \mathbb{I}_\xi = \mathbb{I}_{\xi'},$$

that is the thesis of Proposition 11. \square

A decomposition example. Remark 14 provides an example of an X in $L^2[\Omega; \mathbb{F}]$ with $\mathbf{Ste}(X) = 0$ for which $\mathbb{E}[X] \neq H_X^\perp$ and for which H_X^\perp is not necessarily \mathbb{I}_0 ; i.e., in terms of Proposition 12, X is not a translation but its deterministic part H_X^\perp in the decomposition (10) is not just reduced to the origin.

Remark 14 Let $\mathbb{R}^d = \mathbb{R}$, $(\Omega = [0, 1], \mathcal{B}_{[0,1]}, \mathbb{P})$ where $\mathcal{B}_{[0,1]}$ denotes the Borel σ -algebra on $[0, 1]$ w.r.t. the euclidean metric and $\mathbb{P} = \mu$ is the Lebesgue measure. Let X be the FRV defined by $X := \mathbb{I}_{[-\omega, \omega]}$, for any $\omega \in [0, 1]$ (actually X is a random sets, but similar arguments can be repeated either for the triangular FRV $Y = (-\omega, 0, \omega)$ or for the trapezoidal FRV $X \oplus Y = (-2\omega, -\omega, \omega, 2\omega)$). Clearly $X \in L^2[\Omega; \mathbb{F}]$ and $\mathbf{Ste}(X) = 0$. Moreover,

$$f_m(\omega) := \min X_1(\omega) = -\omega \quad \text{and} \quad f_M(\omega) := \max X_1(\omega) = \omega$$

are integrable selections of the 1-level X_1 . Obviously, any other integrable selection f of X_1 satisfies

$$f_m(\omega) \leq f(\omega) \leq f_M(\omega), \quad \text{for each } \omega \in [0, 1].$$

Then

$$-\frac{1}{2} = \mathbb{E}f_m \leq \mathbb{E}f \leq \mathbb{E}f_M = \frac{1}{2},$$

and, by the convexity of Aumann expectation and because $X_1 = X_\alpha$ for any $\alpha \in [0, 1]$, $\mathbb{E}X_1 = [-\frac{1}{2}, \frac{1}{2}] = \mathbb{E}X_\alpha$, that is $\mathbb{E}X = \mathbb{I}_{[-\frac{1}{2}, \frac{1}{2}]}$.

We prove that $\mathbb{E}X \notin \Theta_X$ and hence, by Proposition 12, X is not a FRV translation. In fact, note that

$$X \ominus_H \mathbb{E}X = \mathbb{I}_{[-\omega, \omega]} \ominus_H \mathbb{I}_{[-\frac{1}{2}, \frac{1}{2}]} = \begin{cases} \mathbb{I}_{[-\omega+\frac{1}{2}, \omega-\frac{1}{2}]}, & \omega > \frac{1}{2}, \\ \mathbb{I}_0, & \omega = \frac{1}{2}, \\ \text{it does not exist,} & \omega < \frac{1}{2}, \end{cases}$$

implies

$$\mathbb{P}(\mathbb{E}X \in \theta_X) = \mathbb{P}(\text{there exists } X \ominus_H \mathbb{E}X) = \mathbb{P}\left(\omega \geq \frac{1}{2}\right) = \frac{1}{2},$$

and hence $\mathbb{E}X \notin \Theta_X$.

Actually we can show that $\Theta_X = \{\mathbb{I}_0\}$ and hence $H_X^\perp = \mathbb{I}_0$. In fact, by absurd let $\eta \in \Theta_X$ with $\eta \neq \mathbb{I}_0$, then there exists $\alpha \in [0, 1]$ such that $\eta_\alpha = [a, b]$ with $a < b$ and there exists $X_\alpha \ominus_H \eta_\alpha$, here \ominus_H is considered as the Hukuhara difference for subsets in \mathbb{R} . On the other hand

$$[-\omega, \omega] \ominus_H [a, b] = \begin{cases} [-\omega - a, \omega - b], & \omega - b > -\omega - a, \\ \{-\frac{b+a}{2}\}, & \omega = \frac{b-a}{2}, \\ \text{it does not exist,} & \omega < \frac{b-a}{2}, \end{cases}$$

and, as consequence,

$$\mathbb{P}([- \omega, \omega] \ominus_H [a, b] \text{ does not exist}) = \mu\left[\left(-\infty, \frac{b-a}{2}\right) \cap [0, 1]\right] = \frac{b-a}{2} > 0$$

(recall that, by hypothesis, $b > a$). This is an absurd since $\eta \in \Theta_X$ by hypothesis. Thus $\Theta_X = \{\mathbb{I}_0\} \neq \mathbb{E}X = \mathbb{I}_{[-\frac{1}{2}, \frac{1}{2}]}$.

Finally, in order to introduce a more general example, let us consider

$$X = \mathbb{I}_{[-\omega, \omega]} \oplus \mathbb{I}_{[-\frac{1}{2}, \frac{1}{2}]} = \mathbb{I}_{[-\omega - \frac{1}{2}, \omega + \frac{1}{2}]}$$

so that, from Corollary 8, we immediately obtain that

$$\mathbb{I}_{[-1, 1]} = \mathbb{E}X \neq H_X^\perp = \mathbb{I}_{[-\frac{1}{2}, \frac{1}{2}]}.$$

Note that, this is a case in which H_X^\perp is different from \mathbb{I}_0 . □

5 Conclusion

In this paper, we have proven that any square integrable FRV can be decomposed as $X = H_X^\perp \oplus Y \oplus \mathbb{I}_{\mathbf{Ste}(X)}$, where H_X^\perp is a unique deterministic fuzzy convex compact set (i.e. in \mathbb{F}) and Y is an element of $L^2[\Omega; \mathbb{F}]$ with $\mathbf{Ste}(Y) = 0$. The method used to obtain H_X^\perp is a particular case where the problem posed in [7, p.174–175] is well-solved by defining the Hukuhara set Θ_X . In this view, H_X^\perp may be interpreted as an expectation for X that satisfies some of the properties, listed in [7, p.190] for random closed sets but trivially extendible in the fuzzy case, of a “reasonable” expectation of X .

The decomposition theorem proposed in Section 4 could not be compared with the fuzzy regression problem stated in [13]. In fact, in that paper, the authors look for the best linear approximation function of a given square integrable FRV Y by another square integrable FRV X , studying the minimization problem

$$\inf_{a \in \mathbb{R}, \eta \in \mathbb{F}} \mathbb{E}[d_2(Y, aX \oplus \eta)^2].$$

Besides the open problems exposed in Section 4 (concerning the use of different metrics), future works may consider the possibility to relax some hypothesis. As example, replacing \mathbb{R}^d with an Hilbert or a Banach space (problems may arise considering the embedding j and hence the closure of the Hukuhara set Θ_X), or dropping convexity hypothesis and hence stating a decomposition theorem for a fuzzy random element whose level sets are not necessarily convex. Finally, note that we restricted our studies to the existence of a such H_X^\perp . However, it is certainly interesting to establish whenever H_X^\perp could be explicitly computed or approximated, though even in particular cases.

Acknowledgements

The authors would like to thank Prof. V. Capasso for the helpful discussions during the preparation of this paper. The authors are also very grateful to the anonymous referees for the valuable comments that helped to improve the paper, especially for suggesting the use of the D_θ^φ metric and for having pointed out the possibility of an extension to the unbounded support case.

References

- [1] V. N. Bobylev. Support function of a fuzzy set and its characteristic properties. *Mathematical Notes*, 37, 281–285, 1985.
- [2] E. G. Bongiorno. A Note on Fuzzy Set-Valued Brownian Motion. *Statistics and Probability Letters*, 82, 827-832, 2012.
- [3] A. Colubi, J. S. Dominguez-Menchero, M. Lopez-Diaz, D. A. Ralescu. A $D_E[0, 1]$ representation of random upper semicontinuous functions. *Proceedings of the American Mathematical Society*, 130, 3237–3242, 2002.
- [4] P. Diamond, P. Kloeden. *Metric spaces of fuzzy sets: theory and applications*. Wiley, 1994.
- [5] G. González-Rodríguez, A. Colubi, M. A. Gil. Fuzzy data treated as functional data: A one-way ANOVA test approach. *Computational Statistics and Data Analysis*, 56, 943–955, 2012.

- [6] S. Li, Y. Ogura, V. Kreinovich. *Limit Theorems and Applications of Set-Valued and Fuzzy Set-Valued Random Variables*. Kluwer Academic Publishers Group, Dordrecht, 2002.
- [7] I. Molchanov. *Theory of random sets*. Springer. (2005)
- [8] W. Näther. Linear Statistical Inference for Random Fuzzy Data. *Statistics*, 29:3, 221–240, 1997.
- [9] W. Näther. On random fuzzy variables of second order and their application to linear statistical inference with fuzzy data. *Metrika*, 51, 201–221, 2000.
- [10] A. B. Ramos-Guajardo, A. Colubi, G. González-Rodríguez, M. A. Gil. One-sample tests for a generalized Fréchet variance of a fuzzy random variable. *Metrika*, 71, 185–202, 2010.
- [11] W. Trutschnig, G. González-Rodríguez, A. Colubi, M. A. Gil. A new family of metrics for compact, convex (fuzzy) sets based on a generalized concept of mid and spread. *Information Sciences*, 179, 3964–3972, 2009.
- [12] T. Vetterlein, M. Navara. Defuzzification using Steiner points. *Fuzzy Sets and Systems*, 157, 1455–1462, 2006.
- [13] A. Wünsche, W. Näther. Least-squares fuzzy regression with fuzzy random variables. *Fuzzy Sets and Systems*, 130, 43–50, 2002.