



A novel CD46 mutation in a patient with microangiopathy clinically resembling thrombotic thrombocytopenic purpura and normal ADAMTS13 activity

by Raffaella Rossio, Luca Andrea Lotta, Silvia Pontiggia, Nicolo' Borsa Ghiringhelli, Isabella Garagiola, Gianluigi Ardissino, Danijela Mikovic, Massimo Cugno, and Flora Peyvandi

Haematologica 2014 [Epub ahead of print]

Citation: Rossio R, Lotta LA, Pontiggia S, Borsa Ghiringhelli N, Garagiola I, Ardissino G, Mikovic D, Cugno M, and Peyvandi F. A novel CD46 mutation in a patient with microangiopathy clinically resembling thrombotic thrombocytopenic purpura and normal ADAMTS13 activity. Haematologica. 2014; 99:xxx doi:10.3324/haematol.2014.111062

Publisher's Disclaimer.0

E-publishing ahead of print is increasingly important for the rapid dissemination of science. Haematologica is, therefore, E-publishing PDF files of an early version of manuscripts that have completed a regular peer review and have been accepted for publication. E-publishing of this PDF file has been approved by the authors. After having E-published Ahead of Print, manuscripts will then undergo technical and English editing, typesetting, proof correction and be presented for the authors' final approval; the final version of the manuscript will then appear in print on a regular issue of the journal. All legal disclaimers that apply to the journal also pertain to this production process.

A novel *CD46* mutation in a patient with microangiopathy clinically resembling thrombotic thrombocytopenic purpura and normal ADAMTS13 activity

Raffaella Rossio*¹, Luca Andrea Lotta*¹, Silvia Pontiggia¹, Nicolò Borsa Ghiringhelli², Isabella Garagiola¹, Gianluigi Ardissino³, Danijela Mikovic⁴, Massimo Cugno⁵ and Flora Peyvandi¹

1Angelo Bianchi Bonomi Hemophilia and Thrombosis Center, Fondazione IRCCS Ca' Granda - Ospedale Maggiore Policlinico, Department of Pathophysiology and Transplantation, University of Milan and Luigi Villa Foundation, Milan, Italy

2Laboratory of Medical Genetics, Center for HUS Control Fondazione IRCCS Ca' Granda - Ospedale Maggiore Policlinico, Milan, Italy

3Center for HUS Control, Fondazione IRCCS Ca' Granda - Ospedale Maggiore Policlinico, Milan, Italy

4Blood Transfusion Institute of Serbia, Svetog Save 39, Belgrade, Serbia

5Internal Medicine, Fondazione IRCCS Ca' Granda - Ospedale Maggiore Policlinico, Department of Pathophysiology and Transplantation, University of Milan, Milan, Italy

*RR and LAL contributed equally to this manuscript

Running Title: Complement abnormalities in TTP.

Correspondence

Flora Peyvandi, MD, PhD, Angelo Bianchi Bonomi Hemophilia and Thrombosis Center, Fondazione IRCCS Ca' Granda - Ospedale Maggiore Policlinico, Department of Pathophysiology and Transplantation, Università degli Studi di Milano and Luigi Villa Foundation. Italy.
E-Mail: flora.peyvandi@unimi.it

Acknowledgments

LAL is recipient of the 2012 Bayer Hemophilia Award – Early Career Investigator Award and of the Raffaello De Biasi Prize (Associazione Italiana Centri Emofilia, AICE). FP is recipient of the 2011 Bayer Hemophilia Award – Special Project Award. The study was supported by Fondazione Cariplo (Grant n. 2011-0524), Italo Monzino Foundation, Italian Ministry of Health (Ricerca finalizzata RF-2009-1530493), Regione Lombardia – Direzione Generale di Sanità (Decreto n. 13848, 11/12/2009) and by Fondazione Fiera Milano. The authors would like to thank R. Donadelli and R. Piras (Unit of Genetics and Molecular Basis of Renal Diseases, Mario Negri Institute for Pharmacological Research, Villa Camozzi, Via Camozzi 3, Ranica Bergamo, Italy) for their help with MCP expression measurements.

The differential diagnosis and etiologic classification of the two main forms of thrombotic microangiopathy (TMA) —thrombotic thrombocytopenic purpura (TTP) and atypical haemolytic uraemic syndrome (aHUS)— remain challenging¹. The terms TTP and HUS are used to describe the clinical presentation of these diseases. HUS has prominent renal involvement, whereas neurological manifestations are common in TTP. The distinction is not always reliable; neurologic complications can be present in patients with aHUS and renal failure not requiring dialysis can be present in patients affected by TTP.

aHUS is characterised by hyperactivation of the alternative complement pathway^{2,3}, whilst TTP is characterised by the severe deficiency of the von Willebrand factor (VWF) cleaving protease ADAMTS13⁴. The prevalence of severe ADAMTS13 deficiency (i.e. activity below 10%) in TTP is high in patients with idiopathic disease and none or minimal renal involvement. However in a proportion of patients with idiopathic TTP and minor renal involvement at acute disease presentation, ADAMTS13 activity may be only slightly reduced or even normal. The pathophysiological mechanisms in these patients are unknown. In order to clarify the etiology in a patient (27 year old female) with acute TTP and normal ADAMTS13 activity, we investigated the complement system-, ADAMTS13-, and VWF-related biochemical properties^{5,6} at acute phase and during remission. Concentration of the complement system components C3, C4 and factor H were measured by radial immunodiffusion using commercial methods. C3 and C4 levels were measured by C3 and C4 NOR Partigen (Siemens, Marburg, Germany). Complement system activity was evaluated by ELISA (Wieslab complement assay, EuroDiagnostica, Malmö, Sweden). The wells of microtitre strips were coated with specific activators of the classical, alternative or mannose-binding lectin pathway, and the serum was diluted in a buffer containing specific blockers in order to ensure that only one pathway was activated during incubation. The wells were then washed and membrane attack complex (C5b-9) was detected. Anti-factor H antibodies were assayed by an enzyme-linked immunosorbent assay (ELISA) that used purified factor H for capture and anti-human immunoglobulin G (IgG), A (IgA) and M (IgM) for detection⁷.

We performed exome sequencing to identify potentially causal mutations. Three first-degree relatives of the patient, including the patient's mother and her two sons, were also studied. The study was approved by the institutional review board of the Fondazione IRCCS Ca' Granda – Ospedale Maggiore Policlinico. All participants signed informed consent.

Our patient presented to her local hospital with fever, respiratory tract infection, abdominal pain, and headache. She had no past medical history and didn't take any drugs. Laboratory investigations revealed thrombocytopenia (platelet counts: $14 \times 10^9/L$; normal values [nv]: $150-450 \times 10^9/L$) and microangiopathic hemolytic anemia (hemoglobin: 8.9 g/dL; nv: 11.5-16.0 g/dL) with negative Coombs test and evidence of schistocytes in peripheral blood smear and reduced haptoglobin levels (34 mg/dl n.v. 45-164). Total and indirect bilirubin (43.6 and 38.2 $\mu\text{mol/l}$; n.v.< 20 $\mu\text{mol/l}$ and n.v.< 7 $\mu\text{mol/l}$) and lactate dehydrogenase (4559 U/l; n.v.: 160-320 U/l) were increased. Serum creatinine was slightly increased (117 $\mu\text{mol/L}$; nv: 53-106 $\mu\text{mol/L}$), but the patient did not have renal failure according to the RIFLE (Risk, Injury, Failure, Loss, ESRD) criteria. Creatinine levels returned to normal levels (89 $\mu\text{mol/L}$) within 48 hours. A diagnosis of TTP was made and plasma exchange (PEX) commenced. Remission was achieved after 7 plasma exchange procedures. The patient initiated regular follow-up and remained relapse-free seven years after the episode. She had two pregnancies without complications. Consumption of VWF multimers of large and intermediate size was observed. The finding of reduced ULVWF ratio in acute phase could suggest a TTP-like TMA, despite normal ADAMTS13 levels, with the proposed mechanism: the excess of ULVWF (possibly due to endothelial cells activation and ULVWF release) overwhelmed ADAMTS13 cleaving activity, with VWF-mediated platelet aggregation and ULVWF consumption in the thrombi. Recently Cataland et al reported that patients with aHUS did not showed mid- to large- sized VWF multimer in acute phase ⁸.

Low C3 and C4 levels as well as low activity of the classical and alternative complement pathways may reflect a consumption of complement factors due to activation. This explanation is supported by the observation that during remission complement levels normalized. A selective activation of

alternative pathway is typical of aHUS, but a degree of classical pathway activation is described⁹. Not only complement parameters, but also all the other measurements were normal during remission (Table 1).

We sequenced the entire protein coding area of the genome by Illumina HiSeq 2000. In the exome of the patient, we identified 25068 SNVs and 209 open reading frame indels. A total of 12 SNVs in 6 genes were found. We focused on complement factor and complement regulation genes (*CFH*, *CFI*, *CD46*, *CFB*, *THBD* and *C3* genes). Sequencing identified 9 single nucleotide variants (SNVs; 6 missense or splice-site SNVs and 3 additional coding synonymous SNVs reported in Supplemental Table 2). We did not detect short insertions or deletions (indels) in these genes. Neither *C3*, *CFB*, *CFHR3*, *CFHR4*, nor *THBD* had functional variants. Of all variants found in the protein-coding area of thrombotic-microangiopathy-associated genes, one was novel: i.e. not present in dbSNP139 (URL: <http://www.ncbi.nlm.nih.gov/SNP/http://www.1000genomes.org/faq/how-do-i-cite-1000-genomes-project>) or in 1000 Genomes (URL <ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/>).

This novel variant was a heterozygous splice-site SNV of *CD46* (c.475+1G>A), encoding the complement protein MCP. Notably, we did not find the variant in 90 Italian controls by PCR and Sanger sequencing. Association of *CD46* splice-site variants with aHUS have been reported previously^{10,11}. The *CD46* variant is predicted by NetGene2 and Spliceport to abolish the natural donor splice site of exon 4, creating an alternative splice-site into the exon 4. The activation of the new splice-site is predicted to result in the deletion of 21 nucleotides (c.631_652delTAAGCCCCCAATATGTGAAA), determining the lack of 6 aminoacids (p.G152_C157del). These analyses suggest that the variant may be causally implicated in the TMA of our patient. In order to support this hypothesis, we screened for the variant in the family members of the patient, and also made measurements related to the complement activation. Genetic studies on first degree relatives have potential psychologic and financial implications for them, but the identification of carriers of mutations may allow adequate follow-up of at-risk subjects,

particularly during triggering events such as infections and pregnancy. In fact it is known that aHUS penetrance is incomplete.

To this end, we studied the circulating CD46 mRNA by reverse transcription PCR (RT-PCR) and found that both sons of the patient had the c.475+1G>A variant (in heterozygous form), whereas her mother did not. Meanwhile, complement system measurements showed activation of the alternative complement pathway in the younger of the two sons (Son 2) and no alteration of complement activation in the older son (Son 1), nor in the mother of the patient. Real time PCR and Sanger sequencing of the CD46 mRNA spanning exons 3-6 confirmed the presence of the altered CD46 transcript bearing the deletion of 21 nucleotides in the patient and in both sons. Expression of MCP was analysed on granulocytes from the patient and a control subject using a FACSAria cytometer (BD Biosciences) and showed a 50% reduction of this protein.

The differential diagnosis of TMAs is challenging. TTP is accompanied by severe deficiency of ADAMTS13 activity, whilst aHUS is characterized by complement system abnormalities due to genetic mutations. Severe ADAMTS13 deficiency has been described in a proportion of patients with a clinical diagnosis of aHUS previously,¹² suggesting that these disorders could overlap not only for their clinical characteristics, but also for their pathophysiological mechanisms.

In this patient with an isolated episode of TMA and minimal renal impairment, a detailed phenotypic characterization followed by exome-wide genetic analysis identified a novel *CD46* mutation as a likely cause for the TMA. Interestingly, the patient did not show signs of prominent renal involvement and was diagnosed with TTP. The novel finding of our report is that a patient with TMA clinically resembling TTP and minor renal involvement may carry a rare, disruptive genetic variant of complement regulator gene, such as *CD46*.

This case of TMA at presentation did not meet a clear-cut criteria for diagnosis of either TTP or aHUS, because the clinical presentation is TTP-like. However, non-deficient ADAMTS13 activity at presentation and in remission, evidence of complement activation at presentation and the finding of a mutation of *CD46* collectively support the hypothesis that the most appropriate diagnosis

would be that of a complement-mediated TMA/aHUS rather than a diagnosis of TTP. This case report emphasizes the need for a new classification of different types of TMA based not only on clinical presentation, but also on disease mechanism and molecular evaluation. A new classification of TMA is important because the therapeutic approach to TTP and aHUS is evolving and more etiology-based. New tailored therapies are being introduced and used in combination with PEX. These include anti-CD20 antibodies¹³, inhibitors of VWF-platelet interactions¹⁴ and “terminal complement” pathway inhibitors¹⁵ like eculizumab. As the therapeutic approach to different forms of TMA diverges, it is becoming more important to establish the underlying etiology. This paper confirms the utility of the analysis of both the ADAMTS13 activity and complement evaluation in patients with acute TMA.

Authorship and Disclosures

RR, LAL, and FP designed the study; DM recruited the patient and family members; RR, LAL, SP, NGB, IG, and GA extracted or generated clinical or experimental data; LAL, RR, and MC analyzed the data; RR, LAL, MC, and FP interpreted the results; RR and LAL wrote the manuscript; all authors read and approved the final version of the manuscript.

GA is a member of SAB of the aHUS Global Registry supported by Alexion. FP has received honoraria for participating as a speaker at satellite symposia and educational meetings organized by Novo Nordisk, CSL Behring, LFB, Grifols, Bayer and Baxter, and she has received research grant funding from Novo Nordisk, Biotest and Kedrion Biopharma. All other authors report no conflicts of interest.

References

- 1-Mannucci PM. Thrombotic thrombocytopenic purpura and the hemolytic uremic syndrome: much progress and many remaining issues. *Haematologica*. 2007;92(7):878-80.
- 2-Stuhlinger W, Kourilsky O, Kanfer A, Sraer JD. Haemolytic-uraemic syndrome: evidence for intravascular C3 activation. *Lancet* 1974;2(7883):788-9.
- 3-Noris M, Remuzzi G. Atypical hemolytic-uremic syndrome. *N Engl J Med*. 2009;361(17):1676-87.
- 4-George JN. Clinical practice. Thrombotic thrombocytopenic purpura. *N Engl J Med*. 2006;354(18):1927-35.
- 5- Tsai HM, Lian EC. Antibodies to von Willebrand factor-cleaving protease in acute thrombotic thrombocytopenic purpura. *N Engl J Med*. 1998;339(22):1585-94.
- 6-Peyvandi F, Lavoretano S, Palla R, Feys HB, Vanhoorelbeke K, Battaglioli T et al ADAMTS13 and anti-ADAMTS13 antibodies as markers for recurrence of acquired thrombotic thrombocytopenic purpura during remission. *Haematologica*. 2008;93(2):232-9.
- 7- Cugno M, Gualtierotti R, Possenti I, Testa S, Tel F, Griffini S et al. Complement Functional Tests For Monitoring Eculizumab Treatment In Patients With Atypical Hemolytic Uremic Syndrome. *J Thromb Haemost*. 2014 May 23 [Epub ahead of print]
- 8- Cataland SR, Holers VM, Geyer S, Yang S, Wu HM. Biomarkers of terminal complement activation confirm the diagnosis of aHUS and differentiate aHUS from TTP. *Blood* 2014;123(24):3733-8
- 9- Noris M, Caprioli J, Bresin E, , Mossali C, Pianetti G, Gamba S et al. Relative role of genetic complement abnormalities in sporadic and familial aHUS and their impact on clinical phenotype. *Clin J Am Soc Nephrol*. 2010;5(10):1844-59
- 10-Fremeaux-Bacchi V, Moulton EA, Kavanagh D, Dragon-Durey MA, Blouin J, Caudy A et al Genetic and functional analyses of membrane cofactor protein (CD46) mutations in atypical hemolytic uremic syndrome. *J Am Soc Nephrol*. 2006;17(7):2017-25.
- 11-Richards A, Kathryn Liszewski M, Kavanagh D, Fang CJ, Moulton E, Fremeaux-Bacchi V, et al Implications of the initial mutations in membrane cofactor protein (MCP; CD46) leading to atypical hemolytic uremic syndrome. *Mol Immunol*. 2007 ;44(1-3):111-22.
- 12-Remuzzi G, Galbusera M, Noris M, Canciani MT, Daina E, Bresin E, et al Von Willebrand factor cleaving protease (ADAMTS13) is deficient in recurrent and familial thrombotic thrombocytopenic purpura and hemolytic uremic syndrome *Blood* 2002; 100(3): 778-785
- 13-Scully M, McDonald V, Cavenagh J, Hunt BJ, Longair I, Cohen H et al A phase 2 study of the safety and efficacy of rituximab with plasma exchange in acute acquired thrombotic thrombocytopenic purpura. *Blood*. 2011;118(7):1746-53.

14-Cataland SR, Peyvandi F, Mannucci PM, Lämmle B, Kremer Hovinga JA, Machin SJ et al Initial experience from a double-blind, placebo-controlled, clinical outcome study of ARC1779 in patients with thrombotic thrombocytopenic purpura. *Am J Hematol.* 2012;87(4):430-2.

15- Fakhouri F, Frémeaux-Bacchi V, Loirat C. Atypical hemolytic uremic syndrome: From the rediscovery of complement to targeted therapy. *Eur J Intern Med.* 2013;24(6): 492-5


Table 1. ADAMTS13-, VWF- and complement system-related plasmatic measurements at acute disease and remission in the patient.

Parameter	Normal values	Acute disease	Remission
ADAMTS13 activity, %	46-160	47	100
ADAMTS13 antigen, %	40-155	71	86
Anti-ADAMTS13 antibodies ^a	Absent	Absent	Absent
VWF antigen, %	55-165	194	150
ULVWF ratio	0.85-1.21	0.50	1.01
Classical complement pathway activity, %	69-129	56	79
Alternative complement pathway activity, %	30-113	24	60
Mannose binding lectin complement pathway activity, %	0-125	45	74
C3, %	70-130	69	114
C4, %	70-130	58	77
FH antigen, %	66-122	62	94
Anti-FH antibodies, Unit/mL	Less than 5.20	1.55	0.09

VWF, von Willebrand factor; ULVWF, ultra-large forms of von Willebrand factor; C3, complement system component 3; C4, complement system component C4, FH, complement system factor H.

^aAnti-ADAMTS13 antibodies were screened by western blotting.

Figure 1. Reverse transcription PCR analysis of the CD46 transcripts in the patient and her family.



The figure shows the results of southern blotting of *CD46* cDNA (upper left), the family tree (upper right), and the Sanger sequencing traces (bottom). Family tree: I, II, and III indicate the generations. Sanger traces: A, mother who carried wild-type *CD46* alleles; B, patient; C, first son; D, the second son; B, C, and D carried the *CD46* c.475+1G>A heterozygote variant.