


Accepted Manuscript

Title: Adropin and apelin fluctuations throughout a season in professional soccer players: Are they related with performance?

Author: Fabian Sanchis-Gomar Rafael Alis Ermanno Rampinini Andrea Bosio Davide Ferioli Antonio La Torre Jincheng Xu Veronica Sansoni Silvia Perego Marco Romagnoli Giovanni Lombardi



PII: S0196-9781(15)00146-1
DOI: <http://dx.doi.org/doi:10.1016/j.peptides.2015.05.001>
Reference: PEP 69473

To appear in: *Peptides*

Received date: 30-3-2015
Revised date: 4-5-2015
Accepted date: 5-5-2015

Please cite this article as: Sanchis-Gomar F, Alis R, Rampinini E, Bosio A, Ferioli D, La Torre A, Xu J, Sansoni V, Perego S, Romagnoli M, Lombardi G, Adropin and apelin fluctuations throughout a season in professional soccer players: Are they related with performance?, *Peptides* (2015), <http://dx.doi.org/10.1016/j.peptides.2015.05.001>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

32 **Abstract**

33 Myokines are likely to be involved in the whole-body metabolic adaptive changes that
34 occur in response to regular exercise. We aimed to investigate the association of the two
35 myokines (adropin and apelin) with physical performance in professional soccer
36 players. To this purpose, we analyzed the fluctuations of circulating levels of both
37 adropin and apelin in professional soccer players during a season and evaluated the
38 possible association of these myokines with the performance level. *Creatine kinase*
39 (CK) and *lactate dehydrogenase* (LDH) activity as well as iron, transferrin and high-
40 sensitivity C-Reactive protein (hsCRP), ferritin, soluble transferrin receptor (sTfR), free
41 testosterone/cortisol ratio (FTCR), total iron binding capacity (TIBC) were also
42 determined. Fifteen male professional soccer players from an Italian Serie A team were
43 included in this study. Regarding the results of the biochemical analyses, the patterns of
44 changes in the biomarkers of fatigue and inflammation, i.e., HsCRP, CK and LDH
45 reflected the effects of the training throughout the season. No significant changes were
46 observed in adropin, while apelin exhibited variations that seem not to be related with
47 performance. In addition, both adropin and apelin did not represent valuable strategy to
48 assist in the performance assessment of professional soccer players.

49 **Keywords:** skeletal muscle; performance; cytokines.

50

51

52

53

54

55

56 **Introduction**

57 *Myokines* are cytokines produced by skeletal muscles, especially induced by exercise,
58 modulating different metabolic processes [6]. By influencing metabolism locally in the
59 muscles, myokines are thought to be involved in the whole-body metabolic adaptive
60 changes that occur in response to regular exercise like, for example, attenuation of fat
61 accumulation [2]. Skeletal muscle and pancreas act in a synergistic manner to monitor
62 systemic glucose homeostasis, and it has been suggested that myokines mediate the
63 cross-talk between insulin-sensitive tissues [17]. Striated skeletal muscle is one of the
64 body's largest tissues. However, it is unclear how contracting skeletal muscles transmit
65 metabolic positive effects on health. One of the possible explanations for the health
66 benefit of exercise can be that regular muscle contractions produce important
67 messengers such as myokines [5]. Released circulating myokines may explain how
68 normal muscle activity influences mood, physical performance and cognitive function
69 [14].

70 It has been shown that exercise up-regulates the expression of the newly described
71 myokine apelin in patients with type 2 diabetes [11]. In addition, apelin expression is
72 induced by exercise and secreted in vitro in human primary myotubes, and may behave
73 as a novel exercise-regulated myokine with autocrine/paracrine action [4]. Apelin is also
74 up-regulated by insulin, contributing thus to glucose homeostasis [19]. Finally, apelin is
75 highly implicated in cardiovascular function [10].

76 Adropin is also a recently described myokine involved in the regulation of lipid
77 metabolism. It was first isolated in 2008 by Kumar et al. in liver and brain tissues [12].
78 In mice, adropin regulates physical activity (locomotion and coordination) via the *NB-*
79 *3/Notch* signaling pathway [20]. Elevated circulating levels of adropin reduce insulin

80 resistance and glucose intolerance that arise in response to metabolic stress [7]. In this
81 case, there is no clear evidence about whether exercise can regulate circulating levels of
82 this myokine.

83 Therefore, because myokines are clearly involved in exercise-associated metabolic and
84 cardiac changes, and hence could be potentially implicated in performance
85 improvements throughout a soccer season, we aimed to analyze the fluctuations of
86 circulating apelin and adropin levels in professional soccer players during a season. In
87 addition, we also evaluated the possible association of both myokines with the
88 performance level.

89 **Material and methods**

90 *Subjects*

91 Fifteen male professional soccer players from an Italian Serie A team (age (mean \pm SD)
92 27 \pm 5 years, weight 76.9 \pm 4.1 kg, height 1.82 \pm 0.05 m, body fat 8.7 \pm 2.4 %) were included
93 in this study. Goalkeepers were not considered in this study since their physical load
94 during soccer games is different from the other field players and as such their training
95 programs are also different. All participants were informed of the purpose, protocol, and
96 procedures of the study before agreeing to participate. The study complies with the
97 World Medical Association Declaration of Helsinki regarding ethical conduct of
98 research involving human subjects and/or animals and was approved by the ethics
99 committee of University of Valencia, and by the soccer clubs involved.

100 *Experimental Protocol*

101 The players were sampled 3 times during the last part of a competitive season (in
102 January, in March, in May). The competitive season finished at the end of May.

103 Thereafter, all players took a vacation and returned to the team discipline at the
104 beginning of July, when the players included in the study were sampled again (just
105 before preseason training beginning). At all sampling points, both mokines were
106 assessed along with the physical performance determinations. An extensive biochemical
107 and hematological profile study was also performed at all time-points but at May.

108 *Physical performance determinations*

109 At all time-points, the players were subjected to three physical performance tests, a
110 continuous running test (Mognoni's Test) [8], a high-intensity intermittent test (HIT)
111 [18] and a counter-movement jump (CMJ) test.

112 During the Mognoni's Test, blood lactate (La^-) concentration was determined
113 immediately after a single 6-min run at $13.5 \text{ km}\cdot\text{h}^{-1}$ while the mean heart rate (HR) of
114 the last minute of running was considered for the analysis. After 10-min of passive
115 recovery, subjects completed, following an acoustic signal, a HIT protocol (total
116 duration = 5 min) consisting of 10 x 10 s shuttle running at $18 \text{ km}\cdot\text{h}^{-1}$ over a 25-m
117 course with an 180° direction change from run to run and 20s of passive recovery
118 between runs. Immediately after the HIT protocol, blood La^- concentration was
119 determined and the mean HR of 5-min run was considered for the analysis.

120 During both tests, blood La^- accumulation was measured using a portable amperometric
121 microvolume lactate analyzer (Lactate Plus, Nova Biomedical, Waltham, MA, USA).
122 Capillary blood samples ($0.7 \mu\text{l}$) were collected from the earlobe. Before the tests, the
123 analyzer was calibrated following the instructions of the manufacturer. HR data was
124 collected using Polar Team² Pro system (Kempele, Finland).

125 The CMJ test was performed using a portable force platform (Quattro Jump, Kistler,
126 Switzerland). In short, after a standardized warm-up, the subjects performed 6 single
127 CMJ and jump height and peak power output (PPO), averaged from the 3 best jumps
128 were recorded.

129 In addition, the body fat percentage was estimated at all time points by the skin-fold
130 technique, based on the Jackson and Pollock formula [9].

131 *Blood sampling*

132 Blood collection and sample management before analysis was carried out strictly
133 following the good laboratory practice for pre-analytical phase of sports biochemistry
134 and hematology tests [3]. The samples were drawn by venipuncture in the antecubital
135 vein in fasting conditions. For hematological determinations, blood was collected in
136 K₃EDTA vacuum tubes (Vacutainer, BD, Franklin Lakes, NJ, USA) and for
137 biochemical tests, in vacuum plain tubes without additives (Vacutainer). The former
138 were allowed to coagulate, then centrifuged at 3000 g for 10 min at room temperature
139 and after centrifugation, serum was separated into aliquots. Whole blood tubes were
140 immediately kept refrigerated at 4° C and assayed within 24 hours. Serum aliquots were
141 frozen at -80 °C until the assay.

142 *Laboratory Methods*

143 Adropin and apelin were determined in serum samples using available commercial
144 competitive enzyme-linked immunosorbent assay kits (CSB-EL007669HU, Cusabio,
145 Wuhan, China and EIA-APC, RayBiotech, Norcross, GA, USA; respectively). Both
146 assays were performed in duplicate following manufacturer's instructions. Intra-assay
147 coefficients of variation were 13.92% for adropin and 4.52% for apelin determinations.

148 Full blood cell count was carried out on the automated analyzer XE-2100L (Sysmex,
149 Kobe, Japan). *Creatine kinase* (CK) and *lactate dehydrogenase* (LDH) activity as well
150 as iron, transferrin and high-sensitivity C-Reactive protein (hsCRP) concentration were
151 determined on the automated clinical chemistry platform ADVIA 1800 (Siemens
152 Healthcare Diagnostics, Erlangen, Germany), employing proprietary reagents. Cortisol
153 and testosterone were immunoassayed on the automated analyzer Elecsys 1010 (Roche
154 Diagnostics, Mannheim, Germany) using the dedicated electro-chemiluminescence
155 immunoassay kits. Ferritin was assayed on automated immunoassay system (ADVIA
156 Centaur) and soluble transferrin receptor (sTfR) on a fully automated
157 immunonephelometer (BN ProSpec), both provided by Siemens Healthcare Diagnostics
158 and by using proprietary immunoassays. For calculation of the testosterone/cortisol ratio
159 (FTCR), free testosterone was assumed as 2% of the total testosterone, and the formula
160 previously validated was adopted [1]. Also the total iron binding capacity (TIBC) was
161 estimated by applying the formula [TIBC ($\mu\text{g/dL}$) = transferrin (g/L) x 140] derived
162 from the stoichiometric relationship between divalent transferrin and iron.

163 *Statistical analysis*

164 All data were analyzed for normality by Shapiro-Wilk test. Since the majority of the
165 variables were not normally distributed, non-parametric tests were adopted. The effect
166 of training and detraining (sampling time: January, March, May and July) on the
167 parameters tested was analyzed with the Friedman's test (χ^2) and paired comparisons
168 were performed with the Wilcoxon's test (z). The Spearman's coefficient (ρ) was used
169 to explore the correlation between adropin and apelin levels as well as with the other
170 parameters determined. The statistical analyses were performed using SPSS, version 21
171 (IBM Corporation, Armonk, NY, USA). The results were considered statistically
172 significant at $p \leq 0.05$. Data were expressed as median (10th-90th percentile).

173 **Results**

174 The aerobic endurance, assessed by post-exercise La^- levels in Mogroni's and HIT tests
175 changed at the end part of the competitive season and by the detraining period
176 [Mogroni $\chi^2(3)=19.53$, $p<0.001$; HIT $\chi^2(3)=15.53$, $p=0.001$; see **Figure 1**]. In the
177 Mogroni's test, La^- levels in January [3.50(2.55-6.82) mM] were higher than in March
178 [2.85(2.02-5.33) mM, $z=-2.552$, $p=0.011$] and in May [3.02(2.08-4.66) mM, $z=-2.601$,
179 $p=0.009$]. In the same line, La^- levels in the HIT test in March [2.47(1.37-6.21) mM]
180 were lower than in January [4.40(1.65-8.35) mM, $z=-2.045$, $p=0.041$] and May
181 [3.65(1.52-6.07) mM, $z=-2.090$, $p=0.002$]. This data would indicate a progressive
182 adaptation with training. However, the July La^- values were higher in both tests (see
183 **Figure 1**) which would reflect the detraining occurring during the vacation period
184 between the end of the season and the beginning of the next pre-season. Accordingly,
185 the cardiovascular implication was higher after the detraining period in both Mogroni's
186 [$\chi^2(3)=13.00$, $p=0.005$] and HIT [$\chi^2(3)=11.44$, $p=0.010$] tests (see **Figure 1**).
187 Nevertheless, we failed to find any significant effect of training and detraining on either
188 jump height [$\chi^2(3)=5.23$, $p=0.156$] or PPO [$\chi^2(3)=1.46$, $p=0.692$] in the CMJ test.

189 **Figure 2** shows a significant increase in the apelin concentration from 341.8(283.0-
190 444.8) ng/mL in January to 433.3(373.4-677.5) ng/mL in March. Nonetheless, no
191 statistically significant changes were found in either May or July compared to the
192 previous time points. No significant changes were observed in adropin at any time.

193 A full blood cell panel along with several biochemical parameters was performed in
194 January, March and July sampling times, but it could not be performed in May. These
195 data are displayed on **Table 1**. The detraining period induced some alterations in
196 erythrocyte indices. The mean corpuscular volume decreased while mean corpuscular

197 hemoglobin increased in July in comparison with the previous sampling points (**Table**
198 **1**). The red blood distribution width was slightly lower in July compared with January
199 levels ($z=-2.284$, $p=0.022$, **Table 1**). HsCRP, an inflammatory marker, was found to be
200 lower in July compared to January ($z=-2.528$, $p=0.011$). After detraining, levels of both
201 biomarkers of muscle damage CK and LDH, were significantly lower compared with
202 the previous time points (see **Table 1**).

203 Players' weight [January 77.0 (69.8-82.3) kg, March 77.0(69.0-83.5) kg, May
204 76.5(68.0-84.8) kg and July 77.6(70.1-86.9) kg] and percentage of body fat [January
205 8.4(5.8-13.1) %, March 8.1(6.5-12.0) %, May 8.2(5.5-12.0) %, July (9.2(7.0-12.4) %]
206 did not significantly change at the end of the competitive season or after the detraining
207 period [$\chi^2(3)=6.126$, $p=0.106$; $\chi^2(3)=4.576$, $p=0.206$; respectively].

208 Finally, no correlation was found between adropin or apelin concentrations and
209 performance parameters at all time points measured (**Supplementary Table 1**). On the
210 other hand, significant correlations were found between adropin and apelin levels and
211 other hematological and biochemical parameters measured (**Supplementary Table 2**),
212 although those correlations did not provide additional insights.

213 **Discussion**

214 No significant changes were observed in adropin levels, while apelin exhibited
215 variations that seem not to be related with performance. On the other hand, the patterns
216 of changes in the biomarkers of fatigue and inflammation, i.e., HsCRP, CK and LDH
217 reflected the effects of the training throughout the season.

218 While apelin levels showed a significant increase only in the first time point, possibly
219 linked to an increased effort, fluctuations in adropin levels did not reach statistical

220 significance. In both cases, however, the distribution widths within the study cohort
221 were large: this was particularly true in the case of adropin but was also evident for
222 apelin at the end of the season and after the rest period. It thus seems that the use of
223 these two markers is not useful to assist in the performance assessment of professional
224 soccer players. It should be also mentioned that, while other myokines seem to be
225 relatively stable within person [13-16] thereby allowing longitudinal analyses, there is
226 paucity of such data on apelin and adropin.

227 The main limitation of our study is the low number of subjects included on each
228 experimental group that can decrease the power of the statistical analyses performed.
229 Since season openings in Europe are regularly at mid/late August, we did not have a
230 first season's start sample to compare. In addition, the specific design of this study with
231 only professional athletes recruited limits the generalizability of its results in less
232 trained individuals. Moreover, due to a problem of samples conservation, the
233 hematological and biochemical panel could not be determined at May. Nevertheless,
234 this study needs to be evaluated in both larger and different cohorts before they can be
235 translated into clinical practice.

236 In conclusion, a one-season follow-up of professional soccer player's training did not
237 lead to observe changes in circulating apelin and adropin levels. Accordingly, neither
238 apelin nor adropin were associated with changes in performance level.

239 **Competing financial interests**

240 The authors declare no competing financial interests.

241

242 **Acknowledgements**

243 This research has been supported by grant DEP2012-37494 from the Spanish
244 Government and by grants 2013-168-002 and 2012-011-001 from Catholic University
245 of Valencia. RA is predoctoral fellow of Catholic University of Valencia. A special
246 thanks to the athletes involved in the study, to Cedal laboratory and Dr Alberto Dolci
247 for the support in the data collection.

248 **References**

249 [1] Adlercreutz H, Harkonen M, Kuoppasalmi K, Naveri H, Huhtaniemi I, Tikkanen H, et al.
250 Effect of training on plasma anabolic and catabolic steroid hormones and their response during
251 physical exercise. *Int J Sports Med.* 1986;7 Suppl 1:27-8.

252 [2] Aydin S. Three new players in energy regulation: preptin, adropin and irisin. *Peptides.*
253 2014;56:94-110.

254 [3] Banfi G, Dolci A. Preanalytical phase of sport biochemistry and haematology. *The Journal*
255 *of sports medicine and physical fitness.* 2003;43:223-30.

256 [4] Besse-Patin A, Montastier E, Vinel C, Castan-Laurell I, Louche K, Dray C, et al. Effect of
257 endurance training on skeletal muscle myokine expression in obese men: identification of apelin
258 as a novel myokine. *Int J Obes (Lond).* 2013.

259 [5] Bruunsgaard H. Physical activity and modulation of systemic low-level inflammation. *J*
260 *Leukoc Biol.* 2005;78:819-35.

261 [6] Febbraio MA, Pedersen BK. Contraction-induced myokine production and release: is
262 skeletal muscle an endocrine organ? *Exerc Sport Sci Rev.* 2005;33:114-9.

- 263 [7] Ganesh Kumar K, Zhang J, Gao S, Rossi J, McGuinness OP, Halem HH, et al. Adropin
264 deficiency is associated with increased adiposity and insulin resistance. *Obesity (Silver Spring)*.
265 2012;20:1394-402.
- 266 [8] Impellizzeri FM, Rampinini E, Marcora SM. Physiological assessment of aerobic training in
267 soccer. *Journal of sports sciences*. 2005;23:583-92.
- 268 [9] Jackson AS, Pollock ML. Generalized equations for predicting body density of men. *Br J*
269 *Nutr*. 1978;40:497-504.
- 270 [10] Kadoglou NP, Lampropoulos S, Kapelouzou A, Gkontopoulos A, Theofilogiannakos EK,
271 Fotiadis G, et al. Serum levels of apelin and ghrelin in patients with acute coronary syndromes
272 and established coronary artery disease--KOZANI STUDY. *Transl Res*. 2010;155:238-46.
- 273 [11] Kadoglou NP, Vrabas IS, Kapelouzou A, Lampropoulos S, Sailer N, Kostakis A, et al. The
274 impact of aerobic exercise training on novel adipokines, apelin and ghrelin, in patients with type
275 2 diabetes. *Med Sci Monit*. 2012;18:CR290-5.
- 276 [12] Kumar KG, Trevaskis JL, Lam DD, Sutton GM, Koza RA, Chouljenko VN, et al.
277 Identification of adropin as a secreted factor linking dietary macronutrient intake with energy
278 homeostasis and lipid metabolism. *Cell Metab*. 2008;8:468-81.
- 279 [13] Panagiotou G, Mu L, Na B, Mukamal KJ, Mantzoros CS. Circulating irisin, omentin-1, and
280 lipoprotein subparticles in adults at higher cardiovascular risk. *Metabolism: clinical and*
281 *experimental*. 2014;63:1265-71.
- 282 [14] Pedersen BK, Febbraio M. Muscle-derived interleukin-6--a possible link between skeletal
283 muscle, adipose tissue, liver, and brain. *Brain Behav Immun*. 2005;19:371-6.
- 284 [15] Pedersen BK, Febbraio MA. Muscle as an endocrine organ: focus on muscle-derived
285 interleukin-6. *Physiol Rev*. 2008;88:1379-406.
- 286 [16] Pedersen BK, Febbraio MA. Muscles, exercise and obesity: skeletal muscle as a secretory
287 organ. *Nat Rev Endocrinol*. 2012;8:457-65.
- 288 [17] Pedersen L, Hojman P. Muscle-to-organ cross talk mediated by myokines. *Adipocyte*.
289 2012;1:164-7.

290 [18] Rampinini E, Sassi A, Azzalin A, Castagna C, Menaspa P, Carlomagno D, et al.
291 Physiological determinants of Yo-Yo intermittent recovery tests in male soccer players. *Euro J*
292 *Appl Physiol.* 2010;108:401-9.

293 [19] Soriguer F, Garrido-Sanchez L, Garcia-Serrano S, Garcia-Almeida JM, Garcia-Arnes J,
294 Tinahones FJ, et al. Apelin levels are increased in morbidly obese subjects with type 2 diabetes
295 mellitus. *Obes Surg.* 2009;19:1574-80.

296 [20] Wong CM, Wang Y, Lee JT, Huang Z, Wu D, Xu A, et al. Adropin is a brain membrane-
297 bound protein regulating physical activity via the NB-3/Notch signaling pathway in mice. *J Biol*
298 *Chem.* 2014;289:25976-86.

299

Table and Figures Legends

Table 1. Players' hematological and biochemical parameters in January, March, and July (i.e., at the beginning of the next pre-season).

Figure 1. Blood lactate (La^-) and heart rate (HR) in the Mogroni and HIT protocols, and height jump and peak power output in the counter-movement jump test (CMJ) during the competitive season (January, March and May) and in the next pre-season stage (July). During the Mogroni's Test, La^- concentration was determined immediately after the test, while the mean HR of the test's last minute of running was considered. During the HIT protocol La^- concentration was determined just after the test and the mean HR of 5-min run was considered. CMJ height and peak power output was averaged from the 3 best of 6 jump repetitions. Data represented as median (horizontal line), 1st to 3rd quartile (box) and 10th to 90th percentile (whiskers). Significant comparisons are indicated.

Figure 2. Apelin and adropin concentrations during the competitive season (January, March and May) and in the next pre-season stage (July). Data represented as median (horizontal line), 1st to 3rd quartile (box) and 10th to 90th percentile (whiskers). Significant comparisons are indicated.

Table 1.

	In-season January	In-season March	Pre-season July	Friedman's p-value
RBC ($\times 10^6/\mu\text{L}$)	5.03(4.60-5.73)	5.12(4.78-5.45)	5.11(4.72-5.87)	0.247
Hemoglobin (g/dL)	15.1(13.8-16.0)	15.3(14.3-16.2)	15.1(13.7-16.7)	0.302
Hematocrit (%)	44.0(41.0-47.3)	44.5(42.0-46.8)	44.9(40.1-47.7)	0.591
MCV (fL)	87.2(81.9-89.6) ^{***}	85.9(83.0-91.2) ^{***}	84.7(80.9-88.4)	<0.001
MCH (pg)	30.0(27.9-31.2) [*]	30.1(27.9-31.1) ^{**,#}	29.5(27.8-30.9)	<0.001
MCHC (g/dL)	34.3(33.1-35.5)	34.5(33.2-35.9)	34.5(33.2-36.6)	0.581
RDW (%)	13.0(12.5-13.5) [*]	12.9(12.4-13.5)	12.8(12.3-13.6)	0.011
Reticulocytes (%)	0.67(0.51-1.02) ^{**}	0.66(0.43-0.88) ^{**}	0.82(0.57-1.33)	0.005
IRF (%)	3.2(1.44-5.98)	2.8(0.94-6.28)	3.2(1.26-5.78)	0.721
Fe ($\mu\text{g/dL}$)	82(60-117)	96(58-131)	84(67-145)	0.627
Ferritin (ng/mL)	125.7(69.1-246.7)	130.2(48.7-244.7)	113.4(51.6-217.6)	0.070
Transferrin (mg/dL)	249(213-273)	242(211-290)	259(207-297)	0.085
TIBC ($\mu\text{g/dL}$)	349(299-383)	339(296-405)	363(290-416)	0.085
Transferrin saturation (%)	25(18-32)	26(18-41)	26(18-39)	0.349
sTfR (mg/L)	1.30(0.84-1.63)	1.15(0.90-1.49)	1.19(0.83-1.62)	0.591
WBC ($\times 10^3/\mu\text{L}$)	4.97(4.01-6.52)	4.70(3.89-7.5)	5.01(4.25-7.76)	0.516
Lymphocytes ($\times 10^3/\mu\text{L}$)	2.2(1.4-3.3)	2.1(1.5-2.9)	2.3(1.6-3.0)	0.272
Neutrophils ($\times 10^3/\mu\text{L}$)	2.3(1.4-2.9)	2.4(1.4-4.1)	2.3(1.5-3.9)	0.179
Monocytes ($\times 10^3/\mu\text{L}$)	0.4(0.3-0.6)	0.4(0.3-0.6)	0.4(0.3-0.6)	0.971
Basophils ($\times 10^3/\mu\text{L}$)	0.02(0.01-0.03)	0.02(0.01-0.04)	0.02(0.01-0.03)	0.928
Eosinophils ($\times 10^3/\mu\text{L}$)	0.1(0.1-0.4) ^{**}	0.1(0.1-0.5)	0.2(0.1-0.5)	0.009
hsCRP (mg/L)	0.97(0.40-8.22) [*]	0.71(0.16-2.25)	0.30(0.13-2.49)	0.022
Creatine kinase (U/L)	392(240-2053) ^{**}	382(168-713) [*]	294(125-925)	<0.001
Lactate dehydrogenase (U/L)	205(73-271) [*]	190(148-231) ^{*,#}	158(127-217)	<0.001
Cortisol (ng/mL)	238.0(190.3-277.4)	214.9(183.9-271.7)	214.1(135.2-260.1)	0.165
Testosterone (ng/mL)	7.21(4.15-9.57)	7.61(5.39-9.34)	7.16(5.39-9.95)	0.766
FTCR	0.74(0.47-0.95)	0.82(0.58-0.99)	0.91(0.61-1.57)	0.241

RBC: red blood cell, MCV: mean corpuscular volume, MCH: mean corpuscular hemoglobin, MCHC: mean corpuscular hemoglobin concentration, RDW: red blood cell distribution width, IRF: non-mature reticulocyte fraction, sTfR: soluble transferrin receptor, TIBC: total iron-binding capacity, WBC: white blood cell, hsCRP: high-sensitivity C-Reactive protein, FTCR: free-testosterone to cortisol ratio. Data as median (10th-90th percentile).

*p<0.05, **p<0.01 and ***p<0.001 vs July. #p<0.05, ##p<0.01 and ###p<0.01 vs January.

Highlights

- Myokines are involved in metabolic adaptive changes induced by regular exercise.
- We investigated the association of two myokines (adropin and apelin) with physical performance.
- No significant changes were observed in adropin.
- Apelin exhibited variations that seem not to be related with performance.
- Apelin and adropin levels are not related to performance in professional soccer players.

