



New insight on crystal and spot development in hard and extra hard cheeses: association of spots with incomplete aggregation of curd granules

Journal:	<i>Journal of Dairy Science</i>
Manuscript ID	JDS-16-11050.R1
Article Type:	Research
Date Submitted by the Author:	n/a
Complete List of Authors:	D'Incecco, Paolo; University of Milan, Department of Food, Environmental and Nutritional Sciences Limbo, Sara; University of Milan, Department of Food Environmental and Nutritional Sciences Faoro, Franco; University of Milan, Department of Agricultural and Environmental Sciences - Production, Landscape, Agroenergy Hogenboom, John; University of Milan, Department of Food Environmental and Nutritional Sciences Rosi, Veronica; University of Milan, Department of Food Environmental and Nutritional Sciences Morandi, Stefano; National Research Council , ISPA Pellegrino, Luisa; University of Milan, Department of Food Environmental and Nutritional Sciences
Key Words:	hard and extra hard cheeses, cheese ultrastructure, calcium phosphate crystal, free amino acids

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11

INTERPRETATIVE SUMMARY

New insight on crystal and spot development in hard and extra hard cheeses: association of spots with incomplete aggregation of curd granules

D’Incecco

The study was carried out with a multidisciplinary approach, using techniques such as confocal, Raman and electron microscopy, to shed light on the phenomenon of speck and spot blowing in hard cheese. The obtained information on structure and chemical composition of these particles, allowed us to formulate an original theory for their genesis. Furthermore, these results provide useful information to be applied in cheese making technology in order to contain the incidence of the speck and spot phenomenon.

12 **RUNNING HEAD: CRYSTALS AND SPOTS IN HARD AND EXTRA HARD CHEESES**

13 **New insight on crystal and spot development in hard and extra hard cheeses: association of**
14 **spots with incomplete aggregation of curd granules**

15 **P. D’Incecco,^{*1} S. Limbo,* F. Faoro,[†] J. Hogenboom,* V. Rosi,* S. Morandi,[‡] L. Pellegrino***

16

17 * Department of Food Environmental and Nutritional Sciences, State University of Milan, 20133
18 Milan, Italy.

19

20 † Department of Agricultural and Environmental Sciences - Production, Landscape, Agroenergy,
21 State University of Milan, 20133 Milan, Italy.

22

23 ‡ Institute of Sciences of Food Production–Italian National Research Council, 20133 Milan, Italy.

24

25 ¹Corresponding author: P. D’Incecco

26 Department of Food, Environmental and Nutritional Sciences, University of Milan, Via Celoria 2,
27 20133 Milan, Italy

28 Telephone: (+39) 0250316679

29 Fax: (+39) 0250316672

30 E-mail: paolo.dincecco@unimi.it

31

32 **ABSTRACT**

33 Chemical composition and structure of different types of macro- (specks, spots) and micro particles
34 (microcrystals) present in hard and extra hard cheeses were investigated. Light microscopy revealed
35 that the small, hard specks had the structure of crystalline tyrosine, as confirmed by amino acid
36 analysis. Spots showed a complex structure, including several curd granules, cavities, and
37 microcrystals, and were delimited by a dense protein layer. Spots contained less moisture and ash
38 than the adjacent cheese area, and more protein, including significantly higher contents of valine,
39 methionine, isoleucine, leucine, tyrosine and phenylalanine. Microcrystals were observed by light
40 and electron microscopy and analysed by confocal micro Raman. Among others, calcium phosphate
41 crystals appeared to consist of a central star-shaped structure immersed in a matrix of free fatty
42 acids besides leucine and phenylalanine, in free form or in small peptides. A hypothetical
43 mechanism for the formation of these structures has been formulated.

44

45 **Keywords:** hard and extra hard cheeses, cheese ultrastructure, calcium phosphate crystal, free
46 amino acids, electron microscopy

47

48

INTRODUCTION

49 Hard and Extra-Hard are attributes used to define cheeses having a firm and brittle body texture
50 (Codex Alimentarius, 1978). Hard and extra-hard cheeses share low moisture content, close
51 structure, and a long ripening period. During ripening, many chemical, biochemical and
52 microbiological phenomena take place. The biochemical changes are very important for the
53 development of the flavour and texture of these cheeses and are characteristic of the different
54 varieties. Proteolysis is the most relevant among the biochemical phenomena due to its complexity
55 and final impact on the cheese taste. In fact, casein breakdown progressively brings to large and
56 medium peptides, then to small peptides and free amino acids (FAA). Since FAA are rather stable,
57 they tend to accumulate with the ripening time and may reach up to 20-24% on cheese protein basis
58 in 10-12 months old extra-hard cheeses (Masotti et al., 2010). Proteinases and peptidases that
59 catalyse proteolysis in cheese originate from different sources, namely milk, rennet, starter and non-
60 starter lactic acid bacteria (LAB). LAB have complex enzyme patterns that release peptides and
61 amino acids from the proteins into the cheese environment to satisfy their own nutritional
62 requirements (Gatti et al., 2014). After vat processing, the loss of water, diffusion of salt, and
63 formation of soluble molecules, such as FAA and lactate, are factors concurring to the increase of
64 solute concentration and concomitant decrease of water activity (a_w) in cheese throughout the whole
65 ripening period. Beside these main events, minor changes contribute to lower the water activity in
66 cheese, like changes in water binding by new carboxylic and amino groups formed on protein
67 hydrolysis (McSweeney, 2004). The moisture content and a_w are strongly correlated in cheese
68 throughout ripening (Marcos, 1993).

69 The increasing solute (salt, ions, FAA) concentration in cheese water phase may give rise to
70 aggregation and crystallization phenomena that result in different types of structures observed by
71 some authors in the interior and on the surface of different cheese varieties (Bianchi et al., 1974;
72 Agarwal et al., 2006; Tansman et al., 2015). Although earliest studies date back to the 1900s

73 (Babcock et al., 1903; Tuckey et al., 1938), a clear and unambiguous characterization of these
74 structures has not been achieved yet. Moreover, an univocal association between the terminology
75 (e.g. crystals, specks, dots, granules, spots, pearls) and the appearance of these structures is still
76 lacking.

77 In long ripened extra-hard cheeses, having a thick dry rind, these structures develop inside the
78 cheese becoming visible when the wheel is cut. Typically, two different types of structures visible
79 to the naked eye can be observed, referred to as specks and spots in this article. Specks look bright
80 white and firmer against the cheese matrix, and are usually smaller than 2-3 mm. Consumers
81 (cheese lovers) appreciate the crispness of the specks while chewing the cheese and their
82 contribution to the overall cheese taste. Previous studies about specks reported them to contain
83 clusters of tyrosine, cysteine as well as other FAA, calcium lactate and magnesium (Shock et al.
84 1948) or tyrosine and phenylalanine (Giolitti and Mascherpa, 1970). More recently, Bottazzi et al.,
85 (1994) and Tansman et al., (2015) converged on identifying them as tyrosine crystals in extra hard
86 cheeses.

87 Spots are spherical and paler than the cheese, and can grow up to 4-5 mm. They appear to be
88 amorphous and firmer with respect to the surrounding cheese matrix, and become visible after 10-
89 12 months of ripening. **Spots can become so numerous and flashy that may influence the visual**
90 **appeal of the cheese.** Spots have been very little studied, moreover without achieving consistent
91 results (Giolitti and Mascherpa, 1970; Bianchi et al., 1974; Tansman et al., 2015).

92 Besides specks and spots, extra-hard cheeses contain microscopic crystals, mostly investigated in
93 Cheddar cheese. However, some authors generically referred to crystals, without distinguishing
94 between the microscopic ones and those visible to the naked eye (Kalab, 1980; Bottazzi et al., 1982;
95 Washam et al., 1985; Bottazzi et al., 1994).

96 The aim of the present work was to shed light on the nature and origin of specks, spots and
97 microscopic crystals in extra-hard cheeses by a multidisciplinary approach. Due to the effectiveness
98 in cheese structure studies (Ong et al., 2010; Schrader et al., 2012; D'Incecco et al., 2015), various

99 microscopy techniques and different dyes (light and fluorescence, confocal, confocal micro Raman
100 and transmission electron microscopy) were used in combination with chemical data to achieve an
101 unambiguous characterization of these particles in cheese. Our ultimate goal was to formulate an
102 hypothesis on the origin of these structures as they appear in hard and extra-hard cheeses. This
103 knowledge will contribute useful information to understanding the bioavailability of selected
104 minerals and nutrients in cheese. Furthermore, this knowledge could provide insights into the nature
105 of these structures that might lead to new manufacturing strategies to control the formation of spots
106 in commercial cheeses.

107 MATERIALS AND METHODS

108 **Cheese samples and collection of specks and spots.** Eleven extra-hard cheeses, ripened for 18-20
109 months, were kindly provided by two dairies producing Grana Padano (7 cheeses) and Parmigiano-
110 Reggiano (4 cheeses) respectively. Specks were harvested from the cheese using a pin, while spots
111 were collected from the cheese using a spatula and then gently brushed to remove the cheese matrix
112 on the surface. Specks and spots were separately collected from individual cheese samples in a
113 sufficient amount (20-22 g) to conduct all the analyses. Equivalent cheese amount was taken from
114 the portion (0.5-cm thick) immediately surrounding the single spot, as shown in Figure 1, as used as
115 a term of reference. When necessary, a slice representative of the whole cheese was taken as well.
116 Additional cheese portions were taken as required for microscopy investigations with various
117 techniques. In particular, 20 spots of different size and taken from different cheeses were observed
118 for their structural characterization.

119 Chemicals.

120 Glutaraldehyde, paraformaldehyde, cacodylate buffer, and osmium tetroxide were purchased from
121 Agar Scientific (Stansted, UK). Toluidine blue, rhodamine and single amino acids were purchased
122 from Sigma Aldrich (Milan, Italy). Ninhydrin was purchased from Biochrom Ltd (Cambridge, UK).
123 Water purified with Milli-Q system (Millipore Corp., Bedford, MA) was used.

124 Composition analyses.

125 The ISO Standard methods for cheese were used to determine the content of protein (ISO
126 27871:2011), fat (ISO 1735:2004), ash (ISO 5545:2008) and moisture (ISO5534:2004),
127 respectively. Content of calcium and phosphorus were determined by ICP-MS spectrometer
128 (Agilent Technologies, Milan, Italy).

129 The pattern of FAA was determined on the various cheese portions (including pecks and spots)
130 using the method described by Masotti et al. (2010). Briefly, the cheese portion was solubilized
131 with sodium citrate, homogenized and deproteinized with sulphosalicylic acid. The extract was
132 diluted (1:1) with lithium citrate buffer at pH 2.2, filtered and analysed by ion exchange
133 chromatography. The chromatographic separation was carried out on a Biochrom 30+ (Biochrom
134 Ltd, Cambridge, UK) amino acid analyser operated under the conditions provided by the
135 manufacturer. These employ an eight-step elution program with lithium citrate buffers of increasing
136 pH and ionic strength, post-column derivatisation with ninhydrin, and detection at 440 and 570 nm.
137 The quantification was carried out using four-level calibration lines of the 21 amino acids in the
138 range 0.75-22.5 mg/L and using norleucine (Sigma Aldrich) as an internal standard. Repeatability
139 values of ISO Standard 13903:2005 were fulfilled.

140 **Amino acid diffusion trial.**

141 To confirm the different diffusivity of individual FAA within the cheese, 0.3 mL of an aqueous
142 FAA solution having three times the concentration of the cheese water phase was injected into the
143 core of a spot-free cheese portion (a disk of 10 g) using a microsyringe. The cheese portion was
144 kept in a forced-ventilation thermostatic oven at 18 °C for 18 days and then sampled as described in
145 the Results. The FAA pattern and the moisture content were determined in each sampled portion.

146 **Light and Confocal Microscopy.**

147 Specimens for light and fluorescence microscopy observations were thin sections obtained from
148 resin-embedded cheese samples prepared as below described for transmission electron microscopy
149 (TEM). Thin sections (4-5 per sample) were directly dried on the microscope slide, stained and
150 subsequently washed. Two different staining were performed, (i) toluidine blue (1% in water, w/v)

151 for 5 min at room temperature, to visualize the overall protein structure by light microscopy, and (ii)
152 rhodamine B (0.5% in water, w/v) for 5 min at room temperature. In the latter case, the sample was
153 examined by a Hg lamp, with the following filters: excitation wavelength = 570 nm, emission
154 wavelength = 590 nm. In addition a 5-cm cube of cheese was immersed in ninhydrin solution for 1
155 h and then cut with a blade until a thin section containing a spot was obtained. All samples were
156 examined with an Olympus BX optical microscope (Tokyo, Japan) equipped with Nomarski
157 interference contrast and QImaging Retiga camera (Surrey, BC, Canada). Specimens for confocal
158 microscopy observations were cryo-sectioned by a CM1950 cryostat (Leica, Germany) and stained
159 directly onto the microscope slide with fast green (0.1 % in water, w/v). Sections were examined
160 with a Video confocal microscope, Nikon Vico (Tokyo, Japan).

161 **Transmission Electron Microscopy.**

162 Cubes of cheese (1 mm edge length) were fixed in a mixture (w/v) of glutaraldehyde 3% and
163 paraformaldehyde 2% in cacodylate buffer for 2 h at 4 °C, then washed with cacodylate buffer for 1
164 h and post-fixed in osmium tetroxide (1% in water, w/v) for 2 h. After the dehydration in an ethanol
165 series, the samples were embedded in London Resin White™ resin and cured at 60 °C for 24 h.
166 Ultrathin sections (50 to 60 nm thick) were stained with uranyl acetate and lead citrate and
167 examined with a Philips E208 transmission electron microscope (Aachen, Germany).

168 **Confocal Micro Raman.**

169 The Raman spectral data were collected in the range from 3200 to 200/cm Raman shift using a
170 confocal DXR Raman Microscope (Thermo Scientific, Waltham, MA, USA). An Olympus 50X
171 objective (numerical aperture 0.75) with a 50 µm confocal pinhole was used to collect the Raman
172 signal directly from a flat area of the sample (cut using a sharp knife) with a spatial resolution lower
173 than 1 µm, without any preparation of the sample. A laser with an excitation wavelength of 780 nm
174 with a low energy power (5-10 mW) to avoid overheating and a 400 lines/mm grating was used to
175 record Raman spectra over the focalized area. A photobleaching time equal to 1 min was set up. For
176 the specks, spectra were collected individually while for the microcrystals a selected area was

177 analysed collecting around 70 spectra over the entire surface, using 10 μm as the interval between
178 positions. In particular, each sample was placed on an automated x,y mapping stage and Raman
179 spectra were obtained at different points of the selected surface, by moving it under the microscope
180 objective. Autofocus at each map point was applied. Omnic Atlus software (Thermo Fisher
181 Scientific, Madison, WI, USA) was used to obtain Raman spectra, perform spectrometer operations
182 and process data. All spectra were corrected for background contributions and an automated
183 subtraction of cosmic ray peaks was employed.

184 **Statistical analysis**

185 Statistical treatment of data was performed by means of SPSS Win 12.0 program (SPSS Inc.,
186 Chicago, IL, USA). Data were analysed by Student's t-test and one way Anova. A $P < 0.05$ was
187 assumed as significance limit.

188 **RESULTS**

189 **Speck characterization.** Attempts to obtain a section of specks or to embed them in resin failed
190 because they were too hard and packed. In contrast, specks *in toto* could be directly observed by
191 light microscopy after isolation by cheese and showed the characteristic structure of crystalline
192 tyrosine (Figure 2). The FAA pattern (Figure 2) indicated that specks were indeed tyrosine crystals
193 of $> 95\%$ purity. The peak of ammonia in the chromatogram derived from the elution buffers and
194 thus was ignored in purity calculation. The Raman spectrum also confirmed the nature of the
195 specks: the doublet Raman bands at 828 and 848/cm due to the Fermi resonance between ring
196 fundamental and overtone were strongly evident, as was the ring-O stretching vibration located at
197 1263/cm (Culka et al., 2010). Also the highly resolved bands found in the finger print region (1614-
198 250/cm) were associated to the signals collected for the solid and pure (98-99%) crystalline form of
199 L-tyrosine (spectrum not shown).

200 **Spot characterization.**

201 Spot structure was firstly examined by light microscopy. A cheese portion was immersed in the
202 protein-staining solution (ninhydrin) and then sectioned up to reveal a spot inside which appeared

203 surrounded by an highly dense layer (Figure 3, arrowhead) that impaired staining permeation inside.
204 However, some stain could permeate through preferential micro pathways (junctions, indicated by
205 arrows in Figure 3), making the structure visible. Thin sections of resin-embedded spots, stained
206 with toluidine blue, gave more insight of the inner structure (Figure 4a). The spot appeared as made
207 of several curd particles having clean-cut irregular shapes and size up to 0.5 mm. The darker lines,
208 corresponding to the junctions among curd particles, were richer in protein than the particle body.
209 Several openings, including a large hole collecting the junctions, were visible in the section (Figure
210 4b). Although the cheese around the spot showed the same composite structure (not shown), large
211 cavities were observed only inside spots. With respect to the curd particles, the junctions appeared
212 as thick protein strings almost free of fat (Figure 4c) and rich in microcrystals (arrows).
213 Microcrystals were also observed in the cheese matrix outside the spot, as further discussed.
214 The ultrastructure of the spot, examined by TEM, proved to be remarkably different compared to
215 that of the surrounding cheese (Figure 5). In particular, the interface between protein (Figure 5, in
216 grey) and fat (in white) was more irregular and fringed by crystal-like particles (Figure 5a) in
217 respect with the cheese surrounding the spot (Figure 5b).
218 The chemical composition of the spots was compared to that of the cheese portion just around them
219 and to that of the whole cheese. Five different cheeses were individually analysed (Table 1). Spots
220 proved to be significantly richer in protein ($p < 0.00$) and poorer in moisture ($p < 0.00$) and ash
221 ($p < 0.00$) with respect to the surrounding cheese portion, which did not differ from the whole
222 cheese. Fat content was not different ($p < 0.17$). The contents of both calcium and phosphorus were
223 10% lower in the spot than in the cheese, with a Ca/P molar ratio of 1 in both zones (not shown).
224 The protein fraction was characterized by capillary zone electrophoresis as described in our
225 previous work (Masotti et al., 2010). No difference could be evidenced, neither in casein nor in
226 peptide patterns, between the spots and the rest of the cheese (not shown), indicating that the
227 primary proteolysis had proceeded to the same extent within and outside the spot. On the contrary,
228 significant differences were found in the FAA patterns of the two zones.

229 In order to compare data of cheese portions (i.e. spots and surrounding cheese) having different
230 moisture contents, and considering that FAA are soluble molecules, values were expressed on the
231 respective moisture content. Furthermore, data from both Grana Padano and Parmigiano-Reggiano
232 were pooled to increase statistical significance. In particular, spots contained significantly higher
233 amounts (from 4 to 12 times) of six FAA, i.e. valine ($p<0.00$), methionine ($p<0.00$), isoleucine
234 ($p<0.00$), leucine ($p<0.00$), tyrosine ($p<0.00$) and phenylalanine ($p<0.00$) (Figure 6). It is worthy of
235 remark that the content of the other FAA was the same as in the cheese portion around the spot.
236 Nevertheless, this last portion had the usual FAA pattern we observed for the two target cheeses in
237 previous studies (Cattaneo et al., 2008; Masotti et al., 2010). To achieve direct confirmation of such
238 a different composition in individual FAA within the cheese, a simple experiment was carried out.
239 A water solution having the approximate concentration of FAA in cheese water phase was injected
240 into a spot-free cheese slice. After a resting time suitable to allow diffusion of the solution, the
241 cheese was sampled taking three distinct portions: one circular, corresponding to the injection point
242 (mimicking the spot), and two concentric rings around it, as shown in Figure 7 (inset). Despite of
243 the approximate experimental conditions and sampling procedure, the obtained data confirmed that
244 the same six FAA were retained in the zone where the mixture was injected, while the others
245 diffused to the surrounding portions largely reaching an equilibrium (Figure 7).

246 **Microscopic crystals.**

247 Microscopic crystals were investigated through various microscopy techniques. Observations by
248 fluorescence microscopy of cheese semi-thin sections, after resin embedding and staining with
249 rhodamine B, allowed to see a huge number of microscopic crystals in bright red (Figure 8a).
250 Confocal microscopy showed that crystals had different shapes, i.e. circular, oval or kidney-shaped,
251 and their core was not fluorescent, indicating a different composition in respect to the peripheral
252 zone (Figure 8b). The number of crystals ranged from 30 to 100 crystals/mm² in 18-20 months
253 ripened cheeses and apparently was not different between cheese and inside spots. Crystals with the

254 same structure were detected in younger (three and six months of ripening) extra hard cheeses,
255 although in a lower amount (not shown).

256 Microscopic crystals observed by TEM showed a complex ultrastructure (Figure 8c). Three main
257 zones could be outlined: (i) a central star-shaped crystal, (ii) an intermediate zone and (iii) a
258 peripheral compact shell at the interface with the protein matrix. Figures 8d shows both the external
259 shell and the intermediate zone to be constituted by fibrils with prismatic morphology, radially
260 ordered and packed around the central crystal.

261 Confocal Raman microscopy was used to obtain information on the chemical composition since this
262 technique does not require any sample preparation. Area maps of the crystals, detected both inside
263 and outside the spot, confirmed the presence of distinct regions, as observed by TEM. Spectra of the
264 central star shaped crystal matched the calcium phosphate spectrum, dominated by the very strong
265 band at 986/cm and the medium band at 878/cm (Figure 9) that derived from the symmetric
266 stretching mode of the phosphate group (Sauer et al., 1994). The band at 1081/cm corresponded to
267 the stretching vibration (ν_3) of PO_4^{3-} , while the band at 588/cm to the P-O and O-P-O stretching and
268 bending modes (ν_4) of the same group. In calcium phosphate crystals, the minerals can be identified
269 by the position and shape of the main bands. Raman shifts and assignment for some calcium
270 phosphate minerals were studied by Koutsopoulos (2002). The Savitsky-Golay second derivative of
271 the spectra highlighted the presence of other weak bands attributable to the dibasic calcium
272 phosphate dehydrate form. No calcium phosphate was detected in other parts of the complex crystal
273 structure. In fact, in the outer zone, appearing as a dark area in the optical magnification used for
274 the Raman acquisition, spectra presented bands that arose from both free fatty acids and proteins
275 (Figure 10a). The spatial distribution of fatty acid/protein with respect the calcium phosphate within
276 the crystal area is shown in Figure 10b (in blue). It is evident that the crystal is immersed in the
277 cheese matrix. A deeper analysis of the spectra of this portion provided a major characterization of
278 the protein structure contribution. Usually, the amide I and III peaks in a protein are less sharply
279 resolved if compared with the signals of small peptides (Jenkins et al., 2005). In other words,

280 Raman spectral signatures of the single amino acids are retained in the peptide or protein, being
281 largely derived from their side chain or backbone. The higher the resolution and intensity of the
282 bands in a complex spectrum, the more probable the contribution of amino acids and/or small
283 peptides. Thus, in the original spectrum collected from the cheese matrix around the phosphate
284 crystal, some peculiar bands of amino acids were evident. In particular, a spectral subtraction of an
285 unsaturated free fatty acid spectrum (considered as a background) from the original spectrum of the
286 external region returned the profile of leucine (match higher than 60% with the pure spectrum of L-
287 leucine) and phenylalanine (match equal to 53% with the pure spectrum of L-phenylalanine).
288 Leucine was mainly characterized by the bands at 1237/cm due to the twisting of CH₂, and by the
289 bands at 1187 and 1132/cm due to the rocking of NH₃⁺, whilst phenylalanine showed a very intense
290 band around 1000/cm. These results indicated that the protein component around the crystal was
291 mainly due to these two amino acids, present in free form or in small peptides.

292 Other crystalline structures were randomly detected within the cheese and analysed by means of
293 micro Raman spectroscopy. Only calcium carbonate crystals were identified so far. They appeared
294 spheroidal in shape and translucent: the peaks at 1083, 1410, 713 and 284/cm confirmed their
295 nature (Tlili et al., 2001) (data not shown). The presence of calcium carbonate in cheese seems to be
296 attributable to the microbial metabolism that produces CO₂ (Gaucheron et al. 1999).

297 DISCUSSION

298 Speck characterization

299 By evaluating the confocal micro Raman spectrum and FAA composition of specks we obtained a
300 tyrosine purity >95% confirming previous reports indicating the presence of this amino acid in
301 these structures (Bottazzi et al., 1994; Tansman et al., 2015). Free tyrosine concentration increases
302 in the whole cheese throughout the ripening process, like for all other FAA (Pellegrino et al., 1997).
303 In our samples of Grana Padano and Parmigiano-Reggiano, free tyrosine concentration was
304 approximately 0.8 g / 100 g of water phase (Figure 6), i.e. about ten times higher than its water
305 solubility at room temperature (Grosse Daldrup et al., 2010). Therefore, formation of crystals

306 spread within the cheese would suggest that progress of proteolysis is not homogeneous in the
307 matrix thus leading to FAA accumulation, including tyrosine, preferentially into micro openings
308 until saturation. Furthermore, we have obtained by light microscopy further structural details of
309 tyrosine crystals from cheese (Figure 2), showing that the former reported description (Tansman et
310 al., 2015) is due to micro spike of the crystalized amino acid. When observed by atomic force
311 microscopy in protein hydrolysates (McPherson et al., 2012), tyrosine crystals appeared to be
312 covered by a stable layer of 3-nm particles, likely represented by micellar arrangement of small
313 peptides present in the medium and able to prevent re-solubilisation of crystals once formed.
314 Considered the remarkable content of peptides in ripened cheeses, this aspect would be worthy of
315 investigation.

316 **Spot characterization**

317 Contrary to the hard specks, spots were easily resin embedded and cut into thin sections that
318 allowed a deeper characterization by various microscopy techniques. Previous studies on spots in
319 cheese did not investigate their structure and ultrastructure (Giolitti and Mascherpa, 1970; Bianchi
320 et al. 1974; Tansman et al., 2015). Unexpectedly, when the cheese was directly stained with
321 ninhydrin and observed by light microscopy, the original grains of curd were still visible. During
322 cheese manufacturing, curd grains (20-50 mm), obtained by cutting the rennet gel, are let in hot
323 whey for 50-60 min to settle and aggregate at the bottom of the vat. Further fusing and shrinking of
324 curd grains are induced by the subsequent acidification of the cheese loaf when kept in mould for
325 48 hours. Since Grana Padano and Parmigiano-Reggiano loaves are not pressed, high temperature
326 and low pH (5.2-5.3) play the major role in promoting whey draining (Pellegrino et al., 1997) and
327 the tight aggregation of curd grains. However, as said above, these grains appeared as separate units
328 even several months later. The junctions between contiguous grains are low in fat because many fat
329 globules escape the protein network at the surface of the grains before they stick together. For those
330 grains having either irregular shape or different size their aggregation during settling forms an
331 internal hole, connected with several openings and radial junctions. This pattern was frequently

332 observed within the spots (Figure 3b). In the young cheese, even before the spot formation, these
333 hollow cavities likely represent preferential zones where whey stagnates and entrapped bacterial
334 cells find substrates for growth (Le Boucher et al., 2016). Growth of bacterial colonies as affected
335 by local concentration of substrates in cheese is receiving increasing attention (Silva et al., 2013;
336 Jeanson et al., 2015). Although no intact bacterial cells were detected by TEM in cavities within the
337 spot, as well as in the surrounding cheese, likely due to the prolonged ripening process, an indirect,
338 preliminary confirmation of this hypothesis came out by measuring the amount of total DNA
339 extracted from the spot and the surrounding cheese following the procedure of (Cremonesi et al.,
340 2007). In fact, DNA amount was more than three times higher in the spots than in the surrounding
341 cheese (72.94 μg vs 22.41 $\mu\text{g}/\text{mg}$).

342 Spots proved to be significantly more dry, with respect to the surrounding cheese portion, and to
343 contain more protein and less ash (Table 1), confirming the data reported by Tansman et al. (2015).
344 In addition, we showed the cheese portion just around the spot had the same chemical composition
345 as the rest of the cheese, confirming that the spot is a fully isolated unit. The dense layer we
346 observed around it by light microscopy is likely responsible for this (Figure 3, arrowhead). Overall,
347 these findings point to a migration of whey, with consequent draining of solutes, occurring locally
348 within the cheese where the spot would originate later.

349 Spots were also reported to have a different FAA composition compared to the whole cheese.
350 Bianchi et al. (1974) found spots in 18- and 25-month ripened Grana Padano cheeses to contain
351 (g/100 g): leucine (9.86), isoleucine (4.96), methionine (1.64), valine (1.52), glutamic acid (1.84),
352 and asparagine (1.63), as dominant FAA. However, these authors did not notice that the last two
353 FAA were equally abundant in the whole cheese. Recently, Tansman et al. (2015) detected only 1%
354 of free leucine in spots from a 24-month old Parmigiano-Reggiano cheese **but they did not give**
355 **explanation for the discrepancy of this result with those of literature.** We obtained FAA data similar
356 to those of Bianchi et al. (1974), however we put forward a different interpretation. The key to
357 understand the genesis of the spot relays in the chemical properties of the different FAA present in

358 the spot and in the cheese regardless of their concentration. In fact, the six FAA we detected at
359 much higher levels in the spot (i.e. leucine, isoleucine, methionine, valine, phenylalanine and
360 tyrosine) (Figure 6) are all hydrophobic because of either the aromatic or branched chain structure,
361 and thus insoluble in the water phase of cheese. This shared FAA characteristic leads us to
362 hypothesise that local water movements are responsible for their different distribution and,
363 consequently, for the spot blowing in hard and extra hard cheeses. This hypothesis was confirmed
364 by our experiment mimicking the migration of FAA within the cheese.

365 As already mentioned, in young cheese the cavities in which the residual whey stagnates could be
366 the sites where bacterial colonies develop. During cheese ripening, the water phase in these sites
367 becomes more concentrated in solutes due to lactic acid bacteria metabolism and, at a later stage, to
368 cell lysis. Progressively, the water phase migrates through the curd grain junctions, dragging solutes
369 including, preferentially, polar FAA over the hydrophobic ones. Consequently, less water soluble
370 molecules concentrate in a restricted area, that evolves into a spot, from which water moves away
371 radially (Figure 7) dragging away also salts. For this reason, many crystals were visible in the
372 junctions (Figure 4c).

373 **Microscopic crystal characterization**

374 Microcrystals appeared to be spread within the whole cheese. Among these crystals we have up to
375 now identified calcium phosphate by micro Raman, directly on the cheese, without any sample
376 preparation. The central star shaped crystals observed by TEM (Figure 8c) were clearly assigned to
377 selected types of calcium phosphate. Gaucheron et al. (1999) demonstrated that the supersaturation
378 of calcium phosphate salts increases strongly during cheese ripening due to the rise of pH,
379 explaining the salt precipitation. Unfortunately, the few references that discuss crystals
380 ultrastructure in cheese give no indication about the nature of the calcium phosphate. In our study,
381 the micro Raman analysis produced sufficient information about their morphological features and
382 complex ultrastructure. In fact, this technique allowed to evidence for the first time that the
383 fibrillary layer surrounding the central crystal contains leucine and, to a lower amount,

384 phenylalanine. These hydrophobic amino acids are reported to limit growth of calcium phosphate
385 crystals due to their absorption on the crystal surface that blocks the active growth sites (Dalas et
386 al., 2008). Although calcium lactate could form in cheese, due to lactic acid fermentation, the
387 chelation of calcium by the phosphates should greatly prevail (Heertje et al., 1981), in agreement
388 with the relevant number of calcium phosphate crystals we detected. Arkwall et al. (2006) reported
389 that calcium lactate crystals only form when pH is higher than 5.1. We could not find calcium
390 lactate crystals, although we cannot exclude their presence in extra hard cheeses, as different
391 unidentified crystalline structures were observed by TEM. Some researcher detected calcium lactate
392 crystals in cheeses other than Grana Padano and Parmigiano-Reggiano (Washam et al., 1982;
393 Tansman et al., 2014), whereas only Bottazzi et al. (1982) detected them in 14-month ripened Grana
394 cheese indicating their low incidence, i.e. 2-3 per square mm.

395 Presence of leucine crystals is likely within the spot where free leucine showed a concentration in
396 the water phase ten times higher than in the cheese (Figure 6). Crystalline leucine was also detected
397 by Tansman et al. (2015) in pearls collected from Parmigiano-Reggiano cheese.

398 CONCLUSIONS

399 In conclusion, combining the background knowledge about extra-hard cheese manufacturing and
400 ripening with the new information about composition and structure of spots achieved in this study,
401 we can hypothesize an incomplete aggregation of curd granules and a consequent local whey
402 stagnation to be at the origin of spot development. Growth and lysis of bacterial cells entrapped in
403 these micro cavities influence both metabolite availability and micro environmental conditions, that
404 in turn regulate diffusion and crystallization of solutes locally. Therefore, all practices finalized to
405 the syneresis improvement, such as fine-tuning of temperature control and curd grain size, would
406 greatly reduce spot number and the occurrence of other confined phenomena.

407 REFERENCES

- 408 Agarwal, S., J. R. Powers, B. G. Swanson, S. Chen, and S. Clark. 2006. Cheese pH, protein
409 concentration, and formation of calcium lactate crystals. *J. Dairy Sci.* 89: 4144-4155.
- 410
- 411 Agarwal, S., K. Sharma, B. G. Swanson, G. U. Yüksel, and S. Clark. 2006. Nonstarter lactic acid
412 bacteria biofilms and calcium lactate crystals in Cheddar cheese. *J. Dairy Sci.* 89: 1452–
413 1466.
- 414
- 415 Babcock, S. M., L. H. Russel, A. Vivian, and U. S. Baer. 1903. Condition affecting the
416 development of white specks in cold-curd cheese. Pages 180-183 in *Wis. Agric. Expt. Sta.*
417 19th Annu. Rep.
- 418
- 419 Bianchi, A.; G. Beretta, G. Caserio, and G. Giolitti. 1974. Amino acid composition of granules and
420 spots in Grana Padano cheeses. *J. Dairy Sci.* 57: 1504–1508.
- 421
- 422 Bottazzi, V., B. Battistotti, and F. Bianchi. 1982. The microscopic crystalline inclusions in Grana
423 cheese and their x-ray microanalysis. *Milchwissenschaft Milk Sci. Internat.* 37: 577–580.
- 424
- 425 Bottazzi, V., F. Lucchini, A. Rebecchi, and G. L. Scolari. 1994. I cristalli del formaggio grana
426 (Crystals present in Grana cheese). *Sci. Tecn. Latt.-Cas.* 45: 7–14.
- 427
- 428 Cattaneo, S., J. A. Hogenboom, F. Masotti, V. Rosi, L. Pellegrino, and P. Resmini. 2008. Grated
429 Grana Padano cheese: new hints on how to control quality and recognize imitations. *Dairy*
430 *Sci. Technol.* 88: 595-605.

431

432 Codex General Standard for Cheese. Codex Standard 283-1978.

433

434 Cremonesi, P., G. Perez, G. Pisoni, P. Moroni, S. Morandi, M. Luzzana, M. Brasca, and B.
435 Castiglioni. 2007. Detection of enterotoxigenic *Staphylococcus aureus* isolates in raw milk
436 cheese. *Lett. Appl. Microbiol.* 45: 586–591.

437

438 Culka, A., J. Jehlicka, and H. G. M. Edwards. 2010. Acquisition of Raman spectra of amino acids
439 using portable instruments: Outdoor measurements and comparison. *Spectrochim. Acta A.*
440 77: 978–983.

441

442 Dalas, E., P. Malkaj, Z. Vasileiou, and D. G. Kanellopoulou. 2008. The effect of Leucine on the
443 crystal growth of calcium phosphate. *J. Mater. Sci.: Mater. Med.* 19: 277-282.

444

445 D’Incecco, P., F. Faoro, T. Silvetti, K. Schrader, and L. Pellegrino. 2015. Mechanisms of
446 *Clostridium tyrobutyricum* removal through natural creaming of milk: A microscopy study.
447 *J. Dairy Sci.* 98: 5164-5172.

448

449 Gatti, M., B. Bottari, C. Lazzi, E. Neviani, and G. Mucchetti. 2014. Invited review: Microbial
450 evolution in raw-milk, long-ripened cheeses produced using undefined natural whey
451 starters. *J. Dairy Sci.* 97: 573-591.

452

- 453 Gaucheron, F., Y. Le Graet, F. Michel, V. Briard and M. Piot. Evolution of various salt
454 concentrations in the moisture and in the outer layer and centre of a model cheese during its
455 brining and storage in ammoniacal atmosphere. *Lait*. 79: 553-566.
- 456
- 457 Giolitti, G. and G. F. Mascherpa. 1970. La formazione di depositi di tirosina in formaggi a lunga
458 stagionatura. *Ind. Latte*. 6: 83-85.
- 459
- 460 Grosse Daldrup, J. B., C. Held, F. Ruether, G. Schembecker, and G. Sadowski. 2009. Measurement
461 and modeling solubility of aqueous multisolute amino-acid solutions. *Ind. Eng. Chem.* 49:
462 1395–1401.
- 463
- 464 Heertje, I., M. J. Boskamp, F. Van Kleef, and F. H. Gortemaker. 1981. The microstructure of
465 processed cheese. *Neth. Milk Dairy J.* 35, 177.
- 466
- 467 Jeanson, S., J. Floury, V. Gagnaire, S. Lortal, and A. Thierry. 2015. Bacterial colonies in solid
468 media and foods: a review on their growth and interactions with the micro-environment.
469 *Front. Microbiol.* 6: 1284.
- 470
- 471 Jenkins, A. L., R. A. Larsen, and T. B. Williams. 2005. Characterization of amino acids using
472 Raman spectroscopy. *Spectrochim. Acta A.* 61: 1585–1594.
- 473
- 474 Kaláb, M., J. Yun, and S. H. Yiu. 1987. Textural properties and microstructure of process cheese
475 food rework. *Food Microbiol.* 6: 181-192.

476

477 Koutsopoulos, S. 2002. Synthesis and characterization of hydroxyapatite crystals: a review study on
478 the analytical methods. *J. Biomed. Mater. Res.*, 62: 600-612.

479

480 Le Boucher, C., V. Gagnaire, V. Briard-Bion, J. Jardin, M. B. Maillard, G. Dervilly-Pinel, B. Le
481 Bizec, S. Lortal, S. Jeanson, and A. Thierry. 2016. Spatial distribution of *Lactococcus lactis*
482 colonies modulates the production of major metabolites during the ripening of a model
483 cheese. *Appl. Environ Microbiol.* 82: 202–210.

484

485 Marcos, A. 1993. Water activity in cheese in relation to composition, stability and safety. Pages
486 439-469 in *Cheese: Chemistry, Physics and Microbiology*, Vol. 1, ed.; Fox, P. F.; Chapman
487 & Hall, London.

488

489 Masotti, F., J. A. Hogenboom, V. Rosi, I. De Noni, and L. Pellegrino. 2010. Proteolysis indices
490 related to cheese ripening and typicalness in PDO Grana Padano cheese. *Int. Dairy J.* 20:
491 352-359.

492

493 McPherson, A., S. B. Larson, and Y. G. Kuznetsov. 2012. Tyrosine Microcrystals Produced by
494 Digestion of Proteins with Pancreatic Enzymes. *Crystal Growth & Design.* 12: 3594-3602.

495

496 McSweeney, P. L. H. 2004. Biochemistry of cheese ripening. *Int. J. Dairy Technol.* 57:127–144.

497

498 Ong, L., R. R. Dagastine, S. E. Kentish, and S. L. Gras. 2010. Transmission electron microscopy
499 imaging of the microstructure of milk in cheddar cheese production under different
500 processing conditions. *J. Food Sci.* 75: 135–145.

501

502 Pellegrino, L. G. Battelli, P. Resmini, P. Ferranti, F. Barone, and F. Addeo. 1997. Effects of heat
503 load gradient occurring in moulding on characterization and ripening of Grana Padano. *Lait.*
504 77: 217-228.

505

506 Sauer, G. R., W. B. Zunic, J. R. Durig, and R. E. Wuthier. 1994. Fourier transform Raman
507 spectroscopy of synthetic and biological calcium phosphates. *Calcif. Tissue Int.* 54: 414-
508 420.

509

510 Schrader, K. 2012. Structural analysis of milk and milk products. *Strukturanalyse von Milch und*
511 *Milchprodukten.* 133: 30-34.

512

513 Shock, A. A., W. J. Harper, A. M. Swanson, and H. H. Sommer. 1948. What's in those "white
514 specks" on Cheddar. *Wisconsin Agric. Experiment. Station. Bull.* 474: 31–32.

515

516 Silva, J. V. C., P. D. S. Peixoto, S. Lortal, and J. Floury. 2013. Transport phenomena in a model
517 cheese: the influence of the charge and shape of solutes on diffusion. *J. Dairy Sci.* 96: 6186–
518 6198.

519

- 520 Tansman, G., P. S. Kindstedt, and J. M. Hughes. 2014. Powder x-ray diffraction can differentiate
521 between enantiomeric variants of calcium lactate pentahydrate crystal in cheese. *J. Dairy*
522 *Sci.* 97: 7354–7362.
- 523
- 524 Tansman, G., P. S. Kindstedt, and J. M. Hughes. 2015. Crystal fingerprinting: elucidating the
525 crystals of Cheddar, Parmigiano-Reggiano, Gouda, and soft washed-rind cheeses using
526 powder x-ray diffractometry. *Dairy Sci. Technol.* 95: 651–664.
- 527
- 528 Tlili, M. M., M. Ben Amor, C. Gabrielli, S. Joiret, G. Maurin, and P. Rousseau. 2001.
529 Characterization of CaCO₃ hydrates by micro-Raman Spectroscopy. *J. Raman Spectrosc.*
530 33: 10–16.
- 531
- 532 Tuckey, S. L., H. A. Ruehe, and G. L. Clark. 1938. X-ray diffraction analysis of white specks in
533 Cheddar cheese. *J. Dairy Sci.* 21: 161.
- 534
- 535 Washam, C. J., T. J. Kerr, V. J. Hurst, and W. E. Rigsby. 1985. A scanning electron microscopy
536 study of crystalline structures on commercial cheese. *Dev. Ind. Microbiol.* 26: 749–761.

537

TABLES

538 **Table 1. Chemical composition of spot, cheese portion around it and whole cheese.***

	MOISTURE	FAT	PROTEIN	ASH
SPOT	24.2 ^a ± 1.2	27.5 ^a ± 2.6	37.6 ^a ± 1.3	3.9 ^a ± 0.1
CHEESE AROUND THE SPOT	29.5 ^b ± 1.0	29.6 ^a ± 1.2	33.5 ^b ± 0.6	4.4 ^b ± 0.3
WHOLE CHEESE	29.8 ^b ± 1.0	29.9 ^a ± 1.1	33.5 ^b ± 0.2	4.4 ^b ± 0.2

539

540 * Data were mean ± standard deviation based on duplicate analyses of five cheeses.

541 ^{a, b} Means in the same column with different letters are significantly different (p= 0.05).

542

For Peer Review

543

FIGURE CAPTIONS

544 **Figure 1.** Photograph of extra hard cheese showing speaks (SK) and spots (ST). (CS) cheese
545 portion surrounding a spot, sampling mode. Inset, isolated spots from the cheese.

546

547 **Figure 2.** Free amino acid analysis and (inset) light microscopy of a speck isolated from cheese.
548 Bar=200 μm .

549

550 **Figure 3.** Hand-made section of Grana Padano cheese including a spot: light microscopy after
551 ninhydrin staining revealed a dense layer limiting the spot (arrowhead) and the junctions among
552 curd granules (arrows). Bar=200 μm .

553

554 **Figure 4.** Light microscopy of spot semi-thin section (2-5 μm) stained with toluidine blue. (a) the
555 curd junctions are visible as darker lines; bar=200 μm . (b): Detail of the large hole in which
556 junctions converge; bar=50 μm . (c) Microcrystals (arrows) along a curd junction; bar=10 μm .

557

558 **Figure 5.** Ultrathin section of spot ultrastructure (a) in comparison with the surrounding cheese (b):
559 the network of electron dense proteins shows a more irregular profile in the spot, particularly at the
560 interface with the electron transparent fat matrix; bar=1 μm .

561

562 **Figure 6.** Free amino acid content of the spot and the cheese portion around it. Data were mean of
563 duplicate analyses of eleven cheeses. (*) Values statistically different ($p < 0.05$).

564

565 **Figure 7.** Free amino acid content of the portions 1-3 taken from cheese after the diffusion of 0.3
566 mL aqueous amino acid solution injected in 1 (inset). Data were mean of duplicate analyses.

567

568 **Figure 8.** Calcium phosphate crystals in cheese. (a) Fluorescence microscopy of crystals stained
569 with rodhamine B. (b) Video confocal microscopy of crystals stained with fast green and observed
570 with TRITC and FITC filters, revealing a non-fluorescent core. (c) Ultrathin section of a crystal
571 showing a star shaped zone in the center; at the interface with protein matrix, the fibrillar structure
572 of the crystal is visible (d).

573

574 **Figure 9.** Raman spectrum of calcium phosphate crystal in cheese. Stars represent typical signals of
575 the salt.

576

577 **Figure 10.** Raman spectra (a) and images (b) of spatial distribution of lipid/protein components
578 (red/green areas) around the crystal (covered by the blue area). High, medium and very low
579 component concentrations are indicated as red, green and blue respectively.

580

1 **FIGURE CAPTIONS (printed and on-line version)**

2 **Figure 1.** Photograph of extra hard cheese showing speaks (SK) and spots (ST). (CS) cheese
3 portion surrounding a spot, sampling mode. Inset, isolated spots from the cheese.

4

5 **Figure 2.** Free amino acid analysis and (inset) light microscopy of a speck isolated from cheese.
6 Bar=200 μm .

7

8 **Figure 3.** Hand-made section of Grana Padano cheese including a spot: light microscopy after
9 ninhydrin staining revealed a dense layer limiting the spot (arrowhead) and the junctions among
10 curd granules (arrows). Bar=200 μm .

11

12 **Figure 4.** Light microscopy of spot semi-thin section (2-5 μm) stained with toluidine blue. (a) the
13 curd junctions are visible as darker lines; bar=200 μm . (b): Detail of the large hole in which
14 junctions converge; bar=50 μm . (c) Microcrystals (arrows) along a curd junction; bar=10 μm .

15

16 **Figure 5.** Ultrathin section of spot ultrastructure (a) in comparison with the surrounding cheese (b):
17 the network of electron dense (grey) proteins shows a more irregular profile in the spot, particularly
18 at the interface with the electron transparent (white) fat matrix; bar=1 μm .

19

20 **Figure 6.** Free amino acid content of the spot and the cheese portion around it. Data were mean of
21 duplicate analyses of eleven cheeses. (*) Values statistically different ($p < 0.05$).

22

23 **Figure 7.** Free amino acid content of the portions 1-3 taken from cheese after the diffusion of 0.3
24 mL aqueous amino acid solution injected in 1 (inset). Data were mean of duplicate analyses.

25

26 **Figure 8.** Calcium phosphate crystals in cheese. (a) Fluorescence microscopy of crystals stained
27 with rodhamine B. (b) Video confocal microscopy of crystals stained with fast green and observed
28 with TRITC and FITC filters, revealing a non-fluorescent core. (c) Ultrathin section of a crystal
29 showing a star shaped zone in the center; at the interface with protein matrix, the fibrillar structure
30 of the crystal is visible (d).

31


32 **Figure 9.** Raman spectrum of calcium phosphate crystal in cheese. Stars represent typical signals of
33 the salt.

34

35 **Figure 10.** Raman spectra (a) and images (b) of spatial distribution of lipid/protein components
36 around the crystal (central dark grey area).


37

38




D’Incecco Figure 1.

For Peer Review




D'Incecco Figure 2.

For Peer Review




D’Incecco Figure 3.

For Peer Review




D'Incecco Figure 4.




D’Incecco Figure 5.


Peer Review




D'Incecco Figure 6.




D'Incecco Figure 7.



D'Incecco Figure 8.



D'Incecco Figure 9.



D'Incecco Figure 10.