



## POOR EARLY GROWTH AND HIGH SALT INTAKE IN INDIAN INFANTS

Journal:	<i>International Journal of Food Sciences and Nutrition</i>
Manuscript ID	CIJF-2016-0672.R1
Manuscript Type:	Research Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Genovesi, Simonetta; University of Milano-Bicocca , Department of Medicine and Surgery; Azienda Ospedaliera San Gerardo Antolini, Laura; University of Milano-Bicocca, Department of Medicine and Surgery, Centre of Biostatistics for Clinical Epidemiology Orlando, Antonina; University of Milano-Bicocca Brahmocharry Mandal , Sujit; Institute for Indian Mother and Child De Servi, Alessandra; Project for People Capelli, Silvia; Project for People Giussani, Marco; Family Paediatrician Nava, Elisa; University of Milano-Bicocca Agostoni, Carlo; University of Milan, Clinical Sciences and Community Health; IRCCS Ca' Granda, Ospedale Maggiore Policlinico, Pediatric Clinic 2 Gallieni, Maurizio; ASST Santi Paolo e Carlo, University of Milano, Nephrology and Dialysis Unit
Keywords:	Salt, India, Growth, hypertension, Infants, Nutrition

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60**1 POOR EARLY GROWTH AND HIGH SALT INTAKE IN INDIAN INFANTS**

2 Genovesi Simonetta<sup>a,b</sup>, Antolini Laura<sup>c</sup>, Orlando Antonina<sup>a</sup>, Brahmochary Sujit<sup>d</sup>, De Servi  
3 Alessandra<sup>e</sup>, Capelli Silvia<sup>e</sup>, Giussani Marco<sup>f</sup>, Nava Elisa<sup>a</sup>, Agostoni Carlo<sup>g</sup>, Gallieni Maurizio<sup>e,h</sup>.

4  
5  
6 Affiliations

7 <sup>a</sup> Department of Medicine and Surgery, University of Milano-Bicocca and Nephrology Unit, San  
8 Gerardo Hospital, Monza, Italy

9 <sup>b</sup> Department of Cardiovascular, Neural and Metabolic Sciences, Istituto Auxologico Italiano,  
10 Milan, Italy

11 <sup>c</sup> Department of Medicine and Surgery, Centre of Biostatistics for Clinical Epidemiology,  
12 University of Milano-Bicocca, Monza, Italy

13 <sup>d</sup> Institute for Indian Mother and Child (IIMC), Kolkata, India

14 <sup>e</sup> Project for People, Milan, Italy

15 <sup>f</sup> Family Paediatrician, Milano, Italy

16 <sup>g</sup> Pediatric Clinic, Department of Clinical Sciences and Community Health, University of Milan,  
17 Fondazione IRCCS Ca' Granda - Ospedale Maggiore Policlinico, Milan, Italy

18 <sup>h</sup> Nephrology and Dialysis Unit, Ospedale San Carlo Borromeo, ASST Santi Paolo e Carlo,  
19 Department of Biomedical and Clinical Sciences 'L. Sacco', University of Milan, Italy

20  
21 Email Addresses: Simonetta Genovesi [simonetta.genovesi@unimib.it](mailto:simonetta.genovesi@unimib.it); Laura Antolini  
22 [laura.antolini@unimib.it](mailto:laura.antolini@unimib.it); Sujit Brahmochary [iimcmmission@gmail.com](mailto:iimcmmission@gmail.com); Alessandra De Servi  
23 [alessandra\\_deservi@alice.it](mailto:alessandra_deservi@alice.it); Silvia Capelli [silvi.capelli@yahoo.it](mailto:silvi.capelli@yahoo.it); Marco Giussani [abrjg@tin.it](mailto:abrjg@tin.it);  
24 Antonina Orlando [antonina.orlando@unimib.it](mailto:antonina.orlando@unimib.it); Elisa Nava [elisa.nava4@virgilio.it](mailto:elisa.nava4@virgilio.it);  
25 Carlo Agostoni [carlo.agostoni@unimi.it](mailto:carlo.agostoni@unimi.it); Maurizio Gallieni [maurizio.gallieni@fastwebnet.it](mailto:maurizio.gallieni@fastwebnet.it)

26  
27 **Corresponding author:** Dr. Maurizio Gallieni

28 Nephrology and Dialysis Unit, Ospedale San Carlo Borromeo, ASST Santi Paolo e Carlo,  
29 Univeristy of Milano

30 Via Pio II, 3 – 20153 Milano, Italy

31 tel: +39 0240222445 – ORCID 0000-0002-2011-2160

32 Email: [maurizio.gallieni@fastwebnet.it](mailto:maurizio.gallieni@fastwebnet.it)

33  
34  
35 **Keywords:** Infant, Growth, Nutrition, Hypertension, Salt, India.

1  
2  
3 37 **ABSTRACT**

4 38 The influence of feeding patterns on the growth of infants and how salt is included in the diet are  
5 39 unknown in the area of West Bengal, India. A cross-sectional study was carried on 517 infants  
6 40 (median age 6.5 months). Negative Z-scores were observed for all anthropometric parameters.  
7  
8 41 72.7% of infants aged 0-6 months received exclusive breastfeeding. In the 6-12 month-old group  
9 42 (n=235), 91.5% had salt added to foods. In a regression model adjusted for age, a low salt diet  
10 43 resulted a significant factor in increasing weight-for-length and BMI-for-age z-scores, with  
11 44 increments equal to 0.637 SD (p=0.037) and 0.650 SD (p=0.036), respectively.

12  
13  
14  
15  
16 45 In West Bengal infants showing poor growth, breastfeeding was associated with better  
17 46 anthropometric indexes, but early in life salt is added to their diet. Early life low weight coupled  
18 47 with high salt intake may be a risk factor for arterial hypertension in Indian children.  
19  
20  
21 48

## 49 INTRODUCTION

50 According to the World Health Organization, non-communicable chronic diseases, including  
51 hypertension and kidney disease, are the most relevant public health problems worldwide, also in  
52 developing countries (WHO/FAO, 2002). Recent studies from our group showed a high prevalence  
53 of hypertension within an Indian population living in West Bengal, both in adults (Gallieni et al.  
54 2013) and children (Genovesi et al. 2011). The causes of these findings remain uncertain,  
55 considering that the association with increased body weight, even if significant, is rather modest.  
56 Further studies were therefore undertaken through a collaboration of two Italian Universities, the  
57 Indian non-governmental organization (NGO) Institute for Indian Mother and Child (IIMC),  
58 operating in West Bengal, and the Italian NGO Project for People (Gallieni et al. 2014). IIMC  
59 health workers follow and support pregnant women in district 24 of South Parganas, a  
60 rural/suburban area south of the city of Kolkata, which allowed us to recruit mothers for the present  
61 study.

62 The impact of early nutrition in regulating the expression of chronic degenerative disease later in  
63 life has been receiving increasing attention in the last decades (Agostoni et al. 2013). The incidence  
64 and outcomes of several metabolic diseases, such as hypertension and cardiovascular disorders,  
65 have been associated with birth weight, growth and feeding patterns in early childhood (Wu et al.  
66 2009). In addition, low birth weight may be a marker of impaired nephrogenesis, which may be  
67 causally related to hypertension and chronic kidney disease (White et al. 2009). Epidemiological  
68 and clinical evidence indicates that salt intake is a key regulator of blood pressure in children  
69 (Cooper et al. 1980, Geleijnse et al. 1997). Strengthening the possible role of salt intake in infancy,  
70 several clinical studies (Whitten et al. 1980, Hofman et al. 1983, Pomeranz et al. 2002) and meta-  
71 analyses of controlled trials (He et al. 2006) established a clear beneficial effect of the reduction in  
72 salt intake on blood pressure in children.

73 The purpose of this cross-sectional study was to analyze the influence of feeding patterns, from  
74 lactation to weaning, on the growth of infants up to their first year of life. In addition, the timing of  
75 salt intake in their diets was also investigated. Such information could be useful in better  
76 understanding the possible link between salt intake in infants and the possible effects of this habit  
77 on the higher prevalence of hypertension in pediatric age observed in this geographic area (Gallieni  
78 et al. 2013, Genovesi et al. 2011).

## 80 METHODS

81 IIMC health care workers and international volunteers collected data on the health status of infants  
82 and mothers. We studied 517 babies from 11 centers of aforementioned area, from January to

1  
2  
3 83 March 2013. Their median age was 6.5 months (interquartile range = 3.1-9.2, range =1-12), and  
4 84 48.6% were females babies. A team composed of doctors, nurses, health workers and international  
5 85 medical students participated in the screening procedures. Visits and measurements were carried out  
6 86 in IIMC owned medical ambulatory structures or schools, adapted as day-clinics (Figure 1). The  
7 87 children nutritional status was assessed by a dietary questionnaire, administered to mothers by  
8 88 experienced nutritionist at the time of children visit, and consisting of two parts. The first part  
9 89 related to children demographic data (name, surname, gender and date of birth). The second part  
10 90 related to two periods of child feeding: lactation and complementary feeding. Information about  
11 91 lactation concerned breastfeeding and its duration and the possible intake of other types of milk  
12 92 (cow's milk or formula). Information about introduction of complementary foods concerned type  
13 93 and time of introduction. Specific questions about adding further salt to food and "adult food"  
14 94 intake were also provided. Anthropometric characteristics of the sample were assessed by  
15 95 measuring length and weight with an infantometer and mobile digital scale (SECA, Vogel & Halke  
16 96 GmbH, Hamburg, Germany).

17 97 No formal evaluation and approval of the study protocol was obtained by an ethics committee, for  
18 98 the following reasons: the study was conducted in a very poor suburban/rural area in Kolkata  
19 99 suburbs as part of the regular volunteer activities of IIMC in support of poor mothers and children  
20 100 without access to medical care; IIMC is a NGO institution without an internal ethics committee; the  
21 101 study was observational and no drugs or diagnostic tests were administered to the studied subjects.  
22 102 Nevertheless, mothers gave their consent for the study participation and the study was conducted  
23 103 following the principles and guidelines in the World Medical Association (2000) Declaration of  
24 104 Helsinki on medical research involving human subjects  
25 105 (<http://www.wma.net/en/30publications/10policies/b3/>). No human subject included in this study  
26 106 can be identified via the paper; data were fully anonymized.

27 107 The study population was partitioned into 3 age groups: 0 (1day)-3 completed months, 3 (3 months  
28 108 + 1 day)- 6 completed months, 6 (6 months +1 day)-12 completed months. The impact on growth of  
29 109 exclusive breastfeeding versus not exclusive breastfeeding was evaluated in the first two age  
30 110 groups. Babies aged 6-12 completed months were analyzed to evaluate the impact of solid food  
31 111 introduction.

32 112 Raw anthropometric measures were processed by calculating the gender-specific z-score of length-  
33 113 for-age, weight-for-length, body mass index (BMI)-for-age, and weight-for-age, according to the  
34 114 World Health Organization charts (de Onis and Yip, 1996). Calculating z-score converts  
35 115 anthropometric measurements in the number of standard deviations the observed value is far from  
36 116 the expected median value in the reference population. Z-score values were subsequently

1  
2  
3 117 categorized into <-3 standard deviations, within -3 and -2 standard deviations, within -2 and -1  
4 118 standard deviations for the definition of poor growth (WHO, 2009) as follows:  
5  
6 119 • severe, moderate and mild stunting when based on length-for-age,  
7  
8 120 • severe, moderate and mild wasting when based on weight-for-length or BMI-for age.  
9  
10 121 • severe, moderate and mild under-weight when based on weight-for-age.  
11  
12

### 13 *Statistical analysis*

14 124 Anthropometric measures and z-score distributions were described graphically by box-plot graphs  
15 125 and by the summary measures mean, standard deviation, median and the 0.25 and 0.75 quantiles.  
16 126 Distributions were compared across age groups by F test and Bonferroni correction for post-hoc  
17 127 pairwise comparisons. Percentages of z-score <-3, within -3 and -2, within -2 and -1 standard  
18 128 deviations were represented graphically by bar-plot graphs. Percentages of z-score <-1, as overall  
19 129 percentages of poor growth, were compared across age groups by Chi-square test. The dependence  
20 130 between the z-score distributions and exclusive vs non exclusive breastfeeding, salt consumption vs  
21 131 non consumption, adult food consumption vs non consumption were investigated in regression  
22 132 models adjusted for age.  
23  
24  
25  
26  
27  
28  
29

### 30 133

## 31 134 **RESULTS**

32 135 Table 1 describes anthropometrics characteristics according to age group, which are represented by  
33 136 box-plots in Figure 2. Length-for-age z-score showed, in all age groups, negative mean values with  
34 137 a mild, non-significant worsening trend over time. When considering standardized measures of  
35 138 weight with respect to the achieved length, namely weight-for-length z-score and the BMI-for-age  
36 139 z-score, negative mean values were observed in all age groups. For both, weight-for-length z-score  
37 140 and BMI-for-age z-score, there was a sensible difference across age groups with a tendency to  
38 141 improve growth from 0-3 months to 3-6 months. A subsequent flexion at 6-12 months was observed  
39 142 only for weight-for-length z-score. Negative averages were observed in all age groups for weight-  
40 143 for-age z-score.  
41  
42  
43  
44  
45  
46  
47

48 144 Table 2 describes the percentages of severe, moderate and mild poor growth showing that about  
49 145 50% of children aged 0-3 months showed poor growth for all parameters. For stunting (length-for-  
50 146 age z-score) and wasting (weight-for-length z-score) no significant differences were observed  
51 147 across age groups, whereas the percentage of moderate and severe BMI-for-age z-score alterations  
52 148 significantly decreased with increasing age. A similar trend was observed for under weight (weight-  
53 149 for-age z-score) from 0-3 months to 3-6 months. When considering only severe and moderate poor  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 150 growth, the overall percentages were about 20% for all parameters with the exception of wasting  
4 151 (weight-for-length z-score), which was more modest.

5  
6 152 The analysis of dietary habits showed that 72.7% of the 238 babies aged 0-6 months received  
7  
8 153 exclusive breastfeeding, a further 25.2% received mixed feeding, and thus only 2.1% artificial  
9  
10 154 feeding. When contrasting mixed breastfeeding versus exclusive breastfeeding a modest non-  
11  
12 155 significant tendency to increase was observed for all parameters (i.e. observed averages for weight-  
13  
14 156 for-length z-score were -0.69 and -0.62,  $p=0.676$ ), even when adjusting for age.

15 157 Considering the 6-12 months group ( $n=235$  babies), we observed that 91.5% ( $n=215$  babies) had  
16 158 salt added to foods and the majority of them ( $n=209/215$ , 97.2%) started at six completed months.  
17  
18 159 In a regression model adjusted for age, not having added salt resulted a significant factor in  
19  
20 160 increasing weight-for-length z-score and BMI-for-age z-score, with increments equal to 0.637 SD  
21  
22 161 ( $p=0.0373$ ) and 0.650 SD ( $p=0.0360$ ) respectively (Table 3). No significant relationships were  
23  
24 162 observed for length-for-age z-score and weight-for-age z-score.

25 163 Among infants fed adult foods, most ingested “kitchuri”, consisting of rice, lentils, vegetables,  
26 164 chicken or fish with added salt, cornmeal, sugar and fried oil. The percentage of infants fed adult  
27  
28 165 foods starting at 6 months of life was 79.2%. A regression model adjusted for age showed that  
29  
30 166 infants not assuming adult food had higher weight-for-length and BMI-for-age z-scores, with  
31  
32 167 increment equal to 0.560 SD ( $p= 0.0071$ ) and 0.589 SD ( $p= 0.0051$ ) (Table 4).

33 168 Further differences were not detectable in infants fed other types of solids, such as flour, biscuits,  
34  
35 169 vegetables, fruits, meat and fish.

36 170

## 37 38 171 **DISCUSSION**

39 172 In this population, there was a high prevalence of malnutrition, in particular in the first three months  
40  
41 173 of life, as suggested by anthropometric parameters. Unfortunately, the analysis could not be  
42  
43 174 adjusted for infants' birth weight since the delivery occurred at home where weighing the newborn  
44  
45 175 is still an unusual practice. However, considering the rates of malnutrition observed in the youngest  
46  
47 176 group, a high prevalence of low birth weight for gestational age may be hypothesized. A low  
48  
49 177 intrauterine growth would lead to increased insulin resistance, according to Barker hypothesis,  
50  
51 178 (Barker et al 2010). In addition, insulin increase is associated with higher blood pressure values also  
52  
53 179 in children (Genovesi et al. 2012). We found a high percentage of breastfeeding mothers and the  
54  
55 180 breastfeeding practice was associated with an improvement of all anthropometric indexes of infants,  
56  
57 181 underscoring the importance of encouraging breastfeeding even in low income countries. The  
58  
59 182 complementary feeding period shows some peculiar characteristics. A high percentage of infants  
60  
183 have salt added to foods quite early. Salt intake starting in the pediatric age is associated to a high

1  
2  
3 184 prevalence of arterial hypertension (Whitten et al. 1980, Hofman et al. 1983, Pomeranz et al. 2002,  
4 185 He et al. 2006, Strazzullo et al. 2012) and we speculate that this evidence could at least partly  
5 186 explain the high prevalence of hypertension observed in our previous investigations in the West  
6 187 Bengal region (Gallieni et al. 2013, Genovesi et al. 2011). Infants who received added salt showed  
7 188 also lower body weight measures standardized for height than those who did not. This association  
8 189 was observed even in those 6 to 12 month-old infants fed the kitchuri food. While the biologic  
9 190 plausibility of high salt intake and lower body weight is unclear, it has already been observed both  
10 191 experimentally (Coelho et al. 2006) and clinically in children (Stein et al. 2006), suggesting some  
11 192 hormone-mediated mechanisms. This unusual salt consumption coupled with early low weight  
12 193 conditions may contribute to the early development of arterial hypertension (Calkins et al. 2011).  
13 194 However, because the results of this study do not include a long term evaluation of blood pressure,  
14 195 the effects of increased salt intake during infancy on the prevalence of hypertension in Indian  
15 196 children remain hypothetical.

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

197 In conclusion, the nutritional pattern of West Bengal babies can be considered adequate only in the  
198 0-6 months period, when exclusive breastfeeding is common. Breastfeeding practice prevents the  
199 development of infections and dysentery causing underweight or mortality, and is therefore  
200 particularly important. However, the complementary feeding period in this population ensues in a  
201 critical time, because the early intake of added salt and adult food may have a negative effect on  
202 growth and later health, if coupled with the major risk of unfavorable programming associated with  
203 intrauterine growth retardation (Keijzer et al. 2005). The results of this study may be used for  
204 informing mothers of the importance of continuing breastfeeding as well as retarding the  
205 introduction of salt in their babies' diet.

#### 206 207 **Geolocation information.**

208 This study was undertaken in West Bengal, India

#### 209 210 **Acknowledgments**

211 Funding: this research was carried on without specific funding. The authors acknowledge all the  
212 Indian and International volunteers working at the Institute for Indian Mother and Child, Kolkata,  
213 India, who contributed to the collection of data.

#### 214 215 **Declaration of interests statement**

216 The authors report no conflicts of interest.



218 **References**

- 219 Agostoni C, Baselli L, Mazzoni MB. 2013. Early nutrition patterns and diseases of adulthood: A  
220 plausible link? *Eur J Intern Med.* 24:5-10.
- 221 Barker DJ, Gelow J, Thornburg K, Osmond C, Kajantie E, Eriksson JG. 2010. The early origins  
222 of chronic heart failure: impaired placental growth and initiation of insulin resistance in  
223 childhood. *Eur J Heart Failure.* 12: 819–825.
- 224 Calkins K, Devaskar S. 2011. Fetal origins of adult disease. *Curr Probl Pediatr Adolesc Health*  
225 *Care.* 41:158-176.
- 226 Coelho MS, Passadore MD, Gasparetti AL, Bibancos T, Prada PO, Furukawa LL, Furukawa  
227 LN, Fukui RT, Casarini DE, Saad MJ, et al. 2006. High- or low-salt diet from weaning to  
228 adulthood: effect on body weight, food intake and energy balance in rats. *Nutr Metab*  
229 *Cardiovasc Dis.* 16:148-155.
- 230 Cooper R, Soltero I, Liu K, Berkson D, Levinson S, Stamler J. 1980. The association between  
231 urinary sodium excretion and blood pressure in children. *Circulation.* 62:97–104.
- 232 de Onis M, Yip R. 1996. The WHO growth chart: historical considerations and current scientific  
233 issues. *Bibl Nutr Dieta.* 53:74-89.
- 234 He FJ, MacGregor GA. 2006. Importance of salt in determining blood pressure in children:  
235 meta-analysis of controlled trials. *Hypertension.* 48:861-869.
- 236 Hofman A, Hazebroek A, Valkenburg HA. 1983. A randomized trial of sodium intake and  
237 blood pressure in newborn infants. *JAMA.* 250:370–373.
- 238 Gallieni M, Ene-Iordache B, Aiello A, Tucci B, Sala V, Brahmochary Mandal SK, Doneda A,  
239 Stella A, Carminati S, Perico N, Genovesi S. 2013. Hypertension and kidney function in an  
240 adult population of West Bengal, India: role of body weight, waist circumference, proteinuria  
241 and rural area living. *Nephrology.* 18:798-807.
- 242 Gallieni M, Aiello A, Tucci B, Sala V, Brahmochary Mandal SK, Doneda A, Genovesi S. 2014  
243 The burden of hypertension and kidney disease in Northeast India: the Institute for Indian  
244 Mother and Child noncommunicable diseases project. *Scientific World Journal.* 2014:320869.
- 245 Genovesi S, Antolini L, Gallieni M, Aiello A, Mandal SK, Doneda A, Giussani M, Stella A,  
246 Tucci B, Valsecchi MG. 2011. High prevalence of hypertension in normal and underweight  
247 Indian children. *J Hypertens.* 29:217-221.
- 248 Genovesi S, Brambilla P, Giussani M, Galbiati S, Mastriani S, Pieruzzi F, Stella A, Valsecchi  
249 MG, Antolini L. 2012. Insulin resistance, pre-hypertension, hypertension and blood pressure  
250 values in paediatric age. *J Hypertension.* 30:327-335.

- 1  
2  
3 251 Geleijnse JM, Hofman A, Witteman JC, Hazebroek AA, Valkenburg HA, Grobbee DE 1997.  
4 252 Long-term effects of neonatal sodium restriction on blood pressure. *Hypertension*. 29:913–917.  
5 253 Keijzer-Veen MG, Schrevel M, Finken MJ, Dekker FW, Nauta J, Hille ET, Frölich M, van der  
6 254 Heijden BJ; Dutch POPS-19 Collaborative Study Group. 2005. Microalbuminuria and lower  
7 255 glomerular filtration rate at young adult age in subjects born very premature and after  
8 256 intrauterine growth retardation. *J Am Soc Nephrol*. 16:2762-2768.  
9 257 Pomeranz A, Dolfin T, Korzets Z, Eliakim A, Wolach B. 2002. Increased sodium  
10 258 concentrations in drinking water increase blood pressure in neonates. *J Hypertens*. 20:203–207.  
11 259 Stein LJ, Cowart BJ, Beauchamp GK. 2006. Salty taste acceptance by infants and young  
12 260 children is related to birth weight: longitudinal analysis of infants within the normal birth  
13 261 weight range. *Eur J Clin Nutr*. 60:272-279.  
14 262 Strazzullo P, Campanozzi A, Avallone S. 2012. Does salt intake in the first two years of life  
15 263 affect the development of cardiovascular disorders in adulthood? *Nutr Metab Cardiovasc Dis*.  
16 264 22:787-792.  
17 265 White SL, Perkovic V, Cass A, Chang CL, Poulter NR, Spector T, Haysom L, Craig JC, Salmi  
18 266 IA, Chadban SJ, Huxley RR. 2009. Is low birth weight an antecedent of CKD in later life? A  
19 267 systematic review of observational studies. *Am J Kidney Dis*. 54:248-261.  
20 268 Whitten CF, Stewart RA. 1980. The effect of dietary sodium in infancy on blood pressure and  
21 269 related factors. Studies of infants fed salted and unsalted diets for five months at eight months  
22 270 and eight years of age. *Acta Paediatr Scand Suppl*. 279:1–17.  
23 271 WHO/FAO (2002) Diet, nutrition and the prevention of chronic diseases: report of a joint  
24 272 WHO/FAO expert consultation. Available from:  
25 273 <http://www.who.int/dietphysicalactivity/publications/trs916/en/> (accessed June 2016).  
26 274 WHO (2009) Child Growth Standards and the Identification of Severe Acute Malnutrition in  
27 275 Infants and Children: A Joint Statement by the World Health Organization and the United  
28 276 Nations Children's Fund. 2009 Geneva. Available from:  
29 277 <http://www.ncbi.nlm.nih.gov/books/NBK200775/> (accessed June 2016).  
30 278 Wu TC, Chen PH. 2009. Health consequences of nutrition in childhood and early infancy.  
31 279 *Pediatr Neonatol*. 50:135-142.

280

281

282

283 Table 1. Descriptive statistics of anthropometric characteristics (raw and z-score) according to age  
 284 group  
 285

Continuous variable	0-3 months (n=124;24.0%)					3-6 months (n=131;25.3%)					6-12 months (n=262;50.7%)					p-value
	mean	SD	median	q25	q75	mean	SD	median	q25	q75	mean	SD	median	q25	q75	
Length (cm)	54.5	3.5	54.5	52.0	57.0	61.1	3.4	61.0	59.5	63.0	68.2	4.1	68.0	65.0	71.0	<0.0001
Weight (kg)	4.2	0.8	4.2	3.6	4.7	6.0	1.0	5.9	5.4	6.6	7.5	1.1	7.4	6.7	8.1	<0.0001
BMI (kg/m <sup>2</sup> )	13.8	1.6	13.9	12.5	15.1	16.0	1.6	16.0	14.9	16.8	16.0	1.8	15.8	14.9	16.9	<0.0001
Length for age (z-score)	-1.01	1.43	-1.02	-1.89	-0.16	-1.09	1.38	-1.14	-1.86	-0.29	-1.32	1.43	-1.21	-2.09	-0.52	0.0864
Weight for length (z-score) * ^	-0.88	1.24	-0.97	-1.54	-0.11	-0.43	1.13	-0.31	-1.21	0.27	-0.71	1.28	-0.79	-1.44	0.05	0.0115
BMI for age (z-score)*	-1.32	1.13	-1.18	-1.89	-0.66	-0.72	1.16	-0.55	-1.39	-0.04	-0.73	1.29	-0.81	-1.47	0.01	<0.001
Weight for age (z-score)*	-1.17	1.27	-1.09	-1.88	-0.48	-0.81	1.23	-0.77	-1.28	-0.02	-0.90	1.14	-0.88	-1.67	-0.11	0.0474

286  
 287  
 288 0-3 months = from 1 day to 3 completed months, 3-6 months = from 3 months + 1 day to 6  
 289 completed months, 6-12 months = from 6 months + 1 day to 12 completed months, BMI = body  
 290 mass index; SD=standard deviation; q25 = 0.25 quantile; q75 = 0.75 quantile; \* = significant post-  
 291 hoc comparison 0-3 months vs 3-6 months; ^ = significant post-hoc comparison 3-6 months vs 6-12  
 292 months.  
 293

294 Table 2. Growth status according to age group

295

Categorical variable	0-3 months (n=124;24.0%)						3-6 months (n=131;25.3%)						6-12 months (n=262;50.7%)						p value
	severe		moderate		mild		severe		moderate		mild		severe		moderate		mild		
	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	
Stunting (Length for age)	6	4.8	20	16.1	38	30.7	8	6.1	20	15.3	39	29.8	30	11.5	40	15.3	75	28.6	0.662
Wasting (Weight for length)	2	1.6	4	3.2	40	32.3	1	0.8	6	4.6	31	23.7	6	2.3	14	5.3	72	27.5	0.345
Wasting (BMI for age)*	14	11.3	13	10.5	46	37.1	3	2.3	12	9.2	36	27.5	10	3.8	26	9.9	73	27.9	0.002
Under weight (Weight for age)*	11	8.9	16	12.9	43	34.7	6	4.6	13	9.9	28	21.4	10	3.8	35	13.4	73	27.9	0.004

296

297

298 0-3 months = from 1 day to 3 completed months, 3-6 months = from 3 months + 1 day to 6  
 299 completed months, 6-12 months = from 6 months + 1 day to 12 completed months, severe = below -  
 300 3 SD, moderate = between -3 and -2 SD, mild = between -2 and -1 SD, BMI = body mass index;  
 301 SD=standard deviation; \* = significant post-hoc comparison 0-3 months vs 3-6 months.

302

303

304 Table 3. Effect of salt consumption on growth indexes (model adjusted for age, n=235, 6-12  
 305 months)  
 306

Variable	Intercept				Salt				Age (months)			
	<i>b</i>	(95% CI)	<i>p</i> -value	<i>b</i>	(95% CI)	<i>p</i> -value	<i>b</i>	(95% CI)	<i>p</i> -value			
Length for age (z-score)	-0.717	-1.718 0.284	0.162	0.271	-0.381 0.922	0.416	-0.087	-0.181 0.007	0.072			
Weight for length (z-score)	-0.217	-1.133 0.699	0.643	-0.637	-1.232 -0.041	0.037	0.007	-0.079 0.093	0.872			
BMI for age (z-score)	-0.710	-1.637 0.218	0.135	-0.650	-1.253 -0.046	0.036	0.060	-0.027 0.148	0.175			
Weight for age (z-score)	-0.467	-1.263 0.329	0.251	-0.427	-0.945 0.091	0.107	-0.002	-0.077 0.073	0.958			

307  
 308  
 309

Or Peer Review Only

310 Table 4. Effect of adult food consumption on growth indexes (model adjusted for age, n=235, 6-12  
 311 months).  
 312

Variable	Intercept				Adult food				Age (months)			
	<i>b</i>	(95% CI)		<i>p</i> -value	<i>b</i>	(95% CI)		<i>p</i> -value	<i>b</i>	(95% CI)		<i>p</i> -value
Length for age (z-score)	-0.757	-1.680	0.165	0.109	0.404	-0.038	0.846	0.075	-0.090	-0.183	0.003	0.059
Weight for length (z-score)	-0.342	-1.186	0.501	0.427	-0.560	-0.964	-0.156	0.007	0.006	-0.079	0.091	0.898
BMI for age (z-score)	-0.827	-1.681	0.026	0.058	-0.589	-0.998	-0.180	0.005	0.059	-0.027	0.145	0.177
Weight for age (z-score)	-0.649	-1.389	0.091	0.087	-0.202	-0.557	0.152	0.264	-0.007	-0.082	0.067	0.851

313  
 314  
 315  
 316

or Peer Review Only

1  
2  
3 317 **Figure legend**

4 318

5 319 **Figure 1. A room of an elementary school from a remote village turned into a day-clinic for**  
6 **data collection.** On the front-left the infantometer used for measurements of babies. On the right a  
7 320 group of mothers.  
8 321  
9 322

10  
11 323 **Figure 2. Graphical representation of anthropometric characteristics (z-scores) according to**  
12 **age group.** Boxplot explanation: upper horizontal line of box = 75th percentile; lower horizontal  
13 324 line of box = 25th percentile; horizontal bar within box = median, square within box = mean;  
14 325 vertical lines out of the box = minimum and maxim.  
15 326  
16 327

17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 1. Descriptive statistics of anthropometric characteristics (raw and z-score) according to age group

Continuous variable	0-3 months (n=124;24.0%)					3-6 months (n=131;25.3%)					6-12 months (n=262;50.7%)					
	mean	SD	median	q25	q75	mean	SD	median	q25	q75	mean	SD	median	q25	q75	p-value
Length (cm)	<b>54.5</b>	3.5	<b>54.5</b>	52.0	<b>57.0</b>	<b>61.1</b>	3.4	<b>61.0</b>	59.5	<b>63.0</b>	<b>68.2</b>	4.1	<b>68.0</b>	65.0	<b>71.0</b>	<0.0001
Weight (kg)	<b>4.2</b>	0.8	<b>4.2</b>	3.6	<b>4.7</b>	<b>6.0</b>	1.0	<b>5.9</b>	5.4	<b>6.6</b>	<b>7.5</b>	1.1	<b>7.4</b>	6.7	<b>8.1</b>	<0.0001
BMI (kg/m <sup>2</sup> )	<b>13.8</b>	1.6	<b>13.9</b>	12.5	<b>15.1</b>	<b>16.0</b>	1.6	<b>16.0</b>	14.9	<b>16.8</b>	<b>16.0</b>	1.8	<b>15.8</b>	14.9	<b>16.9</b>	<0.0001
Length for age (z-score)	<b>-1.01</b>	1.43	<b>-1.02</b>	-1.89	-0.16	<b>-1.09</b>	1.38	<b>-1.14</b>	-1.86	<b>-0.29</b>	<b>-1.32</b>	1.43	<b>-1.21</b>	-2.09	<b>-0.52</b>	0.0864
Weight for length (z-score) * ^	<b>-0.88</b>	1.24	<b>-0.97</b>	-1.54	-0.11	<b>-0.43</b>	1.13	<b>-0.31</b>	-1.21	<b>0.27</b>	<b>-0.71</b>	1.28	<b>-0.79</b>	-1.44	<b>0.05</b>	0.0115
BMI for age (z-score)*	<b>-1.32</b>	1.13	<b>-1.18</b>	-1.89	-0.66	<b>-0.72</b>	1.16	<b>-0.55</b>	-1.39	<b>-0.04</b>	<b>-0.73</b>	1.29	<b>-0.81</b>	-1.47	<b>0.01</b>	<0.001
Weight for age (z-score)*	<b>-1.17</b>	1.27	<b>-1.09</b>	-1.88	-0.48	<b>-0.81</b>	1.23	<b>-0.77</b>	-1.28	<b>-0.02</b>	<b>-0.90</b>	1.14	<b>-0.88</b>	-1.67	<b>-0.11</b>	0.0474

0-3 months = from 1 day to 3 completed months, 3-6 months = from 3 months + 1 day to 6 completed months, 6-12 months = from 6 months + 1 day to 12 completed months, BMI = body mass index; SD=standard deviation; q25 = 0.25 quantile; q75 = 0.75 quantile; \* = significant post-hoc comparison 0-3 months vs 3-6 months; ^ = significant post-hoc comparison 3-6 months vs 6-12 months.



Table 2. Growth status according to age group.

Categorical variable	0-3 months (n=124;24.0%)						3-6 months (n=131;25.3%)						6-12 months (n=262;50.7%)						p value
	severe		moderate		mild		severe		moderate		mild		severe		moderate		mild		
	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	
Stunting (Length for age)	6	4.8	20	16.1	38	30.7	8	6.1	20	15.3	39	29.8	30	11.5	40	15.3	75	28.6	0.662
Wasting (Weight for length)	2	1.6	4	3.2	40	32.3	1	0.8	6	4.6	31	23.7	6	2.3	14	5.3	72	27.5	0.345
Wasting (BMI for age)*	14	11.3	13	10.5	46	37.1	3	2.3	12	9.2	36	27.5	10	3.8	26	9.9	73	27.9	0.002
Under weight (Weight for age)*	11	8.9	16	12.9	43	34.7	6	4.6	13	9.9	28	21.4	10	3.8	35	13.4	73	27.9	0.004

0-3 months = from 1 day to 3 completed months, 3-6 months = from 3 months + 1 day to 6 completed months, 6-12 months = from 6 months + 1 day to 12 completed months, severe = below -3 SD, moderate = between -3 and -2 SD, mild = between -2 and -1 SD, BMI = body mass index; SD=standard deviation; \* = significant post-hoc comparison 0-3 months vs 3-6 months.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

Table 3. Effect of salt consumption on growth indexes( model adjusted for age, n=235, 6-12 months)

Variable	Intercept			Salt			Age (months)		
	<i>b</i>	(95% CI)	<i>p</i> -value	<i>b</i>	(95% CI)	<i>p</i> -value	<i>b</i>	(95% CI)	<i>p</i> -value
Length for age (z-score)	-0.717	-1.718 0.284	0.162	0.271	-0.381 0.922	0.416	-0.087	-0.181 0.007	0.072
Weight for length (z-score)	-0.217	-1.133 0.699	0.643	-0.637	-1.232 0.041	0.037	0.007	-0.079 0.093	0.872
BMI for age (z-score)	-0.710	-1.637 0.218	0.135	-0.650	-1.253 0.046	0.036	0.060	-0.027 0.148	0.175
Weight for age (z-score)	-0.467	-1.263 0.329	0.251	-0.427	-0.945 0.091	0.107	-0.002	-0.077 0.073	0.958

Table 4. Effect of adult food consumption on growth indexes. (model adjusted for age, n=235, 6-12 months).

Variable	Intercept				Adult food				Age (months)			
	<i>b</i>	(95% CI)		<i>p</i> -value	<i>b</i>	(95% CI)		<i>p</i> -value	<i>b</i>	(95% CI)		<i>p</i> -value
Length for age (z-score)	-0.757	-1.680	0.165	0.109	0.404	-0.038	0.846	0.075	-0.090	-0.183	0.003	0.059
Weight for length (z-score)	-0.342	-1.186	0.501	0.427	-0.560	-0.964	-0.156	0.007	0.006	-0.079	0.091	0.898
BMI for age (z-score)	-0.827	-1.681	0.026	0.058	-0.589	-0.998	-0.180	0.005	0.059	-0.027	0.145	0.177
Weight for age (z-score)	-0.649	-1.389	0.091	0.087	-0.202	-0.557	0.152	0.264	-0.007	-0.082	0.067	0.851

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Figure 1. A room of an elementary school from a remote village turned into a day-clinic for data collection. On the front-left the infantometer used for measurements of babies. On the right a group of mothers.

Figure 1  
245x164mm (300 x 300 DPI)

View Only




Figure 2. Graphical representation of anthropometric characteristics (z-scores) according to age group. Boxplot explanation: upper horizontal line of box = 75th percentile; lower horizontal line of box = 25th percentile; horizontal bar within box = median, square within box = mean; vertical lines out of the box = minimum and maximum.

Figure 2  
508x281mm (72 x 72 DPI)