

1 **Published in Veterinary Immunology and Immunophatology,**

2 **DOI: 10.1016/j.vetimm.2016.10.007**

3

4 **Original Article**

5

6 **In vitro assessment of the effects of temperature on phagocytosis, reactive oxygen species**
7 **production and apoptosis in bovine polymorphonuclear cells**

8

9

10 Cristina Lecchi^{*a}, Nicola Rota^b, Andrea Vitali^c, Fabrizio Ceciliani^a, Nicola Lacetera^c

11 ^aDipartimento di Medicina Veterinaria, Università di Milano, Milano, Italy

12 ^bDipartimento di Scienze Veterinarie per la Salute, la Produzione Animale e la Sicurezza
13 Alimentare, Università di Milano, Milano, Italy

14 ^cDipartimento di Scienze Agrarie e Forestali, Università della Tuscia, Viterbo, Italy

15

16 * Corresponding author. Tel.: +39 0250318100.

17 E-mail address: cristina.lecchi@unimi.it (C. Lecchi).

18

19 **Abstract**

20 Heat stress exerts a direct negative effect on farm animal health, triggering physiological
21 responses. Environmental high temperature induces immunosuppression in dairy cows, increasing
22 the risk of mastitis and milk somatic cell counts. The influence of heat stress on leukocytes
23 activities have not been fully elucidated. The present *in vitro* study was aimed at assessing whether
24 the exposure to temperature simulating conditions of severe whole body hyperthermia affects
25 defensive functions of bovine blood polymorphonuclear cells.

26 Blood was collected from seven clinically healthy, multiparous, late lactating Holstein cows.
27 After isolation, PMN were incubated at either 39 or 41 °C. Phagocytosis, respiratory burst and
28 apoptosis were then investigated. The selected temperatures of 39 °C or 41 °C mimicked conditions
29 of normothermia or severe heat stress, respectively.. Phagocytosis assay was carried out by
30 measuring the fluorescence of phagocytosed fluorescein-labelled *E. coli* bioparticles. The modulation
31 of oxidative burst activity was studied by the cytochrome C reduction method. Apoptosis was
32 determined by measuring the activities of two enzymes that play an effector role in the process,
33 namely Caspase-3 and Caspase-7. Statistical analyses were performed using SPSS 22.0. A Student
34 t-test for paired samples and a Generalised Estimating Equation were used based on data
35 distribution.

36 , The phagocytosis rate was reduced (-37%, $P < 0.01$) when PMN were incubated for 2 h at
37 41°C, when compared to phagocytosis rate measured at 39 °C. The oxidative burst, as determined
38 by extracellular production of reactive oxygen species (ROS), was also reduced by the exposure of
39 cells to 41°C compared to 39 °C. Such reduction ranged between -2 and -21% ($P < 0.05$). Apoptosis
40 rate was not affected by different temperatures.

41 The results reported in this study suggest that phagocytosis and ROS production in PMN
42 exposed to severe high temperature are impaired, partially explaining the higher occurrence of
43 infections during periods of hot weather.

44

45 *Key words:* bovine; polymorphonuclear cells; high temperature; apoptosis; ROS

46

47 **1. Introduction**

48 Heat stress (HS) induced immunosuppression is indicated as one of the mechanisms through
49 which climate changes, more specifically the increase of air temperature, are expected to exert a
50 direct negative effect on human and animal health in the next decades (Gaughan et al., 2009; Martin
51 et al., 2010; Dang et al., 2012).

52 Studies focusing on the relationships between environmental temperature and health
53 problems in dairy cows pointed out higher risks of mastitis (Giesecke 1985; Smith et al. 1985;
54 Waage et al. 1998; Olde Riekerink et al., 2007) and an increase of milk somatic cell counts
55 (Bertocchi et al., 2014) during periods of hot weather. The mechanisms behind the higher risk of
56 mastitis during periods of HS have not been elucidated although it a negative action of HS on
57 defensive mechanisms has been suggested as one of the most probable mechanism (Giesecke 1985;
58 Bertocchi et al., 2014).

59 The effect of HS on cattle immune functions (enhancement, suppression, or no effect)
60 depends on several factors, which include breed, genotype, age, acclimation level, intensity and
61 duration of the exposure to high temperatures, recovery opportunities, the specific immune
62 parameter taken into account, the experimental models adopted (*in vivo*, *ex vivo*, and *in vitro*) and
63 their interactions (Lacetera, 2012). Severe HS has been shown to impair immune functions in dairy
64 cows. In particular, both *ex vivo* and *in vitro* studies focusing on proliferation of mitogens
65 stimulated peripheral blood mononuclear cells (PBMC) pointed out that high temperatures alter the
66 ability of PBMC to proliferate (Lacetera et al., 2005, 2009). Furthermore, preliminary results from
67 ongoing *in vitro* studies indicate that incubation temperatures simulating whole body severe
68 hyperthermia (41 °C and above) impair the secretion of TNF-alpha and IFN-gamma from PBMC
69 isolated from dairy cows (Lacetera et al., unpublished).

70 The effects of high temperatures on functions of bovine polymorphonuclear cells (PMN)
71 received only limited attention, despite their involvement in the protection against mastitis and
72 metritis (Mehrzhad et al., 2010; LeBlanc et al., 2011). Elvinger and co-workers (1991) reported that
73 incubation temperature of 42 °C inhibited certain response of bovine PMN even if results from
74 different experiments were not clear and conclusive. Do Amaral and co-workers (2011) reported
75 that summer cooling may exert a positive effect on phagocytosis and oxidative burst of PMN
76 isolated from heat stressed dairy cows.

77 The apoptosis of PMN plays important roles in promoting resolution of the acute
78 inflammatory response (Headland and Norling, 2015). The impact of HS on PMN apoptosis has
79 been investigated in humans (Callahan et al., 1999; Kettritz et al., 2006; Nagarsekar et al., 2008;
80 Bzowska et al., 2011; Boyko et al., 2014). On the contrary, no information is available in bovine
81 species. Studies referred to humans focused on several aspects of PMN functions, which included
82 oxidants' production, heat shock response and apoptosis.

83 The present *in vitro* study was aimed at assessing whether exposure to temperature
84 simulating conditions of severe whole body hyperthermia affects functions of PMN in dairy cattle.

85

86 **2. Materials and Methods**

87 *2.1 Materials*

88 HEPES, RPMI-1640, HBSS with Mg⁺⁺ and Ca⁺⁺, Red Blood Cell Lysing Buffer Hybri-Max,
89 PMA (phorbol 12-myristate 13-acetate), Cytochrome *c*, Percoll, trypan blue, PBS without Ca⁺⁺ and
90 Mg⁺⁺ and endotoxin-free water were purchased from Sigma-Aldrich Co. (St.Louis, MO, USA).
91 Fetal Bovine Serum was provided by Biochrom AG (Berlin, Germany) and Apo-ONE[®]
92 Homogeneous Caspase-3/7 Assay from Promega Corporation (Milano, Italy). Ninety-six wells
93 sterile cell culture plates and 384 wells black sterile cell culture plates were purchased from Nunc

94 (Rochester, NY, USA). *Escherichia coli* (K-12 strain) BioParticles[®], fluorescein conjugate, was
95 provided by Molecular Probes (Invitrogen, San Giuliano Milanese (Milano), Italy).

96 2.2 Source and PMN isolation

97 The experiment was carried out on cells which were used for *in vitro* studies. Blood was
98 collected from seven clinically healthy multiparous (2nd and 3rd calving), pregnant and late
99 lactating (265 ± 50 days in milk) Holstein Friesian cows. Animals used in this study were managed
100 according to the National Law for Animal Welfare and Protection (Italy). Late lactating cows were
101 selected in order to avoid conditions of negative energy balance, which are known to interfere with
102 immune functions. Cows used in the study did not suffer from clinical diseases in the last month
103 before bleeding and somatic cell count at the time of blood collection was lower than 200,000
104 cells/ml. The experiment was carried out in March when cows were not exposed to heat stress
105 conditions. In detail, during the sampling period average daily temperature was 8 ± 1.2 °C with 0
106 and 15 °C as minimum and maximum temperatures, respectively. Cows were housed and fed in free
107 stalls, had free access to water, and were milked twice daily at 0600 and 1800 h.

108 Blood was collected by jugular venipuncture into sterile tubes containing 3.2% sodium
109 citrate as anticoagulant and was processed within 1h of collection. Bovine PMN were isolated as
110 described by Smits and coworkers (2000) with only minor modifications. Samples containing less
111 than 5% of eosinophils were used. Blood was centrifuged at 1000 x g for 30 min at 4°C: The
112 plasma, buffy coat and top one-third of the red blood cell pellet were removed. The remainder of
113 pellet containing PMN was diluted 1:3 vol:vol in sterile cold PBS, gently layered on 10 ml Percoll
114 1.087 g/ml and centrifuged at 400 x g for 40 min at room temperature. After the removal of Percoll,
115 the remaining red blood cells were removed by hypotonic lysis (Pisani et al., 2009) and PMN were
116 washed four times with sterile cold PBS. All functional assays were performed simultaneously on
117 the same day for any cow included in the present study.

118 2.3 Phagocytosis assay

119 Opsonisation of fluorescein-labelled *Escherichia coli* (K-12 strain) bioparticles was carried
120 out by incubating the bacterial suspension (6×10^8) with 20% autologous serum for 30 min on a
121 rocking roller at 37°C. The suspension was centrifuged at 800 x g for 15 min and suspended in
122 sterile HBSS. Opsonised *E. coli* were stored at -20°C until use.

123 The concentration of PMN was adjusted to 3×10^6 /ml and 100µl of this solution (3×10^5 cells)
124 were suspended in complete RPMI-1640 (RPMI 1640 with 20mM HEPES, 10% heat inactivated
125 FBS, 100 IU penicillin/ml, 100 µcells) and left at 39°C in humidified atmosphere at 5% CO₂ for 1h
126 to restore resting conditions. The PMN were then incubated for 1h at 39°C or 41°C in humidified
127 atmosphere of 5% CO₂. The temperature of 41 °C for this and for the experiments described below
128 was selected to simulate conditions of whole body hyperthermia, which can be detected in severely
129 heat stressed dairy cows (Silanikove, 2000). All the experiments were carried out in triplicate. The
130 cells were washed twice with sterile HBSS and fluorescein-labelled *E. coli* bioparticles (Pisani et
131 al., 2009) with a ratio of 42 particles/cell were added; PMN were incubated for 1h at 39°C or 41°C
132 in humidified atmosphere of 5% CO₂. Phagocytosis assay was carried out by measuring the
133 fluorescence of fluorescein-labelled *E. coli* bioparticles. Cells were washed twice with sterile HBSS
134 and incubated with 50 µL trypan blue for 1 min at room temperature to quench non-internalised
135 fluorescent bacteria. The cells were washed twice with HBSS. Fluorescence intensity was measured
136 using a fluorescence plate reader (Fluoroscan Ascent). Laser excitation was set at 485 nm and
137 fluorescence emission was collected using a 530 nm band pass filter.

138 2.4 Determination of Reactive Oxygen Species (ROS) production

139 The modulation of oxidative burst activity of isolated bovine PMN was studied by
140 cytochrome *c* reduction assay as previously described (Rinaldi et al., 2008). All the experiments
141 were carried out in duplicate. Aliquots of 2×10^5 PMN were suspended in 100 µl of HBSS with
142 Mg⁺⁺ and Ca⁺⁺ in two 96-well sterile plates and incubated at 39°C in humidified atmosphere of 5%

143 CO₂ for 1h to restore resting conditions. The PMN were then incubated for 1h at 39°C or 41°C in
144 humidified atmosphere of 5% CO₂. At the end of the incubation period, 10 µl of cytochrome *c* (1
145 mM) and 50 µl of HBSS or PMA (2.5 µg/mL final concentration) were added. The final volume of
146 all wells was adjusted to 200 µl with HBSS. Absorbance was measured on a plate reader (Bio-Tec
147 Instruments Inc., Winooski, VT, USA). The optical density (OD) was measured at a wavelength of
148 550 nm for 210 minutes and at 30 min intervals.

149 *2.5 PMN apoptosis measurement*

150 The concentration of PMN was adjusted to 2x10⁶/ml and 25µl of this solution (5x10⁴ cells),
151 seeded in two 384 wells black sterile cell culture plates and incubated for 1h at 39°C in humidified
152 atmosphere of 5% CO₂ to restore resting conditions. The PMN were then incubated for 2h at either
153 39°C or 41°C in humidified atmosphere of 5% CO₂. All the experiments were carried out in
154 triplicate. Apoptosis was determined by measuring the activities of two enzymes that play effector
155 roles in cows' apoptosis: caspase-3 and caspase-7 (Ceciliani et al., 2007). Briefly, to each wells
156 containing 5x10⁴ PMN in 25µl was added the same volume of Apo-ONE[®] Homogeneous Caspase-
157 3/7 Reagent, previously diluted 1:100 in the reaction buffer. Fluorescent intensity was measured
158 using a fluorescence plate reader Fluoroscan Ascent. Laser excitation was set at 485nm and
159 fluorescence emission was collected at 530nm band pass filter.

160 *2.6 Statistical analysis*

161 Statistical analyses were carried out using SPSS 22.0 for Windows (IBM, SPSS Inc, USA).
162 Descriptive statistics of different observed parameters are expressed as the mean (±SEM). For
163 phagocytosis, after assessment of normal distribution of data by Shapiro-Wilk Test, treated samples
164 were compared with control samples through a Student's t-Tests for paired samples. For apoptosis
165 and ROS, since the data were not normally distributed (Shapiro-Wilk Test), they were compared
166 using a GEE (generalized estimating equation) to take account of replications, in which the
167 dependent variables had an inverse Gaussian distribution, and an identity link function was used.

168 The effect of treatment, time of replications and their interaction were assessed. Goodness of fit was
169 assessed using a quasi-likelihood under independence model criterion (QIC). The threshold for
170 statistical significance was considered to be $P < 0.05$ (*) or $P < 0.01$ (**).

171

172 **3. Results**

173 *3.1 High temperature reduces the phagocytosis capacity of bovine PMN*

174 To explore the capacity of HS to modulate the bovine PMN phagocytosis, the neutrophils
175 were cultured at 39°C or 41°C. Results show that the phagocytosis ability of PMN to engulf
176 fluorescein labelled *E. coli* particles was affected by the exposure to high temperature: the
177 phagocytosis rate was significantly reduced (decrease of 37%, $P < 0.01$) when the cells were
178 incubated for 1 h at 41°C compared to 39°C (Figure 1).

179 *3.2 High temperature attenuates ROS production by resting and PMA-stimulated PMN*

180 In order to study whether HS modulates the generation of extracellular superoxide, PMN
181 were incubated at either 39°C or 41°C in the presence of cytochrome c, the reduction of which
182 occurring specifically in response to the generation of extracellular superoxide anion. Results
183 showed that, starting from 1h after HS, the oxidative burst of PMN, determined by ROS production,
184 was significantly reduced by exposure of the cells at 41°C s compared to 39°C (Figure 2A). In
185 detail the ROS production difference was 2% less after 60 min and the gap increased up to 12%
186 after 210 min. The same experiment was carried out on PMA-stimulated PMN, in order to study
187 whether the oxidative activity is influenced by HS in inflammatory condition (Figure 2B). Results
188 showed that PMN stimulated with PMA for 1h produced higher amount of extracellular ROS as
189 compared with resting PMN at both 39°C or 41°C (Figure 2A and 2B). Furthermore, results
190 confirmed that high temperature (41°C) specifically inhibited PMN ROS production starting from
191 2h after HS (Figure 2B). In details, the ROS production differences between PMN cultured at 39

192 and 41°C were higher than those observed in resting PMN and ranged between 2% at 120 min and
193 21% at 210min.

194 3.3 *High temperature does not affect PMN spontaneous apoptosis*

195 To investigate the apoptosis-modulating activity of HS, the enzymatic activities of caspase-3
196 and caspase-7, the two major executioners of the apoptosis pathway, were measured. Caspase 3/7
197 activity was detected 5 h after PMN exposure at 39°C or 41°C. The results show that the high
198 temperature did not influence the apoptosis rate when cells are exposed at 41°C as compared to
199 39°C (Figure 3).

200

201 **4. Discussion**

202 Results from the present study support the concept that HS, more specifically body
203 hyperthermia consequent to the exposure to high air temperature, is likely to dampen some bovine
204 PMN functions and may thus increase the risk of infections.

205 The findings of this study are in agreement with those already reported for dairy cows or
206 humans. Elvinger et al. (1991) described an impairment of ROS production in PMN exposed in
207 vitro to HS conditions, whereas they did not find any significant effect of HS on phagocytosis.
208 Different experimental conditions (42 instead of 41 °C) and different methods for measuring
209 phagocytosis may partially explain this difference. Furthermore, it has to be noticed that Elvinger
210 and co-workers (1991) cultured PMN at 42 °C, representing a temperature of at least 3 °C above
211 normal body temperature. It must be said that these high temperatures are quite unlikely in under
212 field conditions, and life threatening (Bianca, 1968). More recently, other authors (do Amaral et al.,
213 2011) reported that HS due to lack of cooling decreases ROS production and phagocytosis in
214 neutrophils from transition dairy cows kept in hot environment. The exposure of human PMN to 43
215 °C for 1 hour reduced the net intracellular oxidants production by 46% (Callahan et al., 1999).

216 Furthermore, both macrophages and PMN isolated from healthy humans and exposed *in vitro* to HS
217 conditions showed a significant reduction of NADPH oxidase-mediated O₂⁻ generation (Polla et al.,
218 1995).

219 The mechanisms behind the impairment of phagocytosis and ROS production observed in
220 PMN exposed to 41 °C have not been investigated in the present study. However, results reported
221 elsewhere for human PMN indicated that their exposure to HS conditions elicits a heat shock
222 response (Babcock and Meyer, 1998; Callahan et al., 1999) and that heat shock proteins may exert
223 an anti-inflammatory action. Results referred to immune cells largely differ from those referred to
224 whole body or other cell types (e.g., erythrocytes), which indicate that high temperature stimulates
225 ROS production with consequent higher risk of oxidative stress (Bernabucci et al., 2002; Slimen et
226 al., 2014). The reasons beyond such different behaviour of cells belonging to different
227 compartments are not easy to explain and are beyond the scope of this study.

228 The anti-inflammatory action of HS has been recently described using an inflammatory
229 model in pig and has been associated with physiological adaptation to high temperature (Campos et
230 al., 2014). Decline of ROS production is also in line with previous results relative to bovine PBMC,
231 indicating a reduction of intracellular ROS at 41 °C incubation temperature as a sign of reduction of
232 cellular metabolic rate (Lacetera et al., 2006). Finally, the decline of ROS production in PMN at
233 high temperatures may represent an adaptive mechanisms which reduce the risk of oxidative stress
234 commonly observed in heat stressed dairy cows (Bernabucci et al., 2002).

235 The fine tuning of PMN's lifespan in tissues is essential to reduce inflammatory-derived
236 injuries to the host (Greenlee-Wacker, 2016). We can rule out that the downregulation of ROS
237 production and the decrease of phagocytosis is related to a decrease of cells' number due to
238 apoptosis, given that the apoptosis rate of bovine PMN was not found to be affected by HS.
239 Available data on the effects of HS on apoptosis of human PMN are conflicting, depending on the
240 temperature of HS and time of exposure. Nagarsekar and co-workers (2008) found that culturing

241 human PMN at 39.5 °C for 8h greatly accelerated caspase dependent apoptosis. Exposure of human
242 PMN to 43 °C for 1 hour caused an early increase in the rate of apoptosis (Callahan et al., 1999),
243 whereas in a recent in vitro study carried out by culturing human neutrophils at 37 or 43 °C for 10
244 minutes, no substantial differences in viability were observed (Boyko et al., 2014). Therefore, we
245 may hypothesize that the temperature effects depend on the length of exposure.

246

247 **5. Conclusions**

248 In synthesis, the results of the present in vitro study suggest that some functions of PMN
249 exposed to temperature simulating conditions of severe whole body hyperthermia are impaired and
250 may at least partially explain the higher occurrence of infections during periods of hot weather.

251 Further epidemiology studies are necessary to clarify whether and under which specific
252 circumstances high environmental temperatures are associated with higher incidence of infections.
253 However, the risk of impairment of PMN functions under these conditions support the use of
254 management practices (i.e. cooling), which may help to limit the increase of body temperature and
255 optimization of other environmental features (i.e. general hygienic conditions, nutritional plans,
256 etc.), which may also play a role in causing infections outbreaks in hot environments.

257

258 **Abbreviations**

259 **HS:** heat stress; **PBMC:** peripheral blood mononuclear cells; **PMN:** polymorphonuclear cells; **TNF-**
260 **alpha:** tumor necrosis factor alpha; **INF-gamma:** interferon gamma; **PMA:** phorbol 12-myristate
261 13-acetate; **ROS:** Reactive Oxygen Species; **HBSS:** Hank's Balanced Salt Solution; **OD:** optical
262 density.

263 **Acknowledgements**

264 The authors thank Giorgina Kuzminsky of the DAFNE, Università della Tuscia for help with *in*
265 *vitro* experiments.

266

267 **Competing interest**

268 The authors declare that they have no competing interests.

269

270 **References**

- 271 Babcock GF, Meyer JG. The effects of stress on neutrophil function receptor expression, and
272 apoptosis. *FASEB Journal*. 1998; 12, A1045.
- 273 Bernabucci U, Ronchi B, Lacetera N, Nardone A. Markers of oxidative status in plasma and
274 erythrocytes of transition dairy cows during the hot season. *J Dairy Sci*. 2002; 85, 2173-
275 2179.
- 276 Bertocchi L, Vitali A, Lacetera N, Nardone A, Varisco G, Bernabucci U. Seasonal variations in
277 the composition of Holstein cow's milk and temperature-humidity index (THI)
278 relationship. *Animal* 2014; 8, 667-674.
- 279 Bianca W. Thermoregulation. In: Hafez, E.S.E. (Ed.), *Adaptation of Domestic Animals*. Lea
280 and Febiger, Philadelphia, PA, 1968; pp. 97–118, Chapter 7.
- 281 Boyko AA, Vetchinin SS, Sapozhnikov AM, Kovalenko EI. Changes in the heat shock 70 kDa
282 protein level in human neutrophils induced by heat shock. *Bioorg Khim*. 2014; 40, 488-
283 498.
- 284 Bzowska M, Hamczyk M, Skalniak A, Guzik K. Rapid decrease of CD16 (FcγRIII) expression
285 on heat-shocked neutrophils and their recognition by macrophages. *J Biomed*
286 *Biotechnol*. 2011;2011:284759.
- 287 Callahan TE, Marins J, Welch WJ, Horn JK. Heat shock attenuates oxidation and accelerates
288 apoptosis in human neutrophils. *J Surg Res*. 1999; 85, 317-322.
- 289 Campos PHRF, Merlot E, Damon M, Noblet J, Le Floc'h N. High ambient temperature
290 alleviates the inflammatory response and growth depression in pigs challenged with
291 *Escherichia coli* lipopolysaccharide. *Vet J*. 2014; 200, 404-409.

292 Ceciliani F, Pocacqua V, Miranda-Ribera A, Bronzo V, Lecchi C, Sartorelli P. alpha(1)-Acid
293 glycoprotein modulates apoptosis in bovine monocytes. *Vet Immunol Immunop.* 2007;
294 116, 145-152.

295 Dang VT, Speck P, Benkendorff K. Influence of elevated temperatures on the immune response
296 of abalone, *Haliotis rubra*. *Fish Shellfish Immunol.* 2012; 32, 732-740.

297 do Amaral BC, Connor EE, Tao S, Hayen MJ, Bubolz JW, Dahl GE. Heat stress abatement
298 during the dry period influences metabolic gene expression and improves immune status
299 in the transition period of dairy cows. *J Dairy Sci.* 2011; 94, 86-96.

300 Elvinger F, Hansen PJ, Natzke RP. Modulation of function of bovine polymorphonuclear
301 leukocytes and lymphocytes by high temperature in vitro and in vivo. *Am J Vet Res*
302 1991; 52, 1692-1698.

303 Gaughan JB, Lacetera N, Valtorta SE, Khalifa HH, Hahn GL, Mader TL. Response of domestic
304 animals to climate challenges. In "Biometeorology for adaptation to climate variability
305 and change", Eds., Ebi K.L., Burton I., McGregor G.R., Springer-Verlag, Heidelberg
306 (Germany), 2009; 131-170.

307 GieseckeWH. The effect of stress on udder health of dairy cows. *Onderstepoort J Vet Res.*
308 1985; 52, 175-193.

309 Greenlee-Wacker MC. Clearance of apoptotic neutrophils and resolution of inflammation.
310 *Immunol Rev.* 2016; 273(1), 357-70.

311 Headland SE, Norling LV. The resolution of inflammation: Principles and challenges. *Semin*
312 *Immunol.* 2005; 27, 149-160.

313 Kettritz R, Choi M, Salanova B, Wellner M, Rolle S, Luft FC. Fever-like temperatures affect
314 neutrophil NF-kappaB signaling, apoptosis, and ANCA-antigen expression. *J Am Soc*
315 *Nephrol.* 2006; 17, 1345-1353.

316 Lacetera N, Bernabucci U, Scalia D, Ronchi B, Kuzminsky G, Nardone A. Lymphocyte
317 functions in dairy cows in hot environment. *Int J Biometeorol.* 2005; 50, 105-110.

318 Lacetera N, Bernabucci U, Scalia D, Basiricò L, Morera P, Nardone A. Heat stress elicits
319 different responses in peripheral blood mononuclear cells from brown swiss and holstein
320 cows. *J Dairy Sci.* 2006; 89, 4606-4612.

321 Lacetera N, Bernabucci U, Basiricò L, Morera P, Nardone A.. Heat shock impairs DNA
322 synthesis and down-regulates gene expression for leptin and Ob-Rb receptor in
323 concanavalin A-stimulated bovine peripheral blood mononuclear cells. *Vet Immunol*
324 *Immunop.* 2009; 127, 190-194.

325 Lacetera N. Effect of Environment on Immune Functions. In “Environmental Physiology of
326 Livestock”, Eds., Collier R.J., Collier J.L., Wiley-Blackwell, The Atrium, Southern
327 Gate, Chichester (England), 2012; 165-179.

328 LeBlanc SJ, Osawa T, Dubuc J. Reproductive tract defense and disease in postpartum dairy
329 cows. *Theriogenology* 2011; 76, 1610-1618.

330 Martin LB, Hopkins WA, Mydlarz LD, Rohr JR. The effects of anthropogenic global changes
331 on immune functions and disease resistance. *Ann N Y Acad Sci.* 2010; 1195, 129-148.

332 Mehrzad J, Paape M, Burvenich C.. Role of neutrophils in protection of udder from infection in
333 high yielding dairy cows. *Iranian Journal of Veterinary Research* 2010; 11, 102-118.

334 Nagarsekar A, Greenberg RS, Shah NG, Hasday JD. Febrile-range hyperthermia accelerates
335 caspase-dependent apoptosis in human neutrophils. *J Immunol.* 2008; 181, 2636-2643.

336 Olde Riekerink RG, Barkema HW, Stryhn H. The effect of season on somatic cell count and the
337 incidence of clinical mastitis. *J Dairy Sci.* 2007; 90, 1704-1715.

338 Pisani LF, Lecchi C, Invernizzi G, Sartorelli P, Savoini G, Ceciliani F. In vitro modulatory
339 effect of omega-3 polyunsaturated fatty acid (EPA and DHA) on phagocytosis and ROS
340 production of goat neutrophils. *Vet Immunol Immunopathol.* 2009;131(1-2):79-85.

341 Polla BS, Stubbe H, Kantengwa S, Maridonneau-Parini I, Jacquier-Sarlin MR. Differential
342 induction of stress proteins and functional effects of heat shock in human phagocytes.
343 *Inflammation.* 1995; 19, 363-78.

344 Rinaldi M, Ceciliani F, Lecchi C, Moroni P, Bannerman DD. Differential effects of alpha1-acid
345 glycoprotein on bovine neutrophil respiratory burst activity and IL-8 production. *Vet*
346 *Immunol Immunopathol.* 2008;126(3-4), 199-210.

347 Silanikove N. Effects of heat stress on the welfare of extensively managed domestic ruminants.
348 *Livest Prod Sci.* 2000; 67, 1-18.

349 Slimen IB, Najar T, Ghram A, Dabbebi H, Ben Mrad M, Abdrabbah M. Reactive oxygen
350 species, heat stress and oxidative-induced mitochondrial damage. A review. *Int J*
351 *Hyperthermia.* 2014; 30, 513-523.

352 Smith KL, Todhunter DA, Schoenberger PS. Environmental mastitis: cause, prevalence,
353 prevention. *J Dairy Sci.* 1985; 68, 1531-1553.

354 Smits E, Burvenich C, Guidry AJ, Massart-Leen A. Adhesion receptor CD11b/CD18
355 contributes to neutrophil diapedesis across the bovine blood-milk barrier. *Vet Immunol*
356 *Immunopathol.* 2000; 73, 255-265.

357 Waage S, Sviland S, Odegaard SA. Identification of risk factors for clinical mastitis in dairy
358 heifers. *J Dairy Sci.* 1998; 81, 1275-1284.


359

360

361

362

Figure legends



363


364

365 Fig.1. Phagocytosis of fluorescein labelled *E. coli* bioparticles by bovine PMN.

366 The cells were pre-incubated at 39°C or 41°C and after 1h the *E.coli* bioparticles were added.
367 Phagocytosis was allowed to proceed for 1h at 39°C or 41°C. Each assay was carried out in
368 triplicate. Data are means \pm SEM of seven independent experiments. Significance was declared for
369 $P < 0.05$ (*) and $P < 0.01$ (**).

370


371



372

373 Fig.2. Effect high temperature on bovine PMN generation of extracellular superoxide as measured
 374 by cytochrome c reduction.

375 Resting (A) or PMA-stimulated (B) PMN were exposed to 39°C or 41°C in the presence of
 376 cytochrome c and absorbance values measured every 30 min. Each assay was carried out in
 377 duplicate. Data are means \pm SEM of seven independent experiments. Significance was declared for
 378 $P < 0.05$ (*) and $P < 0.01$ (**).



379

380 Fig.3. The caspase 3/7 enzymatic activity (spontaneous apoptosis) of bovine PMN.

381 PMN were incubation at 39°C or 41°C for 5 h. Each assay was carried out in triplicate. Data are
382 means \pm SEM of seven independent experiments.

383

384