

Lorenzo Rosso, Rosaria Carrinola, Alessandro Palleschi, Davide Tosi, Paolo Mendogni, Ilaria Righi, Mario Nosotti.

Thoracic Surgery and Lung Transplantation Unit,
Fondazione IRCCS Cà Granda Ospedale Maggiore Policlinico, Milan.
University of Milan, Italy.



AUTOFLUORESCENCE BRONCHOSCOPY AS INNOVATIVE MARKER OF AIRWAY COMPLICATIONS AFTER LUNG TRANSPLANTATION

AIRWAY COMPLICATIONS IN LUNG TRANSPLANTATION

- Healing of airway anastomosis is considered the Achilles' heel of lung transplantation.
- Airway complications lead to morbidity, to worsening in QOL and increased health care costs.
- Incidence 10%-15%, related mortality rate 2%-3%.


RISK FACTORS

- Length of mechanical ventilation both in donor (50-70 hrs) and recipient (PGD, acute cellular rejection) - barotrauma, inflammation, infection.
- Height mismatch (gender)
- Length of donor bronchus (secondary carina)
- Corticosteroids (not only adverse effects)
- Ischemic time ?????
- BMI, organ preservation technique, acute injury injury.
- No difference between DCD and DBD.

Machuzak et al


AUTOFLUORESCENCE PRINCIPLES

- Autofluorescence: Tissue stimulated by high energy short wavelength blue light emits light at lower energy frequencies in green and red spectrum.
- Fluorescence reaction is achieved by exciting endogenous fluorophores in the sub-epithelial layer.
- AF is attenuated by thickening of epithelial layer (tumorous tissues) and by hemoglobin.
- Facilitates detections of lung cancer at an early stage.



AUTOFLUORESCENCE PRINCIPLES

- Normal tissues are displayed in green and tumorous lesions in magenta.
- In addition to blue excitation AFI uses green illumination light that is reflected by hemoglobin contained in the living tissues.
- A similar result is obtained in ischemic or infected mucosa (bacterial biofilms).



D. Huttenberger et al.

STUDY AIM


Find a relationship between the degree of bronchial vascularization after lung transplantation and the onset of airway stenosis using autofluorescence bronchoscopy.

METHODS

- From February 2014 and May 2015 all consecutive patients receiving single or double lung transplantation were included.
- Surgical technique was the same in all patients; end to end bronchial anastomosis with two absorbable 4/0 running sutures.
- Patients requiring separate lung ventilation, intensive care unit stay > 7 days, survival < 6 months were excluded.
- Bronchoscopies were scheduled every 7 days for the first month after operation then at 3, 6 and twelve months.
- All procedures were conducted with the same instrument (EVIS Lucera Spectrum AFI, Olympus, Japan) by the same endoscopist and recorded on USB storage device.

METHODS

- Bright pink color was typical of ischemic mucosa.
- Deep green was characteristic for normal mucosa.
- Pictures of the first bronchial carina were analyzed with color histograms.
- The red and green intensity ratio (R/G) was used to grade the pictures.



RESULTS

- 23 patients were eligible for the study and we examined 39 bronchial anastomosis.
- We considered each anastomosis as a unit for statistical purpose.
- We observed 8 bronchial stenosis in 6 patients.
- Airway complications were correlated with ECMO, ischemic time, gender R/D mismatch, PGD.
- R/G ratio was correlated with stenosis onset.
- Statistic analysis: Fischer's exact test, Roc curve, logistic regression.


STUDY POPULATION

39 anastom.	range	mean	50% CI	median	95% CI	SD
AGE R	21-67	43.4	38.1-47.8	41	38.0-48.9	13.43
AGE D	16-67	41.4	36.1-46.7	41	33.0-53.9	16.33
MecVent D	0-8	2.62	1.6-3.5	2	1.0-5.0	2.49
Cold Isc Time	180-1200	474	393.8-553.4	420	193.0-682.4	244
TX	7 SL	16 DL	12 FC	8 IPF	2 BPCO	1 LAM

STENOSIS

	Site	Time	Gender	PGD	ECMO	BAS	MDS	Islet	
1	IPF	L-ant	0 m	FF	-	-	MDD2450	Stent	
2	FC	R-int	6 m	MM	-	-	Pneuma A-Axilla	Pneumato (Stent)	
3	IPF	L-main	3-6 m	MM	-	-	KPC	MDD1450	Stent
4	FC	R-int	6 m	MM	+	Bridge-Post	Pneuma A	MDD2450	Stent
5	FC	L-main	1 m	MM	+	Bridge-Post	Pneuma A	MDD2450	Pneumatic
6	IPF	L-main	1 m	MF	+	-	-	MDD1450	No treat
7	FC	L-main	12 m	FM	+	Post	PhxPC	MDD2450	No treat
8	FC	R-main	12 m	FM	+	Post	PhxPC	MDD2450	No treat


RESULTS



RESULTS


- After Logistic regression we found correlation between stenosis onset and:
 - R/G ratio at 45 days (p=0.0417, OR 37.36)
 - R/G ratio at 3 months (p=0.00231, OR 17.66)
 - R/G ratio at 6 months (p=0.0036, OR 61.98)
 - R/G ratio at 12 months (p=0.0027, OR 533.18)
- Cold ischemia time (p=0.0014, OR 1.00)
- No correlation between stenosis and days of intubation of the donor (p=0.815).

STENOSIS TREATMENT



RESULTS


ROC curve: stenosis and R/G ratio at 3, 6 and 12 months.



Associated criterion R/G > 1.19

RESULTS

ROC curve: stenosis and cold ischemia.



Associated criterion > 840 min

RESULTS

Fisher's exact test

- We found correlation between stenosis and postopECMO (p=0.0105), PGD (p=0.0089).
- No statistical significance with D/R gender mismatch (p=0.087).
- No correlation with ECMO bridge and intraoperative ECMO.

CONCLUSIONS

- Autofluorescence bronchoscopy may allow the assessment of bronchial ischemic mucosa.
- Autofluorescence bronchoscopy may provide a valuable tool as preclinical marker of airway complications after lung transplantation.
- Further studies with larger series are mandatory.