



## New insights on Flavescence dorée phytoplasma ecology in the vineyard agro-ecosystem in southern Switzerland

Journal:	<i>Annals of Applied Biology</i>
Manuscript ID	AAB-2016-0324.R2
Manuscript Type:	Research paper
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Casati, Paola; Università degli Studi di Milano, Dipartimento di Scienze Agrarie e Ambientali – Produzione, Territorio, Agroenergia (DiSAA)</p> <p>Jermi, Mauro; Dipartimento federale dell'economia, della formazione e della ricerca (DEFR) - Agroscope, Istituto delle scienze della produzione vegetale (IPV), Centro di Cadenazzo</p> <p>Quaglino, Fabio; Università degli Studi di Milano, Dipartimento di Scienze Agrarie e Ambientali – Produzione, Territorio, Agroenergia (DiSAA)</p> <p>Corbani, Giacomo; Università degli Studi di Milano, Dipartimento di Scienze Agrarie e Ambientali – Produzione, Territorio, Agroenergia (DiSAA)</p> <p>Schaerer, Santiago; Département fédéral de l'économie, de la formation et de la recherche (DEFR) - Agroscope, Département Protection des végétaux grandes cultures et vigne/Viticulture et oenologie</p> <p>Passera, Alessandro; Università degli Studi di Milano, Dipartimento di Scienze Agrarie e Ambientali – Produzione, Territorio, Agroenergia (DiSAA)</p> <p>Bianco, Piero; Università degli Studi, Dipartimento di Scienze Agrarie e Ambientali – Produzione, Territorio, Agroenergia (DiSAA)</p> <p>Rigamonti, Ivo; Università degli Studi di Milano, Dipartimento di Scienze per gli Alimenti, la Nutrizione, l'Ambiente (DeFENS)</p>
Key Words:	Grapevine yellows, Phytoplasma, Insect vectors, <i>map</i> gene, <i>Orientus ishidae</i> , Hazel, Willow

1  
2  
3 1 **New insights on Flavescence dorée phytoplasma ecology in the**  
4  
5  
6 2 **vineyard agro-ecosystem in southern Switzerland**  
7  
8  
9 3

10  
11 4 Casati P.<sup>1,†</sup>, Jermini M.<sup>2,†</sup>, Quaglino F.<sup>1</sup>, Corbani G.<sup>1</sup>, Schaerer S.<sup>2</sup>, Passera A.<sup>1</sup>,  
12  
13  
14 5 Bianco P.A.<sup>1</sup>, Rigamonti I.E.<sup>3</sup>  
15  
16 6

17  
18 7 <sup>1</sup> Dipartimento di Scienze Agrarie e Ambientali – Produzione, Territorio, Agroenergia (DiSAA), Università  
19  
20 8 degli Studi di Milano, via Celoria 2, 20133 Milan, Italy  
21

22 9 <sup>2</sup> Research Station Agroscope Changins – Wädenswil ACW, Centro di Ricerca di Cadenazzo (TI). A Ramél,  
23  
24 10 18, CH-6593 Cadenazzo, Switzerland  
25

26 11 <sup>3</sup> Dipartimento di Scienze per gli Alimenti, la Nutrizione, l'Ambiente (DeFENS), Università degli Studi di  
27  
28 12 Milano, Via Celoria 2, 20133 Milan, Italy  
29  
30

31 13  
32  
33 14 **Correspondence:** Paola Casati, Dipartimento di Scienze Agrarie e Ambientali – Produzione,  
34  
35 15 Territorio, Agroenergia, Università degli Studi di Milano, via Celoria 2, 20133 Milano, Italy. Tel :  
36  
37 16 +39-02-50316793 ; fax : +39-02-50316781 ; e-mail: [paola.casati@unimi.it](mailto:paola.casati@unimi.it)  
38  
39  
40

41 17  
42  
43 18 **Running title:** Insights on Flavescence dorée ecology  
44  
45 19

46  
47 20 <sup>†</sup> The first two authors contributed equally to the work  
48  
49  
50 21  
51  
52 22  
53  
54 23  
55  
56 24  
57  
58  
59  
60

25 **SUMMARY**

26 Phytoplasmas associated with Flavescence dorée (FDp) grapevine disease are quarantine  
27 pathogens controlled through mandatory measures including the prompt eradication and destruction  
28 of diseased plants, and the insecticide treatments against the insect vector, the ampelophagous  
29 leafhopper *Scaphoideus titanus*. In the present study, a multidisciplinary approach has been applied  
30 to investigate the FDp ecological cycle in a test vineyard agro-ecosystem in Canton Ticino, south  
31 Switzerland. Despite the scarce population density of *S. titanus*, a regular trend of new infections  
32 (3.4% of the total vines) through the years was observed. The leafhopper *Orientalus ishidae* was  
33 found as the most abundant among the captured insect species known as phytoplasma vectors (245  
34 out of 315 specimens). The population of *O. ishidae* was evidenced prevalently (167 specimens) in  
35 the south-western side of the vineyard and within the neighboring forest constituted mainly by hazel  
36 (*Corylus avellana*) and willow (*Salix* spp.). These plant species were found infected by FDp related  
37 strains (30% of analyzed trees) for the first time in this study. Interestingly, *O. ishidae* was found to  
38 harbor FDp related strains in high percentage (26% of the analyzed pools). In addition, 16SrV  
39 phytoplasma group was detected for the first time in the insect *Hyalesthes obsoletus* and a FDp  
40 related strain in *Thamnotettix dilutior*, present in low populations within the test vineyard.  
41 Molecular characterization and phylogenetic analyses of *map* gene sequences of FDp and related  
42 strains, here identified, revealed the great prevalence of the *map*-type FD2 in grapevines (97%) and  
43 in *O. ishidae* pools (72%). Such a *map*-type was found also in hazel and in *T. dilutior*, but not in *S.*  
44 *titanus*. Moreover, *map*-types FD1 and FD3 were identified for the first time in Switzerland in  
45 several host plants and phytoplasma vectors, including grapevine (FD1), *S. titanus* (FD1) and *O.*  
46 *ishidae* (FD1 and FD3). Based on the data obtained in this study, it is reasonable to hypothesize that  
47 the ecological cycle of FDp could be related not exclusively to the grapevine-specific feeding diet  
48 of *S. titanus*, but it could include other insect vector(s) and/or plant host(s). Further studies will be

1  
2  
3 49 needed to prove the role of *O. ishidae* as vector able to transmit FDp from wild plants (e.g. hazel) to  
4  
5 50 grapevine.  
6

7 51

8  
9  
10 52 **Keywords:** Grapevine yellows; Phytoplasma; Insect vectors; *map* gene; *Orientus ishidae*; Hazel;  
11 53 Willow  
12

13 54

## 14 55 INTRODUCTION

15  
16  
17  
18 56 Flavescence dorée (FD) is the most epidemic disease within the grapevine yellows (GY)  
19 57 complex, and causes strong economic damages to viticulture in Europe, both in terms of quality and  
20 58 quantity (Belli *et al.*, 2010). Typical FD symptoms, indistinguishable from those associated with  
21 59 other diseases of the GY complex, include berry shrivel, desiccation of inflorescences, color  
22 60 alterations and curling of the leaves, reduction of growth, and irregular ripening of wood (Belli *et*  
23 61 *al.*, 2010).  
24  
25  
26  
27  
28  
29  
30

31  
32 62 Based on nucleotide sequence analysis of the gene *16S rRNA* and the intergenic spacer  
33 63 between *16S* and *23S rRNA* genes, FD phytoplasmas (FDp) have been classified in two taxonomic  
34 64 subgroups belonging to the 16SrV group (Elm yellows group), 16SrV-C and -D (Lee *et al.*, 1998;  
35 65 Angelini *et al.*, 2001; Davis & Dally, 2001; Lee *et al.*, 2004). As reported for other phytoplasmas  
36 66 belonging to taxonomic group 16SrV (Martini *et al.*, 2002; Navrátil *et al.*, 2009; Malembic-Maher  
37 67 *et al.*, 2011), the variability of the *16S rRNA* gene is insufficient to account for the emerging  
38 68 ecological differences in FDp biological cycles. Thus, further studies focus on finer differentiation  
39 69 of phytoplasmas associated with FD. Molecular and phylogenetic analyses on the genetic locus *map*  
40 70 demonstrate the presence of three *map*-types: FD1 and FD3 (including strains from subgroup  
41 71 16SrV-C), and FD2 (including strains from subgroup 16SrV-D) (Arnaud *et al.*, 2007). Geographic  
42 72 distribution of *map*-types is different: (i) FD1 and FD2 have been identified, respectively, in 17%  
43 73 and 83% of symptomatic grapevines in France, where FD3 is not present (Arnaud *et al.*, 2007); (ii)  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 74 the three *map*-types have been identified in Italy, with a prevalence of FD1 and FD3 in north-  
4  
5 75 western and north-eastern regions, respectively (Arnaud *et al.*, 2007; Foissac & Maixner, 2013);  
6  
7 76 (iii) FD2 and FD3 have been detected in Slovenia and Croatia, with a prevalence of FD3 (Mehle *et*  
8  
9 77 *al.*, 2011; Plavec *et al.*, 2015); (iv) only FD3 has been identified in Serbia (Filippin *et al.*, 2009); (v)  
10  
11 78 in northern Spain, Portugal and Switzerland only FD2 has been reported (Foissac & Maixner,  
12  
13 79 2013).

14  
15  
16 80 Moreover, nucleotide sequences and phylogenetic analyses from multiple genes indicate that  
17  
18 81 FDp, Palatinate grapevine yellows phytoplasmas (PGYp) and Alder yellows phytoplasma (AldYp)  
19  
20 82 are clustered together, supporting their monophyletic origin (Arnaud *et al.*, 2007; Holz *et al.*, 2015).  
21  
22 83 AldYp, widespread in Europe and never detected in America, is transmitted by the insect *Oncopsis*  
23  
24 84 *alni* (Schrank) (Maixner & Reinert, 1999). Such highlights suggest that FDp could have originated  
25  
26 85 in Europe. As a matter of fact, it has been hypothesized that (i) AldYp strains, genetically related to  
27  
28 86 FD1, FD2 and FD3 *map*-types of FDp, are occasionally transmitted to grapevine by *O. alni*  
29  
30 87 (Maixner *et al.*, 2000); (ii) after the entrance of *Scaphoideus titanus* Ball in Europe, such AldYp  
31  
32 88 strains have been transmitted exclusively to grapevine, generating the FD epidemics (Arnaud *et al.*,  
33  
34 89 2007).

35  
36  
37  
38 90 Due to the host plant range and feeding preferences of each phytoplasma vector, the  
39  
40 91 majority of GY (e.g. Bois Noir and Palatinate grapevine yellows) spreads only slowly and,  
41  
42 92 apparently, not from vine to vine. On the other hand, FD is an epidemic disease characterized by its  
43  
44 93 rapid spread within vineyards due to vine-to-vine transmission (EPPO/CABI, 1996). FDp are  
45  
46 94 transmitted by *S. titanus*, a monovoltine leafhopper accidentally introduced to France from North  
47  
48 95 America in the 1950ies. In Europe only species of the *Vitis* genus have been shown to sustain the  
49  
50 96 whole life cycle, from egg to egg, of *S. titanus*; moreover, this leafhopper is considered as  
51  
52 97 oligophagous (Chuche & Thiéry, 2014). Consequently, geographic areas hosting large vector  
53  
54 98 populations can be damaged by strong FD epidemics if FDp are also present. Due to this aspect,  
55  
56  
57  
58  
59  
60

1  
2  
3 99 phytoplasmas associated with FD are quarantine pathogens to be controlled through mandatory  
4  
5 100 measures: (i) promptly eradicating and destroying any plants showing GY type symptoms, even  
6  
7 101 before confirmation of FD infection by laboratory tests where FD is already present, and (ii)  
8  
9 102 compulsory insecticide sprayings to control the *S. titanus* populations. The number of applications  
10  
11 103 varies from one to three per year (targeting nymphs and adults) in commercial vineyards and can be  
12  
13 104 more numerous in nurseries (EFSA, 2014). Even though these measures have reduced the impact of  
14  
15 105 FD, the still broad presence of FDp in different countries indicates that FD epidemic is still  
16  
17 106 important (Botti & Bertaccini, 2007; Jermini *et al.*, 2014).

18  
19  
20  
21 107 Moreover, recent studies report that the FD epidemiological cycle could be more complex  
22  
23 108 than the exclusive FDp transmission from grapevine to grapevine by *S. titanus*; in fact, (i) FDp  
24  
25 109 related strains are also commonly found in other plant hosts such as *Ailanthus*, *Alnus*, and *Clematis*  
26  
27 110 (EFSA, 2014); (ii) several studies reveal that *S. titanus* can feed on other plant species, on which it  
28  
29 111 can only partially complete its biological cycle (Schvester *et al.*, 1962; Caudwell *et al.*, 1970;  
30  
31 112 Trivellone *et al.*, 2013); these host plants could play a role as source of inoculum for FDp  
32  
33 113 transmission to grapevine (Chuche & Thiery, 2014); (iii) other studies reveal that FDp can be  
34  
35 114 transmitted to grapevine by other insects. In detail, Filippin and colleagues (2009) have  
36  
37 115 demonstrated that FDp can be transmitted from *Clematis vitalba* L. to grapevine by the planthopper  
38  
39 116 *Dictyophara europaea* (L.). The frequency of phytoplasma transmission from *Clematis* to grapevine  
40  
41 117 remains to be determined but phytoplasma transmission cannot provoke an FD outbreak in the  
42  
43 118 absence of the leafhopper *S. titanus* (EFSA, 2014). Recently, the leafhoppers *O. alni* (Maixner *et al.*,  
44  
45 119 2000; Arnaud *et al.*, 2007; Mehle *et al.*, 2011) and *Orientalis ishidae* (Matsumura) (Mehle *et al.*,  
46  
47 120 2010, 2011; Gaffuri *et al.*, 2011; Koczor *et al.*, 2013) have been found infected by FDp related  
48  
49 121 strains in Europe; moreover, latest findings revealed the capability of *O. ishidae* to transmit 16SrV  
50  
51 122 group phytoplasmas to grapevine (Lessio *et al.*, 2016).  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 123 Based on such previously described evidences, in the present study carried out in Canton  
4  
5 124 Ticino (Switzerland), a molecular epidemiology approach has been used to test if the FD complex  
6  
7 125 could be considered as an open system by investigating the presence of putative vectors and host  
8  
9 126 plants in the vineyard agro-ecosystem. A special focus has been laid on *O. ishidae*, previously  
10  
11 127 found harboring FDp related strains in the studied area (Trivellone *et al.*, 2016); for the first time,  
12  
13 128 particular attention was paid to investigate the epidemiological role of *O. ishidae* plant hosts in the  
14  
15 129 vineyard and its surroundings.  
16  
17  
18  
19  
20

## 21 131 **MATERIALS AND METHODS**

### 22 132 23 133 **Characteristics of the test vineyard**

24  
25 134 Investigation about the FD epidemiology was carried out in one vineyard, hereby named  
26  
27 135 “the test vineyard”, located in Stabio, a town in the Southern wine-growing region of canton Ticino,  
28  
29 136 Switzerland [WGS 84 (lat/lon) 45.85547, 8.92749; Alt 371 m] and in the surrounding vegetation.  
30  
31 137 The test vineyard consists of 21 rows (2177 vines) of Chardonnay vines grafted on 3309 C  
32  
33 138 rootstock. In this vineyard, grapevines are trained using the Guyot system (distance between rows 2  
34  
35 139 m; plant distance along the row 1 m), occasionally mowing between the rows and chemical weeding  
36  
37 140 along the rows. The vineyard is bordered by woods to the south, meadows to northwest, and  
38  
39 141 vineyards to the east (Figure 1). Based on the directives established by the Cantonal Phytosanitary  
40  
41 142 Service, mandatory destruction of grapevine plants showing GY-typical symptoms are carried out  
42  
43 143 in the vineyard, as well as insecticide treatments against *S. titanus* with the larvicidal insecticide  
44  
45 144 Buprofezin at two time points, at maximum egg hatching and two weeks later, according to  
46  
47 145 monitoring activities and model predictions (Prevostini *et al.*, 2013). Despite these measures,  
48  
49 146 adopted since 2005, and the absence of abandoned vineyards, FD is still present in this area. In  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 147 particular, 3.3% of the grapevine plants in our test vineyard were FDp-positive in 2013 (data from  
4  
5 148 Research Station Agroscope Changins, Switzerland).

6  
7 149

8  
9  
10 150 **Insect monitoring and sampling**

11  
12 151 The activities were carried out from May to November 2014 with the aim to (i) monitor the  
13  
14 152 presence of *S. titanus* and *O. ishidae* nymphs and adults developing on grapevine and wild plants in  
15  
16 153 the test vineyard and in its surroundings, (ii) survey the leafhopper species known as putative  
17  
18 154 phytoplasma vectors inhabiting the vine plants, (iii) determine the flight dynamics of *S. titanus* and  
19  
20 155 *O. ishidae* by monitoring the presence of their adults spreading/dispersing through/across the test  
21  
22 156 vineyard and in its surroundings.

23  
24  
25 157 (i) Three groups each of 30 grapevine plants, randomly selected within the test vineyard,  
26  
27 158 have been sampled weekly from May 13 to September 30 (21 overall samples) with the beating tray  
28  
29 159 method (one hit per plant). Moreover, wild plants hosting known leafhopper phytoplasma vectors,  
30  
31 160 around and within the test vineyard and in neighbouring vineyards, have been occasionally sampled  
32  
33 161 from June 3 to September 2 (38 overall samples) by the beating tray method (up to 4 hits per plant,  
34  
35 162 according to the plant dimension, until reaching 40 hits).

36  
37  
38 163 (ii) Fourteen Aeroxon yellows sticky traps (10 x 25 cm) placed horizontally in grapevine  
39  
40 164 canopy (Jermini *et al.*, 1992a, 1992b) were used to detect the presence of *S. titanus* and other  
41  
42 165 Auchenorrhyncha on the canopy inside the vineyard from July 01 to November 04. Traps were  
43  
44 166 collected and substituted each week.

45  
46  
47 167 These two methods also allowed to simultaneously survey other leafhoppers inhabiting  
48  
49 168 grapevine and wild plants in the test vineyard and in its surroundings.

50  
51  
52 169 (iii) Rebell yellow sticky traps (8 x 16 cm), placed vertically outside the plant canopy, were  
53  
54 170 utilized to capture leafhoppers spreading/dispersing through/across the vineyard and the  
55  
56 171 surrounding vegetation, with the specific objective to determine the flight dynamics of *S. titanus*



1  
2  
3 172 and *O. ishidae*. In detail, six Rebell traps (V1 to V6; V, vineyard) were positioned inside the test  
4  
5 173 vineyard (May 13 - November 04), eight (B1 to B8; B, border) on its borders (May 13 - November  
6  
7 174 04), and nine (W1 to W9; W, woods) outside of it (July 30 - November 04). Particularly, traps W1  
8  
9 175 to W7 were positioned at the external border of the forest neighbouring to the test vineyard, close to  
10  
11 176 other vineyards of the examined area; traps W8 and W9 were positioned inside the forest  
12  
13 177 neighbouring to the test vineyard (Figure 1). Traps were collected and substituted each week.

14  
15  
16 178 Auchenorrhyncha specimens, after being sorted out from the material caught by the traps  
17  
18 179 and the beating tray, were individually identified at the species level with a stereo microscope. All  
19  
20 180 individuals of species belonging to the subfamily Deltocephalinae, considered putative vectors of  
21  
22 181 FDp (Bressan *et al.*, 2006), and other leafhopper and planthopper species, reported in the scientific  
23  
24 182 literature as known vectors of phytoplasmas, were preserved in 70 % alcohol for further molecular  
25  
26 183 analyses.  
27  
28  
29  
30  
31

### 32 185 **Plant sampling**

33  
34 186 During a survey on GY carried out from July to September 2014 in the test vineyard,  
35  
36 187 symptomatic leaves were sampled from 74 grapevine plants, showing GY symptoms for the first  
37  
38 188 time. Furthermore, leaf samples were also collected from symptomless woody and shrubby plants  
39  
40 189 of species (*Betula pendula*, *Carpinus betulus*, *Corylus avellana*, *Salix caprea*, *Salix* sp., *Prunus*  
41  
42 190 *laurocerasus*, *Urtica dioica*) listed in the scientific literature as host plants of *O. ishidae*  
43  
44 191 (<https://gd.eppo.int/reporting/article-4763>; Günthart *et al.*, 2004; Nickel, 2010; Mehle *et al.*, 2011).  
45  
46 192 Finally, leaves were also collected from symptomless plants reported as hosts of phytoplasmas  
47  
48 193 closely related to FD (*Clematis vitalba*) (Filippin *et al.*, 2009). Leaf samples from Madagascar  
49  
50 194 periwinkle (*Catharanthus roseus* L. (G. Don)) plants, maintained in the greenhouse of the  
51  
52 195 Department of Agriculture and Environmental Sciences (University of Milan, Italy), infected by  
53  
54 196 FDp strains FD92 (subgroup 16SrV-D) and FD70 (subgroup 16SrV-C), '*Ca. Phytoplasma solani*'  
55  
56  
57  
58  
59  
60

1  
2  
3 197 strain STOL (subgroup 16SrXII-A), and '*Ca. Phytoplasma asteris*' strain AY1 (subgroup 16SrI-B),  
4  
5 198 were collected and used as controls in molecular analyses.  
6

7 199

8  
9  
10 **Total nucleic acids extraction**

11 201 Total nucleic acids were extracted from pools of the captured insects (each pool includes  
12  
13  
14 202 one to five specimens captured together on the same trap) following a protocol adapted from Alma  
15  
16 203 *et al.* (1997). Briefly, the ethanol-preserved adults were dried onto filter paper, pooled and  
17  
18 204 homogenised in a pre-warmed (65 °C) CTAB-based buffer (2.5% w/v cetyl-trimethyl-ammonium-  
19  
20 205 bromide (CTAB); 100 mM Tris pH 8.0, 1.4 M NaCl; 50 mM EDTA pH8; 1% PVP-40; 0.5%  
21  
22 206 ascorbic acid). After incubation at 65 °C for 20 min, nucleic acids were extracted with one volume  
23  
24 207 of chloroform:isoamyl alcohol 24:1 v/v solution and then precipitated with the addition of one  
25  
26 208 volume of cold isopropanol at -20 °C for 20 min. The obtained total nucleic acids pellet was washed  
27  
28 209 with 70% ethanol, vacuum dried, redissolved in 50 µL TE pH 8.0 and maintained at -20 °C until  
29  
30 210 further use.  
31  
32

33  
34 211 Total nucleic acids were extracted from examined plants using a modified Angelini *et al.*  
35  
36 212 (2001) protocol. Briefly, leaf petioles (1 g) were inserted in extraction bags (12 x 12 cm) (Bioreba,  
37  
38 213 Reinach, Switserland) and ground in 7 ml of pre-warmed (65 °C) 2.5% CTAB-based buffer (see  
39  
40 214 above) using a Tissue Homogenizer (Sediag, Longvic, France). The solution (1 ml) was held at 65  
41  
42 215 °C for 30 min. After incubation, nucleic acids were extracted by adding chloroform:isoamyl alcohol  
43  
44 216 24:1 v/v solution and precipitated by incubation with isopropanol at -20 °C for 10 min. The nucleic  
45  
46 217 acid pellet was washed with 70% ethanol, dried, resuspended in TE buffer, re-precipitated with  
47  
48 218 sodium acetate and absolute ethanol, and re-purified by washing with 70% and 80% ethanol.  
49  
50 219 Obtained nucleic acids pellet was air-dried, suspended in 100 µL of deionised autoclaved water and  
51  
52 220 maintained at -30 °C until further use.  
53  
54

55  
56 221  
57  
58  
59  
60

**222 Detection of Flavescence dorée phytoplasma and related strains**

223 Total nucleic acids extracted from each plant sample and insect pool were used as templates  
224 for real-time PCR assays carried out for the FDp and related strains-specific amplification of the  
225 gene *rplN*, coding the ribosomal protein L14 (Durante *et al.*, 2012), using the commercial kit Real-  
226 time PCR (TaqMan probe and primers) (IPADLAB, Lodi, Italy) and TaqMan Universal PCR  
227 master mix (Applied Biosystems, Monza, Italy), according to the manufacturer's instructions.  
228 Thermocycling was carried out on the StepOnePlus Real-Time PCR System (Applied Biosystems,)  
229 and consisted of an initial denaturation at 95 °C for 10 min followed by 40 cycles of 15 sec at 95 °C  
230 and 1 min at 62 °C. Total nucleic acids extracted from periwinkle plants infected by phytoplasma  
231 strains were used as positive (strains FD92 and FD70) and negative (strains AY1 and STOL)  
232 controls; moreover, total nucleic acids from healthy periwinkle plants and reaction mixture without  
233 template were used as negative controls. Each sample was analyzed in duplicate.

234

**235 Flavescence dorée phytoplasma and related strain typing**

236 FDp and related strains, identified in plants and insect pools by real-time PCR assay, were  
237 typed by PCR-based amplification and sequence analyses of the gene *map* encoding a methionine  
238 aminopeptidase (Arnaud *et al.*, 2007). Precisely, total nucleic acids of infected plants and insect  
239 pools were employed as templates in nested PCR assays performed using primer pair  
240 FD9f5/MAPr1 followed by FD9f6/MAPr2, following the reaction conditions previously described  
241 (Arnaud *et al.*, 2007). All amplifications were performed using the GeneAmp PCR System 9700  
242 (Applied Biosystem). Positive and negative controls employed in nested PCR amplification of *map*  
243 gene were as described above. PCR products were analysed by electrophoresis in 1% agarose gels,  
244 stained with ethidium bromide and observed under UV light.

245 As indicated by Arnaud *et al.* (2007), *map* gene amplicons were typed by restriction  
246 fragment length polymorphism (RFLP) analysis performed through double digestion with the

1  
2  
3 247 enzymes *AluI* and *Eco72I* Fast digest (MBI-Fermentas). Reactions were carried out according to the  
4  
5 248 manufacturer's instructions. RFLP profiles obtained were visualized under UV light in 3% agarose  
6  
7 249 gels stained with ethidium bromide. Attribution of FDp and related strains, identified in the present  
8  
9 250 study, to the *map*-types FD1, 2, and 3 was carried out through the comparison of their *AluI-Eco72I*-  
10  
11 251 RFLP profiles with those described in literature (Arnaud *et al.*, 2007).

12  
13  
14 252 *Map* gene fragments amplified from FDp and related strains, representative of the obtained  
15  
16 253 *AluI-Eco72I*-RFLP profiles and of the phytoplasma hosts, were selected for nucleotide sequence  
17  
18 254 analyses. The *map* gene amplicons were sequenced in both senses (employing primers FD9f6 and  
19  
20 255 MAPr2) by a commercial service (Eurofins, Milan, Italy) to achieve at least 5x coverage per base  
21  
22 256 position. Nucleotide sequence data were assembled by employing the Contig Assembling program  
23  
24 257 of the software BioEdit version 7.0.5 (<http://www.mbio.ncsu.edu/bioedit/bioedit.html>). *Map* gene  
25  
26 258 nucleotide sequences of FDp and 16SrV group phytoplasma strains previously published were  
27  
28 259 retrieved from GenBank (Table S1) and utilized for comparison with the sequences obtained in this  
29  
30 260 study. *Map* gene nucleotide sequences were aligned using the "ClustalW Multiple Alignment"  
31  
32 261 application and analyzed for sequence identity determination using the "Sequence Identity Matrix"  
33  
34 262 application of the software BioEdit.

35  
36  
37  
38 263 Nucleotide sequences of *map* gene of FDp and related strains and 16SrV group phytoplasma  
39  
40 264 strains from the present study and previously described in GenBank were employed for  
41  
42 265 phylogenetic analyses. Minimum-Evolution method was carried out using the Jukes-Cantor model  
43  
44 266 and bootstrap replicated 1000 times with the software MEGA6 to obtain an unrooted phylogenetic  
45  
46 267 tree (Tamura *et al.*, 2013).

47  
48  
49 268

## 50 51 269 **RESULTS**

52  
53  
54 270

### 55 56 271 **Insect monitoring**

1  
2  
3 272 A total of 315 specimens belonging to seven species, previously reported as known or  
4  
5 273 putative phytoplasma vectors, were captured on Rebell and Aeroxon sticky traps positioned inside,  
6  
7 274 on the borders and outside the test vineyard, and by beating tray method. The captured specimens  
8  
9 275 belonged to two Cixiidae species, *Hyalesthes obsoletus* Signoret (15) and *Reptalus cuspidatus*  
10  
11 276 (Fieber) (6), and to five Cicadellidae species, *Orientus ishidae* (245 specimens), *Scaphoideus*  
12  
13 277 *titanus* (45), *Thamnotettix dilutior* (Kirschbaum) (2), *Fieberiella florii* (Stål) (1), and *Anoplotettix*  
14  
15 278 *fuscovenosus* (Ferrari) (1), all belonging to the Deltocephalinae subfamily (Table 1).

16  
17  
18 279 Captures obtained by beating tray method and Aeroxon traps evidenced that three insect  
19  
20 280 species (*S. titanus*, *O. ishidae*, and *H. obsoletus*) are present on the grapevine canopy in the test  
21  
22 281 vineyard. In detail, it was possible to observe on grapevine canopy (i) three nymphs (two at May 27  
23  
24 282 and one at July 1<sup>st</sup>) and 13 adults (two captured by beating tray in August and 11 by Aeroxon trap in  
25  
26 283 August/September) of *S. titanus*, evidencing a sporadic presence of the insect throughout the season  
27  
28 284 (Figure 2A); (ii) 22 adults of *O. ishidae* captured by Aeroxon traps, evidencing a sporadic presence  
29  
30 285 of the insect during the summer and a peak of captures in October (Figure 2A); (iii) three adults of  
31  
32 286 *H. obsoletus* captured in July by Aeroxon traps. Furthermore, from June 21 to July 30, beating tray  
33  
34 287 method allowed the capture of four nymphs of *O. ishidae* from willow (two positive samples out of  
35  
36 288 six), and two nymphs and three adults of *O. ishidae* from hazel tree (three positive samples out of  
37  
38 289 six).

39  
40  
41  
42  
43 290 Captures obtained by Rebell traps highlighted that (i) *O. ishidae* adult population was  
44  
45 291 prevalently spread on the borders (111 out of 214) and on the surrounding vegetation (96 out of  
46  
47 292 214); only seven specimens were captured inside the test vineyard; (ii) the same trend was observed  
48  
49 293 for *S. titanus* (28 specimens out of 29 on the borders and outside the vineyard; only one specimen  
50  
51 294 captured inside the vineyard) (Table 1). Moreover, the obtained results evidenced that specimens of  
52  
53 295 *O. ishidae* (167 out of 214) and *S. titanus* (28 out of 29) were mainly captured on four traps located  
54  
55 296 closely together at the south-western limit of the experimental area. Two traps (B3, B4) were on the  
56  
57  
58  
59  
60

1  
2  
3 297 borders of the test vineyard (103 *O. ishidae* and 7 *S. titanus*), and two (W8, W9) were on the  
4  
5 298 surrounding vegetation, inside the forest where several hazel trees are present (64 *O. ishidae* and 21  
6  
7 299 *S. titanus*) (Figure 1, Table 1). Concerning other planthoppers and leafhoppers, *R. cuspidatus*, *T.*  
8  
9 300 *dilutior* and the majority (9 out of 15) of *H. obsoletus* specimens were captured by traps positioned  
10  
11 301 on the vineyard borders. Moreover, unique specimens of *F. florii* and *A. fuscovenosus* were  
12  
13 302 captured inside and outside the vineyard, respectively (Figure 1, Table 1).

16 303         Given the numbers of captured specimens per species, it was possible to investigate only the  
17  
18 304 flight dynamic of *O. ishidae* and the pattern of *S. titanus* occurrence in the experimental area. In  
19  
20 305 order to compare the results, data were reported as average weekly captures. Inside and on the  
21  
22 306 borders of the test vineyard, dispersing *O. ishidae* adults were captured continuously by Rebell traps  
23  
24 307 (V1 to V6 and B1 to B8) from the end of June until October, with the presence of two periods of  
25  
26 308 higher catches at (i) end of July / beginning of August and (ii) mid October, respectively. The  
27  
28 309 presence of *S. titanus* was negligible with sporadic catches from August to October (Figure 2B).  
29  
30 310 Outside the vineyard, Rebell traps (W1 to W9) confirmed the pattern of *O. ishidae* captures, as  
31  
32 311 observed inside the vineyard from July to September, but were unable to confirm the presence of  
33  
34 312 dispersing adults during October (Figure 2C). The presence of *S. titanus* outside the vineyard was  
35  
36 313 scarce, with low but regular captures from mid September to mid October (Figure 2C).  
37  
38  
39  
40  
41  
42

### 315 **FDp and related strain identification**

45 316         Real-time PCR assays performed to identify FDp and related strains amplified DNA  
46  
47 317 extracted from periwinkle plants infected by FDp strains FD92 and FD70 showed an average Ct  
48  
49 318 (threshold cycle) of 18.05 and 18.88, respectively. No amplification was observed for periwinkle  
50  
51 319 plants infected by phytoplasma strains STOL and AY1, or reaction mixtures without DNA. Only  
52  
53 320 PCR products amplified from insects and plants, with an average Ct < 37, were associated with the  
54  
55 321 presence of FDp and related strains. qPCR assays were carried out on total nucleic acids extracted  
56  
57  
58  
59  
60

1  
2  
3 322 from 119 plant leaf samples and 146 insect pools. FDp and related strain distribution among tested  
4  
5 323 plants is as follows: 70 out of 74 (95%) grapevine samples, 11 out of 31 (35%) hazel tree samples, 3  
6  
7 324 out of 10 (30%) willow samples, and 3 out of 4 (75%) *Clematis vitalba* samples (Table 2).

8  
9 325 Out of 301 leafhoppers captured by sticky traps, 274 specimens were grouped in 146 pools  
10  
11 326 for further molecular analyses. FDp and related strains were detected in 30 out of 146 insect pools  
12  
13 327 (21%). In particular, it was identified in 25 out of 97 (26%) *O. ishidae* pools, 2 out of 29 (7%) *S.*  
14  
15 328 *titanus* pools, 1 out of 11 (9%) *H. obsoletus* pools and 2 out of 2 (100%) *T. dilutior* pools. All  
16  
17 329 examined pools of *F. florii*, *R. cuspidatus*, and *A. fuscovenosus* were negative (Table 1).

18  
19  
20  
21 330

### 22 331 **FDp and related strain typing**

23  
24  
25 332 Nested PCRs carried out using the primer pair FD9f6/MAPr2 amplified DNA from all the  
26  
27 333 plant and insect samples infected by FDp and related strains, with the exception of 5 hazel leaf  
28  
29 334 samples (out of 11) and 2 willow leaf samples (out of 3) (Table 1 and 2). Enzymatic digestions of  
30  
31 335 FD9f6/MAPr2 amplicons with the enzymes *AluI* and *Eco72I* produced three restriction profiles  
32  
33 336 among FDp and related strains identified in the present study (Figure 3). The *AluI-Eco72I*-RFLP  
34  
35 337 patterns obtained in this study were compared to those described in the literature (*i.e.* FD1, FD2 and  
36  
37 338 FD3; Arnaud *et al.*, 2007). Among the FDp and related strains identified from plants and insects in  
38  
39 339 the test vineyard, the *map*-type FD2 was prevalent (89 out of 108, or 82%), while *map*-types FD1  
40  
41 340 and FD3 were identified in 13 (12%) and 6 (6%) instances, respectively (Table 1 and 2). FD2 was  
42  
43 341 detected in grapevine (68 plants), hazel trees (1 plant), *O. ishidae* (18 pools) and *T. dilutior* (2  
44  
45 342 pools). FD1 was identified in grapevine (2 plants), hazel tree (3 plants), willow (1 plant), *O. ishidae*  
46  
47 343 (5 pools), and *S. titanus* (2 pools). FD3 was identified in hazel trees (2 plants), *C. vitalba* (3 plants),  
48  
49 344 *O. ishidae* (2 pools), and *H. obsoletus* (1 pool) (Table 1 and 2).

50  
51  
52 345 Nucleotide sequence analyses, performed on 22 FDp and related strains and 16SrV group  
53  
54 346 phytoplasma strains selected on the basis of both their *map*-type and the phytoplasma hosts (Table  
55  
56  
57  
58  
59  
60

1  
2  
3 347 3), confirmed their assignment to *map*-types as determined by the RFLP assays. As a matter of fact,  
4  
5 348 (i) FDp and related strains with an FD1 RFLP profile shared best *map* gene sequence identity (99.8  
6  
7 349 to 100%) with strain FD70, a FD1 reference strain; (ii) FDp and related strains with an FD2 RFLP  
8  
9 350 profile shared best *map* gene sequence identity (100%) with strain V00-SP5, a FD2 reference strain;  
10  
11 351 (iii) FDp-related and 16SrV group phytoplasma strains with an FD3 RFLP profile shared best *map*  
12  
13 352 gene sequence identity (96.9 to 99.6%) with strain VI04-Toscana1, a FD3 reference strain (Table  
14  
15 353 S2).

16  
17  
18 354 Clustering in a phylogenetic tree built using the *map* gene nucleotide sequences isolated  
19  
20 355 from plant and insect hosts in this study clearly confirms the attribution of the FDp and related  
21  
22 356 strains and 16SrV group phytoplasma strains to *map*-types FD1, FD2 and FD3 (Figure 4).  
23  
24  
25 357

## 26 27 358 **DISCUSSION**

28  
29 359 Due to the mandatory application of insecticides, the population of the FDp vector *S. titanus*  
30  
31 360 within the test vineyard was very low, as shown by the very small number of specimens (three  
32  
33 361 nymphs and two adults) collected by the beating tray method during the growing season and  
34  
35 362 corroborated by the few captures of adults on the Rebell and Aeroxon yellow sticky trap placed  
36  
37 363 inside the test vineyard. This trend confirmed previous evidences by Trivellone *et al.* (2016),  
38  
39 364 reporting high *S. titanus* populations in Swiss localities where its control was not mandatory, and  
40  
41 365 low abundances in vineyards treated with insecticides twice during the growing season. Previous  
42  
43 366 works have demonstrated that *S. titanus* can survive and feed on several plants (Trivellone *et al.*,  
44  
45 367 2013; Chuche & Thiery, 2014), develop on elm (Gibson, 1973) and lay eggs on elder (Gargani *et*  
46  
47 368 *al.*, 2013). In the present study, the majority of *S. titanus* specimens (28 out of 40) was captured on  
48  
49 369 traps B3, B4, W8 and W9 (Figure 1), placed on the borders and outside the test vineyard, close to a  
50  
51 370 wood of mostly hazel and willow trees, with no wild grapevines to be found. Moreover, the two  
52  
53 371 FDp positive *S. titanus* pools, captured on traps W8 and B4, were infected by the *map*-type FD1  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 372 (Table 1). FD1 was never found before in Switzerland neither in vineyards nor in *S. titanus*, but was  
4  
5 373 the most frequent *map*-type on hazel and willow trees in the experimental area, and affected a low  
6  
7 374 minority (two out of 70) of FDP-infected grapevine plants in the test vineyard (Table 2). On the  
8  
9 375 other hand, *S. titanus* was not captured by beating tray from hazel and willow trees in the  
10  
11 376 surroundings of the test vineyard. Based on such evidences, it is reasonable to raise the question if  
12  
13 377 *S. titanus* can live, but not develop, on wild plants long enough to acquire FDP and related strains.  
14  
15

16 378 Despite the compulsory destruction of grapevines with GY typical symptoms and the  
17  
18 379 insecticide treatments, applied at two distinct time points, 74 symptomatic grapevines (3.4%) were  
19  
20 380 identified in the test vineyard in 2014, the same levels as observed in the previous year by  
21  
22 381 Agroscope, suggesting a stable infection rate through the years. This, together with the low  
23  
24 382 populations of *S. titanus* inside and outside the vineyard, as well as the absence of *D. europaea* and  
25  
26 383 *O. alni* [insects known to transmit FDP and related strains from wild plants to grapevine (Chuche &  
27  
28 384 Thiery, 2014)], points at a scenario, in which FDP could be vectored by other phloem-feeding  
29  
30 385 insects. Interestingly, this study revealed an abundant population of the leafhopper *O. ishidae*,  
31  
32 386 recently found able to transmit 16SrV phytoplasmas to vines (Lessio *et al.*, 2016). *O. ishidae* was  
33  
34 387 found to harbor FD related phytoplasma in high percentage (26% of the examined pools),  
35  
36 388 prevalently at the borders of the test vineyard and within the wood. Our data confirms the *O.*  
37  
38 389 *ishidae* contamination rates by FDP related strains recently reported for Canton Ticino, Slovenia  
39  
40 390 and Italy (Mehle *et al.*, 2010; Gaffuri *et al.*, 2011; Trivellone *et al.*, 2016) and its presence in  
41  
42 391 vineyards surrounded by woods (Trivellone *et al.*, 2016).  
43  
44  
45  
46

47 392 Previous studies report *O. ishidae* as polyphagous, mainly on woody plants (Hamilton,  
48  
49 393 1985; Nickel, 2010; Mehle *et al.*, 2011; <https://gd.eppo.int/reporting/article-4763>). Here, as  
50  
51 394 previously reported, nymphs and adults of *O. ishidae* were collected from hazel and willow trees by  
52  
53 395 beating tray method; moreover, the highest density of *O. ishidae* (167 specimens) was observed on  
54  
55 396 traps B3, B4, W8 and W9, which were placed in the southwestern side of the experimental area, at  
56  
57  
58  
59  
60

1  
2  
3 397 the border of the test vineyard and within a wood of mostly hazel (*Corylus avellana*), already  
4  
5 398 known as a plant host for 'Candidatus Phytoplasma asteris' (Cieślińska & Kowalik, 2011), and  
6  
7 399 willow (*Salix* spp.) trees, found infected by FDp related strains (30% of the analyzed samples) for  
8  
9 400 the first time. Moreover, beating tray captures link *O. ishidae* nymphs and adults with hazel and  
10  
11 401 willow, species found infected by FDp related strains in the present study, confirming the role of  
12  
13 402 these plant species as hosts for *O. ishidae* (Lessio *et al.*, 2016). On the other hand, *O. ishidae* was  
14  
15 403 not captured by beating tray on *Clematis vitalba*, a plant rarely found in the examined vineyard and  
16  
17 404 previously reported to host FDp (Filippin *et al.*, 2009). Even more interestingly, even if no *O.*  
18  
19 405 *ishidae* specimen was captured by beating tray on grapevine plants, 22 specimens were captured  
20  
21 406 inside the vineyard, mainly on horizontal Aeraxon traps, which are known to catch insects living on  
22  
23 407 the vine canopies, suggesting that *O. ishidae* adults could be present on grapevine for extended  
24  
25 408 periods.  
26  
27  
28

29 Intriguingly, PCR tests revealed for the first time the presence of 16SrV group phytoplasma  
30  
31 410 in the Cixiidae *Hyalesthes obsoletus*, the vector of 'Candidatus Phytoplasma solani', associated with  
32  
33 411 Bois noir (Quaglino *et al.*, 2013), and FDp related strain in the Cicadellidae *Thamnotettix dilutior*,  
34  
35 412 largely reported as associated with vineyard ecosystems (Bosco *et al.*, 1997; Sanna *et al.*, 2016).  
36  
37 413 Molecular analyses showed that (i) *H. obsoletus* hosts a 16SrV group phytoplasma strain (Ho1),  
38  
39 414 characterized by the typical RFLP profile of the *map*-type FD3, but very divergent from FDp and  
40  
41 415 related strains of *map*-type FD3 based on sequence identity (from 96.9 to 97.3%); in fact, *map* gene  
42  
43 416 nucleotide sequence of strain Ho1 has 15 SNPs compared to the most closely related strain (FDp  
44  
45 417 strain FG1) in FD3 cluster (data not shown); due to the high polyphagy of *H. obsoletus*, it is  
46  
47 418 reasonable to hypothesize that 16SrV group phytoplasma strain Ho1 was acquired by occasional  
48  
49 419 feeding of the insect on infected source plant(s), which should be identified in further studies; (ii) *T.*  
50  
51 420 *dilutior* specimens host FDp related strains indistinguishable from *map*-type FD2 phytoplasma  
52  
53 421 strains (sequence identity 100%) identified in grapevine, hazel, and *O. ishidae* in the test vineyard  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 422 (Figure 4, Table S2). These evidences, along with the occurrence of a low population in the test  
4  
5 423 vineyard, suggest that *T. dilutior*, if confirmed as vector, could play a marginal role in FDp  
6  
7 424 transmission to grapevine.  
8

9  
10 425 Together with ongoing field surveys and molecular analyses, the data of the present study,  
11  
12 426 based on the detection of FDp and related strains in multiple plants and insects inside and outside  
13  
14 427 the test vineyard, (i) confirm the presence in vineyard agro-ecosystems of different leafhopper and  
15  
16 428 plant species infected with FDp and related strains, (ii) confirm the high FDp related strains-  
17  
18 429 infection rate in *O. ishidae* observed in some European Countries (Mehle *et al.*, 2010, 2011; Gaffuri  
19  
20 430 *et al.*, 2011; Koczor *et al.*, 2013; Trivellone *et al.*, 2016), and (iii) show for the first time the high  
21  
22 431 FDp related strain infection rate in hazel, a host plant of *O. ishidae* (Nickel *et al.*, 2010; Lessio *et*  
23  
24 432 *al.*, 2016), very common and abundant in Swiss vineyard agro-ecosystems. It is therefore  
25  
26 433 reasonable to hypothesize that the ecological cycle of FDp may be related not exclusively to the  
27  
28 434 grapevine-specific feeding diet of *S. titanus*, but also to that of other insect vector(s) and/or plant  
29  
30 435 host(s). Bressan *et al.* (2006) tested the specificity of vector transmission of FDp on 15  
31  
32 436 Auchenorrhyncha species, collected in vineyard agro-ecosystem but not developing on grapevine,  
33  
34 437 by abdominal injection of FDp. They demonstrated that three leafhopper species, all belonging to  
35  
36 438 the subfamily Deltocephalinae, transmit FDp. These data suggest that the vector competency of  
37  
38 439 FDp transmission could be widespread and opens the question if in a viticultural area the  
39  
40 440 range/number of effective insect vector species could be determined by the availability of suitable  
41  
42 441 host plant species in a system where FDp strains flow among many different hosts, inside and  
43  
44 442 outside the vineyards.  
45  
46  
47  
48

49  
50 443 During the last decades, the Asian Cicadellidae *O. ishidae* has been reported from several  
51  
52 444 European Countries. This species has also been introduced in North America during the last  
53  
54 445 century. *O. ishidae* is a sap-feeder; although some damage (i.e. uniform browning of apple and  
55  
56 446 hazel foliage) has occasionally been reported in the USA, it is not considered to be a major pest of  
57  
58  
59  
60

1  
2  
3 447 cultivated plants. The main concern raised by the introduction of *O. ishidae* in Europe is the  
4  
5 448 possible transmission of phytoplasma diseases (EPPO, 2015). In fact, experiments conducted in the  
6  
7 449 USA have shown that *O. ishidae* can transmit ‘*Candidatus* Phytoplasma pruni’ to celery plants  
8  
9 450 (Rosenberg *et al.*, 1978). Moreover, the present work and recent studies conducted in several  
10  
11 451 European Countries detected the presence of FDp related strains in specimens of *O. ishidae*  
12  
13 452 collected from diseased vineyards (EPPO, 2015). Interestingly, latest findings revealed the  
14  
15 453 capability of *O. ishidae* to transmit 16SrV phytoplasmas to grapevine (Lessio *et al.*, 2016).  
16  
17

18 454 In this framework, based on the results obtained in the present study, it is reasonable to  
19  
20 455 assume that the settlement of *O. ishidae* in Canton Ticino at the beginning of the 21<sup>st</sup> century could  
21  
22 456 have contributed to the persistence of FDp, despite the application of mandatory insecticide  
23  
24 457 treatments, by transmitting FDp from hazel and willow to grapevine, in a way similar to that  
25  
26 458 hypothesized for *O. alni* (Arnaud *et al.*, 2007; Durante *et al.*, 2012).  
27  
28

29 459 In the last years, the molecular characterization of phytoplasma species showed molecular  
30  
31 460 diversity, representing ecologically separated populations. The investigation of the genetic diversity  
32  
33 461 among phytoplasmas associated with several diseases allows the identification of strain-specific  
34  
35 462 molecular markers, which are useful to improve the understanding of complex phytoplasma  
36  
37 463 ecologies (Davis *et al.*, 2013; Quaglino *et al.*, 2013). Indeed, among grapevine yellows, the  
38  
39 464 knowledge of the biological complexity of the Bois noir disease was improved through the  
40  
41 465 application of a molecular epidemiology approach (Kostadinovska *et al.*, 2014; Kosovac *et al.*,  
42  
43 466 2016; Murolo & Romanazzi, 2015; Quaglino *et al.*, 2016). Concerning FD, molecular markers were  
44  
45 467 produced to distinguish FDp strains in different ecological niches (Martini *et al.*, 2002; Botti &  
46  
47 468 Bertaccini, 2007; Filippin *et al.*, 2009; Quaglino *et al.*, 2010; Malembic-Maher *et al.*, 2011; Durante  
48  
49 469 *et al.*, 2012). The most promising candidate for typing closely related strains within the taxonomic  
50  
51 470 group 16SrV is the *map* gene. It distinguishes three *map*-types (FD1, FD2 and FD3), which are  
52  
53 471 associated with distinct geographic areas (Arnaud *et al.*, 2007) and alder yellows strains in Germany  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 472 (Holz *et al.*, 2015). In this study, data from the molecular characterization and phylogenetic  
4  
5 473 analyses of FDp and related strains reveal the great prevalence of the *map*-type FD2 in grapevines  
6  
7 474 (97%) and in *O. ishidae* pools (72%); FD2 was found also in hazel and in *T. dilutior*, but not in *S.*  
8  
9 475 *titanus*. The *map*-type FD1, identified in few grapevines (3%), was found also in *O. ishidae*, *S.*  
10  
11 476 *titanus*, *C. avellana*, and *Salix* spp.. The *map*-type FD3, not found in grapevine, was identified in *O.*  
12  
13 477 *ishidae*, *C. avellana*, and *C. vitalba*. Together, these data show that in the test vineyard FD is  
14  
15  
16 478 associated prevalently with phytoplasma strains of *map*-type FD2.

17  
18 479 Based on the data obtained in this and in a previous study (Lessio *et al.*, 2016), it is  
19  
20 480 reasonable to hypothesize that in the test vineyard the ecology of FDp is related to grapevine-  
21  
22 481 specific cycle of *S. titanus*, the cycle of other polyphagous vectors (e.g. *O. ishidae*), and the  
23  
24 482 presence of other host plants of FDp and related strains (e.g. hazel and willow). Further studies will  
25  
26 483 be needed to prove the role of *O. ishidae* in FD epidemiology. Considering the new scenario  
27  
28 484 describing the FD ecology as an open system, in which the landscape composition can influence the  
29  
30 485 presence of FDp vectors and host plants impacting on FD epidemics, it appears suitable to re-  
31  
32 486 evaluate the disease control strategies.

33  
34  
35  
36 487

## 37 38 488 **ACKNOWLEDGEMENTS**

39  
40 489 We gratefully thank Prof. Alberto Alma (University of Turin) for sharing important information on  
41  
42 490 *O. ishidae* obtained in his own researches, and Dr. Valeria Trivellone (Research Station Agroscope  
43  
44 491 Changins, Cadenazzo, Switzerland) for the technical assistance in insect identification. Moreover,  
45  
46 492 we are grateful to Mister Piercarlo Saglini, the owner of the test vineyard.

47  
48  
49 493

## 50 51 494 **REFERENCES**

- 1  
2  
3 495 1. Alma A., Bosco D., Danielli A., Bertaccini A., Vibio M., Arzone A. (1997) Identification of  
4  
5 496 phytoplasmas in eggs, nymphs and adults of *Scaphoideus titanus* Ball reared on healthy  
6  
7 497 plants. *Insect Molecular Biology*, **6**, 115–121.
- 8  
9 498 2. Angelini E., Clair D., Borgo M., Bertaccini A., Boudon-Padieu E. (2001) Flavescence  
10  
11 499 dorée in France and Italy. Occurrence of closely related phytoplasma isolates and their near  
12  
13 500 relationships to Palatinate Grapevine yellows and an Alder yellows phytoplasma. *Vitis*, **40**,  
14  
15 501 79–86.
- 16  
17  
18 502 3. Anonymous (2015) *Orientus ishidae*: a potential phytoplasma vector spreading in the EPPO  
19  
20 503 region. *EPPO Reporting Service*, **5**, article 2015/098.
- 21  
22  
23 504 4. Arnaud G., Malembic-Maher S., Salar P., Bonnet P., Maixner M., Marcone C., Boudon-  
24  
25 505 Padieu E., Foissac X. (2007) Multilocus sequence typing confirms the close genetic inter-  
26  
27 506 relatedness between three distinct flavescence dorée phytoplasma strain clusters and group  
28  
29 507 16SrV phytoplasmas infecting grapevine and alder in Europe. *Applied and Environmental*  
30  
31 508 *Microbiology*, **73**, 4001–4010.
- 32  
33  
34 509 5. Belli G., Bianco P.A., Conti M. (2010) Grapevine yellows: past, present and future. *Journal*  
35  
36 510 *of Plant Pathology*, **92**, 303–326.
- 37  
38 511 6. Bosco D., Alma A., Arzone A. (1997) Studies on population dynamics and spatial  
39  
40 512 distribution of leafhoppers in vineyards (Homoptera:Cicadellidae). *Annals of Applied*  
41  
42 513 *Biology*, **130**, 1–11.
- 43  
44  
45 514 7. Botti S., Bertaccini A. (2007) Grapevine yellows in Northern Italy: molecular identification  
46  
47 515 of Flavescence doree phytoplasma strains and of Bois Noir phytoplasmas. *Journal of*  
48  
49 516 *Applied Microbiology*, **103**, 2325–2330.
- 50  
51  
52 517 8. Bressan A., Clair D., Sémétey O., Boudon-Padieu E. (2006) Insect injection and artificial  
53  
54 518 feeding bioassays to test the vector specificity of Flavescence dorée phytoplasma.  
55  
56 519 *Phytopathology*, **96**, 790–796.

- 1  
2  
3 520 9. Caudwell A., Gianotti J., Kuszala C., Larrue J. (1971) Etude du rôle de particules de type  
4  
5 521 mycoplasme dans l'étiologie de la Flavescence dorée de la vigne. Examen cytologique des  
6  
7 522 plants malades et des cicadelles infectieuses. *Annales de Phytopathologie*, **3**, 107–203.  
8  
9 523 10. Caudwell A., Kuszala C., Bachelier J.C., Larrue J. (1970) Transmission de la Flavescence  
10  
11 524 dorée de la vigne aux plantes herbacées par l'allongement du temps d'utilisation de la  
12  
13 525 cicadelle *Scaphoideus littoralis* Ball et l'étude de sa survie sur un grand nombre d'espèces  
14  
15 526 végétales. *Annales de Phytopathologie*, **2**, 415–428.  
16  
17 527 11. Chuche C., Thiery D. (2014) Biology and ecology of the Flavescence dorée vector  
18  
19 528 *Scaphoideus titanus*: a review. *Agronomy for Sustainable Development*, **34**, 381–403.  
20  
21 529 12. Cieślińska M., Kowalik B. (2011) Detection and molecular characterization of 'Candidatus  
22  
23 530 Phytoplasma asteris' in European hazel (*Corylus avellana*) in Poland. *Journal of*  
24  
25 531 *Phytopathology*, **159**, 585–588.  
26  
27 532 13. Davis R.E., Dally E.L. (2001) Revised subgroup classification of group 16SrV phytoplasmas  
28  
29 533 and placement of flavescence dorée-associated phytoplasmas in two distinct subgroups.  
30  
31 534 *Plant Disease*, **85**, 790–797.  
32  
33 535 14. Davis R.E., Zhao Y., Dally E., Lee I.-M., Jomantiene R., Douglas S.M. (2013) 'Candidatus  
34  
35 536 Phytoplasma pruni', a novel taxon associated with X-disease of stone fruits, *Prunus* spp.:  
36  
37 537 multilocus characterization based on 16S rRNA, *secY*, and ribosomal protein genes.  
38  
39 538 *International Journal of Systematic and Evolutionary Microbiology*, **63**, 766–776.  
40  
41 539 15. Durante G., Casati P., Clair D., Quaglino F., Bulgari D., Boudon-Padieu E., Bianco P.A.  
42  
43 540 (2012) Sequence analyses of S10-*spc* operon among 16SrV group phytoplasmas:  
44  
45 541 phylogenetic relationships and identification of discriminating single nucleotide  
46  
47 542 polymorphisms. *Annals of Applied Biology*, **161**, 234–246.  
48  
49 543 16. European and Mediterranean Plant Protection Organization/Commonwealth Agricultural  
50  
51 544 Bureau International. (1996) Grapevine flavescence dorée phytoplasma. In I.M. Smith,

- 1  
2  
3 545 McNamara D.G., Scott P.R., Holderness M. (ed.), Quarantine pests for Europe, 2<sup>nd</sup> edition.  
4  
5 546 Commonwealth Agricultural Bureau International, Wallingford, United Kingdom, pp. 1013–  
6  
7 547 1021.  
8  
9  
10 548 17. European and Mediterranean Plant Protection Organization (2015) *Orientus ishidae*: a  
11  
12 549 potential phytoplasma vector spreading in the EPPO region. *In* EPPO Reporting Service  
13  
14 550 2015 no. 5 - Pest & Diseases, **2015/98**, 8–10.  
15  
16 551 18. European Food Safety Authority (2014). Scientific Opinion on pest categorisation of  
17  
18 552 Grapevine Flavescence dorée. EFSA Panel on Plant Health (PLH). *EFSA Journal*, **12**, 3851.  
19  
20 553 19. Filippin L., Jović J., Cvrković D., Forte V., Clair D., Toševski I., Boudon-Padieu E., Borgo  
21  
22 554 M., Angelini E. (2009) Molecular characteristics of phytoplasmas associated with  
23  
24 555 Flavescence dorée in clematis and grapevine and preliminary results on the role of  
25  
26 556 *Dictyophara europaea* as a vector. *Plant Pathology*, **58**, 826–837.  
27  
28  
29 557 20. Foissac X., Maixner M. (2013) Spread of grapevine phytoplasma diseases. In Torres E.,  
30  
31 558 Laviña A., Jarausch W., Bertaccini A. (ed.), New perspectives in phytoplasma disease  
32  
33 559 management, Book of Abstracts of COST Action FA0807 Workshop; March 22, 2013,  
34  
35 560 Barcelona, Spain.  
36  
37  
38 561 21. Gaffuri F., Sacchi S., Cavagna B. (2011) First detection of the mosaic leafhopper, *Orientus*  
39  
40 562 *ishidae*, in northern Italian vineyards infected by the flavescence dorée phytoplasma. *New*  
41  
42 563 *Disease Reports*, **24**, 22.  
43  
44  
45 564 22. Gargani E., Torrini G., Caradonna S., Bagnoli B. (2013) *Scaphoideus titanus* and *Metcalfa*  
46  
47 565 *pruinosa* egg distribution on different woody parts of Kober 5BB grapevine. *IOBC/wprs*  
48  
49 566 *Bulletin*, **85**, 37–42.  
50  
51  
52 567 23. Gibson L.P. (1973) An annotated list of the Cicadellidae and Fulgoridae of elm. *USDA*  
53  
54 568 *Forest Service Research Paper*, **NE-278**, 1–5.  
55  
56  
57  
58  
59  
60



- 1  
2  
3 569 24. Günthart H., Mühlethaler R., Lauterer P. (2004) Für die Schweiz neue Zikadenarten und  
4  
5 570 Ergänzungen zu bereits bekannten Arten (Hemiptera Auchenorrhyncha). *Mitteilungen der*  
6  
7 571 *Entomologischen Gesellschaft Basel*, **54**, 150–160.
- 8  
9 572 25. Hamilton K.G.A. (1985) Leafhoppers of ornamental and fruit trees in Canada. *Agriculture*  
10  
11 573 *Canada Publication*, **1779E/F**, 1–71.
- 12  
13 574 26. Holz S., Duduk B., Buttner C., Kube M. (2016) Genetic variability of Alder yellows  
14  
15 575 phytoplasma in *Alnus glutinosa* in its natural Spreewald habitat. *Forest Pathology*, **46**, 11–  
16  
17 576 21.
- 18  
19 577 27. Jermini M., D'Adda G., Baumgärtner J., Lozzia G.C., Baillod M. (1992a) Nombre des  
20  
21 578 pièges englués nécessaires pour estimer la densité relative des populations de la cicadelle  
22  
23 579 *Scaphoideus titanus* Ball en vignobles. *Bollettino di Zoologia Agraria e Bachicoltura*, **25**,  
24  
25 580 91–102.
- 26  
27 581 28. Jermini M., Rossi A., Baillod M. (1992b) Etude du piégeage de la cicadelle *Scaphoideus*  
28  
29 582 *titanus* Ball à l'aide de pièges jaunes. *Revue Suisse de Viticulture, Arboriculture,*  
30  
31 583 *Horticulture*, **24**, 235–239.
- 32  
33 584 29. Jermini M., Schaerer S., Johnston H., Colombi L., Marazzi C. (2014) Dix ans de  
34  
35 585 flavescence dorée au Tessin. *Revue Suisse de Viticulture, Arboriculture, Horticulture*, **46**,  
36  
37 586 222–229.
- 38  
39 587 30. Koczor S., Bagarus A.K., Karap A.K., Varga Á., Orosz A. (2013) A rapidly spreading  
40  
41 588 potential pest, *Orientus ishidae* identified in Hungary. *Bulletin of Insectology*, **66**, 221–224.
- 42  
43 589 31. Kosovac A., Radonjić S., Hrnčić S., Krstić O., Toševski I., Jović J. (2016) Molecular tracing  
44  
45 590 of the transmission routes of bois noir in Mediterranean vineyards of Montenegro and  
46  
47 591 experimental evidence for the epidemiological role of *Vitex agnus-castus* (Lamiaceae) and  
48  
49 592 associated *Hyalesthes obsoletus* (Cixiidae). *Plant Pathology*, **65**, 285–298.
- 50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 593 32. Kostadinovska E., Quaglino F., Mitrev S., Casati P., Bulgari D., Bianco P.A. (2014)  
4  
5 594 Multiple gene analyses identify distinct “bois noir” phytoplasma genotypes in the Republic  
6  
7 595 of Macedonia. *Phytopathologia Mediterranea*, **53**, 491–501.  
8  
9 596 33. Lee I.-M., Gundersen D.E., Davis R.E., Bartoszik I.M. (1998) Revised classification scheme  
10  
11 597 of phytoplasmas based on RFLP analyses of 16S rRNA and ribosomal protein gene  
12  
13 598 sequences. *International Journal of Bacteriology*, **48**, 1153–1169.  
14  
15 599 34. Lee I.-M., Martini M., Marcone C., Zhu S.F. (2004). Classification of phytoplasma strains  
16  
17 600 in the elm yellows group (16SrV) and proposal of ‘*Candidatus Phytoplasma ulmi*’ for the  
18  
19 601 phytoplasma associated with elm yellows. *International Journal of Systematic and*  
20  
21 602 *Evolutionary Microbiology*, **54**, 337–347.  
22  
23 603 35. Lessio F., Picciau L., Gonella E., Mandrioli M., Tota F., Alma A. (2016) The mosaic  
24  
25 604 leafhopper *Orientus ishidae*: host plants, spatial distribution, infectivity, and transmission of  
26  
27 605 16SrV phytoplasmas to vines. *Bulletin of Insectology*, **69**, 277–289.  
28  
29 606 36. Linder C., Cavadini M., Schaerer S. (2014) *Dictyophara europaea*: un vecteur potentiel de  
30  
31 607 la flavescence dorée en Suisse? *Revue Suisse de Viticulture, Arboriculture, Horticulture*, **46**,  
32  
33 608 216–219.  
34  
35 609 37. Maixner M., Reinert W. (1999) *Oncopsis alni* (Schrank) (Auchenorrhyncha:Cicadellidae) as  
36  
37 610 a vector of the alder yellows phytoplasma of *Alnus glutinosa* (L.) Gaertn. *European Journal*  
38  
39 611 *of Plant Pathology*, **105**, 87–94.  
40  
41 612 38. Maixner M., Reinert W., Darimont H. (2000) Transmission of grapevine yellows by  
42  
43 613 *Oncopsis alni* (Schrank) (Auchenorrhyncha: Macropsinae). *Vitis*, **39**, 83–84.  
44  
45 614 39. Malembic-Maher S., Salar P., Filippin L., Carle P., Angelini E., Foissac X. (2011) Genetic  
46  
47 615 diversity of European phytoplasmas of the 16SrV taxonomic group and proposal of  
48  
49 616 ‘*Candidatus Phytoplasma rubi*’. *International Journal of Systematic and Evolutionary*  
50  
51 617 *Microbiology*, **61**, 2129–2134.  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 618 40. Martini M., Botti S., Marcone C., Marzachi C., Casati P., Bianco P.A., Benedetti R.,  
4  
5 619 Bertaccini A. (2002) Genetic variability among flavescence dorée from different origins in  
6  
7 620 Italy and France. *Molecular and Cellular Probes*, **16**, 197–208.
- 8  
9 621 41. Mehle N., Ravnikar M., Seljak G., Knapiè V., Dermastia M. (2011) The most widespread  
10  
11 622 phytoplasmas, vectors and measures for disease control in Slovenia. *Phytopathogenic*  
12  
13 623 *Mollicutes*, **1**, 65–76.
- 14  
15 624 42. Mehle N., Rupar M., Seljak G., Ravnikar M., Dermastia M. (2011) Molecular diversity of  
16  
17 625 ‘flavescence dorée’ phytoplasma strains in Slovenia. *Bulletin of Insectology*, **64**, S29– S30.
- 18  
19 626 43. Mehle N., Seljak G., Rupar M., Ravnikar M., Dermastia M. (2010) The first detection of a  
20  
21 627 phytoplasma from the 16SrV (Elm yellows) group in the mosaic leafhopper *Orientalis*  
22  
23 628 *ishidae*. *New Disease Reports*, **22**, 11.
- 24  
25 629 44. Murolo S., Romanazzi G. (2015) In-vineyard population structure of ‘*Candidatus*  
26  
27 630 *Phytoplasma solani*’ using multilocus sequence typing analysis. *Infection, Genetics and*  
28  
29 631 *Evolution*, **31**, 221– 230.
- 30  
31 632 45. Navrátil M., Šafařová D., Válová P., Fránová J., Šimková M. (2009) Phytoplasma associated  
32  
33 633 with witches’-broom disease of *Ulmus minor* MILL. in the Czech Republic: electron  
34  
35 634 microscopy and molecular characterization. *Folia Microbiologica*, **54**, 37–42.
- 36  
37 635 46. Nickel H. (2010) First addendum to the Leafhoppers and Planthoppers of Germany  
38  
39 636 (Hemiptera: Auchenorrhyncha). *Cicadina*, **11**, 107–122.
- 40  
41 637 47. Plavec J., Križanac I., Budinščak Ž., Škorić D., Šeruga Musić M. (2015) A case study of FD  
42  
43 638 and BN phytoplasma variability in Croatia: multigene sequence analysis approach.  
44  
45 639 *European Journal of Plant Pathology*, **142**, 591–601.
- 46  
47 640 48. Prevostini M., Taddeo A.V., Balac K., Rigamonti I., Baumgärtner J., Jermini M. (2013)  
48  
49 641 WAMS - an adaptive system for knowledge acquisition and decision support: the case of  
50  
51 642 *Scaphoideus titanus*. *IOBC-WPRS Bulletin*, **85**, 57–64.

- 1  
2  
3 643 49. Quaglino F., Casati P., Bianco P.A. (2010) Distinct *rpsC* single nucleotide polymorphism  
4  
5 644 lineages of Flavescence dorée subgroup 16SrV-D phytoplasma co-infect *Vitis vinifera* L..  
6  
7 645 *Folia Microbiologica*, **55**, 251–257.  
8  
9 646 50. Quaglino F., Maghradze D., Casati P., Chkhaidze N., Lobjanidze M., Ravasio A., Passera  
10  
11 647 A., Venturini G., Failla O., Bianco P.A. (2016) Identification and characterization of new  
12  
13 648 '*Candidatus* Phytoplasma solani' strains associated with bois noir disease in *Vitis vinifera* L.  
14  
15  
16 649 cultivars showing a range of symptoms severity in Georgia, the Caucasus region. *Plant*  
17  
18 650 *Disease*, **100**, 904–915.  
19  
20 651 51. Quaglino F., Zhao Y., Casati P., Bulgari D., Bianco P.A., Wei W., Davis R.E. (2013)  
21  
22 652 '*Candidatus* Phytoplasma solani', a novel taxon associated with stolbur and bois noir related  
23  
24 653 diseases of plants. *International Journal of Systematic and Evolutionary Microbiology*, **63**,  
25  
26 654 2879–2894.  
27  
28  
29 655 52. Rosenberger D.A., Jones A.L. (1978) Leafhopper vectors of the Peach X disease pathogen  
30  
31 656 and its seasonal transmission from chokecherry. *Phytopathology*, **68**, 782–790.  
32  
33 657 53. Sanna F., Quaglino F., Filisetti S., Casati P., Faccincani M., Bianco P.A., Mori N. (2016)  
34  
35 658 Preliminary results on putative vectors of '*Candidatus* Phytoplasma solani' in Bois noir-  
36  
37 659 affected vineyards in Franciacorta (Lombardy region, North Italy). *Mitteilungen*  
38  
39 660 *Klosterneuburg*, **66**, 28–31.  
40  
41  
42 661 54. Schvester D., Carle P., Moutos G., Bonfils J. (1962) Etudes biologique des cicadelles de la  
43  
44 662 vigne dans le sud-ouest de la France. *Annales des épiphyties*, **13**, 205–237.  
45  
46  
47 663 55. Tamura K., Stecher G., Peterson D., Filipski A., Kumar S. (2013) MEGA6: Molecular  
48  
49 664 Evolutionary Genetics Analysis Version 6.0. *Molecular Biology and Evolution*, **30**,  
50  
51 665 2725–2729.  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 666 56. Trivellone V., Filippin L., Narduzzi-Wicht B., Angelini E. (2016) A regional-scale survey to  
4  
5 667 define the known and potential vectors of grapevine yellow phytoplasmas in vineyards  
6  
7 668 South of Swiss Alps. *European Journal of Plant Pathology*, **145**, 915–927.

8  
9 669 57. Trivellone V., Jermini M., Linder C., Cara C., Delabays N., Baumgärtner J. (2013) Role de  
10  
11 670 la flore du vignoble sur la distribution de *Scaphoideus titanus*. *Revue Suisse de Viticulture,*  
12  
13 671 *Arboriculture, Horticulture*, **45**, 222–228.

14  
15 672

16  
17 673

18  
19 674

20  
21 675

22  
23 676

24  
25 677

26  
27 678

28  
29 679

30  
31 680

32  
33 681

34  
35 682

36  
37 683

38  
39 684

## Supporting Information

Casati *et al.* [Insights on Flavescence dorée ecology]

685

686

687

688 **Table S1** *Map* gene nucleotide sequences of 16SrV group strains retrieved from NCBI GenBank

689 for sequence identity and phylogenetic analyses

Strain	map type	Host	Location	Accession number
FD V00-SP5 and V00-SP9	FD2	<i>Vitis vinifera</i>	Gironde, France	AM384886
FD V02-101 and V01-9	FD1	<i>Vitis vinifera</i>	Gironde, France	AM384887
FD V03-9-2, 9-16, and 9-17	FD1	<i>Vitis vinifera</i>	Pyrénées Atlantiques, France	AM384888
FD V04-11-19, 11-21, and 11-53	FD1	<i>Vitis vinifera</i>	Lot, France	AM384889
FD VI04-C28 and C29	FD3	<i>Vitis vinifera</i>	Veneto, Italy	AM384894
FD VI04-Toscana1	FD3	<i>Vitis vinifera</i>	Toscana, Italy	AM384895
FD VI04-188-04 and 248-04	FD3	<i>Vitis vinifera</i>	Piemonte, Italy	AM384896
FD FD 70	FD3	<i>Catharanthus roseus</i>	France	AM238512
PGY PGY-A		<i>Vitis vinifera</i>	Rheinland-Pfalz, Germany	AM384892
PGY PGY-B Yellows (B type)		<i>Vitis vinifera</i>	Rheinland-Pfalz, Germany	AM384893
PGY PGY-C EY 38		<i>Vitis vinifera</i>	Rheinland-Pfalz, Germany	AM384891
PGY V04-11-01		<i>Vitis vinifera</i>	Haut-Rhin, France	AM384890
EY1 E04-D714		<i>Ulmus glabra</i>	Haute-Vienne, France	AM384901
AldY AI04-3-13 and AI04-3-7		<i>Alnus glutinosa</i>	Basilicata, Italy	AM384884
Ald YWJ1444-32		<i>Alnus glutinosa</i>	Pyrénées Orientales, France	AM384897
AldY ALY		<i>Catharanthus roseus</i>	Basilicata, Italy	AM384885
AldY A06-30-3	FD1	<i>Alnus glutinosa</i>	Gironde, France	FN561864
AldY Ag30 MAC		<i>Alnus glutinosa</i>	Macedonia	KJ605451
AldY 74-08-MNE		<i>Alnus glutinosa</i>	Montenegro	KC188998
AldY SW7		<i>Alnus glutinosa</i>	Germany	KP238304
AldY SW38		<i>Alnus glutinosa</i>	Germany	KP238315
AldY SW1		<i>Alnus glutinosa</i>	Germany	KP238302
AldY 75-08-MNE		<i>Alnus glutinosa</i>	Montenegro	KC188999
AldY Ag26 MAC		<i>Alnus glutinosa</i>	Macedonia	KJ605448
AldY Ag27 MAC		<i>Alnus glutinosa</i>	Macedonia	KJ605449
AldY Ag29 MAC		<i>Alnus glutinosa</i>	Macedonia	KJ605450
AldY A06-30-25		<i>Alnus glutinosa</i>	Gironde, France	FN561865
AldY SW13		<i>Alnus glutinosa</i>	Germany	KP238307
AldY SW23		<i>Alnus glutinosa</i>	Germany	KP238310
AldY A06-30-20		<i>Alnus glutinosa</i>	Gironde, France	FN561863
AldY SW4		<i>Alnus glutinosa</i>	Germany	KP238303
AldY SW8		<i>Alnus glutinosa</i>	Germany	KP238305
AldY SW17		<i>Alnus glutinosa</i>	Germany	KP238309
FD FG 1	FD3	<i>Vitis vinifera</i>	Serbia	KP238300

690

# Supporting Information

Casati *et al.* [Insights on Flavescence dorée ecology]

**Table S2** Sequence identity of *map* gene of FDp and related strains identified in the present study and 16SrV phytoplasma strains available at NCBI GenBank

#	Type	Strain	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	
1	FD1	Vv128	ID																									
2	FD1	Vv129	1	ID																								
3	FD1	Ca249	1	1	ID																							
4	FD1	Ca253	1	1	1	ID																						
5	FD1	Sa252	0,998	0,998	0,998	0,998	ID																					
6	FD1	Oi99	0,998	0,998	0,998	0,998	0,996	ID																				
7	FD1	Oi116	1	1	1	1	0,998	0,998	ID																			
8	FD1	St121	1	1	1	1	0,998	0,998	1	ID																		
9	FD1	St142	1	1	1	1	0,998	0,998	1	1	ID																	
10	FD2	Vv178	0,98	0,98	0,98	0,98	0,982	0,978	0,98	0,98	0,98	ID																
11	FD2	Vv220	0,98	0,98	0,98	0,98	0,982	0,978	0,98	0,98	0,98	1	ID															
12	FD2	Ca158	0,98	0,98	0,98	0,98	0,982	0,978	0,98	0,98	0,98	1	1	ID														
13	FD2	Oi12	0,98	0,98	0,98	0,98	0,982	0,978	0,98	0,98	0,98	1	1	1	ID													
14	FD2	Oi32	0,98	0,98	0,98	0,98	0,982	0,978	0,98	0,98	0,98	1	1	1	1	ID												
15	FD2	Td8	0,98	0,98	0,98	0,98	0,982	0,978	0,98	0,98	0,98	1	1	1	1	1	ID											
16	FD2	Td31	0,98	0,98	0,98	0,98	0,982	0,978	0,98	0,98	0,98	1	1	1	1	1	1	ID										
17	FD3	Ca254	0,984	0,984	0,984	0,984	0,985	0,982	0,984	0,984	0,984	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	1	ID		
18	FD3	Ca257	0,984	0,984	0,984	0,984	0,985	0,982	0,984	0,984	0,984	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	1	ID		
19	FD3	Ca256	0,984	0,984	0,984	0,984	0,985	0,982	0,984	0,984	0,984	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	1	1	ID	
20	FD3	Oi102	0,982	0,982	0,982	0,982	0,984	0,98	0,982	0,982	0,982	0,98	0,98	0,98	0,98	0,98	0,98	0,98	0,98	0,998	0,998	0,998	ID					
21	FD3	Oi115	0,984	0,984	0,984	0,984	0,985	0,982	0,984	0,984	0,984	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	1	1	1	0,998	ID			
22	FD3	Ho1	0,957	0,957	0,957	0,957	0,959	0,955	0,957	0,957	0,957	0,955	0,955	0,955	0,955	0,955	0,955	0,955	0,955	0,973	0,973	0,973	0,971	0,973	ID			
23	FD1	FD70	1	1	1	1	0,998	0,998	1	1	1	0,98	0,98	0,98	0,98	0,98	0,98	0,98	0,984	0,984	0,984	0,982	0,984	0,957	ID			
24	FD2	V00-SP5	0,98	0,98	0,98	0,98	0,982	0,978	0,98	0,98	0,98	1	1	1	1	1	1	1	0,982	0,982	0,982	0,98	0,982	0,955	0,98	ID		
25	FD3	VI04-Toscana1	0,984	0,984	0,984	0,984	0,985	0,982	0,984	0,984	0,984	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,982	0,996	0,996	0,996	0,994	0,996	0,969	0,984	0,982	ID

**Table 1** Insect vector specimens captured by yellow sticky traps in the test vineyard from May to November 2014, and FDp and related strains molecular identification and characterization in pooled insects

Trap	Species <sup>b</sup>																											
	<i>S. titanus</i>					<i>O. ishidae</i>					<i>T. dilutior</i>				<i>R. cuspidatus</i>			<i>F. florii</i>			<i>H. obsoletus</i>				<i>A. fuscovenosus</i>			
	Cpt	And	Pls	Inf	map	Cpt	And	Pls	Inf	map	Cpt	Pls	Inf	map	Cpt	Pls	Inf	Cpt	Pls	Inf	Cpt	Pls	Inf	map	Cpt	Pls	Inf	
Rebell inside the vineyard	V1																1	1	0									
	V2	1	1	1	0																							
	V3					5	5	4	0												1	1	1	FD3(1)				
	V4																											
	V5																					2	1	0				
	V6						2	2	2	1	FD2(1)																	
Rebell on the borders	B1																3	3	0									
	B2					1	1	1	0								2	1	0									
	B3	1	1	1	0	88	88	26	10	FD1(2) FD2(8)		1	1	1	FD2(1)	1	1	0										
	B4	6	6	4	1	FD1(1)	15	15	8	2	FD2(2)																	
	B5						2	2	1	0																		
	B6						1	1	1	0		1	1	1	FD2(1)													
	B7						4	4	3	0											5	4	0					
	B8																				4	2	0					
Rebell outside the vineyard <sup>a</sup>	W1					15	15	8	1	FD3(1)																		
	W2					4	4	3	0																			
	W3																											
	W4					4	4	4	2	FD1(1) FD2(1)																		
	W5					4	4	4	2	FD1(1) FD2(1)																		
	W6					2	2	1	0																			
	W7					3	3	3	0																			
	W8	15	15	15	1	FD1(1)	29	29	12	3	FD1(1) FD2(2)														1	1	0	
	W9	6	6	6	0		35	35	12	4	FD2(3) FD3(1)																	
Aerixon		11	2	2	0	22	4	4	0												3	3	0					
		40	31	29	2	FD1(2)	236	218	97	25	FD1(5) FD2(18) FD3(2)	2	2	2	FD2(2)	6	5	0	1	1	0	15	11	1	FD3(1)	1	1	0

<sup>a</sup> traps W1 to W7 positioned at the external border of the forest neighboring other vineyards; traps W8 and W9 positioned within the forest

<sup>b</sup> Cpt, captured specimens; And, analyzed specimens; Pls, pools; Inf, infected pools determined by real-time PCR; map, map-type determined by RFLP analysis of map gene amplified by nested PCR



699 **Table 2** Molecular identification and characterization of FDp and related strains in plants

700

Host	No. of samples	Real-time PCR	map PCR	RFLP map types		
				FD1	FD2	FD3
<i>Vitis vinifera</i>	74	70	70	2	68	0
<i>Corylus avellana</i>	31	11	6	3	1	2
<i>Salix</i> sp.	10	3	1	1	0	0
<i>Clematis vitalba</i>	4	3	3	0	0	3

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721 **Table 3** GenBank accession numbers of *map* gene nucleotide sequences amplified from  
 722 representative FDp, FDp related and 16SrV phytoplasma strains identified in insects and plants  
 723

RFLP <i>map</i> types	Host	No. of strains	Representative strain	Accession Number
FD1	<i>Vitis vinifera</i>	2	Vv128 <sup>a</sup>	KX245031
			Vv129 <sup>a</sup>	KX245030
	<i>Corylus avellana</i>	3	Ca249 <sup>b</sup>	KX245028
			Ca253 <sup>b</sup>	KX245033
	<i>Salix</i> sp.	1	Sa252 <sup>b</sup>	KX245029
	<i>Orientus ishidae</i>	5	Oi99 <sup>b</sup>	KX245034
			Oi116 <sup>b</sup>	KX245032
<i>Scaphoideus titanus</i>	2	St121 <sup>a</sup>	KX245035	
		St142 <sup>a</sup>	KX245036	
FD2	<i>Vitis vinifera</i>	68	Vv178 <sup>a</sup>	KX245026
			Vv220 <sup>a</sup>	KX245025
	<i>Corylus avellana</i>	1	Ca158 <sup>b</sup>	KX245023
	<i>Orientus ishidae</i>	18	Oi12 <sup>b</sup>	KX245021
			Oi32 <sup>b</sup>	KX245022
	<i>Thamnotettix dilutior</i>	2	Td8 <sup>b</sup>	KX245024
			Td31 <sup>b</sup>	KX245027
FD3	<i>Corylus avellana</i>	2	Ca254 <sup>b</sup>	KX245038
			Ca257 <sup>b</sup>	KX245041
	<i>Clematis vitalba</i>	1	Cv256 <sup>b</sup>	KX245042
	<i>Orientus ishidae</i>	2	Oi102 <sup>b</sup>	KX245040
			Oi115 <sup>b</sup>	KX245039
<i>Hyalesthes obsoletus</i>	1	Ho1 <sup>c</sup>	KX245037	

724 <sup>a</sup> FDp strain; <sup>b</sup> FDp related strain; <sup>c</sup> 16SrV phytoplasma strain  
 725  
 726  
 727  
 728  
 729  
 730  
 731  
 732  
 733  
 734  
 735

736 **FIGURE LEGENDS**

737 **Figure 1.** View of the vineyard agro-ecosystem in Stabio, 'Canton Ticino' Region (Switzerland),  
738 and position of the yellow sticky traps (Rebell and Aeroxon) used to capture potential phytoplasma  
739 insect vectors. Rebell traps inside the vineyards: V1 to V6; Rebell traps on the borders: B1 to P8;  
740 Rebell outside the vineyard: W1 to W9. Fourteen aeroxon traps were positioned horizontally on the  
741 grapevine canopy inside the test vineyard.

742 **Figure 2.** Flight dynamics of *Orientus ishidae* (in blue) and *Scaphoideus titanus* (in red) in the  
743 vineyard in Stabio, 'Canton Ticino' Region (Switzerland), determined through the number of  
744 specimens captured inside the vineyard by Aeroxon (A) and Rebell (V1 to V6, B1 to B8) (B) traps,  
745 and outside the vineyard by Rebell (W1 to W9) traps (C).

746 **Figure 3.** Restriction fragment length polymorphism (RFLP) profiles obtained through double  
747 digestion with the enzymes *AluI* and *Eco72I*. RFLP patterns were visualized through  
748 electrophoresis in 3% agarose gel. Attribution of RFLP profiles, identified in the present study, was  
749 carried out through the comparison with the profiles of the *map*-types FD1 (A), 2 (B), and 3 (C)  
750 (Arnaud *et al.*, 2007). Acronyms of the FDp and related strains are reported in Table 3.

751 **Figure 4.** Unrooted phylogenetic tree inferred from analyses of nucleotide sequences of *map* gene.  
752 Minimum-Evolution method was carried out using the Jukes-Cantor model with the software  
753 MEGA6. The reliability of the analyses was subjected to a bootstrap test with 1000 replicates;  
754 bootstrap values lower than 50 are not shown. Phytoplasma strains and their nucleotide sequence  
755 accession numbers are reported in Table 3 and Table S1. Nucleotide sequences from the present  
756 work (Table 3) are written in bold characters.

757

758

759

760

1  
2  
3 761  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43 762  
44  
45 763  
46  
47 764  
48  
49 765  
50  
51 766  
52  
53 767  
54  
55 768  
56  
57 769  
58  
59 770  
60




Figure 1

771

772




Figure 2

773

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789

790

Review

1  
2  
3 791  
4 792  
5  
6 793  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33 794  
34 795  
35 796  
36 797  
37 798  
38 799  
39 800  
40 801  
41 802  
42 803  
43 804  
44 805  
45 806  
46 807  
47 808  
48 809  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 3

810




Figure 4

811