

1 **Characterization of Whole Grain Pasta: Integrating Physical, Chemical, Molecular, and Instrumental Sensory**

2 **Approaches**

3 Alessandra Marti, Stefano Cattaneo, Simona Benedetti, Susanna Buratti, Parisa Abbasi Parizad, Fabio Masotti,

4 Stefania Iametti* and Maria Ambrogina Pagani

5 Department of Food, Environmental and Nutritional Sciences (DeFENS), Università degli Studi di Milano, via G.

6 Celoria 2, 20133 Milan, Italy

7 * Corresponding author: Stefania Iametti; DeFENS, Via G. Celoria 2, 20133 Milan, Italy; Phone +3902-5031-

8 6819; stefania.iametti@unimi.it

9 **Word count of text:**4998

10 **Short version of title:** Features of whole-grain pasta

11 **Choice of journal/section where article should appear:** Food Chemistry

12

13 **ABSTRACT:** The consumption of whole-grain food - including pasta - has been increasing steadily. In the case of
14 whole-grain pasta, given the many different producers, it seems important to have some objective parameters
15 to define its overall quality. In this study, commercial whole-grain pasta samples representative of the Italian
16 market have been characterized from both molecular and electronic-senses (electronic nose and electronic
17 tongue) standpoint in order to provide a survey of the properties of different commercial samples. Only one
18 pasta product showed very low levels of heat damage markers (furosine and **pyrraline**), suggesting that this
19 sample underwent to low temperature dry treatment. In all samples, the furosine content was directly
20 correlated to protein structural indices, since protein structure compactness increased with increasing levels of
21 heat damage markers. Electronic senses were able to discriminate among pasta samples according to the
22 intensity of heat treatment **during the drying step**. Pasta sample with low furosine content was discriminated
23 by umami taste and by sensors responding to aliphatic and inorganic compounds. Data obtained with this
24 multidisciplinary approach **are meant to provide** hints for identifying useful indices for pasta quality.

25 **Keywords:** whole-grain pasta; heat damage; electronic nose; electronic tongue; solubilized proteins

26 **Practical Application:** As observed for semolina pasta, objective parameters based on heat-damage were best
27 suited to define the overall quality of wholegrain pasta, almost independently of compositional differences
28 among commercial samples. Drying treatments of different intensity also had an impact on instrumental
29 sensory traits, that may provide a reliable alternative to analytical determination of chemical markers of heat
30 damage in all cases where there is a need for avoiding time-consuming procedures.

31

32 **Introduction**

33 Given the well-known health benefits of whole grain consumption (Poutanen 2012), consumers are
34 increasingly demanding whole-grain versions of many cereal-based products. Because of its simple formula and
35 worldwide popularity, pasta has been a common product targeted by product developers for offering whole-
36 grain formats. Indeed, a single serving of whole grain pasta – with its 7 g of fiber per 100 g of pasta (Casiraghi
37 and others 2013) - could help in satisfying the recommended daily intake for this component (Fogliano and
38 Vitaglione 2005).

39 Beside the nutritional benefits, there are a spate of sensory (Heiniö and others 2016) and technological
40 (Foschia and others 2013; Rakhesh and others 2015) issues that may become problematic when dealing with
41 whole wheat pasta. Whole-grain pasta appears dark in color, with a bitter taste and with the possibility of
42 occurrence of off-flavors and off-odors during storage. Whole-grain pasta producers are therefore called to
43 meet the demand for a product with high nutritional interest, while paying due attention to the acceptability of
44 the product.

45 In this regard, particular attention should be paid to the choice of the raw material and the
46 technological process. High levels of both damaged starch and amylase activity, typical of whole-meal flour, are
47 responsible for the formation of reducing sugars and for intense Maillard reaction (MR) (De Noni and Pagani
48 2010). The drying step certainly represents the most critical phase in the pasta-making process, because of its
49 impact on the texture, the color, the flavor, and the nutritional properties of the product. Therefore, the
50 definition of the overall quality of pasta cannot neglect the intensity of the MR assessment that is carried out
51 by the determination of appropriate molecular markers. Furosine (ϵ -N-furoylmethyl-L-lysine, FUR) is the most
52 widely used index of heat damage in pasta and in cereal-based products (Resmini and Pellegrino 1994). Heat
53 damage in dried pasta also involves protein glycation and formation of advanced glycation end products (AGEs,
54 e.g. ϵ -pyrrole-lysine (**pyrraline**; ϵ -2-formyl-5-hydroxymethyl-pyrrolaldehyde; ϵ -PL) (Resmini and others 1993) -
55 which may affect protein digestibility (Seiquer and others 2006; Stuknytė and others 2014) and be potentially
56 involved in the onset of some non-communicable diseases (Uribarri and others 2010).

57 In the case of pasta, a relationship has been reported also between the intensity of the heat treatment
58 and the association of proteins through covalent and non-covalent interactions (Bonomi and others 2012). This
59 has been studied also in whole grain pasta (Bock and others 2015), but the system used in those studies was
60 based on common wheat, and therefore different from the semolina-based products that are mandatory in
61 Italy. It seems reasonable that the different availability of water and the different macromolecular organization
62 in whole-grain systems may result – also in this case - in altered patterns for protein-related structural changes
63 and for formation of protein-carbohydrate adducts.

64 Aside from their molecular traits, whole-grain products are worthless unless they have
65 satisfactory sensory traits. Producers have their own panels, but their scores are not always
66 accessible, and are far from being univocal. Therefore, there is a great interest in using electronic senses
67 (electronic nose and electronic tongue) for objective analysis of sensory traits. Electronic nose and electronic
68 tongue (e-nose and e-tongue) allow to evaluate the contribution of different chemical species in determining
69 aroma and tastes in food products (Sliwinska and others 2014). Thus, the application of electronic senses (e-
70 senses) to pasta could represent an attractive – and objective - alternative to sensory tests for quality control
71 and process monitoring.

72 This study **aimed at exploring** the fundamental aspects of structure, flavor, taste, and cooking
73 behavior of whole wheat pasta, and at relating these characteristics to the processing conditions as
74 estimated by heat damage indices. Both chemical and structural indices were taken into account.

75 **Materials and Methods**

76 *Materials*

77 Packages of whole grain pasta manufactured in Italy by different companies were purchased in two Italian
78 supermarkets. A total of 20 spaghetti samples (1.7 mm diameter) were used, from two different production
79 batches of 10 different brands. The samples were labeled with separate codes for identification of the
80 producer and batch, respectively (i.e., a1, a2, b1, b2, etc). Chemical composition of pasta samples was those

81 reported by the manufacturer (see Table 1). No information was available as for the specific materials and
82 processes used by individual producers. When appropriate, pasta samples were ground with a laboratory
83 mill (IKA Universalmühle M20; Janke & Kunkel GmbH & CoKG, IKA Labortechnik, Staufen, Germany), fitted with a
84 water cooling jacket in order to avoid overheating during grinding.

85 *Cooking behavior*

86 Cooking loss was evaluated according to the AACC official method 66-50.01 (AACC, 2001). An aliquot of pasta
87 (10 g) was cooked in boiling water (100 mL) until the optimum cooking time suggested by the
88 producer, as reported on the package with no salt added. After cooking, pasta was drained, cooking water
89 was recovered and its level brought back to the starting volume. An aliquot of water (40 mL) was then dried to
90 constant weight at 105 °C. The residue was weighed and the dry matter reported as percentage of the starting
91 dry material. Results were expressed as grams of solid loss/100 g of dry pasta. Pasta weight increase due to
92 cooking was evaluated by weighing pasta before and after cooking. Each measurement was determined in
93 duplicate.

94 *Evaluation of heat damage*

95 The formation of chemical artefacts potentially capable of affecting protein digestibility was assessed by
96 determining the following molecules arising from MR. The level of FUR and ϵ -PL was determined according to
97 Resmini and others (1990) and Resmini and Pellegrino (1994) respectively. In the case of ϵ -PL, the
98 measurement was carried out only on one batch of pasta, taking into account FUR data performed on all the
99 samples. The analyses were performed in triplicate.

100 *Properties of the protein network*

101 The solubility of proteins in pasta samples was determined in triplicate by using buffers of different
102 composition, as described by lametti and others (2006). Typically, 1 g of finely ground sample was suspended in
103 20 mL of 50 mM sodium phosphate buffer pH 7.0 containing 0.1 M NaCl. After stirring at room temperature for

104 60 min and removal of insoluble materials by centrifugation (2500 x g for 30 min), the protein content in the
105 supernatant was assessed by a dye-binding colorimetric method (Bradford, 1976). Where indicated, the buffer
106 used for protein extraction also contained 8 M urea or 8 M urea and 10 mM dithiothreitol (DTT) as reported by
107 Bonomi and others (2012). A given amount (10 mg) of the milled uncooked pasta was treated with 0.2 ml of
108 SDS-PAGE denaturing buffer (0.125 M Tris-HCl, pH 6.8, 50 % glycerol, 1.7 % SDS; 0.01 % Bromophenol Blue)
109 containing 1% 2-mercaptoethanol when indicated, and heated at 100 °C for 10 min. For comparison, SDS-PAGE
110 analysis was also carried out on the proteins extracted by treating 10 mg of ground pasta samples directly with
111 0.2 ml of denaturing buffer containing 2-mercaptoethanol for 0 min at 100°C. SDS-PAGE was carried out on a
112 fixed porosity gel (12% monomer), using a MiniProtein apparatus (BioRad, Richmond, VA, USA) and gels were
113 Coomassie Blue-stained. Molecular Weight markers (14; 20.1; 30; 45; 67;96 kDa) were from BioRad
114 (Richmond, VA, USA).

115 *E-senses*

116 The volatile profile of the various pasta samples was assessed by a portable e- nose (PEN2, Win Muster
117 Airsense Analytics Inc., Schwerin, Germany). The device consisted of a sampling apparatus, a detecting unit
118 with a sensor array, and an appropriate pattern-recognition software (Win Muster v.1.6) for data recording and
119 elaboration. For e-nose measurements, the sample headspace was exposed to the sensors that provided a
120 signal pattern related to the volatile compounds in the headspace (Benedetti and others 2008; Marengo and
121 others 2017). The sensor array was made up of 10 metal oxide semiconductor–type sensors: W1C (aromatic),
122 W5S (broad range), W3C (aromatics), W6S (hydrogen), W5C (aromatics–aliphatics), W1S (broad methane),
123 W1W (sulfur-containing compounds), W2S (broad alcohol), W2W (sulfur-containing and chlorinated
124 compounds), and W3S (methane-aliphatics).

125 E-nose measurements were performed on both cooked and uncooked pasta samples. For the determination of
126 the aromatic profile, samples were placed in a 40 mL airtight glass vial fitted with a pierceable silicon/Teflon®
127 disk in the cap. The uncooked pasta sample (3 g) was used as such, whereas the cooked pasta sample (5 g of

128 lyophilized material) was wetted with 2 ml of distilled water. After an hour equilibration at room temperature,
129 headspace measurements were performed according to the following conditions: flow rate 300 ml/min,
130 injection time 60 s, flush time 180 s (during which the surface of the sensors was cleaned with air filtered
131 through active carbon). All samples were analyzed in triplicate and the average of sensor responses were used
132 for the subsequent statistical analysis.

133 E-tongue measurements were carried out by a Taste-Sensing System SA 402B (Intelligent Sensor Technology
134 Co. Ltd, Japan). For this study a total of 4 detecting sensors and 2 reference electrodes were used, separated in
135 two arrays according to the membrane charge: hybrid (CT0; AAE) and positive (C00; AE1). Measurements are
136 based on the capability of tasty compounds to modify sensors potential through electrostatic or hydrophobic
137 interactions (Buratti and others 2011). Ten grams of cooked and lyophilized pasta were resuspended in 150 mL
138 of distilled water. Solutions were vortexed for 5 min and centrifuged at 5000 rpm for 10 min at 20° C. After
139 centrifugation, the supernatants were filtered, and placed in 40 mL vessels for e-tongue analysis. Before each
140 measurement, the detecting sensors and the reference electrodes were dipped into a reference solution (30
141 mM potassium chloride, 0.3 mM tartaric acid) and the electric potential was measured for each sensor (V_r).
142 The sensors were then dipped for 30 s into the sample and the potential (V_s) was measured. For each sensor
143 the “relative value” (R_v) was calculated as the difference between the potential of the sample and that of the
144 reference solution ($V_s - V_r$). Sensors were then rinsed with fresh reference solution for 6 s, and dipped again into
145 the reference solution. The new potential of the reference solution was defined as V_r' . For each sensor, the
146 difference between the potential of the reference solution before and after sample measurement ($V_r' - V_r$) is
147 the **CPAV (Change of Membrane Potential caused by Absorption value)** and corresponds to the e-tongue
148 “aftertastes” (Kobayashi and others 2010; Buratti and others 2013). Before a new measurement cycle started,
149 sensors were rinsed for 90 s with washing solutions and then for 180 s with the reference solution. Each
150 sample was analysed in triplicate and sensor outputs were converted to taste information. The “taste values”
151 were calculated by multiplying sensor outputs for appropriate coefficients based on the Weber–Fechner law,

152 which gives the intensity of sensation considering the sensor property for tastes (Kobayashi and others 2010;
153 Buratti and others 2013).

154 *Statistical analysis*

155 Data were elaborated by Principal Component Analysis (PCA) using the MINITAB 14 software package (Minitab
156 Inc., State College, PA, USA, version 12.0). PCA was applied as an exploratory tool to uncover aroma and taste
157 characteristics. Analysis of variance (ANOVA) was performed with Statgraphics XV version 15.1.02 (StatPoint Inc.,
158 Warrenton, VA, USA). Samples were used as factor. When the factor effect was significant ($p \leq 0.05$), differences
159 among the respective means were determined using Fisher's Least Significant Difference (LSD) test.

160 **Results and Discussion**

161 *Cooking behavior*

162 The cooking behavior of commercial whole wheat pasta – evaluated as cooking loss and water absorption - is
163 summarized in Table 2. The amount of material leached into cooking water is frequently used to define the
164 cooking quality of semolina pasta (Marti and others 2016). Indeed, good pasta cooking quality is assured by the
165 formation of a continuous and strengthened network of coagulated gluten proteins, which entraps the starch
166 macromolecules, limiting their swelling and solubilization into the cooking water (Resmini and Pagani 1983).
167 Cooking loss values ranged from 3.38 and 4.73 g/100g pasta for j1 and h2, respectively, but no statistically
168 significant difference was evident when considering this parameter. Although these samples presented the
169 highest and the lowest protein content, no significant correlation ($p > 0.05$) was detected between protein
170 content and cooking loss when all the samples were taken into account.

171 As regards water absorption, values ranged from 107.8 to 155.2 g/100g pasta for g2 and e1,
172 respectively. Despite the contribution of fiber in increasing the amount of water absorbed during cooking, no
173 significant ($p > 0.05$) correlation between fiber content and water absorption was detected. Results on cooking
174 behavior suggest the key role of processing conditions (i.e. drying cycle) in affecting the cooking behavior of

175 whole grain pasta, as already shown in pasta from refined semolina (D'Egidio and others 1990; Marti and
176 others 2013).

177 *Heat Damage*

178 The extent of (MR) in commercial samples of whole wheat pasta, monitored by their FUR and ϵ -PL content, is
179 summarized in **Table 2**. The mean FUR content of the 10 brands is equal to 595 mg/100 g protein, ranging from
180 229 to 836 mg/100 g. The wide range of data herewith recorded suggests high variability in raw material
181 characteristics (i.e. α -amylase activity and reducing sugars content) and/or processing conditions (i.e. drying
182 diagram) among manufacturers. The lowest values of FUR, in the range of 229-262 mg/100 g protein, were
183 found for j1, and j2, respectively. This result suggests the use of medium temperature drying cycle, that results
184 in limited heat damage. Samples j1 and j2 also exhibited the lowest cooking losses (**Table 2**), likely because of
185 the high quality of the raw materials (Marti and others 2013).

186 FUR levels in whole grain pasta below 300 mg/100 g protein are indicative of a mild heat damage
187 (Casiraghi and others 2013). As expected, the whole grain pasta is characterized by higher average FUR values
188 compared to semolina pasta (De Noni and Pagani 2010; Bonomi and others 2012), likely due to the intense
189 amylase activity and the high content in reducing sugars and proteins of whole-meal semolina, that make these
190 products highly susceptible to MR (Resmini and others 1996). **In general, no significant variability was observed**
191 **among pasta batches from the same producer, if not for the “d” brand. This result might be ascribed to the**
192 **differences in the expiration date between these two batches (5 months in the case of samples d1 and d2, vs 2**
193 **months in all other samples),** and by the reported effect of storage time on the FUR content (Cattaneo and
194 others 2008).

195 **In agreement with the FUR data, the lowest value of ϵ -PL was found for “j” brand samples (0.8**
196 **mg/100g proteins), whereas the highest values (>15 mg/100g proteins) were measured in the products of**
197 **brands “a” and “d”. To the best of our knowledge, no ϵ -PL data have been reported for whole-grain pasta. In**
198 **the present work, we found a mean value of 8.5 mg/100 g protein, although this parameter spanned a quite**

199 broad range in the various samples (from 0.8 to 15.8 mg/100g protein). Studies on pasta made from refined
200 semolina found low levels of ϵ -PL (0.58 mg/100g protein) in products dried using a low-temperature cycle,
201 whereas high level of ϵ -PL (3.39 mg/100g protein) were recorded after high temperature drying cycles
202 (Stuknyte and others 2014). It is worth noting that “j” pasta samples exhibited much lower either FUR and ϵ -PL
203 levels than “m” pasta, despite their similar high protein content (14%), suggesting that protein content could
204 not be the only attribute chosen by producers or consumers to define product quality.

205 Indeed, in addition to semolina quality, processing conditions (i.e. drying cycle) play a key role in
206 affecting heat damage and thus pasta nutritional quality. From a nutritional standpoint it is noteworthy that
207 most of the samples from whole semolina here considered showed a level of ϵ -PL higher than values found in
208 high temperature-dried semolina pasta (3.39 g/100 g of protein), for which a decrease in protein *in vitro*
209 digestibility of the cooked product was measured (Stuknyte and others 2014).

210 Finally, the data in Table 2 indicate the absence of a possible connection between either FUR or PYR
211 levels and cooking losses. This confirms that the application of high temperature drying cycles decreased the
212 significance of semolina quality (i.e., protein content and gluten strength indices) in determining the pasta
213 cooking behavior (D’Egidio and others 1990, Cubadda and others 2007, Marti and others 2013).

214 *Overall protein organization*

215 Structural features of proteins in the various pasta sample were investigated by detecting the amount of
216 protein solubilized by buffers of different dissociating ability towards covalent (disulfide) and non-covalent
217 inter-protein bonds. This approach has been shown to provide useful hints for understanding the nature and
218 extent of protein-protein interactions in cereal-based foods including gluten- containing (Bock and others 2015;
219 Bonomi and others 2012) and gluten-free (Cabrera-Chávez and others 2012; Marengo and others 2015) pasta,
220 also in the presence of nutritionally relevant ingredients of non-grain origin (eggs/whey) (Marti and others
221 2014).

222 As shown in Table 3, protein solubility in plain saline buffer was low for all samples, confirming
223 aggregation of otherwise soluble proteins in semolina (albumins and globulins) as a consequence of pasta
224 processing. Addition of urea to the saline extraction buffer resulted in a significant increase in solubilized
225 protein in all samples, suggesting that hydrophobic interactions were very relevant to the structure of
226 whatever protein network is present in these pasta samples (Table 3). When both urea and DTT were present
227 in the extraction medium, the amount of soluble proteins further increased markedly in all samples, suggesting
228 that inter-protein disulfides also play a fundamental role in stabilizing the structure of protein network.

229 By taking into account both the protein solubility data in Table 3 and the indices of thermal damage in
230 Table 2, it becomes evident that the sample with the lowest FUR content (i.e., j1 and j2) also had the highest
231 amount of solubilized protein regardless of the composition of the solubilizing medium. It is worth noting that
232 even a modest increase of thermal damage results in a markedly impaired protein solubility, as exemplified by
233 samples g1 and g2. These findings confirm previous reports on commercial pasta from refined semolina
234 (Bonomi and others 2012). The relationship between conditional proteins solubility and heat damage indices
235 offers hints for a possible relationship between the nature of prevalent interactions among proteins and the
236 type and intensity of processing used for pasta production also when other components are present, as is the
237 case in whole-grain pasta.

238 The nature of the proteins solubilized from the various samples by buffers of different composition was
239 investigated by SDS-PAGE. The resulting electrophoretic profiles were similar in all samples (Figure 1),
240 suggesting that the differences in protein organization observed in the solubility studies may be due to the
241 technological processing rather than to differences in the starting materials. In this frame, it is of interest to
242 note that the intensity of SDS-PAGE traces for proteins in the sample with the lowest level of FUR (Figure 1)
243 confirms the presence of high amount of solubilized proteins also upon treatment with detergents and
244 disulfide reducing agents. In the same “j” samples, at difference with other pasta samples, there is also no
245 evidence for formation of the large aggregates not capable of entering the separating electrophoretic gel
246 (Figure 1).

247 *E-senses*

248 E-senses evaluation of the pasta samples was performed by e-nose before and after cooking, for assessment of
249 the volatiles profile. E-nose data collected on uncooked pasta were elaborated by PCA in combination with
250 heat damage (FUR) and protein solubility (in urea) data in correlation matrix where the first two principal
251 components accounted for 84.8% of the total variance (Figure 2). From the score plot (Fig. 2A) **it can be noticed**
252 **that the pasta samples with the lowest furosine and the highest soluble protein levels (i.e., from the “j” brand,**
253 **Tables 2 and 3) are discriminated from all the other pasta samples, and are located on the right side of the first**
254 **principal component (PC1). As noted above, samples with an intermediate furosine content (g1 and g2) but**
255 **with low protein solubility are not discriminated in Fig. 2A.** By considering the loading plot (Fig. 2B) **the “j”**
256 **samples** are characterized by the highest odor intensity perceived by WW and WS sensors (of broad-range
257 sensitivity and specific for sulfur-containing compounds). In addition, the same “j” samples were characterized
258 by the lowest odor intensity perceived by WC sensors (specific for aromatic compounds). **All the other samples,**
259 **which are located on the left side of the plot, showed high odor intensity perceived by WC sensors.**

260 E-tongue measurements were performed on cooked and lyophilized pasta, after reconstitution in
261 water as done for e-nose measurements on the same cooked pasta samples. E-tongue device is a liquid
262 analytical approach that mimics the taste-sensing mechanism of gustatory system; in this work sensors specific
263 for bitterness, umami and astringency evaluation were applied. The estimated taste values were elaborated
264 together with e-nose data collected on the same pasta samples by PCA and the first two principal components
265 (PC1 and PC2) accounted for 65% of the total variance. Considering the score plot (Fig. 3A) and the loading plot
266 (Fig. 3B), it is possible to observe a clear discrimination of samples on the first two principal components. In
267 particular, the “j1” and “j2” pasta samples, located on the bottom right side of the plot (Fig. 3A), were
268 characterized by WW and WS e-nose sensors and by the umami taste detected by e-tongue (Fig. 3B). Samples
269 “a”, “b” and “d”, located on the left of the plot (Fig. 3A), were discriminated by the **aftertaste-A** and by WC
270 sensors; whereas samples “e”, “f”, and “g” were characterized mostly by **astringency and by bitterness.**

271 **Conclusion**

272 The results provided from this study indicate that the drying process is of paramount relevance to the
273 structural and nutritional quality of whole-grain pasta. As reported for pasta made from refined semolina,
274 wholegrain pasta dried at low temperature had low FUR and ϵ -PL content. Both **these** heat damage indices and
275 indices of protein aggregation and structural compactness suggest that the involved events (i.e., protein
276 glycation and structural rearrangement of gluten proteins) are relatively independent – in pasta – from the
277 amount of available fiber and from water distribution among phases. Protein content in the original material
278 had essentially no impact “per se” on the quality indices mentioned above, and no evidence was found for a
279 selective involvement of specific protein classes or subclasses in these events. The e-senses results
280 demonstrate that the low-temperature dried pasta can be easily discriminated from the high-temperature
281 dried samples. In particular, samples dried at low temperature were characterized by e-tongue sensor specific
282 for umami taste and by e-nose sensors of broad range sensitivity and specific for sulfur-containing compounds.
283 The high temperature treated samples were perceived more bitter and were characterized by e-nose sensors
284 specific for aromatic compounds. In conclusion, this combination of multidisciplinary approaches appears to
285 provide some hints for identifying objective indices for the assessment of whole-grain pasta quality.

286

287 **Acknowledgements**

288 The Authors gratefully acknowledge Mr. Gaetano Cardone (DeFENS, University of Milan, Italy) for technical
289 assistance and F.lli De Cecco (Fara San Martino, Italy) for partial support.

290 **Author Contributions**

291 A. Marti evaluated pasta quality and drafted the manuscript. P. Abbasi-Parizad carried out protein-related
292 studies, that were designed and supervised by S. Iametti, who also wrote the related parts of the manuscript. S.
293 Benedetti carried out e-senses studies, that were designed and supervised by S. Buratti, who also interpreted

294 the results. F. Masotti and S. Cattaneo carried out and interpreted the heat-damage-related studies. S. Iametti
295 was responsible for the coordination of the experimental plan execution. M.A. Pagani conceived the study.

296 **References**

297 Benedetti S, Buratti S, Spinardi A, Mannino S, Mignani I. 2008. Electronic nose as a non-destructive tool
298 to characterise peach cultivars and to monitor their ripening stage during shelf-life. *Postharvest Biol Technol*
299 47:181-188.

300 Bock JE, West R, Iametti S, Bonomi F, Marengo M, Seetharaman K. 2015. Gluten structural evolution
301 during pasta processing of refined and whole grain hard wheat pasta: The influence of mixing, drying, and
302 cooking. *Cereal Chem* 92:460-465.

303 Bonomi F, D'Egidio MG, Iametti S, Marengo M, Marti A, Pagani MA, Ragg EM. 2012. Structure–quality
304 relationship in commercial pasta: A molecular glimpse. *Food Chem* 135: 348-355.

305 Buratti S, Ballabio D, Giovanelli G, Zuluanga Dominguez C. M., Moles A, Benedetti S, Sinelli N. 2011.
306 Monitoring of alcoholic fermentation using near infrared and mid infrared spectroscopies combined with
307 electronic nose and electronic tongue. *Anal Chim Acta* 697:67-74.

308 Buratti S, Casiraghi A, Minghetti P, Giovanelli G. 2013. The joint use of electronic nose and electronic
309 tongue for the evaluation of the sensorial properties of green and black tea infusions as related to their
310 chemical composition. *Food Nutr Sci* 4:605–615.

311 Cabrera-Chávez F, Calderón de la Barca AM, Islas-Rubio AR, Marti A, Marengo M, Pagani MA, Bonomi F,
312 Iametti S. 2012. Molecular rearrangements in extrusion processes for the production of amaranth-enriched,
313 gluten-free rice pasta. *LWT - Food Sci Technol* 47:421-426.

314 Casiraghi MC, Pagani MA, Erba D, Marti A, Cecchini C, D'Egidio MG. 2013. Quality and nutritional
315 properties of pasta products enriched with immature wheat grain. *Int J Food Sci Nutr* 64:544-550.

316 Cattaneo S, Masotti F, Pellegrino L. 2008. Effects of overprocessing on heat damage of UHT milk. Eur
317 Food Res Technol 226:1099–1106.

318 Cubadda RE, Carcea M, Marconi M, Trivisonno MC. 2007. Influence of gluten proteins and drying
319 temperature on the coking quality of durum wheat pasta. Cereal Chem 84:48–55.

320 D’Egidio MG, Mariani BM, Nardi S, Novaro P, Cubadda R. 1990. Chemical and technological variables
321 and their relationships: A predictive equation for pasta cooking quality. Cereal Chem 67:275–281.

322 Fogliano V, Vitaglione P. 2005. Functional foods: planning and development. Mol Nutr Food Res
323 49:256-262.

324 Foschia M, Peressini D, Sensidoni A, Brennan CS. 2013. The effects of dietary fibre addition on the
325 quality of common cereal products. J Cereal Sci 58:216-227.

326 Heiniö RL, Noort MWJ, Katina K, Alam SA, Sozer N, de Kock HL, Hersleth M, Poutanen K. 2016. Sensory
327 characteristics of wholegrain and bran-rich cereal foods—A review. Trends Food Sci Technol 47:25-38.

328 Iametti S, Bonomi F, Pagani MA, Zardi M, Cecchini C, D’Egidio MG. 2006. Properties of the protein and
329 carbohydrate fractions in immature wheat kernels. J Agric Food Chem 54:10239-10244.

330 Kobayashi Y, Habara M, Ikezaki H, Chen R, Naito Y, Toko K. 2010. Advanced taste sensors based on
331 artificial lipids with global selectivity to basic taste qualities and high correlation to sensory scores. Sensors
332 10:3411-3443.

333 Marengo M, Bonomi F, Marti A, Pagani MA, Elkhailifa AEO, Iametti S. 2015. Molecular features of
334 fermented and sprouted sorghum flours relate to their suitability as components of enriched gluten-free pasta.
335 LWT - Food Sci Technol 63:511-518.

336 Marengo M, Baffour LC, Buratti S, Benedetti S, Saalia FK, Carpen A, Manful J, Johnson PNT, Barbiroli A,
337 Bonomi F, Pagani MA, Marti A, Iametti S. 2017. Defining the overall quality of cowpea-enriched rice-based
338 breakfast cereals. *Cereal Chem* 94:151-157.

339 Marti A, Seetharaman K, Pagani MA. 2013. Rheological approaches suitable for investigating starch and
340 protein properties related to cooking quality of durum wheat pasta. *J Food Qual* 36:133-138.

341 Marti A, Barbiroli A, Marengo M, Fongaro L, Iametti S, Pagani MA. 2014. Structuring and texturing
342 gluten-free pasta: egg albumen or whey proteins? *Eur Food Res Technol* 238:217-224.

343 Marti A, D'Egidio MG, Pagani MA. 2016. Pasta: Quality testing methods. In: Wrigley C, Corke H,
344 Seetharaman K, Faubion J, editors. *Encyclopedia of Food Grains*, 2nd Edition, volume 4, Oxford: Academic
345 Press. p. 161-165.

346 Poutanen K. 2012. Past and future of cereal grains as food for health. *Trends Food Sci Technol* 25:58-
347 62.

348 Rakshesh N, Fellows CM, Sissons M. 2015. Evaluation of the technological and sensory properties of
349 durum wheat spaghetti enriched with different dietary fibres. *J Sci Food Agr* 95:2-11.

350 Resmini P, Pagani MA. 1983. Ultrastructure studies of pasta: A review. *Food Microstruct* 2:1-12.

351 Resmini P, Pellegrino L, Battelli G. 1990. Accurate quantification of furosine in milk and dairy products
352 by a direct HPLC method. *Ital J Food Sci* 3:173-183.

353 Resmini P, Pagani MA, Pellegrino L, De Noni I. 1993. Formation of 2-acetyl-3-D-glucopyranosylfuran
354 (glucosylisomaltol) from nonenzymatic browning in pasta drying. *Ital J Food Sci* 5:341-353.

355 Resmini P, Pellegrino L. 1994. Occurrence of protein-bound lysylpirrolaldehyde in dried pasta. *Cereal*
356 *Chem* 71:254-262.

357 Resmini P, Pagani MA, Pellegrino L. 1996. Effect of semolina quality and processing conditions on non-
358 enzymatic browning in dried pasta. *Food Aust* 48:362-367.

359 Seiquer I, Díaz-Alguacil J, Delgado-Andrade C, López-Frías M, Hoyos AM, Galdó G, Navarro MP. 2006.
360 Diets rich in Maillard reaction products affect protein digestibility in adolescent males aged 11–14 y. *Am J Clin*
361 *Nutr* 83:1082-1088.

362 Sliwinska M, Wisniewska P, Dymerski T, Namiesnik J, Wardencki W. 2014. Food analysis using artificial
363 senses. *J Agric Food Chem* 62: 1423-1448.

364 Stuknyte M, Cattaneo S, Pagani MA, Marti A, Micard V, Hogenboom J, De Noni I. 2014. Spaghetti from
365 durum wheat: Effect of drying conditions on heat damage, ultrastructure and in vitro digestibility. *Food Chem*
366 149:40-46.

367 Uribarri J, Woodruff S, Goodman S, Cai W, Chen X, Pyzik R, Yong A, Striker GE, Vlassara H. 2010.
368 Advanced glycation end products in foods and a practical guide to their reduction in the diet. *J Am Diet Assoc*
369 110:911-916.

370 Table 1. Chemical parameters for the various pasta samples, as indicated in the label

371

Brand	Ingredients	Proteins (g/100g)	Carbohydrates (g/100g)	Fiber (g/100g)	Lipids (g/100g)
a	Wholegrain organic durum wheat semolina	12.0	67.5	7.0	1.5
b	Wholegrain durum wheat semolina	12.0	65.5	6.0	2.2
c	Wholegrain organic durum wheat semolina	12.2	66.9	6.8	1.7
d	Wholegrain durum wheat semolina, 2% germ	13.0	66.0	7.0	2.5
e	Wholegrain organic durum wheat semolina	13.0	66.0	8.0	2.0
f	Wholegrain durum wheat semolina	13.0	66.7	6.5	2.5
g	Wholegrain organic durum wheat semolina	11.5	64.0	7.0	2.0
h	Wholegrain organic durum wheat semolina	11.0	66.0	7.8	1.9
j	Wholegrain durum wheat semolina	14.0	63.0	7.5	2.5
m	Wholegrain durum wheat semolina	14.0	62.5	7.5	2.5

372

373

374 **Table 2.** Cooking behavior and heat damage of whole grain pasta.

375

Sample	Cooking behaviour		Heat damage	
	Cooking loss	Water absorption	Furosine	Pyrraline
	(g/100g pasta)	(g/100g)	(mg/100 g proteins)	(mg/100 g proteins)
a1	3.75 ± 0.1 ^a	138.2 ± 8.3 ^a	714 ± 9 ^b	15.5 ± 0.6 ^a
a2	4.23 ± 0.4 ^a	124.1 ± 7.4 ^a	742 ± 24 ^b	
b1	3.86 ± 0.4 ^a	129.2 ± 7.8 ^a	737 ± 7 ^b	6.5 ± 0.3 ^c
b2	3.70 ± 0.3 ^a	125.2 ± 7.5 ^a	738 ± 11 ^b	
c1	4.10 ± 0.1 ^a	133.0 ± 8.0 ^a	691 ± 11 ^b	4.8 ± 0.3 ^d
c2	3.57 ± 0.1 ^a	131.9 ± 7.9 ^a	712 ± 4 ^b	
d1	4.25 ± 0.5 ^a	125.1 ± 7.5 ^a	836 ± 22 ^a	15.8 ± 0.4 ^a
d2	3.87 ± 0.7 ^a	128.5 ± 7.7 ^a	618 ± 6 ^c	
e1	4.37 ± 0.1 ^a	155.2 ± 9.3 ^a	742 ± 22 ^b	7.0 ± 0.3 ^c
e2	4.31 ± 0.4 ^a	151.0 ± 9.1 ^a	699 ± 14 ^b	
f1	4.25 ± 0.4 ^a	145.8 ± 8.7 ^a	683 ± 6 ^b	4.6 ± 0.2 ^d
f2	4.32 ± 0.3 ^a	151.2 ± 9.1 ^a	622 ± 19 ^c	
g1	3.81 ± 0.2 ^a	112.2 ± 6.7 ^b	386 ± 16 ^e	2.4 ± 0.2 ^e
g2	3.88 ± 0.1 ^a	107.8 ± 6.5 ^b	350 ± 6 ^e	
h1	4.16 ± 0.8 ^a	144.3 ± 8.7 ^a	582 ± 11 ^d	7.3 ± 0.3 ^c
h2	4.73 ± 0.2 ^a	135.9 ± 8.2 ^a	574 ± 3 ^d	
j1	3.38 ± 0.4 ^b	135.7 ± 8.1 ^a	229 ± 8 ^f	0.8 ± 0.1 ^f
j2	3.63 ± 0.5 ^a	141.0 ± 8.5 ^a	262 ± 3 ^f	

m1	3.75 ± 0.1^a	139.3 ± 8.4^a	620 ± 12^c	8.6 ± 0.4^b
m2	4.01 ± 0.6^a	124.3 ± 7.5^a	677 ± 14^c	

376 Values are means \pm SD. Values with the same superscript letter in a given column are not significantly different
377 ($P \leq 0.05$).

378

379

380 **Table 3.** Protein solubilized from different pasta samples in various media.

381

Sample	50 mM phosphate buffer mg/g sample	+ 6 M urea mg/g sample	+ 6 M urea and 10 mM DTT mg/g sample
a1	1.78 ± 0.08 ^d	8.61 ± 0.54 ^d	28.35 ± 1.32 ^b
a2	1.97 ± 0.07 ^d	8.34 ± 0.53 ^d	32.04 ± 1.42 ^b
b1	3.34 ± 0.13 ^c	15.39 ± 0.95 ^c	33.31 ± 1.13 ^b
b2	3.54 ± 0.18 ^c	16.17 ± 0.98 ^c	32.42 ± 1.15 ^b
c1	6.79 ± 0.31 ^b	27.52 ± 1.50 ^a	39.39 ± 1.42 ^a
c2	6.08 ± 0.24 ^b	27.77 ± 1.04 ^a	37.45 ± 1.51 ^a
d1	2.43 ± 0.13 ^d	12.92 ± 0.97 ^c	25.12 ± 1.16 ^c
d2	3.01 ± 0.21 ^c	13.47 ± 0.98 ^c	23.08 ± 1.87 ^c
e1	4.25 ± 0.26 ^c	17.27 ± 1.41 ^c	34.09 ± 1.98 ^b
e2	5.26 ± 0.28 ^c	21.35 ± 1.82 ^b	38.94 ± 1.86 ^a
f1	5.79 ± 0.32 ^b	20.67 ± 1.12 ^b	40.48 ± 2.10 ^a
f2	5.11 ± 0.33 ^b	20.57 ± 0.99 ^b	37.24 ± 1.89 ^a
g1	3.52 ± 0.16 ^c	20.57 ± 1.05 ^b	32.41 ± 1.79 ^b
g2	4.64 ± 0.18 ^c	19.01 ± 1.11 ^b	31.14 ± 1.68 ^b
h1	3.01 ± 0.13 ^c	19.99 ± 1.23 ^b	35.60 ± 1.78 ^a
h2	3.85 ± 0.12 ^c	15.59 ± 1.14 ^c	39.60 ± 1.86 ^a
j1	7.56 ± 0.35 ^a	30.25 ± 1.71 ^a	43.98 ± 1.85 ^a
j2	6.75 ± 0.29 ^a	33.93 ± 1.87 ^a	42.27 ± 1.96 ^a
m1	4.85 ± 0.18 ^c	20.97 ± 1.23 ^b	33.60 ± 1.24

m2

4.54 ± 0.21^c

20.92 ± 1.25^b

36.66 ± 1.34^a

382 Values are means \pm SD. Values with the same superscript letter in a given column are not significantly different

383 ($P \leq 0.05$).


384

385

386

387

388



389

390

391

392

393

394

395

396

397

398


399

400

401

Figure 1 - SDS-PAGE of proteins in various whole grains pasta. Proper amounts of milled pasta were treated with denaturing buffer containing 1.7 % (w/w) SDS and 1% (w/w) 2-ME and denatured by boiling at 100°C. MW: molecular weight markers (range 14000-96000 Da).


402



403

404

405




406


407 Figure 2 - E-nose PCA score plot (Figure. 2A) and loading plot (Figure 2B) obtained by PCA elaboration of data
408 collected on uncooked pasta in combination with heat damage (FUR) and protein solubility in urea. The data
409 were elaborated in a correlation matrix where the first two principal components accounted for 84.8% of the
410 total variance.

411

412



413



414

415

416 Figure 3 - E-tongue PCA score plot (Figure. 2A) and loading plot (Figure 2B) obtained by PCA elaboration of data
417 collected on cooked pasta in combination with the e-nose data, accounting for 65% of the total variance.

418