

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

# A novel multifunctional cyclometallated iridium(III) complex with interesting second-order nonlinear optical properties and two-photon absorption activity.

Alessia Colombo,<sup>a,b,c</sup> Eleonora Garoni,<sup>a,e</sup> Claudia Dragonetti,<sup>\*a,b,c</sup> Stefania Righetto,<sup>a,b</sup> Dominique Roberto,<sup>a,b,c</sup> Nicolò Baggi,<sup>a</sup> Muriel Escadeillas,<sup>d</sup> Véronique Guerschais,<sup>\*d</sup> and Kenji Kamada<sup>\*e</sup>

<sup>a</sup>Dip. di Chimica dell'Università degli Studi di Milano, <sup>b</sup>UdR di Milano dell'INSTM, <sup>c</sup>Istituto di Scienze e Tecnologie Molecolari del CNR, via Golgi, 19, I-20133 Milano, Italy; <sup>d</sup>Institut des Sciences Chimiques de Rennes UMR 6226 CNRS-Université de Rennes 1, Campus de Beaulieu, 35042 Rennes Cedex, France; <sup>e</sup>IFMRI, National Institute of Advanced Industrial Science and Technology (AIST), 1-8-31 Midorigaoka, Ikeda, Osaka 563-8577, Japan.

Corresponding authors: [claudia.dragonetti@unimi.it](mailto:claudia.dragonetti@unimi.it), [veronique.guerchais@univ-rennes1.fr](mailto:veronique.guerchais@univ-rennes1.fr), [k.kamada@aist.go.jp](mailto:k.kamada@aist.go.jp)

## Abstract

34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

A novel cationic cyclometallated Ir(III) complex, namely [Ir(C<sup>N</sup>-ppy-4-CH=CHC<sub>6</sub>H<sub>4</sub>NEt<sub>2</sub>)<sub>2</sub>(N<sup>N</sup>-bpy-CH=CHC<sub>6</sub>H<sub>4</sub>NEt<sub>2</sub>)] (ppy= 2-phenylpyridine; bpy= 2,2'-bipyridine), was prepared and well characterized. Its good second-order NLO response, as determined in solution by the EFISH technique, and its interesting two-photon absorption activity, as measured by the Z-scan technique in the femtosecond case, make it a good candidate for application in photonics and biomedicine.

53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

*Keywords:* Second-order nonlinear optics; EFISH; two-photon absorption activity; cyclometallated iridium(III) complexes.

## Introduction

Coordination compounds with both luminescent and second-order nonlinear optical (NLO) properties are of great interest as multifunctional materials, as they may offer additional flexibility, when compared to organic NLO chromophores, by introducing new NLO active electronic charge-transfer transitions between the metal and the ligand, tuneable by virtue of the nature, oxidation state and coordination sphere of the metal centre [1,2].

Cyclometallating ligands are often incorporated in Ir(III) complexes for various applications such as in organic light-emitting diodes (OLEDs) [3], light-emitting electrochemical cells (LEECs) [4], and dye-sensitized solar cells (DSSCs) [5]. Whereas cyclometallated Ir(III) complexes have been intensively studied for their interesting luminescent properties, to our knowledge, only a few reports appeared in the literature on their second-order NLO properties [6, 7] or two-photon absorption (TPA) activity [8-11]. Yet, two-photon absorbing materials having large two-photon absorption cross-section ( $\sigma$ ), especially in the near-infrared (NIR) region, are required for photodynamic therapy, bio-imaging, upconversion lasing and drug delivery [9] whereas the design of novel second-order NLO active compounds is crucial for many applications in the fields of optical communication and data processing [2]. Beeby and co-workers reported a series of iridium(III) acetylacetonate complexes, bearing two cyclometallated substituted 2-phenylpyridines, characterized by fair values of TPA cross sections (approximately 20 GM) [10]. Another report concerning iridium complexes of a terpyridyl-stilbene system demonstrated TPA cross-sections in the range of 10-70 GM [11]. More recently, Das and co-workers reported the synthesis of cyclometallated iridium(III) complexes containing two cyclometallated unsubstituted 2-phenylpyridines and a 2,2'-bipyridine bearing highly  $\pi$ -delocalized substituents with TPA cross-sections in the range of 45-196 GM in the femtosecond case [8c]. Besides, some of us reported that iridium(III) acetylacetonate complexes bearing two substituted cyclometallated 4-styryl-2-phenylpyridines are characterized by a good second-order NLO response [7f], as determined by means of the Electric Field Induced Second Harmonic generation (EFISH) technique [12]. Remarkably, the simple luminescent cyclometallated Ir(III) complex  $[\text{Ir}(\text{C}^{\wedge}\text{N-ppy-4-CH=CHC}_6\text{H}_4\text{NEt}_2)_2(\text{acetylacetonate})]$  (ppy = 2-phenylpyridine) [13] shows a  $\mu\beta_{1.907} = -550 \times 10^{-48}$  esu and is a useful tool for the preparation of NLO active polymeric films [7f]. These interesting results prompted us to investigate the effect of the substitution of the acetylacetonate ligand by a  $\pi$ -delocalized bipyridine incorporating electron-donor substituents on the second-order NLO response. Therefore, we prepared the novel cationic cyclometallated

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Ir(III) complex  $[\text{Ir}(\text{C}^{\wedge}\text{N}\text{-ppy-4-CH=CHC}_6\text{H}_4\text{NEt}_2)_2(\text{N}^{\wedge}\text{N}\text{-bpy-CH=CHC}_6\text{H}_4\text{NEt}_2)]$  (**5**) (bpy=bipyridine; Figure 1) and studied its luminescent and NLO properties. Its second-order NLO response was measured in solution through the EFISH technique, working with an incident and non resonant wavelength of 1907 nm, whereas its two-photon absorption activity was evaluated by the Z-scan technique in the femtosecond case.




Figure 1. Investigated cyclometallated Ir(III) complex (**5**)

## Material and methods

### Synthesis

#### General comments

All reagents were purchased from Sigma-Aldrich and were used without further purification. Reactions requiring anhydrous conditions were performed under nitrogen. <sup>1</sup>H NMR spectra were recorded at 400 MHz on a Bruker AVANCE-400 instrument; chemical shifts ( $\delta$ ) are expressed in ppm relative to internal Me<sub>4</sub>Si as standard; signals were abbreviated as s, singlet; bs, broad singlet; d, doublet; t, triplet; q, quartet; m, multiplet. The UV-Vis spectra of the samples solution in dichloromethane were recorded by a Shimadzu UV-3150 spectrometer using 1 cm quartz cells.

**4-methyl-2-phenylpyridine (1).** An excess of lithium (1.82 g, 272 mmol) was suspended in 40 mL of Et<sub>2</sub>O. The suspension was cooled down to 0°C and bromobenzene (21.34 g, 136 mmol) in 50 mL of Et<sub>2</sub>O was added dropwise during 20 minutes. The reaction mixture was stirred overnight at room temperature. The freshly-prepared phenyllithium was added dropwise to a solution of 4-picoline (3.16 g, 34 mmol) in 90 mL toluene. The reaction was stirred at 30°C for 24 hours, distilled water was added dropwise, the residue was extracted with CH<sub>2</sub>Cl<sub>2</sub>, washed with water, dried over MgSO<sub>4</sub> and the solvent was removed under

1 reduced pressure. The product was purified by column chromatography (silica, *n*-  
2 heptane:AcOEt=9:1) to afford **1** as an orange/brown solid (2.44 g). Yield: 42%. <sup>1</sup>H-NMR  
3 (400 MHz, CDCl<sub>3</sub>) δ (ppm): 8.58 (1H, d, *J* = 5.0 Hz), 8.00 (2H, d, *J* = 7.2 Hz), 7.57 (1H, s),  
4 7.49 (2H, t, *J* = 7.4 Hz), 7.43 (1H, t, *J* = 7.2 Hz), 7.08 (1H, d, *J* = 4.9 Hz), 2.44 (3H, s).  
5  
6

7 **4-*N,N*-diethylaminostyryl-2-phenylpyridine (2)**. 4-Methyl-2-phenylpyridine (500 mg, 2.96  
8 mmol), 4-*N,N*-diethylaminobenzaldehyde (620 mg, 3.5 mmol) and *t*-BuOK (700 mg, 6.2  
9 mmol) were dissolved in DMF (20 mL). The reaction mixture was heated at reflux overnight,  
10 then CH<sub>2</sub>Cl<sub>2</sub> (150 mL) was added. The organic phase was washed first with water and then  
11 with an aqueous solution of 1 M KOH, dried with MgSO<sub>4</sub> and the solvent was removed under  
12 reduced pressure. The crude product was purified by column chromatography (silica,  
13 CH<sub>2</sub>Cl<sub>2</sub>:petroleum ether = 7:3) to obtain **2** as a yellow powder (546 mg). Yield: 56%. <sup>1</sup>H  
14 NMR (400 MHz, CDCl<sub>3</sub>) δ (ppm): 8.61 (1H, d, *J* = 5.2 Hz), 8.06 (2H, d, *J* = 7.2 Hz), 7.75  
15 (1H, s) 7.51 (2H, t, *J* = 7.4 Hz), 7.45 (3H, d, *J* = 8.5 Hz), 7.30 (1H, d, *J* = 16.3 Hz), 7.26 (1H,  
16 m), 6.86 (1H, d, *J* = 16.2 Hz), 6.68 (2H, d, *J* = 8.8 Hz), 3.40 (4H, q, *J* = 7.1 Hz), 1.20 (6H, t, *J*  
17 = 7.0 Hz).  
18  
19

20 **[Ir(C<sup>N</sup>-ppy-4-CH=CHC<sub>6</sub>H<sub>4</sub>NEt<sub>2</sub>)<sub>2</sub>Cl]<sub>2</sub> (3)**. IrCl<sub>3</sub> • 3H<sub>2</sub>O (165 mg, 0.55 mmol) and **2** (550  
21 mg, 1.66 mmol) were suspended in a mixture of 2-ethoxyethanol:H<sub>2</sub>O 3:1 (16 mL) in a  
22 Schlenk tube under inert atmosphere. The reaction mixture was heated at reflux for 24 h. The  
23 precipitate was decanted by centrifugation and the solid product was then washed  
24 successively with H<sub>2</sub>O, EtOH and Et<sub>2</sub>O. The solvent was removed with cannula to afford **3** as  
25 an orange powder (290 mg). Yield: 30%. <sup>1</sup>H-NMR (400 MHz, CD<sub>2</sub>Cl<sub>2</sub>) δ (ppm): 9.12 (4H, d,  
26 *J* = 5.9 Hz), 7.94 (4H, s), 7.63 (4H, d, *J* = 7.1 Hz), 7.41 (12H, m), 7.04 (4H, d, *J* = 16 Hz),  
27 6.93 (4H, m), 6.83 (4H, t, *J* = 7.0 Hz), 6.64 (12H, m), 6.05 (4H, d, *J* = 8.0 Hz), 3.48 (16H, q, *J*  
28 = 6.8 Hz), 1.25 (24H, t, *J* = 6.8 Hz).  
29  
30

31 **4,4'-*p*-[(*N,N*-diethyl)aminostyryl]-2,2'-bipyridine (4)**. Diisopropylamine (607.1 mg, 6  
32 mmol) was dissolved in THF (2.5 mL) in a Schlenk tube and the mixture was cooled to 0°C.  
33 A *n*-BuLi solution (1.6 M in hexane, 3.75 mL, 6 mmol) was added slowly. After stirring one  
34 hour, the solution was cooled to -78°C and 4,4'-dimethyl-2,2'-bipyridine (501.1 mg, 2.72  
35 mmol) in THF (13 mL) was added dropwise and the mixture stirred at low temperature for 2  
36 hours. *p*-(*N,N*-diethyl)aminobenzaldehyde (957.1 mg, 5.4 mmol) was dissolved in 6 mL of  
37 THF (6 mL) and added slowly. The resulting mixture was stirred for one hour at low  
38 temperature and overnight at room temperature. After hydrolyzing with distilled water (12  
39 mL), the phases were separated, the organic phase was dried over MgSO<sub>4</sub>, filtered and the  
40 solvent was removed under reduced pressure. The dialcohol was dissolved in toluene (40 mL)  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 in a round-bottomed flask and pyridinium *p*-toluenesulfonate (PPTS, 67.6 mg, 0.27mmol)  
2 was added. The mixture was heated at reflux for 4 hours. After cooling to room temperature,  
3 the solvent was removed under reduced pressure and dichloromethane (16 mL) was added.  
4 The mixture was washed with a saturated solution of NaHCO<sub>3</sub>. The precipitated bipyridine  
5 was filtered, abundantly washed first with H<sub>2</sub>O and then with Et<sub>2</sub>O to obtain **4** as an orange  
6 powder (970 mg). Yield: 71%. <sup>1</sup>H-NMR (400 MHz, CD<sub>3</sub>Cl) δ (ppm): 8.63 (2H, d, *J* = 5.1 Hz),  
7 8.48 (2H, s), 7.47 (4H, d, *J* = 8.9 Hz), 7.42 (2H, d, *J* = 16.2 Hz), 7.36 (2H, dd, *J* = 5.3, 1.8  
8 Hz), 6.92 (2H, d, *J* = 16.2 Hz), 6.71 (4H, d, *J* = 8.9 Hz), 3.43 (8H, q, *J* = 7.1 Hz), 1.22 (12 H,  
9 t, *J* = 7.1 Hz).  
10  
11  
12  
13  
14  
15

16 **[Ir(C<sup>N</sup>-ppy-4-CH=CHC<sub>6</sub>H<sub>4</sub>NEt<sub>2</sub>)<sub>2</sub>(N<sup>N</sup>-bpy-CH=CHC<sub>6</sub>H<sub>4</sub>NEt<sub>2</sub>)] [PF<sub>6</sub>] (**5**)**. A suspension  
17 of **3** (250 mg, 0.14 mmol), **4** (178 mg, 0.35 mmol) and AgPF<sub>6</sub> (88 mg, 0.35 mmol) in 1,2-  
18 dichloroethane (20 mL) was kept at reflux for 2 hours and stirred at room temperature for 16  
19 hours. The solvent was removed under reduced pressure leaving a residue which was washed  
20 with Et<sub>2</sub>O (3 x 20 mL). The compound was precipitated from Et<sub>2</sub>O/CH<sub>2</sub>Cl<sub>2</sub> giving the desired  
21 complex (**5**) as a red solid after filtration (50 mg). Yield: 13%. <sup>1</sup>H NMR (500 MHz, CD<sub>2</sub>Cl<sub>2</sub>) δ  
22 (ppm): 8.32 (2H, s), 7.93 (2H, s), 7.84 (2H, d, *J* = 5.7 Hz), 7.80 (2H, d, *J* = 7.4 Hz), 7.42  
23 (18H, m), 7.08 (2H, t, *J* = 7.7 Hz), 7.02 (2H, d, *J* = 4.5 Hz), 6.95 (4H, m), 6.86 (2H, d, *J* =  
24 16.0 Hz), 6.68 (8H, dd, *J* = 16, 8.9 Hz), 6.46 (2H, d, *J* = 7.9 Hz), 3.41 (16H, m), 1.20 (24H,  
25 m). Elem. Anal. Calcd for C<sub>80</sub>H<sub>84</sub>F<sub>6</sub>IrN<sub>7</sub>P: C, 71.19; H, 6.27; N, 8.30. Found: C, 71.65; H,  
26 6.28; N, 8.33.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36

### 37 EFISH measurements

38  
39 EFISH measurements [12] were carried out at the Dipartimento di Chimica of the Università  
40 di Milano, in CH<sub>2</sub>Cl<sub>2</sub> solutions at a concentration of 1 x 10<sup>-3</sup> M, working with an incident  
41 non-resonant wavelength of 1.907 μm, obtained by Raman-shifting the fundamental 1.064 μm  
42 wavelength produced by a Q-switched, mode-locked Nd<sup>3+</sup>:YAG laser manufactured by  
43 Atalaser. The apparatus for the EFISH measurements is a prototype made by SOPRA  
44 (France). The reported μβ<sub>EFISH</sub> values are the mean values of 16 successive measurements  
45 performed on the same sample. The sign of μβ is determined by comparison with the  
46 reference solvent (CH<sub>2</sub>Cl<sub>2</sub>).  
47  
48  
49  
50  
51  
52  
53  
54  
55


### 56 Two-photon absorption activity measurements

57  
58 All open-aperture Z-scan measurements [13] were performed at AIST in Osaka, for the  
59 solution of the complex **5** held in a 2-mm quartz cuvette. The setup for the Z-scan  
60  
61  
62  
63  
64  
65

1 measurements were reported elsewhere [14]. A femtosecond, wavelength-tunable laser  
 2 (optical parametric amplifier, *SpectraPhysics TOPAS Prime*) was employed as light source to  
 3 obtain the spectrum. Typical pulsewidth was 120 fs and the repetition rate was 1 kHz. First,  
 4 the outline of the TPA spectrum was measured at a single excitation power (0.4 mW,  
 5 corresponding to the on-axis optical intensity at the focal point of 99-150 GW/cm<sup>2</sup>) by  
 6 assuming that the observed open-aperture Z-scan signal originates from the TPA process  
 7 (open dots in Fig. 4). Then the assumption was confirmed and the precise value of the TPA  
 8 cross section ( $\sigma^{(2)}$ ) were obtained by changing the excitation power (filled dots with  
 9 experimental errors in Fig. 4). Finally, the relative TPA spectrum (open dots) were  
 10 superimposed the absolute TPA spectrum (filled dots). For all measurements, in-house  
 11 standard compounds (MPPBT [15] in DMSO) was measured at the same condition and used  
 12 for the spectral correction of  $\sigma^{(2)}$ .  
 13  
 14  
 15  
 16  
 17  
 18  
 19  
 20  
 21  
 22  
 23

## 24 Results and Discussion


25 The complex  $[\text{Ir}(\text{C}^{\wedge}\text{N-ppy-4-CH=CHC}_6\text{H}_4\text{NEt}_2)_2(\text{N}^{\wedge}\text{N-bpy-CH=CHC}_6\text{H}_4\text{NEt}_2)][\text{PF}_6]$   
 26 (**5**) was prepared as shown in Figure 2.  
 27  
 28  
 29  
 30  
 31  
 32  
 33



**Figure 2.** (i) *t*-BuOK, DMF, 155°C, 15 h; (ii) IrCl<sub>3</sub>·3H<sub>2</sub>O, 2-ethoxyethanol/H<sub>2</sub>O, 100°C, 24 h; (iii) a) *n*-BuLi, iPr<sub>2</sub>NH, *p*-(*N,N*-diethyl)aminobenzaldehyde, THF, 0°C → -78°C → rt, 15 h; b) PPTS, toluene, 110°C, 4 h; (iv) AgPF<sub>6</sub>, 1,2-dichloroethane, 84°C, 2 h → rt, 15 h.

1  
2 The pro-ligand 4-(*N,N*-diethyl)aminostyryl-2-phenylpyridine (**2**) was readily prepared upon  
3 deprotonation of the methyl group of 4-methyl-2-phenylpyridine, obtained by reaction of  
4 phenyllithium with 4-picoline, followed by condensation with *p*-(*N,N*-  
5 diethyl)aminobenzaldehyde, as previously reported [16]. Heating at reflux this pro-ligand  
6 with IrCl<sub>3</sub> • 3H<sub>2</sub>O in a 2-ethoxyethanol:H<sub>2</sub>O 3:1 mixture for 24 h afforded the related  $\mu$ -  
7 dichloro-bridged cyclometallated iridium dimer intermediate (**3**) [16]. The ancillary ligand  
8 4,4'-*p*-[(*N,N*-diethyl)aminostyryl]-2,2'-bipyridine (**4**) was prepared like **2** but starting from  
9 4,4'-dimethyl-2,2'-bipyridine [17]. The desired complex (**5**) was finally obtained by refluxing  
10 in 1,2-dichloroethane for 2 hours this  $\pi$ -delocalized substituted bipyridine with the  $\mu$ -  
11 dichloro-bridged cyclometallated iridium dimer. It was obtained as a dark red powder and  
12 well characterized (see Material and methods).  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

23 The absorption spectrum of complex **5** was registered both in dichloromethane and  
24 dimethyl sulfoxide (DMSO, Figure 3a) whereas the photophysical data are listed in Table 1.  
25 By analogy with other cationic iridium(III) complexes bearing two cyclometallated  
26 phenylpyridines and a bipyridine [6e], the absorption bands under 350 nm can reasonably be  
27 assigned to <sup>1</sup>( $\pi$ - $\pi^*$ ) transitions of the styrylphenylpyridine and styrylbipyridine ligands, the  
28 intense band between 350 and 550 nm can be assigned to an Intra-Ligand Charge Transfer  
29 (ILCT) transition in which the pyridyl moiety acts as a  $\pi^*$ -acceptor group hiding the (d(Ir)  $\rightarrow$   
30  $\pi^*(ppy/bpy)$ ) MLCT band, while the weak band centered at 620 nm may be due to direct  
31 population of triplet charge transfer states [6e]. A pronounced solvatochromic behavior is  
32 observed going from DMSO to CH<sub>2</sub>Cl<sub>2</sub>, typical of electronic transitions with an appreciable  
33 degree of charge-transfer character; the blue shift of the band with increasing solvent polarity  
34 suggests a lower dipole moment in the excited state than in the ground state [18].  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



**Figure 3.** **a)** Superposition of absorption spectra of complex **5** in DMSO (red) and CH<sub>2</sub>Cl<sub>2</sub> (blue); **b)** Excitation spectrum at 77 K (blue line), emission spectrum at 298 K (black line) and emission spectrum at 77 K (red line), all in a degassed 5 × 10<sup>-5</sup> M THF solution of **5** (λ<sub>exc</sub> = 450 nm).

TI K

are shown in Figure 3b, and the data are summarized in Table 1. The novel complex is only very weakly emissive in fluid solution at room temperature (Figure 3b), because the radiative decay is quenched by the known competitive non-radiative *trans-cis* isomerization process of the C-C double bond, triggered by excitation with light before emission, of bipyridines linked to an Ir(III) center [6e, 19].

$\lambda_{\text{abs}} / \text{nm}$ ( $\epsilon \times 10^3 / \text{M}^{-1}\text{cm}^{-1}$ ) <sup>a</sup>	$\lambda_{\text{em}} / \text{nm}$ <sup>b</sup>	$\tau / \mu\text{s}$ <sup>b</sup>	$\mu\beta_{1.907} / \times 10^{-48} \text{esu}^c$
454 (7.54), 621 (0.59)	669, 736	57	-960

**Table 1.** Photophysical data for complex **5**. [a] In CH<sub>2</sub>Cl<sub>2</sub> solution at 298 K; in DMSO: λ<sub>abs</sub> = 440 nm, ε × 10<sup>3</sup> = 5.37 M<sup>-1</sup>cm<sup>-1</sup>. [b] In THF at 77 K (λ<sub>exc</sub> = 460 nm). [c] In CH<sub>2</sub>Cl<sub>2</sub> solution (10<sup>-3</sup>M); uncertainty in EFISH measurements is ±10%

On the other hand, **5** is emissive at 77 K in THF, the luminescence spectrum is highly structured and the emission is located in the red region of the visible spectrum, as predicted upon incorporation of extended π-conjugated substituents in the ligands. With reference to previously reported Ir(III) complexes bearing two cyclometallated phenylpyridines and one bipyridine [6e], considering the highly structured spectrum and the long lifetime (57 μs), luminescence can be assigned to radiative decay from a triplet state predominantly localized on the styryl-substituted ligands, rather than from the MLCT state. In any case the



1 participation of the metal in the emission is necessary to promote the population of the  
2 formally forbidden triplet emissive state.


3 The quadratic hyperpolarizability of complex **5** was measured by the solution phase  
4 direct current EFISH generation method, which can provide direct information on the intrinsic  
5 molecular NLO properties through eq 1:  
6  
7

$$\gamma_{\text{EFISH}} = (\mu\beta_{\lambda}/5kT) + \gamma(-2\omega; \omega, \omega, 0) \quad (1)$$

8  
9  
10  
11  
12  
13 where  $\mu\beta_{\lambda}/5kT$  is the dipolar orientational contribution and  $\gamma(-2\omega; \omega, \omega, 0)$  is the third-order  
14 polarizability corresponding to the mixing of two optical fields at  $\omega$  and the dc poling field at  
15  $\omega=0$ . This latter term is often referred as the electronic cubic contribution to  $\gamma_{\text{EFISH}}$ , which is  
16 usually negligible.  $\beta_{\lambda}$  is the projection along the dipole moment ( $\mu$ ) axis of the vectorial  
17 component of the tensor of the quadratic hyperpolarizability at the incident wavelength  $\lambda$ . All  
18 EFISH measurements were carried out working in  $\text{CH}_2\text{Cl}_2$  solutions at a concentration of  $10^{-3}$   
19 M, with a non resonant incident wavelength of 1.907  $\mu\text{m}$ , obtained by Raman-shifting in a  
20 high pressure  $\text{H}_2$  of the fundamental 1.064  $\mu\text{m}$  wavelength produced by a Q-switched, mode-  
21 locked  $\text{Nd}^{3+}$ :YAG laser.  
22  
23  
24  
25  
26  
27  
28  
29

30 It turned out that complex **5** is characterized by a negative value of  $\mu\beta_{1.907}$  ( $-960 \times 10^{-48}$  esu,  
31 Table 1) in agreement with the observed solvatochromic behavior and with a negative value  
32 of  $\Delta\mu_{\text{eg}}$  (difference of the dipole moment in the excited and ground state) upon excitation [2].  
33 It is worth pointing out that the absolute  $\mu\beta_{1.907}$  value of complex **5** is almost twice that  
34 reported for related  $[\text{Ir}(\text{C}^{\wedge}\text{N}\text{-ppy-4-CH=CHC}_6\text{H}_4\text{NEt}_2)_2(\text{acetylacetonate})]$  (ppy = 2-  
35 phenylpyridine) [7e] suggesting that 4,4'-*p*-[(*N,N*-diethyl)aminostyryl]-2,2'-bipyridine is much  
36 better than acetylacetonate as ancillary ligand in this kind of cyclometallated Ir(III) complexes  
37 for second-order nonlinear optics.  
38  
39  
40  
41  
42  
43  
44

45 Complex **5** was then investigated for its two-photon absorption activity. It was  
46 dissolved in DMSO (0.55 mM) and studied with the z-scan technique in the femtosecond case.  
47 The TPA spectrum is shown in Figure 4.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



**Figure 4.** The OPA (line) and TPA (dots) spectra of complex **5**.

Remarkably, the two-photon absorption cross-sections are quite high, in the range 100-250 GM at 600-1000 nm. Contrarily to previously reported TPA active cationic iridium(III) complexes having two cyclometallated unsubstituted 2-phenylpyridines and a 2,2'-bipyridine bearing highly  $\pi$ -delocalized substituents [8c], the complex investigated here has less extended substituents on the bipyridine but it bears them also on the 2-phenylpyridine ligands. The excellent correspondence of the observed TPA peak with the wavelength-doubled one-photon absorption peak, as shown in Figure 4, suggests that the TPA peak mainly originates from the dipolar contribution [20] governed by asymmetric charge transfer, similar to the EFISH property. In this way, the charge-transfer property and the extended  $\pi$ -conjugation over the whole molecule results in good TPA properties of **5** compared to the previously reported complexes of similar structures. Besides, it is worth pointing out that some iridium(III) complexes bearing two cyclometallated unsubstituted 2-phenylpyridines and a 2-phenylimidazo[4,5-f][1,10]phenanthroline, characterized by two-photon absorption cross-sections in the range 48.8-65.5 GM at 750 nm, were utilized as efficient agents for mitochondrial imaging and tracking in living cells under two-photon (750 nm) excitation [21]. Similarly, complex **5** can find application in biomedical research.

## Conclusion

In conclusion, our work put in evidence how a careful choice of the coordination sphere of cyclometallated iridium(III) complexes can lead to high second-order nonlinear optical properties and two-photon absorption activity, correlated through the charge-transfer character of the complex. Substitution of acetylacetonate by a 4,4'-*p*-[(*N,N*-diethyl)aminostyryl]-2,2'-bipyridine is a simple way to increase the second-order NLO response in this kind of cyclometallated Ir(III) complexes. In addition the use of simple  $\pi$ -delocalized donor substituents on both the cyclometallated 2-phenylpyridine and 2,2'-bipyridine allows to reach high two-photon absorption sections in a wide range of wavelengths. Clearly the novel complex reported here has a great potential for application in photonics and biomedicine.

## Acknowledgements

This work was supported by MIUR, in Italy, and by a Grant-in-Aid for Scientific Research on Innovative Areas "Photosynergetics" (No. JP26107004, K.K.) from MEXT and a Grant-in-Aid for Scientific Research (No. JP25248007, K.K.) from JSPS, Japan. We thank CNRS for a financial support (PICS Rennes-Milan) and l'Università Italo Francese (Progetto Galileo 2015/2016; G15\_50) for mobility support. Regione Lombardia and Fondazione Cariplo are gratefully acknowledged for the use of instrumentation purchased through the SmartMatLab Centre project (2014). We thank Dr. Daniele Marinotto for luminescence and lifetime measurements.

## References

- [1] J. Zyss, *Molecular Nonlinear Optics: Materials, Physics and Devices*, Academic Press, Boston, 1994.
- [2] (a) B.J. Coe, in *Comprehensive Coordination Chemistry II*; Vol. 9, Elsevier Pergamon: Oxford, U.K., 2004. (b) B.J. Coe, N.R.M Curati, *Comments on Inorg. Chem.* 25 (2004) 147-184. (c) O. Maury, H. Le Bozec, *Acc. Chem. Res.* 38 (2005) 691-704. (d) C.E. Powell, M.G. Humphrey, *Coord. Chem. Rev.* 248 (2004) 725-756. (e) E. Cariati, M. Pizzotti, D. Roberto, F. Tessore, R. Ugo, *Coord. Chem. Rev.* 250 (2006) 1210-1233 (f) B. J. Coe, *Acc. Chem. Res.* 39 (2006) 383-393. (g) M.G.Humphrey, M. Samoc, *Adv. Organomet. Chem.* 55 (2007) 61-136.

1 (h) J. Boixel, V. Guerchais, H. Le Bozec, A. Chantzis, D. Jacquemin, A. Colombo, C.  
2 Dragonetti, D. Marinotto, D. Roberto *Chem.Comun.* 51 (2015) 7805-7808.

3 (i) S. Di Bella, C. Dragonetti, M. Pizzotti, D. Roberto, F. Tessore, R. Ugo in *Topics in*  
4 *Organometallic Chemistry* 28. *Molecular Organometallic Materials for Optics*, Vol. 28 (Eds:  
5 H. Le Bozec, V. Guerchais), Springer, 2010, 1-55. (l) O. Maury, H. Le Bozec, in *Molecular*  
6 *Materials*; D.W. Bruce, D. O'Hare and R.I. Walton, Eds.; Wiley: Chichester, 2010, 1-59.

7 [3] (a) R.C. Evans, P. Douglas, C.J. Winscom, *Coord. Chem Rev.* 250 (2006) 2093-2126 and  
8 references therein. (b) L. Flamigni, A. Barbieri, C. Sabatini, B. Ventura, F. Barigelletti, *Top.*  
9 *Curr. Chem.* 281 (2007) 143-203. (c) G. Zhou, W.Y. Wong S. Suo, *J. Photochem. Photobio.*  
10 *C: Photochem. Rev.* 11 (2010) 133-156. (d) W.Y. Wong, C.L. Ho, *Coord. Chem. Rev.* 253  
11 (2009) 1709-1758. (e) W.Y. Wong, C.L. Ho, *J. Mater. Chem.* 19 (2009) 4457-4482. (f) G.  
12 Zhou, W.Y. Wong, X. Yang, *Chem. Asian J.* 6 (2011) 1706-1727. (g) H. Wu, G. Zhou, J.  
13 Zou, C.L. Ho, W.Y. Wong, W. Yang, J. Peng, Y. Cao, *Adv. Mater.* 21 (2009) 4181-4184.

14 [4] (a) J.D. Slinker, J. Rivnay, J.S. Moskowitz, J.B. Parker, S. Bernhard, H.D. Abruna, G.G.  
15 Malliaras, *J. Mater. Chem.* 17 (2007) 2976-2988. (b) E. Zysman-Colman, J.D. Slinker, J.B.  
16 Parker, G.G. Malliaras, S. Bernhard, *Chem. Mater.* 20 (2008) 388-396. (c) H.J. Bolink, L.  
17 Cappelli, E. Coronado, M. Graetzel, E. Ortí, R.D. Costa, M. Viruela, M.K. Nazeeruddin, *J.*  
18 *Am. Chem. Soc.* 128 (2006) 14786-14787. (d) E. Margapoti, V. Shukla, A. Valore, A.  
19 Sharma, C. Dragonetti, C. Kitts, D. Roberto, M. Murgia, R. Ugo M. Muccini, *J. Phys. Chem.*  
20 *C* 113 (2009) 12517-12522. (e) W.Y. Wong, G.J. Zhou, X.M. Yu, H.S. Kwok, Z. Lin, *Adv.*  
21 *Funct. Mater.* 17 (2007) 315-323. (f) E. Margapoti, M. Muccini, A. Sharma, A. Colombo, C.  
22 Dragonetti, D. Roberto, A. Valore, *Dalton. Trans.* 41 (2012) 9227-9231. (g) N.M. Shavaleev,  
23 R. Scopelliti, M. Graetzel, M.K. Nazeeruddin, A. Pertegas, C. Roldan-Carmona, D. Torderab,  
24 H.J. Bolink, *J. Mater. Chem. C*, 1 (2013) 2241-2248. (h) A. F. Henwood, E. Zysman-Colman,  
25 *Top. Curr. Chem.* 374:36 (2016) 1-41. (i) R. D. Costa, E. Ortí, H. J. Bolink, F. Monti, G.  
26 Accorsi, N. Armaroli, *Angew. Chem., Int. Ed.* 51 (2012) 8178-8211.

27 [5] (a) E.I. Mayo, K. Kilsa T. Tirrell, P.I. Djurovich, A. Tamayo, M.E. Thompson, N.S.  
28 Lewis, H.B. Gray, *Photochem. Photobiol. Sci.* 5 (2006) 871-873. (b) E. Baranoff, J.H. Yum,  
29 M. Graetzel, Md.K. Nazeeruddin, *J. Organomet. Chem.* 694 (2009) 2661-2670. (c) E.  
30 Baranoff, J.H. Yum, I. Jung, R. Vulcano, M. Graetzel, Md.K. Nazeeruddin, *Chem. Asian J.* 5  
31 (2010) 496-499. (d) Y. Shinpuku, F. Inui, M. Nakai, Y. Nakabayashi, *J. Photochem. and*  
32 *Photobio. A: Chemistry* 222 (2011) 203-209. (e) C. Dragonetti, A. Valore, A. Colombo, S.  
33 Righetto, V. Trafiletti, *Inorg. Chim. Acta* 388 (2012) 163-167.

1 [6] (a) C. Dragonetti, S. Righetto, D. Roberto, R. Ugo, A. Valore, S. Fantacci, A. Sgamellotti,  
2 F. De Angelis, *Chem. Commun.* (2007) 4116-4118. (b) C. Dragonetti, S. Righetto, D.  
3 Roberto, A. Valore, T. Benincori, F. Sannicolò, F. De Angelis, S. Fantacci, *J Mater Sci: Mater*  
4 *Electron* 20 (2009) S460–S464. (c) C. Dragonetti, S. Righetto, D. Roberto, A. Valore, *Phys.*  
5 *Status Solidi, C6*, No. S1 (2009) S50–S53. (d) A. Valore, E. Cariati, C. Dragonetti, S.  
6 Righetto, D. Roberto, R. Ugo, F. De Angelis, S. Fantacci, A. Sgamellotti, A. Macchioni, D.  
7 Zuccaccia, *Chem. Eur. J.* 16 (2010) 4814-4825. (e) V. Aubert, L. Ordroneau, M. Escadeillas,  
8 J.A.G. Williams, A. Boucekkine, E. Coulaud, C. Dragonetti, S. Righetto, D. Roberto, R. Ugo,  
9 A. Valore, A. Singh, J. Zyss, I. Ledoux-Rak, H. Le Bozec, V. Guerchais, *Inorg. Chem.* 50  
10 (2011) 5027-5038.

11 [7] (a) A. Valore, A. Colombo, C. Dragonetti, S. Righetto, D. Roberto, R. Ugo, F. De Angelis,  
12 S. Fantacci, *Chem. Commun.* 46 (2010) 2414–2416. (b) C. Dragonetti, S. Righetto, D.  
13 Roberto, R. Ugo, A. Valore, F. Demartin, F. De Angelis, A. Sgamellotti, S. Fantacci, *Inorg.*  
14 *Chim. Acta* 361 (2008) 4070-4076. (c) C. Dragonetti, S. Righetto, D. Roberto, R. Ugo, A.  
15 Valore, I. Ledoux-Rak, *Nonlinear Optics, Quantum Optics* 43 (2012) 197-204. (d) M. Zaarour,  
16 V. Guerchais, H. Le Bozec, C. Dragonetti, S. Righetto, D. Roberto, F. De Angelis, S. Fantacci,  
17 M.G. Lobello, *Dalton Trans.* 42 (2013) 155–159. (e) M. Zaarour, A. Singh, C. Latouche,  
18 J.A.G. Williams, I. Ledoux-Rak, J. Zyss, A. Boucekkine, H. Le Bozec, V. Guerchais, C.  
19 Dragonetti, A. Colombo, D. Roberto, A. Valore, *Inorg. Chem.* 52 (2013) 7987–7994. (f) C.  
20 Dragonetti, A. Colombo, D. Marinotto, S. Righetto, D. Roberto, A. Valore, M. Escadeillas, V.  
21 Guerchais, H. Le Bozec, A. Boucekkine, C. Latouche, *J. Organometal. Chem.* 751 (2014)  
22 568-572.

23 [8] (a) Y. Fan, D. Ding, D. Zhao, *Chem. Commun.* 51 (2015) 3446. (b) C.-L. Ho, K.-L. Wong,  
24 H.-K. Kong, Y.-M. Ho, C. T.-L. Chan, W.-M. Kwok, K. S.-Y. Leung, H.-L. Tam, M. H.-W.  
25 Lam, X.-F. Ren, A.-M. Ren, J.-K. Feng, W.-Y. Wong, *Chem. Commun.* 48 (2012) 2525–2527.  
26 (c) M. Sarma, T. Chatterjee, R. Bodapati, K. N. Krishnakanth, S. Hamad, S. V. Rao, S. K.  
27 Das, *Inorg. Chem.* 55 (2016) 3530-3540. (d) W.-J Xu, S.-J. Liu, X. Zhao, N. Zhao, Z.-Q. Liu,  
28 H. Xu, H. Liang, Q. Zhao, X.-Q. Yu, W. Huang, *Chem. Eur. J.* 19 (2013) 621. (e) C. Jin, J.  
29 Liu, Y. Chen, L. Zeng, R. Guan, C. Ouyang, L. Ji, H. Chao, *Chem. Eur. J.* 21 (2015) 12000.

30 [9] (a) M. Pawlicki, H. A. Collins, R. G. Denning, H. L. Anderson, *Angew. Chem. Int. Ed.* 48  
31 (2009) 3244-3266. (b) L. K. McKenzie, I. V. Sazanovich, E. Baggaley, M. Bonneau, V.  
32 Guerchais, J. A. G. Williams, J. A. Weinstein, H. E. Bryant, *Chem. Eur. J.* 23 (2017) 234-238.

33 [10] R. M. Edkins, S.L. Bettington, A.E. Goeta, A. Beeby, *Dalton Trans.* 40 (2011) 12765.

34 [11] L.S. Natrajan, A. Toulmin, A. Chew, S.W. Magennis, *Dalton Trans.* 39 (2010) 10837.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [12] (a) B. F. Levine, C. G. Bethea, *Appl. Phys. Lett.* 24 (1974) 445-447. (b) K. D. Singer, A. F. Garito, *J. Chem. Phys.* 75 (1981) 3572-3580. (c) I. Ledoux, J. Zyss, *J. Chem. Phys.* 73 (1982) 203-213.
- [13] M. Sheik-Bahae, A. A. Said, T.-H. Wei, D. G. Hagan, E. W. van Stryland, *IEEE J. Quant. Electr.* 26 (1990) 760–769.
- [14] K. Kamada, K. Matsunaga, A. Yoshino, K. Ohta, *J. Opt. Soc. Am. B* 20 (2003) 529–537.
- [15] K. Kamada, Y. Iwase, K. Sakai, K. Kondo, K. Ohta, *J. Phys. Chem. C* 113 (2009) 11469–11474.
- [16] M. Lepeltier, T. K.-M. Lee, K. K.-W. Lo, L. Toupet, H. Le Bozec, V. Guerschais, *Eur. J. Inorg. Chem.* (2007) 2734-2747.
- [17] T. Le Bouder, L. Viau, J.-P. Guégan, O. Maury, H. Le Bozec, *Eur. J. Org. Chem.* (2002) 3024-3033
- [18] E. Rossi, A. Colombo, C. Dragonetti, S. Righetto, D. Roberto, R. Ugo, A. Valore, J.A.G. Williams, M.G. Lobello, F. De Angelis, S. Fantacci, I. Ledoux-Rak, A. Singh, *J. Zyss Chem. Eur. J.* 19 (2013) 9875-9883.
- [19] A. Colombo, C. Dragonetti, D. Roberto, A. Valore, C. Ferrante, I. Fortunati, A. L. Picone, F. Todescato, J. A. G. Williams, *Dalton Trans.* 44 (2015) 15712-15720.
- [20] K. Kamada, *Disp. Imag.*,1(2014) 107–130.
- [21] C. Jin, J. Liu, Y. Chen, L. Zeng, R. Guan, C. Ouyang, L. Ji, and H. Chao, *Chem. Eur. J.* 21 (2015) 12000-12010.