

Variation in blood transfusion and coagulation management in Traumatic Brain Injury at the Intensive Care Unit: A survey in 66 neurotrauma centers participating in the Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI) study.

Running title

Transfusion and coagulation management

Table of contents title

Variation in transfusion and coagulation management in European neurotrauma centers

Authors

Jilske A. Huijben, MD.¹, Mathieu van der Jagt, MD, PhD.², Maryse C. Cossen, MSc.¹, Marieke J.H.A. Kruij, MD, PhD.³, Iain K. Haitsma, MD.⁴, Nino Stocchetti, MD.⁵, Andrew I.R. Maas, MD, PhD.⁶, David K. Menon, MD, PhD.⁷, Ari Ercole, MD, PhD.⁷, Marc Maegele, MD, PhD.⁸, Simon J. Stanworth, MD.⁹, Giuseppe Citerio, MD.¹⁰, Suzanne Polinder, PhD.¹, Ewout W. Steyerberg, PhD.^{1,11} and Hester F. Lingsma, PhD.¹ on behalf of the CENTER-TBI investigators.

Affiliations

1. Center for Medical Decision Sciences, Department of Public Health, Erasmus MC – University Medical Center Rotterdam, Rotterdam, the Netherlands
2. Department of Intensive Care (Office H-611) and Erasmus MC Stroke Center, Erasmus Medical Center Rotterdam, P.O. Box 2040, 3000 CA - University Medical Center Rotterdam, Rotterdam, the Netherlands
3. Department of Hematology, Erasmus MC - University Medical Center Rotterdam, Rotterdam, the Netherlands
4. Department of Neurosurgery, Erasmus MC, 's Gravendijkwal 230, Kamer H-703, 3015, CE - University Medical Center Rotterdam, Rotterdam, the Netherlands

5. Department of Pathophysiology and Transplants, University of Milan, Italy and Fondazione IRCCS Ca' Granda – Ospedale Maggiore Policlinico, Department of Anesthesia and Critical Care, Neuroscience Intensive Care Unit, Milan, Italy
6. Department of Neurosurgery, Antwerp University Hospital and University of Antwerp, Edegem, Belgium
7. Division of Anaesthesia, University of Cambridge, Addenbrooke's Hospital, Cambridge, United Kingdom
8. Department of Traumatology, Orthopedic Surgery and Sportsmedicine, Cologne-Merheim Medical Center (CMMC) and the Institute for Research in Operative Medicine (IFOM), University of Witten/Herdecke, Cologne, Germany
9. NHS Blood and Transplant/Oxford University Hospitals NHS Trust, John Radcliffe Hospital, Oxford, United Kingdom
10. School of Medicine and Surgery, University of Milan-Bicocca, Milan, Italy; Neurointensive care, San Gerardo Hospital, ASST-Monza, Monza, Italy
11. Department of Medical Statistics and Bioinformatics, Leiden University Medical Center, Leiden, the Netherlands

Corresponding author

Jilske Huijben, MD

Full mailing address: P.O. Box 2040, 3000 CA Rotterdam, The Netherlands, internal postal address Na-2223

Contact information: Email: j.a.huijben@erasmusmc.nl, Telephone: 0031 10 703 84 53,
Fax: 0031 107038475

Dr Simon Stanworth

Full mailing address: Headley Way, Headington, Oxford OX3 9DU John Radcliffe Hospital
United Kingdom

Contact information (telephone, fax, and e-mail address): Email:
simon.stanworth@nhsbt.nhs.uk, Telephone: +44 (0)1865, Fax: +44 (0)1865

Dr Giuseppe Citerio

Full mailing address: Dipartimento di Medicina Perioperatoria e Terapie Intensive, Via
Pergolesi 33

P.IVA e C.F. : 09314290967 H San Gerardo - Monza

Contact information (telephone, fax, and e-mail address): Email:
giuseppe.citerio@unimib.it, Telephone: +390392334316, Fax +390392334340

Dr Suzanne Polinder

Full mailing address: P.O. Box 2040, 3000 CA Rotterdam, The Netherlands

Contact information (telephone, fax, and e-mail address): Email:
s.polinder@erasmusmc.nl, Telephone +31 10 704 42 69 or +31 6 26 46 73 38

Prof Ewout W. Steyerberg

Full mailing address: Department of Medical Statistics LUMC PO Box 9600 2300 RC Leiden
The Netherlands

Contact information (telephone, fax, and e-mail address): Email: e.w.steyerberg@lumc.nl

or

E.Steyerberg@ErasmusMC.nl, Telephone: 31 71 5269700

Dr Hester F. Lingsma

Full mailing address: P.O. Box 2040, 3000 CA Rotterdam, The Netherlands

Variation in blood transfusion and coagulation management in Traumatic Brain Injury at the Intensive Care Unit: A survey in 66 neurotrauma centers participating in the Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI) study. (DOI: 10.1089/neu.2017.5194)

This paper has been peer-reviewed and accepted for publication, but has yet to undergo copyediting and proof of correction. The final published version may differ from this proof.

Abstract

Our aim was to describe current approaches and to quantify variability between European intensive care units (ICU)s in patients with TBI. Therefore, we conducted a provider profiling survey as part of the ‘Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury’ (CENTER-TBI) study. The ICU Questionnaire was sent to 68 centers from 20 countries across Europe and Israel. For this study, we used ICU questions focused on 1) hemoglobin target level (Hb-TL), 2) coagulation management, and 3) deep venous thromboembolism (DVT) prophylaxis. Seventy-eight participants, mostly intensivists and neurosurgeons of 66 centers completed the ICU questionnaire. For ICU-patients, half of the centers (N=34; 52%) had a defined Hb-TL in their protocol. For patients with TBI, 26 centers (41%) indicated a Hb-TL between 70 and 90 g/l and 38 centers (59%) above 90 g/l. To treat trauma related hemostatic abnormalities the use of fresh frozen plasma (N=48; 73%) or platelets (N=34; 52%) was most often reported, followed by the supplementation of vitamin K (N=26; 39%). Most centers reported using DVT prophylaxis with anticoagulants frequently or always (N=62; 94%). In the absence of hemorrhagic brain lesions, 14 centers (21%) delayed DVT prophylaxis until 72 hours after trauma. If hemorrhagic brain lesions were present, the number of centers delaying DVT prophylaxis for 72 hours increased to 29 (46%). Overall, a lack of consensus exists between European ICUs on blood transfusion and coagulation management. The results provide a baseline for the CENTER-TBI study and the large between-center variation indicates multiple opportunities for comparative effectiveness research.

Keywords: intensive care unit; traumatic brain injury; coagulopathy; transfusion; Europe

Introduction

The management of hemorrhage and disordered coagulation is a common and critically important challenge in trauma patients. This is particularly the case for patients with severe traumatic brain injury (TBI) where physicians have to balance the risks of progressive hemorrhage in the brain against secondary thrombotic complications including deep venous thrombosis (DVT). Many controversies continue to exist regarding the appropriate management for optimizing blood and coagulation status.

Transfusion thresholds for anaemia are a particularly controversial area in TBI. According to the guidelines^{1, 2}, transfusion in general critically ill patients is recommended at a restrictive hemoglobin target level (Hb-TL) of 70 g/l rather than a liberal Hb-TL of 90 g/l or 100g/l. Whether such target levels also apply to patients with TBI is unclear.^{3, 4} Inappropriate use of blood products exposes patients to a number of systemic risks and may even lead to progressive hemorrhagic injury following TBI.³ However, cerebral oxygenation may be improved with higher hemoglobin concentrations^{5, 6} whereas restrictive transfusion thresholds may predispose to brain tissue hypoxia and may increase the risk of early mortality.⁷ On the other hand, a recent large retrospective cohort study indicated that a restrictive blood transfusion policy was not associated with increased mortality and can be cost-effective in patients with TBI.⁸ An additional challenge for the management of both blood - and coagulation status is the presence of coagulopathy.⁹ Both pro- and anticoagulatory abnormalities can be observed after TBI in around one out of three patients.¹⁰⁻¹² Coagulopathy at admission is associated with increased mortality and poor neurological outcome.¹²⁻¹⁴ Coagulopathy may result from defective clot initiation, poor clot formation or hyper fibrinolysis. Acidosis, hypothermia, coagulation factor consumption or dilution, and the more recently described acute coagulopathy of trauma-shock which results from widespread endothelial activation after hypoperfusion may contribute to coagulopathy.¹⁵ Finally, patients with TBI are at increased risk of venous thromboembolism (VTE) (around 20%)¹⁶ compared with general ICU patients (around 6-8%).¹⁷ Here, the balance between the prevention of VTE and the risk of (progressive) hemorrhage of the brain depends largely on the timing of thromboprophylaxis with anticoagulants. However, current Brain Trauma Foundation guidelines do not make clear recommendations on coagulation management.¹⁸

Material and Methods

Participating centers

This study is part of the prospective, longitudinal ‘Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury’ (CENTER-TBI) study in 68 centers from 20 countries across Europe and Israel. The CENTER-TBI investigators and participants are listed in Supplemental Data 1. In 2014, before the start of inclusion of patients, the principle investigators of each center were asked to complete a set of questionnaires on structure and process of care: ‘the Provider Profiling Questionnaires’.¹⁹
²⁰ The questionnaires were about TBI management irrespective of systemic injuries. One of these questionnaires concerned ICU management.

Provider Profiling Questionnaire

The provider profiling questionnaire was developed in a systematic manner. The literature (including guidelines and available surveys) was reviewed and experts of various disciplines (neurosurgeons, (neuro)intensivists, neurologists, emergency department physicians, rehabilitation physicians, medical ethicists, health care economists and epidemiologists) were consulted throughout the different phases in the development process. Preliminary questionnaires were pilot-tested in 16 of the participating centers for unexpected or missing values and ambiguity, and received feedback was incorporated. For more information about the development, administration and content of the total set of provider profiling questionnaires, see Clossen et al., 2016.¹⁹ In this study, we focus on 10 questions (with additional sub questions) on hemoglobin target levels, trauma related coagulation management, and use and timing of thromboprophylaxis (Supplemental Data 2).

Hemoglobin target level and coagulation management

Participants were explicitly asked for their general policy rather than for individual treatment preferences. General policy was defined as ‘the way the large majority of patients (>75%) with a certain indication would be treated’. The ICU questionnaire

that number of centers (N=29; 46%) indicated to wait 72 hours after trauma in the presence of hemorrhagic brain lesions. Low molecular weight heparin was most commonly indicated as the prophylactic drug of choice (N=54; 82%), followed by subcutaneous unfractionated heparin (N=7; 11%) and intravenous heparin (N=1; 2%). (Table 2)

[Insert HuijbenTable2]

Most centers indicated that they would always test a coagulation panel prior to the insertion of a parenchymal sensor (N=45; 69%) or a ventricular catheter (N=46; 71%). The reported minimum platelet count for the insertion of a ventricular catheter was variable: $>100 \times 10^9/L$ (N=30; 46%), $>80 \times 10^9/L$ (N=9; 14%) or $>50 \times 10^9/L$ (N=9; 14%). In most of the remaining centers the minimum platelet count depended on the surgeon (N=13; 20%). Also, the reported minimum International Normalized Ratio (INR) considered safe for placement of a ventricular catheter was variable: <1.4 (N=21; 33%), <1.3 (N=17; 26%) or <1.2 (N=8; 12%). Again, in most of the remaining centers the minimum INR was indicated to depend on surgeon's individual preferences (N=15; 23%). There were no centers that answered 'never' on all questions. (Table 3)

[Insert HuijbenTable3]

Twenty-nine centers indicated identical policies for coagulation management (always using DVT prophylaxis, and always obtaining a coagulation panel prior to insertion of a parenchymal or ventricular catheter). The majority of these centers are located in South and East Europe and Israel (N=13, 56%) versus (N=16, 37%) in North and West Europe and the majority are located in high income countries (N=26, 47%), versus (N=3, 27%) in lower income countries.

Discussion

This study shows large between-center variation in blood transfusion and coagulation-directed policies in critically ill patients with TBI. More centers indicated a restrictive Hb-TL (between 70 g/l and 80 g/L) in general ICU patients compared to patients with TBI. Reported coagulation management was variable regarding timing of deep venous

transfusion strategies to correct coagulopathy in terms of the ratio of packed blood cells, fresh frozen plasma (or similar products) and platelets are still being debated.⁴⁰ This debate pertains both to optimal strategies with regard to reversal of trauma related coagulopathy and management of coagulopathy induced by conventional agents (such as vitamin K antagonists) and newer ones such as direct thrombin inhibitors.^{9, 30, 41} Still, others warn for the use of transfusion considering the possibility of complications of transfusion and unknown effects on (functional) outcome.⁴² Also for coagulation (enhancing) products larger studies are needed to prove a positive balance between the beneficial effects in terms of patient outcome and adverse effects on (thromboembolic) complications.^{27, 28, 42-45} New evidence is clearly needed on these topics, since control of blood and coagulation status could have a large impact on patient outcome, especially in patients with TBI.

Conclusions

In conclusion, we showed substantial variation in blood and coagulation management of patients with TBI at the ICUs in 66 centers in Europe and Israel participating in the CENTER-TBI study. This variation may be largely attributable to the lack of guidelines and high quality evidence on these topics. The large practice variation provides an opportunity to study the effectiveness of different policies in comparative effectiveness research.

10. Laroche, M., Kutcher, M.E., Huang, M.C., Cohen, M.J. and Manley, G.T. (2012). Coagulopathy after traumatic brain injury. *Neurosurgery* 70, 1334-1345.
11. Maegele, M. (2013). Coagulopathy after traumatic brain injury: incidence, pathogenesis, and treatment options. *Transfusion* 53 Suppl 1, 28S-37S.
12. Harhangi, B.S., Kompanje, E.J., Leebeek, F.W. and Maas, A.I. (2008). Coagulation disorders after traumatic brain injury. *Acta Neurochir (Wien)* 150, 165-175; discussion 175.
13. Wafaisade, A., Lefering, R., Tjardes, T., Wutzler, S., Simanski, C., Paffrath, T., Fischer, P., Bouillon, B., Maegele, M. and Trauma Registry of DGU. (2010). Acute coagulopathy in isolated blunt traumatic brain injury. *Neurocrit Care* 12, 211-219.
14. Greuters, S., van den Berg, A., Franschman, G., Viersen, V.A., Beishuizen, A., Peerdeman, S.M., Boer, C. and ALARM-BLEEDING investigators. (2011). Acute and delayed mild coagulopathy are related to outcome in patients with isolated traumatic brain injury. *Crit Care* 15, R2.
15. Brohi, K., Cohen, M.J., Ganter, M.T., Schultz, M.J., Levi, M., Mackersie, R.C. and Pittet, J.F. (2008). Acute coagulopathy of trauma: hypoperfusion induces systemic anticoagulation and hyperfibrinolysis. *J Trauma* 64, 1211-1217; discussion 1217.
16. Skrifvars, M.B., Bailey, M., Presneill, J., French, C., Nichol, A., Little, L., Duranteau, J., Huet, O., Haddad, S., Arabi, Y., McArthur, C., Cooper, D.J., Bellomo, R., EPO-TBI investigators and the ANZICS Clinical Trials Group (2016). Venous thromboembolic events in critically ill traumatic brain injury patients. *Intensive Care Med.* 43, 419-428
17. PROTECT Investigators for the Canadian Critical Care Trials Group and the Australian and New Zealand Intensive Care Society Clinical Trials Group., Cook, D., Meade, M., Guyatt, G., Walter, S., Heels-Ansdell, D., Warkentin, T.E., Zytaruk, N., Crowther, M., Geerts, W., Cooper, D.J., Vallance, S., Qushmaq, I., Rocha, M., Berwanger, O. and Vlahakis, N.E. (2011). Dalteparin versus unfractionated heparin in critically ill patients. *N Engl J Med* 364, 1305-1314.
18. Carney, N., Totten, A.M., O'Reilly, C., Ullman, J.S., Hawryluk, G.W., Bell, M.J., Bratton, S.L., Chesnut, R., Harris, O.A., Kissoon, N., Rubiano, A.M., Shutter, L., Tasker, R.C., Vavilala, M.S., Wilberger, J., Wright, D.W. and Ghajar, J. (2016). Guidelines for the Management of Severe Traumatic Brain Injury, Fourth Edition. *Neurosurgery*.
19. Clossen, M.C., Polinder, S., Lingsma, H.F., Maas, A.I., Menon, D., Steyerberg, E.W.,

21

CENTER-TBI Investigators and Participants. (2016). Variation in Structure and Process of Care in Traumatic Brain Injury: Provider Profiles of European Neurotrauma Centers Participating in the CENTER-TBI Study. *PLoS One* 11, e0161367.

20. Maas, A.I., Menon, D.K., Steyerberg, E.W., Citerio, G., Lecky, F., Manley, G.T., Hill, S., Legrand, V., Sorgner, A., Participants, CENTER-TBI Investigators and Participants. (2015). Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI): a prospective longitudinal observational study. *Neurosurgery* 76, 67-80.

21. European Commission. Remuneration of researchers in the public and private sectors Brussels: European Communities; 2007.

22. IBM Corp. Released 2012. IBM SPSS Statistics for Windows [OR Macintosh], Version 21.0. Armonk, NY: IBM Corp.

23. Nyquist, P., Jichici, D., Bautista, C., Burns, J., Chhangani, S., DeFilippis, M., Goldenberg, F.D., Kim, K., Liu-DeRyke, X., Mack, W. and Meyer, K. (2017). Prophylaxis of Venous Thrombosis in Neurocritical Care Patients: An Executive Summary of Evidence-Based Guidelines: A Statement for Healthcare Professionals From the Neurocritical Care Society and Society of Critical Care Medicine. *Crit Care Med* 45, 476-479.

24. Carson, J.L., Stanworth, S.J., Roubinian, N., Fergusson, D.A., Triulzi, D., Doree, C. and Hebert, P.C. (2016). Transfusion thresholds and other strategies for guiding allogeneic red blood cell transfusion. *Cochrane Database Syst Rev* 10, CD002042.

25. Robertson, C.S., Hannay, H.J., Yamal, J.M., Gopinath, S., Goodman, J.C., Tilley, B.C., Epo Severe TBI Trial Investigators, Baldwin, A., Rivera Lara, L., Saucedo-Crespo, H., Ahmed, O., Sadasivan, S., Ponce, L., Cruz-Navarro, J., Shahin, H., Aisiku, I.P., Doshi, P., Valadka, A., Neipert, L., Waguspak, J.M., Rubin, M.L., Benoit, J.S. and Swank, P. (2014). Effect of erythropoietin and transfusion threshold on neurological recovery after traumatic brain injury: a randomized clinical trial. *JAMA* 312, 36-47.

26. McIntyre, L.A., Fergusson, D.A., Hutchison, J.S., Pagliarello, G., Marshall, J.C., Yetisir, E., Hare, G.M. and Hebert, P.C. (2006). Effect of a liberal versus restrictive transfusion strategy on mortality in patients with moderate to severe head injury. *Neurocrit Care* 5, 4-9.

27. Yuan, Q., Wu, X., Du, Z.Y., Sun, Y.R., Yu, J., Li, Z.Q., Wu, X.H., Mao, Y., Zhou, L.F. and Hu, J. (2015). Low-dose recombinant factor VIIa for reversing coagulopathy in patients with isolated traumatic brain injury. *J Crit Care* 30, 116-120.

28. Etemadrezaie, H., Baharvahdat, H., Shariati, Z., Lari, S.M., Shakeri, M.T. and Ganjeifar, B. (2007). The effect of fresh frozen plasma in severe closed head injury. *Clin Neurol Neurosurg* 109, 166-171.
29. Batchelor, J.S. and Grayson, A. (2013). A meta-analysis to determine the effect of preinjury antiplatelet agents on mortality in patients with blunt head trauma. *Br J Neurosurg* 27, 12-18.
30. Pakraftar, S., Atencio, D., English, J., Corcos, A., Altschuler, E.M. and Stahlfeld, K. (2014). Dabigatran etixilate and traumatic brain injury: Evolving anticoagulants require evolving care plans. *World J Clin Cases* 2, 362-366.
31. Abdel-Aziz, H., Dunham, C.M., Malik, R.J. and Hileman, B.M. (2015). Timing for deep vein thrombosis chemoprophylaxis in traumatic brain injury: an evidence-based review. *Crit Care* 19, 96.
32. Phelan, H.A., Eastman, A.L., Madden, C.J., Aldy, K., Berne, J.D., Norwood, S.H., Scott, W.W., Bernstein, I.H., Pruitt, J., Butler, G., Rogers, L. and Minei, J.P. (2012). TBI risk stratification at presentation: a prospective study of the incidence and timing of radiographic worsening in the Parkland Protocol. *J Trauma Acute Care Surg* 73, S122-127.
33. Phelan, H.A., Wolf, S.E., Norwood, S.H., Aldy, K., Brakenridge, S.C., Eastman, A.L., Madden, C.J., Nakonezny, P.A., Yang, L., Chason, D.P., Arbique, G.M., Berne, J. and Minei, J.P. (2012). A randomized, double-blinded, placebo-controlled pilot trial of anticoagulation in low-risk traumatic brain injury: The Delayed Versus Early Enoxaparin Prophylaxis I (DEEP I) study. *J Trauma Acute Care Surg* 73, 1434-1441.
34. Schafer, N., Driessen, A., Frohlich, M., Sturmer, E.K., Maegele, M. and TACTIC partners (2015). Diversity in clinical management and protocols for the treatment of major bleeding trauma patients across European level I Trauma Centres. *Scand J Trauma Resusc Emerg Med* 23, 74.
35. Albrecht, V., Schafer, N., Sturmer, E.K., Driessen, A., Betsche, L., Schenk, M. and Maegele, M. (2015). Practice management of acute trauma haemorrhage and haemostatic disorders across German trauma centres. *Eur J Trauma Emerg Surg*.
36. Driessen, A., Schafer, N., Albrecht, V., Schenk, M., Frohlich, M., Sturmer, E.K., Maegele, M. and TACTIC partners. (2015). Infrastructure and clinical practice for the detection and management of trauma-associated haemorrhage and coagulopathy. *Eur J Trauma Emerg*

Surg 41, 413-420.

37. Hamada, S.R., Gauss, T., Pann, J., Dunser, M., Leone, M. and Duranteau, J. (2015). European trauma guideline compliance assessment: the ETRAUSS study. *Crit Care* 19, 423.

38. Sena, M.J., Rivers, R.M., Muizelaar, J.P., Battistella, F.D. and Utter, G.H. (2009). Transfusion practices for acute traumatic brain injury: a survey of physicians at US trauma centers. *Intensive Care Med* 35, 480-488.

39. Badenes, R., Oddo, M., Suarez, J.I., Antonelli, M., Lipman, J., Citerio, G. and Taccone, F.S. (2017). Hemoglobin concentrations and RBC transfusion thresholds in patients with acute brain injury: an international survey. *Crit Care* 21, 159.

40. Hallet, J., Lauzier, F., Mailloux, O., Trottier, V., Archambault, P., Zarychanski, R. and Turgeon, A.F. (2013). The use of higher platelet: RBC transfusion ratio in the acute phase of trauma resuscitation: a systematic review. *Crit Care Med* 41, 2800-2811.

41. Gaist, D., Garcia Rodriguez, L.A., Hellfritsch, M., Poulsen, F.R., Halle, B., Hallas, J. and Pottegard, A. (2017). Association of Antithrombotic Drug Use With Subdural Hematoma Risk. *JAMA* 317, 836-846.

42. Reddy, G.D., Gopinath, S. and Robertson, C.S. (2015). Transfusion in Traumatic Brain Injury. *Curr Treat Options Neurol* 17, 46.

43. Joseph, B., Hadjizacharia, P., Aziz, H., Kulvatunyou, N., Tang, A., Pandit, V., Wynne, J., O'Keeffe, T., Friese, R.S. and Rhee, P. (2013). Prothrombin complex concentrate: an effective therapy in reversing the coagulopathy of traumatic brain injury. *J Trauma Acute Care Surg* 74, 248-253.

44. Schochl, H., Schlimp, C.J. and Maegele, M. (2014). Tranexamic acid, fibrinogen concentrate, and prothrombin complex concentrate: data to support prehospital use? *Shock* 41 Suppl 1, 44-46.

45. Schochl, H., Voelckel, W., Maegele, M., Kirchmair, L. and Schlimp, C.J. (2014). Endogenous thrombin potential following hemostatic therapy with 4-factor prothrombin complex concentrate: a 7-day observational study of trauma patients. *Crit Care* 18, R147.

Table 1. Red blood cell transfusion policy ^a

Items	Number	N	(%)
questionnaire	completed		
Protocol at the ICU			
Protocol	65		
- Presence of a protocol with a Hb-TL		34	(52%)
- Absence of a protocol with a Hb-TL		31	(48%)
Transfusion at Hb-TL in protocol (open question)			
	29	1	(3%)
- 110 g/L		8	(28%)
- 100g/L		4	(14%)
- 90 g/L		9	(31%)
- 80 g/L		5	(18%)
- 70 g/L		1	(3%)
- 80-100 g/L		1	(3%)
- 70-80 g/L			
In non-neurological critically ill patients			
Transfusion at Hb-TL	63		
- > 100 g/L		1	(2%)
- Between 90 g/l and 100g/L		6	(9%)
- Between 80 g/l and 90 g/L		21	(33%)
- Between 70 g/l and 80 g/L		35	(56%)
In patients with TBI ^b			
Transfusion at Hb-TL	64		
- > 100 g/L		18	(28%)
- Between 90 g/l and 100g/L		20	(31%)
- Between 80 g/l and 90 g/L		16	(25%)
- Between 70 g/l and 80 g/L		10	(16%)

Table 2. Coagulation policies, deep venous thrombosis ^a

Items	Number	N	(%)
questionnaire completed			
DVT prophylaxis			
Frequency of DVT prophylaxis	66		
- Never (0-10%)		1	(2%)
- Rarely (10-30%)		0	(0%)
- Sometimes (30-70%)		3	(4%)
- Frequently (70-90%)		18	(27%)
- Always (90-100%)		44	(67%)
Start in the absence of hemorrhagic lesions	65	26	(40%)
- < 24 hours		24	(37%)
- 24-72 hours		14	(21%)
- > 72 hours		1	(2%)
- Never	63		
Start in the presence of hemorrhagic lesions		5	(8%)
- < 24 hours		25	(40%)
- 24-72 hours		29	(46%)
- > 72 hours	64	4	(6%)
- Never		10	(16%)
Start after intracranial surgery		31	(48%)
- < 24 hours		21	(33%)
- 24-72 hours		2	(3%)
- > 72 hours			
- Never			
Pharmacological DVT prophylaxis	66		
- Subcutaneous unfractionated heparin		7	(11%)
		1	(2%)

Table 3. Coagulation policies, ICP monitoring ^a

Items	Number	N	(%)
questionnaire	completed		
Checks prior to insertion of parenchymal sensor for ICP monitoring			
Coagulation panel	65		
- Never (0-10%)		4	(6%)
- Rarely (10-30%)		2	(3%)
- Sometimes (30-70%)		5	(8%)
- Frequently (70-90%)		5	(8%)
- Always (90-100%)		45	(69%)
- Not available ^b		4	(6%)
Checks prior to insertion ventricular catheter for ICP monitoring			
Coagulation panel	65		
- Never (0-10%)		3	(4%)
- Rarely (10-30%)		2	(3%)
- Sometimes (30-70%)		5	(8%)
- Frequently (70-90%)		4	(6%)
- Always (90-100%)		46	(71%)
- Not available ^b		5	(8%)
Minimum platelet count	65		
- >150 x10 ⁹ /L		1	(2%)
- >100 x10 ⁹ /L		30	(46%)
- > 80 x10 ⁹ /L		9	(14%)
- > 50 x10 ⁹ /L		9	(14%)
- Depending on the surgeon		13	(20%)
- No minimum		0	(0%)
- Other		3	(4%)
Minimum INR	65		

-	<1.4	21	(33%)
-	<1.3	17	(26%)
-	<1.2	8	(12%)
-	Depending on the surgeon	15	(23%)
-	No minimum	0	(0%)
-	Other	4	(6%)

Frequencies and percentage of centers with corresponding answers

DVT: deep venous thrombosis, ICP: intracranial pressure, INR: International Normalized Ratio, L: Liter

a) General policy: the way the large majority of patients >75% with a certain indication would be treated at the intensive care b) Centers that did not have this technique

Supplemental data 1. Center-TBI investigators and participants

Adams Hadie ¹, Alessandro Masala ², Allanson Judith ³, Amrein Krisztina ⁴,
Andaluz Norberto ⁵, Andelic Nada ⁶, Andrea Nanni ², Andreassen Lasse ⁷, Anke Audny ⁸,
Antoni Anna ⁹, Ardon Hilko ¹⁰, Audibert Gérard ¹¹, Auslands Kaspars ¹², Azouvi Philippe ¹³,
Baciu Camelia ¹⁴, Bacon Andrew ¹⁵, Badenes Rafael ¹⁶, Baglin Trevor ¹⁷, Bartels Ronald ¹⁸,
Barzó Pál ¹⁹, Bauerfeind Ursula ²⁰, Beer Ronny ²¹, Belda Francisco Javier ¹⁶,
Bellander Bo-Michael ²², Belli Antonio ²³, Bellier Rémy ²⁴, Benali Habib ²⁵, Benard Thierry ²⁴,
Berardino Maurizio ²⁶, Beretta Luigi ²⁷, Beynon Christopher ²⁸, Bilotta Federico ¹⁶,
Binder Harald ⁹, Biqiri Erta ¹⁴, Blaabjerg Morten ²⁹, Borgen Lund Stine ³⁰, Bouzat Pierre ³¹,
Bragge Peter ³², Brazinova Alexandra ³³, Brehar Felix ³⁴, Brorsson Camilla ³⁵, Buki Andras ³⁶,
Bullinger Monika ³⁷, Bučková Veronika ³³, Calappi Emiliana ³⁸, Cameron Peter ³⁹,
Carbayo Lozano Guillermo ⁴⁰, Carise Elsa ²⁴, Carpenter K. ⁴¹, Castaño-León Ana M. ⁴²,
Causin Francesco ⁴³, Chevallard Giorgio ¹⁴, Chierigato Arturo ¹⁴, Citerio Giuseppe ^{44, 45},
Cnossen Maryse ⁴⁶, Coburn Mark Coburn ⁴⁷, Coles Jonathan ⁴⁸, Cooper Jamie D. ⁴⁹,
Correia Marta ⁵⁰, Covic Amra ⁵¹, Curry Nicola ⁵², Czeiter Endre ⁵³, Czosnyka Marek ⁵⁴,
Dahyot-Fizelier Claire ²⁴, Damas François ⁵⁵, Damas Pierre ⁵⁶, Dawes Helen ⁵⁷,
De Keyser Véronique ⁵⁸, Della Corte Francesco ⁵⁹, Depreitere Bart ⁶⁰, Ding Shenghao ⁶¹,
Dippel Diederik ⁶², Dizdarevic Kemal ⁶³, Dulière Guy-Loup ⁵⁵, Dzeko Adelaida ⁶⁴,
Eapen George ¹⁵, Engemann Heiko ⁵¹, Ercole Ari ⁶⁵, Esser Patrick ⁵⁷, Ezer Erzsébet ⁶⁶,
Fabricius Martin ⁶⁷, Feigin Valery L. ⁶⁸, Feng Junfeng ⁶¹, Foks Kelly ⁶², Fossi Francesca ¹⁴,
Francony Gilles ³¹, Frantzén Janek ⁶⁹, Freo Ulderico ⁷⁰, Frisvold Shirin ⁷¹, Furmanov Alex ⁷²,
Gagliardo Pablo ⁷³, Galanaud Damien ²⁵, Gao Guoyi ⁷⁴, Geleijns Karin ⁴¹,
Ghuysen Alexandre ⁷⁵, Giraud Benoit ²⁴, Glocker Ben ⁷⁶, Gomez Pedro A. ⁴²,
Grossi Francesca ⁵⁹, Gruen Russell L. ⁷⁷, Gupta Deepak ⁷⁸, Haagsma Juanita A. ⁴⁶,
Hadzic Ermin ⁶⁴, Haitsma Iain ⁷⁹, Hartings Jed A. ⁸⁰, Helbok Raimund ²¹, Helseth Eirik ⁸¹,
Hertle Daniel ²⁸, Hill Sean ⁸², Hoedemaekers Astrid ⁸³, Hofer Stefan ⁵¹,
Hutchinson Peter J. ¹, Håberg Asta Kristine ⁸⁴, Jacobs Bram ⁸⁵, Janciak Ivan ⁸⁶,
Janssens Koen ⁵⁸, Jiang Ji-yao ⁷⁴, Jones Kelly ⁸⁷, Kalala Jean-Pierre ⁸⁸,
Kamnitsas Konstantinos ⁷⁶, Karan Mladen ⁸⁹, Karau Jana ²⁰, Katila Ari ⁶⁹, Kaukonen Maija ⁹⁰,
Keeling David ⁵², Kerforne Thomas ²⁴, Ketharanathan Naomi ⁴¹, Kettunen Johannes ⁹¹,

Journal of Neurotrauma
Variation in blood transfusion and coagulation management in Traumatic Brain Injury at the Intensive Care Unit: A survey in 66 neurotrauma centers participating in the Collaborative European Neuro Trauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI) study. (DOI: 10.1089/neu.2017.5194)

This paper has been peer-reviewed and accepted for publication, but has yet to undergo copyediting and proof correction. The final published version may differ from this proof.

Kivisaari Riku⁹⁰, Koliás Angelos G.¹, Kolumbán Bálint⁹², Kompanje Erwin⁹³,
Kondziella Daniel⁶⁷, Koskinen Lars-Owe³⁵, Kovács Noémi⁹², Kálovits Ferenc⁹⁴,
Lagares Alfonso⁴², Lanyon Linda⁸², Laureys Steven⁹⁵, Lauritzen Martin⁶⁷, Lecky Fiona⁹⁶,
Ledig Christian⁷⁶, Lefering Rolf⁹⁷, Legrand Valerie⁹⁸, Lei Jin⁶¹, Levi Leon⁹⁹,
Lightfoot Roger¹⁰⁰, Lingsma Hester⁴⁶, Loeckx Dirk¹⁰¹, Lozano Angels¹⁶,
Luddington Roger¹⁷, Luijten-Arts Chantal⁸³, Maas Andrew I.R.⁵⁸, MacDonald Stephen¹⁷,
MacFayden Charles⁶⁵, Maegele Marc¹⁰², Majdan Marek³³, Major Sebastian¹⁰³,
Manara Alex¹⁰⁴, Manhes Pauline³¹, Manley Geoffrey¹⁰⁵, Martin Didier¹⁰⁶,
Martino Costanza², Maruenda Armando¹⁶, Maréchal Hugues⁵⁵, Mastelova Dagmara⁸⁶,
Mattern Julia²⁸, McMahon Catherine¹⁰⁷, Melegh Béla¹⁰⁸, Menon David⁶⁵,
Menovsky Tomas⁵⁸, Morganti-Kossmann Cristina¹⁰⁹, Mulazzi Davide³⁸,
Mutschler Manuel¹⁰², Mühlhan Holger¹¹⁰, Negru Ancuta¹¹¹, Nelson David⁸²,
Neugebauer Eddy¹⁰², Newcombe Virginia⁶⁵, Noirhomme Quentin⁹⁵, Nyirádi József⁴,
Oddo Mauro¹¹², Oldenbeuving Annemarie¹¹³, Oresic Matej¹¹⁴, Ortolano Fabrizio³⁸,
Palotie Aarno^{91, 115, 116}, Parizel Paul M.¹¹⁷, Patruno Adriana¹¹⁸, Payen Jean-François³¹,
Perera Natascha¹¹⁹, Perlberg Vincent²⁵, Persona Paolo¹²⁰, Peul Wilco¹²¹,
Pichon Nicolas¹²², Piilgaard Henning⁶⁷, Piippo Anna⁹⁰, Pili Floury Sébastien¹²³,
Pirinen Matti⁹¹, Ples Horia¹¹¹, Polinder Suzanne⁴⁶, Pomposo Inigo⁴⁰, Psota Marek³³,
Pullens Pim¹¹⁷, Puybasset Louis¹²⁴, Ragauskas Arminas¹²⁵, Raj Rahul⁹⁰,
Rambadagalla Malinka¹²⁶, Rehorčíková Veronika³³, Rhodes Jonathan¹²⁷,
Richardson Sylvia¹²⁸, Ripatti Samuli⁹¹, Rocka Saulius¹²⁵, Rodier Nicolas¹²², Roe Cecilie¹²⁹,
Roise Olav¹³⁰, Roks Gerwin¹³¹, Romegoux Pauline³¹, Rosand Jonathan¹³²,
Rosenfeld Jeffrey¹⁰⁹, Rosenlund Christina¹³³, Rosenthal Guy⁷², Rossaint Rolf⁴⁷,
Rossi Sandra¹²⁰, Rostalski Tim¹¹⁰, Rueckert Daniel⁷⁶, Ruiz de Arcaute Felix¹⁰¹,
Rusnák Martin⁸⁶, Sacchi Marco¹⁴, Sahakian Barbara⁶⁵, Sahuquillo Juan¹³⁴,
Sakowitz Oliver^{135, 136}, Sala Francesca¹¹⁸, Sanchez-Pena Paola²⁵, Sanchez-Porrás Renan^{28,}
¹³⁵, Sandor Janos¹³⁷, Santos Edgar²⁸, Sasse Nadine⁵¹, Sasu Luminita⁵⁹, Savo Davide¹¹⁸,
Schipper Inger¹³⁸, Schlößer Barbara²⁰, Schmidt Silke¹¹⁰, Schneider Annette⁹⁷,
Schoechl Herbert¹³⁹, Schoonman Guus¹³¹, Schou Rico Frederik¹⁴⁰,
Schwendenwein Elisabeth⁹, Schöll Michael²⁸, Sir Özcan¹⁴¹, Skandsen Toril¹⁴²,
Smakman Lidwien¹⁴³, Smeets Dirk¹⁰¹, Smielewski Peter⁵⁴, Sorinola Abayomi¹⁴⁴,

Stamatakis Emmanuel ⁶⁵, Stanworth Simon ⁵², Stegemann Katrin ¹¹⁰, Steinbüchel Nicole ¹⁴⁵,
Stevens Robert ¹⁴⁶, Stewart William ¹⁴⁷, Steyerberg Ewout W. ⁴⁶, Stocchetti Nino ¹⁴⁸,
Sundström Nina ³⁵, Synnot Anneliese ^{149, 150}, Szabó József ⁹⁴, Söderberg Jeannette ⁸²,
Taccone Fabio Silvio ¹⁶, Tamás Viktória ¹⁴⁴, Tanskanen Päivi ⁹⁰, Tascu Alexandru ³⁴,
Taylor Mark Steven ³³, Te Ao Braden ⁶⁸, Tenovuo Olli ⁶⁹, Teodorani Guido ¹⁵¹,
Theadom Alice ⁶⁸, Thomas Matt ¹⁰⁴, Tibboel Dick ⁴¹, Tolia Christos ¹⁵²,
Tshibanda Jean-Flory Luaba ¹⁵³, Tudora Cristina Maria ¹¹¹, Vajkoczy Peter ¹⁵⁴,
Valeinis Egils ¹⁵⁵, Van Hecke Wim ¹⁰¹, Van Praag Dominique ⁵⁸, Van Roost Dirk ⁸⁸,
Van Vlierberghe Eline ¹⁰¹, Vande Vyvere Thijs ¹⁰¹, Vanhauzenhuyse Audrey ^{25, 95},
Vargiolu Alessia ¹¹⁸, Vega Emmanuel ¹⁵⁶, Verheyden Jan ¹⁰¹, Vespa Paul M. ¹⁵⁷, Vik Anne ¹⁵⁸,
Vilcinis Rimantas ¹⁵⁹, Vizzino Giacinta ¹⁴, Vleggeert-Lankamp Carmen ¹⁴³, Volovici Victor ⁷⁹,
Vulekovic Peter ⁸⁹, Vámos Zoltán ⁶⁶, Wade Derick ⁵⁷, Wang Kevin K.W. ¹⁶⁰, Wang Lei ⁶¹,
Wildschut Eno ⁴¹, Williams Guy ⁶⁵, Willumsen Lisette ⁶⁷, Wilson Adam ⁵, Wilson Lindsay ¹⁶¹,
Winkler Maren K.L. ¹⁰³, Ylén Peter ¹⁶², Younsi Alexander ²⁸, Zaaroor Menashe ⁹⁹,
Zhang Zhiqun ¹⁶³, Zheng Zelong ²⁸, Zumbo Fabrizio ², de Lange Stefanie ⁹⁷,
de Ruitter Godard C.W. ¹⁴³, den Boogert Hugo ¹⁸, van Dijck Jeroen ¹⁶⁴,
van Essen Thomas A. ¹²¹, van Heugten Caroline ⁵⁷, van der Jagt Mathieu ¹⁶⁵, van der
Naalt Joukje ⁸⁵

- ¹ Division of Neurosurgery, Department of Clinical Neurosciences, Addenbrooke's Hospital & University of Cambridge, Cambridge, UK
- ² Department of Anesthesia & Intensive Care, M. Bufalini Hospital, Cesena, Italy
- ³ Department of Clinical Neurosciences, Addenbrooke's Hospital & University of Cambridge, Cambridge, UK
- ⁴ János Szentágotthai Research Centre, University of Pécs, Pécs, Hungary
- ⁵ University of Cincinnati, Cincinnati, Ohio, United States
- ⁶ Division of Surgery and Clinical Neuroscience, Department of Physical Medicine and Rehabilitation, Oslo University Hospital and University of Oslo, Oslo, Norway
- ⁷ Department of Neurosurgery, University Hospital Northern Norway, Tromsø, Norway
- ⁸ Department of Physical Medicine and Rehabilitation, University hospital Northern Norway
- ⁹ Trauma Surgery, Medical University Vienna, Vienna, Austria
- ¹⁰ Department of Neurosurgery, Elisabeth-Tweesteden Ziekenhuis, Tilburg, the Netherlands
- ¹¹ Department of Anesthesiology & Intensive Care, University Hospital Nancy, Nancy, France
- ¹² Riga Eastern Clinical University Hospital, Riga, Latvia
- ¹³ Raymond Poincaré hospital, Assistance Publique – Hôpitaux de Paris, Paris, France
- ¹⁴ NeuroIntensive Care, Niguarda Hospital
- ¹⁵ Neurointensive Care, Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, UK
- ¹⁶ Department Anesthesiology and Surgical-Trauma Intensive Care, Hospital Clinic Universitari de Valencia, Spain
- ¹⁷ Cambridge University Hospitals, Cambridge, UK
- ¹⁸ Department of Neurosurgery, Radboud University Medical Center
- ¹⁹ Department of Neurosurgery, University of Szeged, Szeged, Hungary
- ²⁰ Institute for Transfusion Medicine (ITM), Witten/Herdecke University, Cologne, Germany
- ²¹ Department of Neurocritical care, Innsbruck Medical University, Innsbruck, Austria

- 41 Intensive Care and Department of Pediatric Surgery, Erasmus Medical Center, Sophia Children's Hospital, Rotterdam, The Netherlands
- 42 Department of Neurosurgery, Hospital Universitario 12 de Octubre, Madrid, Spain
- 43 Department of Neuroscience, Azienda Ospedaliera Università di Padova, Padova, Italy
- 44 NeuroIntensive Care, Azienda Ospedaliera San Gerardo di Monza, Monza, Italy
- 45 School of Medicine and Surgery, Università Milano Bicocca, Milano, Italy
- 46 Department of Public Health, Erasmus Medical Center-University Medical Center, Rotterdam, The Netherlands
- 47 Department of Anaesthesiology, University Hospital of Aachen, Aachen, Germany
- 48 Department of Anesthesia & Neurointensive Care, Cambridge University Hospital NHS Foundation Trust, Cambridge, UK
- 49 School of Public Health & PM, Monash University and The Alfred Hospital, Melbourne, Victoria, Australia
- 50 Radiology/MRI department, MRC Cognition and Brain Sciences Unit, Cambridge, UK
- 51 Institute of Medical Psychology and Medical Sociology, Universitätsmedizin Göttingen, Göttingen, Germany
- 52 Oxford University Hospitals NHS Trust, Oxford, UK
- 53 Department of Neurosurgery, University of Pecs and MTA-PTE Clinical Neuroscience MR Research Group and Janos Szentagothai Research Centre, University of Pecs, Hungarian Brain Research Program (Grant No. KTIA 13 NAP-A-II/8), Pecs, Hungary
- 54 Brain Physics Lab, Division of Neurosurgery, Dept of Clinical Neurosciences, University of Cambridge, Addenbrooke's Hospital, Cambridge, UK
- 55 Intensive Care Unit, CHR Citadelle , Liège, Belgium
- 56 Intensive Care Unit, CHU , Liège, Belgium
- 57 Movement Science Group, Faculty of Health and Life Sciences, Oxford Brookes University, Oxford, UK
- 58 Department of Neurosurgery, Antwerp University Hospital and University of Antwerp, Edegem, Belgium
- 59 Department of Anesthesia & Intensive Care, Maggiore Della Carità Hospital, Novara, Italy
- 60 Department of Neurosurgery, University Hospitals Leuven, Leuven, Belgium

Journal of Neurotrauma
Variation in blood transfusion and coagulation management in Traumatic Brain Injury at the Intensive Care Unit: A survey in 66 neurotrauma centers participating in the Collaborative European Neuro Trauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI) study. (DOI: 10.1089/neu.2017.5194)

This paper has been peer-reviewed and accepted for publication, but has yet to undergo copyediting and proof correction. The final published version may differ from this proof.

- ⁶¹ Department of Neurosurgery, Renji Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China
- ⁶² Department of Neurology, Erasmus MC, Rotterdam, the Netherlands
- ⁶³ Department of Neurosurgery, Medical Faculty and clinical center University of Sarajevo, Sarajevo, Bosnia Herzegovina
- ⁶⁴ Department of Neurosurgery, Regional Medical Center dr Safet Mujić, Mostar, Bosnia Herzegovina
- ⁶⁵ Division of Anaesthesia, University of Cambridge, Addenbrooke's Hospital, Cambridge, UK
- ⁶⁶ Department of Anaesthesiology and Intensive Therapy, University of Pécs, Pécs, Hungary
- ⁶⁷ Departments of Neurology, Clinical Neurophysiology and Neuroanesthesiology, Region Hovedstaden Rigshospitalet, Copenhagen, Denmark
- ⁶⁸ National Institute for Stroke and Applied Neurosciences, Faculty of Health and Environmental Studies, Auckland University of Technology, Auckland, New Zealand
- ⁶⁹ Rehabilitation and Brain Trauma, Turku University Central Hospital and University of Turku, Turku, Finland
- ⁷⁰ Department of Medicine, Azienda Ospedaliera Università di Padova, Padova, Italy
- ⁷¹ Department of Anesthesiology and Intensive care, University Hospital Northern Norway, Tromsø, Norway
- ⁷² Department of Neurosurgery, Hadassah-hebrew University Medical center, Jerusalem, Israel
- ⁷³ Fundación Instituto Valenciano de Neurorehabilitación (FIVAN), Valencia, Spain
- ⁷⁴ Department of Neurosurgery, Shanghai Renji hospital, Shanghai Jiaotong University/school of medicine, Shanghai, China
- ⁷⁵ Emergency Department, CHU , Liège, Belgium
- ⁷⁶ Department of Computing, Imperial College London, London, UK
- ⁷⁷ Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore; and Monash University, Australia
- ⁷⁸ Department of Neurosurgery, Neurosciences Centre & JPN Apex trauma centre, All India Institute of Medical Sciences, New Delhi-110029, India
- ⁷⁹ Department of Neurosurgery, Erasmus MC, Rotterdam, the Netherlands

- ⁸⁰ Department of Neurosurgery, University of Cincinnati, Cincinnati, Ohio, USA
- ⁸¹ Department of Neurosurgery, Oslo University Hospital, Oslo, Norway
- ⁸² Department of Physiology and Pharmacology, Section of Perioperative Medicine and Intensive Care, Karolinska Institutet, Stockholm, Sweden
- ⁸³ Department of Intensive Care Medicine, Radboud University Medical Center
- ⁸⁴ Department of Medical Imaging, St. Olavs Hospital and Department of Neuroscience, Norwegian University of Science and Technology, Trondheim, Norway
- ⁸⁵ Department of Neurology, University Medical Center Groningen, Groningen, Netherlands
- ⁸⁶ International Neurotrauma Research Organisation, Vienna, Austria
- ⁸⁷ National Institute for Stroke & Applied Neurosciences of the AUT University, Auckland, New Zealand
- ⁸⁸ Department of Neurosurgery, UZ Gent, Gent, Belgium
- ⁸⁹ Department of Neurosurgery, Clinical centre of Vojvodina, Novi Sad, Serbia
- ⁹⁰ Helsinki University Central Hospital
- ⁹¹ Institute for Molecular Medicine Finland, University of Helsinki, Helsinki, Finland
- ⁹² Hungarian Brain Research Program - Grant No. KTIA 13 NAP-A-II/8, University of Pécs, Pécs, Hungary
- ⁹³ Department of Intensive Care and Department of Ethics and Philosophy of Medicine, Erasmus Medical Center, Rotterdam, The Netherlands
- ⁹⁴ Department of Neurological & Spinal Surgery, Markusovszky University Teaching Hospital, Szombathely, Hungary
- ⁹⁵ Cyclotron Research Center, University of Liège, Liège, Belgium
- ⁹⁶ Emergency Medicine Research in Sheffield, Health Services Research Section, School of Health and Related Research (SCHARR), University of Sheffield, Sheffield, UK
- ⁹⁷ Institute of Research in Operative Medicine (IFOM), Witten/Herdecke University, Cologne, Germany
- ⁹⁸ VP Global Project Management CNS, ICON, Paris, France
- ⁹⁹ Department of Neurosurgery, Rambam Medical Center, Haifa, Israel
- ¹⁰⁰ Department of Anesthesiology & Intensive Care, University Hospitals Southampton NHS Trust, Southampton, UK

- ¹⁰¹ icoMetrix NV, Leuven, Belgium
- ¹⁰² Cologne-Merheim Medical Center (CMMC), Department of Traumatology, Orthopedic Surgery and Sportmedicine, Witten/Herdecke University, Cologne, Germany
- ¹⁰³ Centrum für Schlaganfallforschung, Charité – Universitätsmedizin Berlin, Berlin, Germany
- ¹⁰⁴ Intensive Care Unit, Southmead Hospital, Bristol, Bristol, UK
- ¹⁰⁵ Department of Neurological Surgery, University of California, San Francisco, California, USA
- ¹⁰⁶ Department of Neurosurgery, CHU , Liège, Belgium
- ¹⁰⁷ Department of Neurosurgery, The Walton centre NHS Foundation Trust, Liverpool, UK
- ¹⁰⁸ Department of Medical Genetics, University of Pécs, Pécs, Hungary
- ¹⁰⁹ National Trauma Research Institute, The Alfred Hospital, Monash University, Melbourne, Victoria, Australia
- ¹¹⁰ Department Health and Prevention, University Greifswald, Greifswald, Germany
- ¹¹¹ Department of Neurosurgery, Emergency County Hospital Timisoara , Timisoara, Romania
- ¹¹² Centre Hospitalier Universitaire Vaudois
- ¹¹³ Department of Intensive Care, Elisabeth-Tweesteden Ziekenhuis, Tilburg, the Netherlands
- ¹¹⁴ Department of Systems Medicine, Steno Diabetes Center, Gentofte, Denmark
- ¹¹⁵ Analytic and Translational Genetics Unit, Department of Medicine; Psychiatric & Neurodevelopmental Genetics Unit, Department of Psychiatry; Department of Neurology, Massachusetts General Hospital, Boston, MA, USA
- ¹¹⁶ Program in Medical and Population Genetics; The Stanley Center for Psychiatric Research, The Broad Institute of MIT and Harvard, Cambridge, MA, USA
- ¹¹⁷ Department of Radiology, Antwerp University Hospital and University of Antwerp, Edegem, Belgium
- ¹¹⁸ NeuroIntensive Care Unit, Department of Anesthesia & Intensive Care Azienda Ospedaliera San Gerardo di Monza, Monza, Italy
- ¹¹⁹ International Projects Management, ARTTIC, Munchen, Germany

- ¹²⁰ Department of Anesthesia & Intensive Care, Azienda Ospedaliera Università di Padova, Padova, Italy
- ¹²¹ Dept. of Neurosurgery, Leiden University Medical Center, Leiden, The Netherlands and Dept. of Neurosurgery, Medical Center Haaglanden, The Hague, The Netherlands
- ¹²² Intensive Care Unit, CHU Dupuytren, Limoges, France
- ¹²³ Intensive Care Unit, CHRU de Besançon, Besançon, France
- ¹²⁴ Department of Anesthesiology and Critical Care, Pitié -Salpêtrière Teaching Hospital, Assistance Publique, Hôpitaux de Paris and University Pierre et Marie Curie, Paris, France
- ¹²⁵ Department of Neurosurgery, Kaunas University of technology and Vilnius University, Vilnius, Lithuania
- ¹²⁶ Rezekne Hospital, Latvia
- ¹²⁷ Department of Anaesthesia, Critical Care & Pain Medicine NHS Lothian & University of Edinburg, Edinburgh, UK
- ¹²⁸ Director, MRC Biostatistics Unit, Cambridge Institute of Public Health, Cambridge, UK
- ¹²⁹ Department of Physical Medicine and Rehabilitation, Oslo University Hospital/University of Oslo, Oslo, Norway
- ¹³⁰ Division of Surgery and Clinical Neuroscience, Oslo University Hospital, Oslo, Norway
- ¹³¹ Department of Neurology, Elisabeth-TweeSteden Ziekenhuis, Tilburg, the Netherlands
- ¹³² Broad Institute, Cambridge MA Harvard Medical School, Boston MA, Massachusetts General Hospital, Boston MA, USA
- ¹³³ Department of Neurosurgery, Odense University Hospital, Odense, Denmark
- ¹³⁴ Department of Neurosurgery, Vall d'Hebron University Hospital, Barcelona, Spain
- ¹³⁵ Klinik für Neurochirurgie, Klinikum Ludwigsburg, Ludwigsburg, Germany
- ¹³⁶ University Hospital Heidelberg, Heidelberg, Germany
- ¹³⁷ Division of Biostatistics and Epidemiology, Department of Preventive Medicine, University of Debrecen, Debrecen, Hungary
- ¹³⁸ Department of Traumasurgery, Leiden University Medical Center, Leiden, The Netherlands
- ¹³⁹ Department of Anaesthesiology and Intensive Care, AUVA Trauma Hospital, Salzburg, Austria

Deep venous thrombosis (DVT) prophylaxis

The responses to the following questions should represent, as best as practicable, a general consensus on treatment at your centre, rather than individual management preferences.

	Never (0-10%)	Rarely (10-30%)	Sometimes (30-70%)	Frequently (70-90%)	Always (90-100%)
53. How often is DVT prophylaxis used?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
54. If you use DVT prophylaxis, when is DVT prophylaxis initiated?					
	<input type="checkbox"/> < 24 hrs	<input type="checkbox"/> 24-72 hrs	<input type="checkbox"/> < 72 hrs	<input type="checkbox"/> Never	
In the absence of hemorrhagic lesions	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
In the presence of hemorrhagic lesion	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
After intracranial surgery	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
55. In patients who receive DVT prophylaxis, what medication is given?					
<input type="checkbox"/> Subcutaneous unfractionated heparin					
<input type="checkbox"/> Low-molecular weight heparin					
<input type="checkbox"/> Other, please specify.....					

Journal of Neurotrauma
 Variation in blood transfusion and coagulation management in Traumatic Brain Injury at the Intensive Care Unit: A survey in 66 neurotrauma centers participating in the Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI) study. (DOI: 10.1089/neu.2017.5194)
 This paper has been peer-reviewed and accepted for publication, but has yet to undergo copyediting and proof correction. The final published version may differ from this proof.

56. Coagulopathy related to the trauma is treated with :

	Never (0-10%)	Rarely (10-30%)	Sometimes (30-70%)	Frequently (70-90%)	Always (90-100%)
Fresh Frozen plasma (FFP)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Platelets	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Fibrinogen	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Novo 7 (recombinant factor VII)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Vitamin K	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
PCC (Prothrombin Complex Concentrate)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Other, please specify...	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Journal of Neurotrauma
Variation in blood transfusion and coagulation management in Traumatic Brain Injury at the Intensive Care Unit: A survey in 66 neurotrauma centers participating in the Collaborative European Neuro Trauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI) study. (DOI: 10.1089/neu.2017.5194)
This paper has been peer-reviewed and accepted for publication, but has yet to undergo copyediting and proof correction. The final published version may differ from this proof.

General ICU treatments / protocols

Red blood cell policy

- 61. Does the Intensive Care Unit (ICU) protocol specify a target goal for hemoglobin concentration?
 - No
 - Yes, please specify

- 62. Do you have a transfusion target in patients with Traumatic Brain Injury (TBI) in the acute phase?
 - >100 g/l or 6 mmol/l
 - Between 90g/l or 5.5 mmol/l and 100 g/l or 6 mmol/l
 - between 80 g/l or 5 mmol/l and 90 g/l or 5.5 mmol/l
 - Between 70 g/l or 4.0 mmol/l and 80 g/l or 5 mmol/l

- 63. What is your transfusion target in patients with non-neurological critical illness?
 - >100 g/l or 6 mmol/l
 - between 90g/l or 5.5 mmol/l and 100 g/l or 6 mmol/l
 - between 80 g/l or 5 mmol/l and 90 g/l or 5.5 mmol/l
 - Between 70 g/l or 4.0 mmol/l and 80 g/l or 5 mmol/l