

SCIENTIFIC OPINION

ADOPTED: 21 April 2016

doi: 10.2903/j.efsa.2016.4479

Safety and efficacy of *Lactobacillus plantarum* DSM 29025 as a silage additive for all animal species

EFSA Panel on Additives and Products or Substances used in Animal Feed (FEEDAP)

Abstract

Lactobacillus plantarum is a technological additive intended to improve the ensiling process at a minimum proposed dose of 5.0×10^7 colony-forming units (CFU)/kg fresh material. The bacterial species *L. plantarum* is considered by EFSA to be suitable for the qualified presumption of safety approach to safety assessment. As the identity of the strain has been clearly established and as no antibiotic resistance of concern was detected, the use of the strain as a silage additive is considered safe for livestock species, for consumers of products from animals fed the treated silage and for the environment. In the absence of data, no conclusion can be drawn on the skin and eye irritancy of the additive. The additive should be considered to have the potential to be a respiratory sensitiser. Three studies with laboratory-scale silos were made using samples of forage of differing dry matter and water-soluble carbohydrate content. In each case, replicate silos containing treated forage were compared with identical silos containing the same but untreated forage. The results showed that the additive has the potential to improve the production of silage from easy, moderately difficult and difficult to ensile material by improving the preservation of nutrients. This was shown at the proposed application rate of 5×10^7 CFU/kg forage.

© 2016 European Food Safety Authority. *EFSA Journal* published by John Wiley and Sons Ltd on behalf of European Food Safety Authority.

Keywords: technological additive, silage additive, *Lactobacillus plantarum*, safety, QPS, efficacy

Requestor: European Commission

Question number: EFSA-Q-2015-00652

Correspondence: feedap@efsa.europa.eu

Panel members: Gabriele Aquilina, Giovanna Azimonti, Vasileios Bampidis, Maria de Lourdes Bastos, Georges Bories, Andrew Chesson, Pier Sandro Cocconcelli, Gerhard Flachowsky, Jürgen Gropp, Boris Kolar, Maryline Kouba, Secundino López Puente, Marta López-Alonso, Alberto Mantovani, Baltasar Mayo, Fernando Ramos, Guido Rychen, Maria Saarela, Roberto Edoardo Villa, Robert John Wallace and Pieter Wester

Acknowledgements: The Panel wishes to thank the members of the Working Group on Microorganisms for the preparatory work on this scientific opinion.

Suggested citation: EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2016. Scientific opinion on the safety and efficacy of *Lactobacillus plantarum* DSM 29025 as a silage additive for all animal species. EFSA Journal 2016;14(6):4479, 10 pp. doi: 10.2903/j.efsa.2016.4479

ISSN: 1831-4732

© 2016 European Food Safety Authority. *EFSA Journal* published by John Wiley and Sons Ltd on behalf of European Food Safety Authority.

This is an open access article under the terms of the [Creative Commons Attribution-NoDerivs License](https://creativecommons.org/licenses/by-nd/4.0/), which permits use and distribution in any medium, provided the original work is properly cited and no modifications or adaptations are made.



The EFSA Journal is a publication of the European Food Safety Authority, an agency of the European Union.



Summary

Following a request from the European Commission, the Panel on Additives and Products or Substances used in Animal Feed (FEEDAP Panel) was asked to deliver a scientific opinion on the safety for target animals, consumers, users and for the environment, and on the efficacy of a specific strain of *Lactobacillus plantarum* when used as a technological additive intended to improve the ensiling process at a minimum proposed application rate of 5.0×10^7 colony-forming units (CFU)/kg fresh material.

The bacterial species *L. plantarum* is considered by the European Food Safety Authority (EFSA) to be suitable for the qualified presumption of safety approach to safety assessment. Therefore, no specific demonstration of safety required other than confirmation of the absence of resistance to antibiotics of human and veterinary clinical significance. As the identity of the strain has been clearly established, and as no antibiotic resistance was detected, the use of the strain as a silage additive is presumed safe for livestock species, consumers of products from animals fed the treated silage and for the environment.

In the absence of data, no conclusion can be drawn on the skin and eye irritancy of the additive. The additive should be considered to have the potential to be a respiratory sensitiser.

Three studies with laboratory-scale silos, each lasting 90 days, were made using samples of forage of differing dry matter and water-soluble carbohydrate contents representing material considered easy, moderately difficult and difficult to ensile. In each case, replicate silos containing treated forage were compared with identical silos containing the same but untreated forage. The FEEDAP Panel concluded that the additive has the potential to improve the production of silage from easy, moderately difficult and difficult to ensile material by improving the preservation of nutrients. This was shown at the proposed application rate of 5×10^7 CFU/kg forage.

Table of contents

Abstract.....	1
Summary.....	3
1. Introduction.....	5
1.1. Background and Terms of Reference.....	5
1.2. Additional information.....	5
2. Data and methodologies.....	5
2.1. Data.....	5
2.2. Methodologies.....	5
3. Assessment.....	6
3.1. Characterisation.....	6
3.1.1. Characterisation of the active agent.....	6
3.1.2. Manufacturing process and characterisation of the product.....	6
3.1.3. Shelf life.....	7
3.1.4. Stability in water.....	7
3.1.5. Conditions of use.....	7
3.2. Safety.....	7
3.2.1. Safety for the target species, consumers and environment.....	7
3.2.2. Safety for the user.....	7
3.3. Efficacy.....	8
4. Conclusions.....	8
Documentation provided to EFSA.....	9
References.....	9
Abbreviations.....	9
Annex A – Executive Summary of the Evaluation Report of the European Union Reference Laboratory for Feed Additives on the Method(s) of Analysis for <i>Lactobacillus plantarum</i> DSM 29025.....	10

1. Introduction

1.1. Background and Terms of Reference

Regulation (EC) No 1831/2003¹ establishes the rules governing the Community authorisation of additives for use in animal nutrition. In particular, Article 4(1) of that Regulation lays down that any person seeking authorisation for a feed additive or for a new use of a feed additive shall submit an application in accordance with Article 7.

The European Commission received a request from Microferm Limited² for the authorisation of *Lactobacillus plantarum* DSM 29025, when used as a feed additive for all animal species (category: Technological additive; functional group: Silage additive).

According to Article 7(1) of Regulation (EC) No 1831/2003, the Commission forwarded the application to the European Food Safety Authority (EFSA) as an application under Article 4(1) (authorisation of a feed additive or new use of a feed additive). EFSA received directly from the applicant the technical dossier in support of this application. The particulars and documents in support of the application were considered valid by EFSA as of 21 January 2016.

According to Article 8 of Regulation (EC) No 1831/2003, EFSA, after verifying the particulars and documents submitted by the applicant, shall undertake an assessment in order to determine whether the feed additive complies with the conditions laid down in Article 5. EFSA shall deliver an opinion on the safety for the target animals, consumer, user and the environment and on the efficacy of the product *Lactobacillus plantarum* DSM 29025, when used under the proposed conditions of use (see Section 3.1.5).

1.2. Additional information

The additive is a preparation containing viable cells of *L. plantarum* DSM 29025. It has not been previously authorised as a feed additive in the European Union (EU).

The species *L. plantarum* is considered by EFSA to be suitable for the Qualified Presumption of Safety (QPS) approach to safety assessment (EFSA, 2007, EFSA BIOHAZ Panel 2013). This approach requires the identity of the strain to be conclusively established and evidence that the strain does not show resistance to antibiotics of human and veterinary importance.

2. Data and methodologies

2.1. Data

The present assessment is based on data submitted by the applicant in the form of a technical dossier³ in support of the authorisation request for the use of *L. plantarum* DSM 29025 as a feed additive. The technical dossier was prepared following the provisions of Article 7 of Regulation (EC) No 1831/2003, Regulation (EC) No 429/2008⁴ and the applicable EFSA guidance documents.

EFSA has verified the European Union Reference Laboratory (EURL) report as it relates to the methods used for the control of the active agent in animal feed. The Executive Summary of the EURL report can be found in Annex A.

2.2. Methodologies

The approach followed by the FEEDAP Panel to assess the safety and the efficacy of *L. plantarum* DSM 29025 is in line with the principles laid down in Regulation (EC) No 429/2008 and the relevant guidance documents: Guidance on technological additives (EFSA FEEDAP Panel, 2012a), Technical guidance: Tolerance and efficacy studies in target animals (EFSA FEEDAP Panel, 2011) Guidance for establishing the safety of additives for the consumer (EFSA FEEDAP Panel, 2012b), Guidance on studies concerning the safety of use of the additive for users/workers (EFSA FEEDAP Panel, 2012c),

¹ Regulation (EC) No 1831/2003 of the European Parliament and of the Council of 22 September 2003 on additives for use in animal nutrition. OJ L 268, 18.10.2003, p. 29.

² Microferm Limited, Spring Lane North, Malvern Link WR141BU Worcestershire United Kingdom.

³ FEED dossier reference: FAD-2015-0035.

⁴ Commission Regulation (EC) No 429/2008 of 25 April 2008 on detailed rules for the implementation of Regulation (EC) No 1831/2003 of the European Parliament and of the Council as regards the preparation and the presentation of applications and the assessment and the authorisation of feed additives. OJ L 133, 22.5.2008, p. 1.

and Guidance on the assessment of bacterial susceptibility to antimicrobials of human and veterinary importance (EFSA FEEDAP Panel 2012d).

3. Assessment

The additive is a preparation of viable cells of a single strain of *L. plantarum* DSM 29025 intended for use as a technological additive (silage additive) for all animal species.

3.1. Characterisation

3.1.1. Characterisation of the active agent

The strain was isolated from grass. It is deposited in the Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ) with the accession number DSM 29025.⁵ It has not been genetically modified.

Species identity was established by the phenotypic properties and by the nearly complete 16S rRNA gene sequence, which, by comparison with sequences recorded in databases, enabled the strain to be unambiguously identified as *L. plantarum*. Multilocus sequence typing based on sequencing four specific genes (*rpoA*, *pheS*, *atpA* and *dnaK*) was proposed as a means of strain-specific detection.⁶ Although the method is suitable for the discrimination of closely related strains, its effectiveness depends on the selection of sequences to be compared. No data were provided to illustrate that comparison of the four gene fragments chosen in this case is able to distinguish between DSM 29025 and other *L. plantarum* strains.⁷

The genetic stability was examined by comparing working cultures with culture collection stock using randomly amplified polymorphic DNA-polymerase chain reaction amplification (RAPD-PCR).⁸ No differences in the resultant patterns were observed.

The bacterial strain was tested for antibiotic susceptibility using broth microdilution techniques. The battery of antibiotics used included those recommended by EFSA (EFSA FEEDAP Panel, 2012d).⁹ All the minimum inhibitory concentration values were below the corresponding EFSA cut-off values except for chloramphenicol (MIC = 16 mg/L, cut-off value = 8 mg/L) which exceeded by one dilution. This is within the normal variation around the mean, and thus, does not raise concerns for safety.

3.1.2. Manufacturing process and characterisation of the product¹⁰

The manufacturing process is detailed in the dossier. The product consists of approximately 38% cells, 2% spent medium and 60% cryoprotectants. The additive is produced with a minimum declared content of 8×10^{10} colony-forming units (CFU)/g.¹¹ Material safety datasheets are provided for all medium components and cryoprotectants but no purity criteria are included.¹²

The strain is also intended for use in grow-up formulations in which numbers of bacteria are increased by incubation before application to forage. As the growth of the strain is encouraged, the product is also available in a formulation which contains (feed grade) nitrogen sources and buffer salts.

Analysis of five freeze-dried cell batches made showed a mean value of 5.1×10^{11} CFU/g additive (range $4.1\text{--}6.2 \times 10^{11}$ CFU/g additive).

Microbial contamination is routinely monitored at various points in the manufacturing process and in the final product. Limits are set for yeasts and filamentous fungi (< 10 CFU/g), presumptive coliforms, and *Escherichia coli* (< 10 CFU/g) and *Salmonella* spp. (absent in 25 g). Compliance with specifications was proved in five batches, except in one case that showed levels of presumptive coliforms of 40 CFU/g.¹³ Given the nature of the fermentation medium and the excipients, the probability of contamination with heavy metals or mycotoxins is considered to be low and consequently not included in routine monitoring of batches. Three batches of corn steep liquor powder (medium component) and five batches of *L. plantarum* (excipient not given) were tested for

⁵ Technical dossier/Section II/Annex_II_8_safedeposit_29025.

⁶ Technical dossier/Section II/Annex_II_2_5_ID_29024.

⁷ Technical dossier/Section II/Annex_II_2_5_ID_29025.

⁸ Technical dossier/Section II/Annex_II_2_genetic_stability_29025.

⁹ Technical dossier/Section II/Annex_II_1_antibioticresistance_29025.

¹⁰ This section has been amended following the confidentiality claims made by the applicant.

¹¹ Technical dossier/Supplementary information April 2016.

¹² Technical dossier/Section III/Annex MSDS Raw materials.

¹³ Technical dossier/Section II/Annex_II_4_contamination.

heavy metals (lead, cadmium and mercury), arsenic and aflatoxins (B1, B2, G1 and G2).¹⁴ Aflatoxin G2 was not detected ($< 0.01 \mu\text{g}/\text{kg}$), levels of aflatoxin B1 were $\leq 0.03 \mu\text{g}/\text{kg}$, of B2 were $\leq 0.05 \mu\text{g}/\text{kg}$ and of G1 were not detected or $\leq 0.04 \mu\text{g}/\text{kg}$. Contamination with heavy metals and arsenic was low and of no concern (lead $< 0.1 \text{ mg}/\text{kg}$, cadmium $< 0.1 \text{ mg}/\text{kg}$, mercury $< 0.01 \text{ mg}/\text{kg}$ and arsenic $< 0.1 \text{ mg}/\text{kg}$).

No specific data were provided on the particle size distribution or dusting potential of the additive under assessment.

3.1.3. Shelf life

Three batches of the product standardised with maltodextrin to give a count of 1×10^{11} CFU/g and another three with dextrose to a level of 2.5×10^{10} CFU/g were stored in sealed aluminium foil bags at ambient temperature.¹⁵ Viability losses were insignificant for both formulations over 6 months but reached up to 13% after 12 months in maltodextrin formulations and 8% in the dextrose formulations.

3.1.4. Stability in water

A batch of product was standardised to give a count of 1×10^{11} CFU/g using dextrose and ammonium and potassium phosphates as buffer salts. An experiment was designed to mirror practical conditions in which, typically, 10 g of product would be dissolved in 2 L of water and applied to 1 tonne of forage to deliver 1×10^9 CFU/kg.¹⁶ Three replicates of *L. plantarum* in solution were stored at room temperature and samples removed over 7 days. Viable cell counts made indicated that the strain was fully stable for at least 3 days under these conditions. Viability losses (up to approximately 25%) were observed at 7 days.

3.1.5. Conditions of use

The additive is intended for use with all forages and for all animal species at a proposed minimum concentration of 5×10^7 CFU/kg forage if applied with other microorganisms or 1×10^8 CFU/kg, if applied alone. It is to be applied as an aqueous suspension.

3.2. Safety

3.2.1. Safety for the target species, consumers and environment

In the view of the FEEDAP Panel, the antibiotic resistance qualification has been met and the identity of the strain established. Consequently, *L. plantarum* DSM 29025 is considered to be suitable for the QPS approach to safety assessment and, consequently, is presumed safe for the target species, consumers of products from animals fed treated silage and the environment.

3.2.2. Safety for the user

No specific data on skin/eye irritation or skin sensitisation were provided for the additive under application. Therefore, no conclusions can be drawn on the skin and eye irritancy of the additive. Given the proteinaceous nature of the active agent, the additive should be considered to have the potential to be a respiratory sensitiser.

Once an active agent has been authorised as a silage additive, different formulations can be placed on the market with reference to that authorisation. The applicant listed several cryoprotectants and carriers which would allow multiple formulations of the additive to be produced, and consequently, not all forms can be directly tested for user safety. However, for assessing the safety for the user of the additive, the active agent is the principal concern provided that other components do not introduce safety issues. For this specific product, the excipients used in the preparation of the final formulation do not introduce additional risks.

¹⁴ Technical dossier/Section II/Annex_II_6_mycotoxins_heavymetals.

¹⁵ Technical dossier/Section II/2.4.1.1.

¹⁶ Technical dossier/Section II/2.4.1.2.

3.3. Efficacy

Three laboratory experiments were made with different forage samples. The duration of the experiments was 90 days. In all the studies, forage was ensiled in mini-silos with a capacity of 4.5 L. All the silos were fitted with air-locks to vent gas. The ambient temperature during ensiling was controlled at $20 \pm 2^\circ\text{C}$. The additive was dissolved in water and sprayed on the forage at an intended concentration of 5×10^7 CFU/kg fresh matter (not confirmed by analysis). Forage for the control silos were sprayed with an equal volume of water, but without the additive. Four replicate silos were prepared for each experimental treatment (with or without the additive). The forages used were mixtures of clover and grasses with different botanical composition and different dry matter (DM) and water-soluble carbohydrate (WSC) content (see Table 1) to represent material easy to ensile (study 1), moderately difficult to ensile (study 2) and difficult to ensile (study 3), as specified by Regulation (EC) No 429/2008.

Table 1: Characteristics of the forage samples used in the five ensiling experiments

Study	Test material	Dry matter content (%)	Water-soluble carbohydrate content (% fresh matter)
1 ^(a)	Red clover, timothy, meadow fescue	43.4	3.4
2 ^(b)	Red clover, timothy, meadow fescue, meadow foxtail, perennial ryegrass	40.8	2.3
3 ^(c)	Alfalfa, red clover, white clover, meadow fescue	21.8	1.2

(a): Annexes IV.1 and IV.4.

(b): Annexes IV.1 and IV.3.

(c): Annexes IV.1 and IV.2.

Silos were opened after 90 days and the contents were analysed by conventional methods to determine silage dry matter (DM) content, pH, lactic and volatile fatty acids concentrations, ethanol, ammonia and total nitrogen. DM loss during ensiling was calculated.

Statistical evaluation of data was by a non-parametric test (Wilcoxon Kruskal–Wallis test), comparing treated versus control silos. Significance was declared at $p < 0.05$.

Table 2: Summary of the analysis of ensiled material recovered at the end of the ensiling period (90 days) with *Lactobacillus plantarum* DSM 29025

Study	Application rate (CFU/kg forage)	Dry matter loss (%)	pH	Lactic acid (% dry matter)	Acetic acid (% dry matter)	Ammonia-N (% total N)
1	0	2.2	4.6	7.1	1.6	6.4
	5×10^7	1.7*	4.4*	6.8	1.0*	4.6*
2	0	1.7	4.8	4.5	1.1	7.3
	5×10^7	1.4*	4.3*	6.6*	0.9*	4.2*
3	0	3.9	4.6	7.8	3.6	8.8
	5×10^7	3.6*	4.6	8.1	3.3*	8.4

*: Significantly different from the control value at $p < 0.05$.

The addition of *L. plantarum* DSM 29025 at 5×10^7 CFU/kg fresh material decreased DM loss and acetic acid concentration in all forage materials (Table 2). With easy and moderately difficult to ensile clover–grass mixtures, the additive also decreased ammonia-N in silage implying a better preservation of proteins.

The additive has the potential to improve the preservation of nutrients in silage prepared from easy, moderately difficult and difficult to ensile material.

4. Conclusions

As the identity of *L. plantarum* DSM 29025 has been established and no antibiotic resistance of concern has been detected, following the QPS approach to safety assessment, the use of this strain as a silage additive is considered safe for the target species, consumers of products from animals fed treated silage and the environment.

In the absence of data, no conclusion can be drawn on the skin and eye irritancy of the additive. The additive should be considered to have the potential to be a respiratory sensitiser.

The additive has the potential to improve the production of silage from easy, moderately difficult and difficult to ensile material by improving the preservation of nutrients. This was shown at the proposed application rate of 5×10^7 CFU/kg forage.

Documentation provided to EFSA

- 1) *Lactobacillus plantarum* (DSM 29025) October 2015. Submitted by Microferm Limited.
- 2) *Lactobacillus plantarum* (DSM 29025). Supplementary information February 2016. Submitted by Microferm Limited.
- 3) Evaluation report of the European Union Reference Laboratory for Feed Additives on the Methods(s) of Analysis for *Lactobacillus plantarum* DSM 29025.
- 4) Comments from Member States.

References

- EFSA (European Food Safety Authority), 2007. Opinion of the Scientific Committee on a request from EFSA on the introduction of a Qualified Presumption of Safety (QPS) approach for assessment of selected microorganisms referred to EFSA. EFSA Journal 2007;5(12):587, 16 pp. doi:10.2903/j.efsa.2007.587
- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), 2013. Scientific Opinion on the maintenance of the list of QPS biological agents intentionally added to food and feed (2013 update). EFSA Journal 2013;11(11):3449, 108 pp. doi:10.2903/j.efsa.2013.3449
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2011. Technical guidance: tolerance and efficacy studies in target animals. EFSA Journal 2011;9(5):2175, 15 pp. doi:10.2903/j.efsa.2011.2175
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2012a. Guidance for the preparation of dossiers for technological additives. EFSA Journal 2012;10(1):2528, 23 pp. doi:10.2903/j.efsa.2012.2528
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2012b. Guidance for establishing the safety of additives for the consumer. EFSA Journal 2012;10(1):2537, 12 pp. doi:10.2903/j.efsa.2012.2537
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2012c. Guidance on studies concerning the safety of use of the additive for users/workers. EFSA Journal 2012;10(1):2539, 5 pp. doi:10.2903/j.efsa.2012.2539
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2012d. Guidance on the assessment of bacterial susceptibility to antimicrobials of human and veterinary importance. EFSA Journal 2012;10(6):2740, 10 pp. doi:10.2903/j.efsa.2012.2740

Abbreviations

CFU	colony-forming unit
DM	dry matter
EC	European Commission
EURL	European Union Reference Laboratory
FEEDAP Panel	EFSA Panel on Additives and Products or Substances used in Animal Feed
MIC	minimum inhibitory concentration
PFGE	pulsed field gel electrophoresis
QPS	Qualified Presumption of Safety
RAPD-PCR	randomly amplified polymorphic DNA-polymerase chain reaction amplification
WSC	water-soluble carbohydrate

Annex A – Executive Summary of the Evaluation Report of the European Union Reference Laboratory for Feed Additives on the Method(s) of Analysis for *Lactobacillus plantarum* DSM 29025¹⁷

In the current application authorisation is sought under Article 4(1) for *Lactobacillus plantarum* DSM 29025 under the category/functional group 1(k) “technological additives”/“silage additives”, according to Annex I of Regulation (EC) No 1831/2003. Authorisation is sought for the use of the feed additive for all animal species. According to the Applicant, the active substance in the feed additive consists in viable cells of the non-genetically modified strain *Lactobacillus plantarum* DSM 29025. The feed additive is to be marketed as a powder containing a minimum *Lactobacillus plantarum* DSM 29025 concentration of 8×10^{10} Colony Forming Unit (CFU)/g. The feed additive is intended to be added to silage at a minimum dose of 5×10^7 CFU/kg fresh silage.

For the identification of *Lactobacillus plantarum* DSM 29025, the EURL recommends for official control Pulsed Field Gel Electrophoresis (PFGE), a recognised standard methodology for genetic identification. This methodology for microbial identification is currently being evaluated by the CEN Technical Committee 327 to become a European Standard.

For the enumeration of *Lactobacillus plantarum* DSM 29025, the Applicant submitted the ring-trial validated spread plate method EN 15787 which was already evaluated by EURL in the frame of previous *Lactobacillus plantarum* dossiers. Based on the performance characteristics available, the EURL recommends for official control this ring-trial validated EN 15787 method for the enumeration of *Lactobacillus plantarum* DSM 29025 in the feed additive per se.

The Applicant did not provide any data or experimental method for the determination of *Lactobacillus plantarum* DSM 29025 in silage, since the unambiguous determination of the content of *Lactobacillus plantarum* DSM 29025 added to silage is not achievable by analysis. Therefore, the EURL cannot evaluate nor recommend any method for official control to determine *Lactobacillus plantarum* DSM 29024 in silage.

Further testing or validation of the methods to be performed through the consortium of National Reference Laboratories as specified by Article 10 (Commission Regulation (EC) No 378/2005) is not considered necessary.

¹⁷ The full report is available on the EURL website: https://ec.europa.eu/jrc/sites/default/files/finrep_fad_2015_0035_lactob_plantarum.pdf