

1 **Nutrients, phytochemicals and botanical origin of commercial bee pollen**
2 **from different geographical areas**

3
4 Claudio Gardana^{a,*}, Cristian Del Bo^{'a}, Marta C. Quicazán^b, Ana Ruby Corrrea^b and Paolo
5 Simonetti^a

6
7
8
9 ^a Università degli Studi di Milano, DeFENS – Department of Food, Environmental and
10 Nutritional Sciences. Via Celoria 2, 20133 Milano,

11 ^b Universidad Nacional de Colombia, Institute of Food Science and Technology, Bogotá,
12 Colombia.

13
14
15 **Corresponding author.* Dr. Claudio Gardana. Tel.: +39-02-50316722; Fax: +39-02-
16 50316721; E-mail: claudio.gardana@unimi.it

17
18 **Abbreviations**

19 AA: amino acid; BP: bee pollen; DBP: dried bee pollen; MUFA: monounsaturated fatty acids;
20 PA: polyamine; SFA: saturated fatty acids; TL: total lipid; UFA: unsaturated fatty acids

21 **ABSTRACT**

22

23 This work evaluated the nutritional, phytochemical composition and botanical origin of
24 commercial bee pollen from three different countries. Fructose (17-23%) was the most
25 abundant sugar in all samples, followed by glucose (14-16 %) and sucrose (5-6%). The
26 protein content in Colombian (24%) and Italian (22%) pollen was higher compared to Spanish
27 sample (14%). The total lipid contents were higher for the Spanish (6%) and Colombian
28 pollens (6%) than the Italian sample (2.5%). Twenty-one fatty acids were identified, and the
29 most abundant were palmitic, α -linolenic, linoleic and oleic acid. Colombian pollen was rich
30 in n-3 fatty acids, while Italian and Spanish samples contained high amounts of n-6 fatty
31 acids. Polyphenols and carotenoids were identified by UHPLC-DAD-Orbitrap mass
32 spectrometry detection. Thirty-nine polyphenols were identified, and the dominant
33 compounds were tri-caffeoyl- and caffeoyl-di-*p*-coumaroyl spermidine derivatives. Di-lauryl-
34 zeaxanthin was the main carotenoid detected in all the samples analysed. Colombian pollen
35 contained traces of lutein, zeaxanthin, β -carotene and phytoene, while only β -carotene was
36 present in the Spanish and Italian samples. After saponification, the average total amount of
37 carotenoids was 57, 25 and 221 $\mu\text{g g}^{-1}$ in pollen from Spain, Italy and Colombia, respectively.
38 The free proline to total free amino acid ratio was 53, 59 and 78 for pollen from Spain, Italy
39 and Colombia, respectively.

40

41 *Keywords*

42 Food analysis, Food composition, Pollen, Mass spectrometry, Nutrients, Polyamines,
43 Polyphenols, Carotenoids

44 **1. Introduction**

45

46 The chemical composition of bee pollen depends on several factors, such as plant source,
47 geographical origin, seasonal conditions and bee activities (Nogueira et al., 2012).

48 Furthermore, the content of nutrients can change based on the processing methods, steps and
49 storage conditions (Bogdanov, 2004).

50 Pollen contains proteins, amino acids, carbohydrates, lipids, vitamins and minerals (Yang
51 et al., 2013), polyphenols (Mihajlovic et al., 2015), phytosterols (Wang et al., 2015) and
52 carotenoids (Bunea et al., 2014). In particular, pollen contains 23 % of protein on average,
53 including 10 % of essential amino acids (AA) such as methionine, lysine, threonine, histidine,
54 leucine, isoleucine, valine, phenylalanine, and tryptophan (Almeida-Muradian et al., 2005).
55 The total lipid (TL) content of bee pollen (BP) is typically lower than 10% of the dry weight
56 (DW) (Nicolson, 2011) and the most abundant are linoleic (18:2n-6), α -linolenic (18:3n-3)
57 and palmitic (16:0) acids (Szczęśna, 2006). Sugar occurs in pollen loads in the amount of 40
58 % on average, ranging from 15 to 24% for fructose, 11-18% for glucose and 4-9% for
59 sucrose. Other sugars such as arabinose, ribose, isomaltose and melibiose accounted for about
60 1% (Szczęśna et al., 2002).

61 The yellow-red appearance of pollen loads, which are formed by the aggregation of
62 microscopic granules, is determined mainly by the presence of pigments, such as flavonoids
63 and carotenoids (Stanley and Linskens, 1974).

64 The principal carotenoids found in bee pollen after saponification are lutein and β -
65 cryptoxanthin, while β -carotene is detected in small or trace amounts (Mărgăoan et al., 2014).
66 Carotenoids have important roles in human health. β -Carotene, for example, constitutes the
67 principal source of vitamin A, and its dietary intake could lower the risk of various types of
68 degenerative diseases (Milani et al., 2017).

69 The glycosides of quercetin, kaempferol and isorhamnetin are the predominant flavonoids
70 found in BP. The total amount ranges between 0.3–1.1% (Bonhevi et al., 2001; Han et al.,
71 2012).

72 Bee pollen also contains polyamines (PAs) and PAs linked to cinnamic acid derivatives,
73 such as *p*-coumaric acid, caffeic acid and ferulic acid. The biosynthesis of PAs occurs by the
74 addition of one or two aminopropyl groups to putrescine, to form spermidine and spermine,
75 respectively. Covalent binding of PAs to *p*-coumaric acid, caffeic acid and ferulic acid,
76 abundant in many plant families, gives rise to hydroxyl-cinnamic acid (HCA) amides (Aloisi
77 et al., 2016). These are involved in the organization of the cell wall, and most of them are
78 associated with fertility (Grienenberger et al., 2009). Moreover, the conjugated PAs are non-
79 peptide antagonists of tachykinin neurokinin1 receptors and may benefit depression and
80 anxiety (Yamamoto et al., 2002).

81 Bee pollen may exert a wide range of biological activities, including antifungal,
82 antimicrobial, antiviral, immunostimulating, local analgesic, hypolipidaemic, anti-
83 atherosclerotic, anti-inflammatory and cerebrovascular (Denisow and Denisow-Pietrzyk,
84 2016; Pascoal et al., 2014).

85 The use of the geographical indication for bee products has been proposed because of a
86 higher demand from consumers for traced and well-characterized products. Thus, the
87 evaluation of the nutritional composition and quality parameters of bee pollen from different
88 geographical areas could lead useful information to both producers and consumers. Also in
89 view of the increased number of studies that have highlighted bee pollen as a functional food
90 (Komosinska-Vassev et al., 2015).

91 Given this considerations, the main objective of this work was to determine and
92 compare the nutritional and phytochemical composition of commercial BP produced in

93 apiaries from Colombia, Italy and Spain. Regarding nutritional parameters, the moisture, ash,
94 lipid, carbohydrate, amino acid and protein contents were determined.

95 The Proline Index, the ratio of proline/total free amino acids, has been determined as an
96 indicator of the freshness of the pollen. For fresh products, the proline index must be less than
97 80 % (Serra Bonvehí and Escolá Jorda, 1997).

98 As phytochemicals, the profiles of the carotenoids, polyphenols and PAs were evaluated
99 by ultra-performance liquid chromatography coupled to diode-array detector and high-
100 resolution mass spectrometer.

101 Principal component analysis (PCA) was carried out to check for similarity between
102 samples according to the nutrients, phytochemicals and geographical area of origin.

103 Finally, botanical origin were identified through palynological analysis.

104

105 **2. Materials and methods**

106

107 *2.1. Chemicals*

108

109 Astaxanthin, lutein, zeaxanthin, β -cryptoxanthin, violaxanthin, echinenone (internal
110 standard), α -carotene and β -carotene were provided by Hoffmann-La Roche (Basel,
111 Switzerland). The fatty acid methyl esters (FAME) were purchased from Supelco (Bellefonte,
112 PA, USA). Fructose, glucose, sucrose, acetonitrile, methanol, formic acid, chloroform,
113 ethanol, tetrahydrofuran (THF), butylated hydroxytoluene, hexane, norvaline (IS), glutamine,
114 cysteine, asparagine and tryptophan were procured from Sigma-Aldrich (St. Louis, MO,
115 USA). ACCQ*TAG Ultra Derivatization kit for the AA determination was purchased from
116 Waters (Milford, USA). The kit contained AA standard (Alanine, arginine, glycine, histidine,
117 isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tyrosine,

118 valine, aspartic acid and glutamic acid), derivatising reagent (6-aminoquinolyl-N-
119 hydroxysuccinimidyl carbamate), eluents and a sub-2 µm column for AA analysis. Water was
120 supplied through a Milli-Q apparatus (Millipore, Milford, MA, USA).

121

122 2.2. *Samples*

123

124 In this study commercial bee pollen samples from Italy (n=1), Spain (n=1) and
125 Colombia (n=1) were evaluated for their content in nutrients, phytochemicals and botanical
126 origin. Blended BP from southern Spain was harvested by beekeepers in the province of
127 Córdoba (37°50' N, 4°45' W) between April and July 2017 and stored at -20 °C.

128 Italian sample resulted from the blend of pollen loads collected by the beekeeper in the
129 municipality of Montaldeo (44°40'05 N, 8°43'52 E) in Alessandria province (Piedmont)
130 between April and July 2017 and immediately frozen.

131 No information is available regarding the technological treatment carried out on Italian
132 and Spanish pollen. The samples were vacuum-packed and available in 100 g packs.

133 Colombian pollen comes from the native forest of the Colombian Cundi-boyacence high
134 pleateau zone at about 2200m above sea level (5°55'45 N, 75°59'02 W), where about 90% of
135 the bee-pollen domestic production is concentrated. The bee pollen was gathered, sun
136 dehydrated for 2 days and packed. The drying of the pollen by directly exposing to the sun is
137 the most used method by farmers because Cundi-boyacence high plateau is one of the zones
138 with highest solar radiation in Colombia.

139 *2.3. Botanical origin of the pollen loads*

140

141 Approximately 2 g of each pollen load was grouped into subsamples, according to their
142 colour and each subsample was weighed and analysed by a Nikon Eclipse 50i optic
143 microscope at 1000 and 400× for identification and counting, respectively. The total
144 percentage of each pollen was catalogued as described by Barth (2004) into dominant pollen
145 (PD, >45% of the total), secondary pollen (PS, 16–45%), important pollen (PI, 3–15%) and
146 occasional pollen (PO, <3%). A reference collection of the Fojanini Foundation (Sondrio, IT)
147 and various pollen morphology guides were used to identify the pollen types.

148

149 *2.4. Moisture and ash determination*

150

151 The vacuum packs of pollen, 100g, have been opened and aliquots of 3g were weighed
152 and heated at 65 °C for 24 h in a ventilated stove. The samples were then cooled in the drier
153 to laboratory temperature, about 24 °C, and weighed. Moisture content was obtained by
154 difference. Ash determination was made through gravimetry after incineration in a muffle
155 furnace at 600°C until constant weight (AOAC, 2005).

156

157 *2.5. Determination of the TL and fatty acids*

158

159 The Soxhlet extraction was achieved with 5 g of BP powder using a Soxtec HT 1043
160 system (Foss, Milan, IT) containing 180 mL of a solution ethyl ether:petroleum ether (2:1,
161 v/v). The BP was extracted at 140°C for 6h, followed by a 30 min solvent rinse and solvent
162 evaporation. The weighed residue was considered as the lipid content (% DW). Subsequently,
163 the fatty acids were analysed by GC-FID. Chromatographic separations were achieved using

164 an Omegawax 320 capillary column (30 m × 0.32 mm i.d.; Supelco, Milan, IT), under the
165 following conditions: initial isotherm, 140°C for 5 min; temperature gradient, 2°C min⁻¹ to
166 210°C; final isotherm, 210°C for 20 min. The injector temperature was 250°C. The injection
167 volume was 1 µL, with a 1/100 split ratio and the FID temperature was 250°C. The carrier
168 and makeup gas were H₂ and N₂, respectively. Fatty acid retention times were obtained by
169 injecting the Omegawax test mix as the standard. The fatty acid content was expressed as a
170 percentage of the total fatty acids using the following formula:

171 $FA \% = (\text{Peak area} / \text{Total area}) \times 100$, where Peak area= area of the fatty acid and Total
172 area= sum of the areas of the individual fatty acids detected in the chromatogram.

173

174 *2.6. Protein determination*

175

176 The nitrogen (N) content was determined by conventional acid hydrolysis and Kjeldahl
177 digestion, using a copper catalyst in 2 g of pulverised pollen. The ammonia was distilled and
178 collected in a solution of boric acid, which was then titrated against standard acid. Digestion
179 and distillation were carried out using a Kjeltec 1002 apparatus (Foss, Milan, IT). Protein
180 content was calculated as total N×5.6.

181

182 *2.7. Carotenoid determination*

183

184 *2.7.1. Extraction*

185

186 Carotenoids were extracted from 10 g of BP with 30 mL of hexane. Extraction was
187 conducted under agitation for 2 min and diminished light. The procedure was repeated until
188 the BP became colourless. The extracts were combined and dried. The residue was dissolved

189 in 50 mL of methanol:THF (80:20, v/v). An aliquot was then used for the determination of
190 free carotenoids by LC-DAD.

191

192 2.7.2. Saponification

193

194 Hydrolysis of esterified carotenoids was carried out according to Riso and Porrini
195 (1997), with slight modifications. Briefly, 40 mL of each extract was saponified with 10 %
196 methanolic KOH (9:1, v/v), at room temperature, in the dark, under an N₂ stream for 2 h, with
197 a 5-mL aliquot removed every 20 min. For the removal of soaps and alkalis, petroleum ether
198 was added, and the solution washed with a saturated NaCl solution to achieve neutrality. The
199 organic layer containing carotenoids was dried over anhydrous sodium sulphate, evaporated
200 to dryness and the resultant samples were then kept under N₂ at -80°C until utilisation.

201

202 2.7.3. Quantitative analysis

203

204 The quantitative determination of the non-esterified carotenoids was performed
205 according to Riso and Porrini (1997), with slight modifications. The chromatographic system
206 was an Alliance 2695 (Waters, Milford, US) coupled to a DAD 2998 (Waters) and a Vydac
207 201TP54 column (C₁₈, 250 × 4.6 mm i.d., 5 µm; Esperia, CA, US). The flow-rate was 1.5 mL
208 min⁻¹, and 50 µL was injected. The column and sample were maintained at 25 and 15 °C,
209 respectively. The eluents were methanol (A) and THF (B). Chromatographic separations were
210 carried out in gradient elution mode, performed as follows: 0 % B for 3 min, 0 % B to 5 % B
211 in 0.1 min, 5% B for 5 min, 5% B to 10 % B in 0.1 min, 10% B for 5 min, 10% B to 20% B in
212 0.1 min and then 20% B for 10 min. Spectra were acquired in the range 220–700 nm and the
213 chromatograms integrated at 445 nm. Acquisition and quantification were performed using

214 Empower software (Waters). Calibration curves for lutein, zeaxanthin, β -cryptoxanthin, α -
215 carotene and β -carotene ranged from 0.1–2.0 $\mu\text{g/mL}$, and the results were expressed as
216 micrograms per gram of DBP. The total carotenoids content was also estimated
217 spectrophotometrically at 450nm from saponified samples and the data reported as μg
218 zeaxanthin/g DBP (zeaxanthin ϵ^{2540} , Lombeida et al., 2016).

219

220 2.7.4. Determination of esterified carotenoids by UHPLC-DAD-Orbitrap mass spectrometry

221

222 The esterified carotenoids were analyzed by using an Acquity UHPLC (Waters),
223 coupled to a DAD eLambda (Waters) and an Exactive Orbitrap (Thermo Scientific, San Jose,
224 CA), equipped with a HESI-II probe for electrospray ionization (ESI). The column was a
225 BEH Shield C_{18} (150 \times 2.1 mm i.d., 1.7 μm ; Waters) and the flow-rate was 0.45 mL min^{-1} .
226 The column and sample were maintained at 45 and 15 $^{\circ}\text{C}$, respectively. Five μL was injected.
227 The eluents were methanol (A) and THF (B). Separations were performed in gradient mode
228 and the profile was: 0% B for 10 min, 0% B to 30% B in 10 min, and then 30% B for 10 min.
229 Data were acquired over the wavelength range 220–700 nm and the chromatograms were
230 plotted in the range 300–550 nm. The operative conditions were as follows: spray voltage +5
231 kV, sheath gas flow-rate 60 (arbitrary units), auxiliary gas flow-rate 20 (arbitrary units),
232 capillary temperature 275 $^{\circ}\text{C}$, capillary voltage +37.5 V, tube lens +125 V, skimmer +26 V,
233 and heater temperature 120 $^{\circ}\text{C}$. The analytes were identified in positive ESI mode by full-
234 scan acquisition (m/z^{+} 200–1000 u), using an isolation window of ± 2 ppm. The automatic gain
235 control (AGC) target, injection time, mass resolution and collision energy were 1×10^6 , 100
236 ms, 50 K and 50 eV respectively. The MS data were processed using Xcalibur software
237 (Thermo Scientific).

238 2.8. Carbohydrate determination

239

240 Approximately 20 mg of BP were dispersed in 10 mL of deionised water, and the
241 suspension was then sonicated for 10 min, centrifuged at 1000 ×g for 5 min, and the
242 supernatant recovered. The residue was extracted with 10 mL of water and treated as
243 described above. The supernatants were combined, and then the final volume was adjusted to
244 50 mL with acetonitrile. The sugar content was assessed by using an Acquity UHPLC
245 (Waters) coupled to an Exactive Orbitrap (Thermo Scientific), equipped with a HESI-II probe
246 for ESI. The column was a BEH amide C₁₈ (150 × 2.1 mm i.d., 1.7 μm; Waters) and the flow-
247 rate was 0.2 mL min⁻¹. One μL was injected. The column and sample were kept at 25 and
248 20°C, respectively. The eluents were 0.02% NH₄OH in water:0.02% NH₄OH in acetonitrile
249 (72:28, v/v). The operative conditions were as follows: spray voltage 3.0 kV, sheath gas flow-
250 rate 40 (arbitrary units), auxiliary gas flow-rate 10 (arbitrary units), capillary temperature
251 300°C, capillary voltage -37.5 V, tube lens -125 V, skimmer -26 V, and heater temperature
252 300°C. The analytes were identified in negative ESI mode by full-scan acquisition (m/z 100–
253 600 u), using an isolation window of ±2 ppm. The AGC target, injection time and mass
254 resolution were 1 × 10⁶, 100 ms and 50 K, respectively. The MS data were processed using
255 Xcalibur software (Thermo Scientific). The peak identity was ascertained by evaluation of
256 both the accurate mass and the retention time. Calibration curves for fructose, glucose and
257 sucrose were constructed in the range 2–50 μg mL⁻¹, and the results were expressed as
258 percentages.

259 The equations of calibration curve for fructose (Fru, 2.0-50.5 μg mL⁻¹), glucose (Glu, 2.0-
260 49.5 μg mL⁻¹) and sucrose (Suc, 2.1-51.5 μg mL⁻¹) were as follows:

261 Fru: $Y = 29.3 X + 5.2$, $R^2=0.997$, $n=5$

262 Glu: $Y = 27.8 X - 3.2$, $R^2=0.995$, $n=5$

263 Suc: $Y = 12.8 X + 2.4$, $R^2=0.995$, $n=5$

264 Where Y =peak area $\times 10^{-3}$, X = $\mu\text{g/mL}$.

265

266 2.9. Polyphenol and PA determination

267

268 The BP samples were finely powdered by milling, and 200 mg extracted with methanol

269 (20 mL) at 70°C for 2 h. The mixture was cooled, filtered and the solid residue was extracted

270 using the same methanol volume. The resulting solutions were mixed and methanol added to

271 adjust the volume to 50 mL. The solution was centrifuged at 1000 $\times g$ for 2 min, and 5 μL

272 injected into the UHPLC system. The analysis was performed on an Acquity UHPLC

273 (Waters) coupled with an eLambda DAD (Waters) and a high-resolution Fourier transform

274 Orbitrap mass spectrometer (Exactive, Thermo Scientific), equipped with a HESI-II probe for

275 ESI and a collision cell (HCD). The operative conditions were as follows: spray voltage -3.0

276 kV, sheath gas flow-rate 55 (arbitrary units), auxiliary gas flow-rate 20 (arbitrary units),

277 capillary temperature 350°C, capillary voltage -37.5 V, tube lens -125 V, skimmer -26 V, and

278 heater temperature 130°C. The injection volume was 5 μL . A BEH Shield C_{18} column (150

279 \times 2.1 mm, 1.7 μm ; Waters) was used for the separation. The column was maintained at 50°C.

280 The flow-rate was 0.45 mL/min, and the eluents were 0.05% formic acid in water (A) and

281 acetonitrile (B). The UHPLC separation was achieved by the following linear elution

282 gradient: 5–35% of B in 10 min, which was then increased from 35 to 80% B in 10 min. The

283 acquisition was made in the full-scan mode in the range (m/z) 80–1000 u, using an isolation

284 window of ± 2 ppm. The AGC target, injection time, mass resolution, energy and gas in the

285 collision cell were 1×10^6 , 100 ms, 50 K, 20-40-60 V and N_2 respectively. The MS data were

286 processed using Xcalibur software (Thermo Scientific). The peak identity was ascertained by

287 evaluating the accurate mass, the fragments obtained in the collision cell and the on-line UV
288 spectra (200–450 nm).

289

290 *2.10. Amino acid determination*

291

292 Free amino acid fraction was obtained as reported by Serra Bonvehí and Escolá Jorda,
293 (1997) with a slight modification. Briefly, 1 g of pulverized pollen was vortexed in 20 mL of
294 water, centrifuged at 6000 x g for 10 min at 4 °C and the supernatant transferred into a 50 mL
295 volumetric flask. The residue was extracted with 20 ml of water, treated a described above
296 and the final volume adjusted to 50 mL with water.

297 Amino acid derivation with AccQ•Tag reagents was conducted according to the
298 manufacturer's protocol. Briefly, amino acids were derivatized by adding 80 µL of AccQ•Tag
299 ultra borate buffer, 10 µL of IS (0.3 mg mL⁻¹) and 20 µL of derivatising reagent to 10 µL of
300 bee pollen extract or standard solution. The mixture was then incubated for 10 min at 55 °C.
301 The 10 min heating was an important step when converted the phenolic side chain of tyrosine
302 to free phenol, so tyrosine could become a major mono-derivatized compound like the rest of
303 the amino acids. Heating the samples has no other significant effect. Amino acid derivatives
304 in room temperature were stable for as long as 1 week.

305 Liquid chromatographic separation was performed on an Acquity UPLC system
306 (Waters) coupled with an eLambda DAD (Waters). The injection volume was 2 µL. The
307 derivatives were separated on an AccQ•Tag Ultra Column (1.7 µm BEH, 100 x 2.1 mm)
308 maintained at 50°C. The flow-rate was 0.7 ml min⁻¹. The eluents and nonlinear separation
309 gradient used were as reported by Armenta et al. (2010). The data were acquired in the range
310 200-450 nm and chromatogram integrated at 260 nm.

311 The stock solution contained a 2.5 mM concentration of each amino acid with the
312 exception of cysteine with the concentration 1.25 mM. Calibration curves were in the range
313 2.2-51 $\mu\text{g mL}^{-1}$.

314

315 *2.11. Statistical analysis*

316

317 Statistical analysis was performed using Statistica software (Statsoft, Tulsa, OK, USA).
318 Results were expressed as mean values \pm standard deviation. Differences between bee pollen
319 samples were analysed by one-way analysis of variance (ANOVA). $P < 0.05$ was considered
320 statistically significant. Principal component analysis (PCA) was performed using R statistical
321 software 3.1.2 by means the function PRCOMP.

322

323 **3. Results and discussion**

324

325 The chemical composition of the pollen samples is given in Table 1. No difference
326 among the groups was found for moisture and ash content ($p > 0.05$). With regard to moisture,
327 fresh bee pollen contains about 20-30% of water, which are favorable conditions for the
328 development of bacteria, yeasts and molds. Thus, for preventing bacterial contaminations and
329 preserve the quality, pollen must be harvested daily and frozen suddenly. Freezing caused no
330 substantial changes in the chemical composition of the pollen loads. After thawing, pollen
331 may be processed by different methods such as desiccation, lyophilization, freeze-drying and
332 microwave-assisted drying. Lyophilisation and drying at 40 °C seem to decrease some
333 nutrients like ascorbic acid (Campos et al., 2005), while microwave-assisted drying seems to
334 reduce some antioxidant compounds such as tocopherols (Conte et al., 2017). After drying,

335 the humidity level should be in the range of 4-8%. In this condition pollen retains its
336 nutritional quality and health safety.

337 In all the samples analyzed the humidity was about 15%. This was most likely due at
338 mild thermal treatment activity of the producers. Despite this relatively high water content,
339 microbiological and chemical analysis have not showed relevant concerns and violations of
340 the safety regulations. In particular, mesophilic microbe content was less than 10^5 cfu g⁻¹. *S.*
341 *aureus*, *E. coli* and *Salmonella spp.* absents and yeasts were 2×10^3 , 5×10^3 and 2×10^4 cfu g⁻¹ in
342 the pollen from Italy, Spain and Colombia, respectively (data not showed). Aflatoxins and
343 ochratoxin were not detected ($<2 \mu\text{g Kg}^{-1}$, data not showed).

344

345 3.1. Nutrients

346

347 Three main sugars were identified and quantified. Fructose was the most abundant,
348 followed by glucose and sucrose (Table 1), and the fructose/glucose ratio varied between 1.20
349 and 1.5. The sugar contents of the various pollen samples were not significantly different and
350 in agreement with literature values (Conti et al., 2016; Szczęśna et al., 2002).

351 The bee pollen protein content ranged from 12.3–21.6 g 100g⁻¹ DBP, evaluated by
352 multiplying the N content by a factor of 5.6. These results were similar to those obtained by
353 Fuenmayor et al. (2014) and Gabriele et al. (2015). The protein contents of the Colombian
354 and Italian pollens were higher compared to Spanish pollen. The differences were likely due
355 to their botanical origins.

356 The content of TL, the relative percentage of saturated fatty acids (SFAs) and
357 unsaturated fatty acids (UFAs), n-3, n-6, and the UFA/SFA ratios of the TLs from the
358 Colombian, Italian and Spanish honeybee-collected pollen is reported in Table 1. The fatty
359 acid compositions are provided in Table 2. The TL content was highly variable and ranged

360 from 2.0–6.5%, with samples from Colombia and Spain containing significant greater TL
361 amounts relative to the Italian bee pollen. Colombian pollen was rich in n-3 fatty acids, while
362 Italian and Spanish samples contained high amounts of n-6 fatty acids.

363 The Colombian pollen has an excellent n-6/n-3 ratio (0.3), which allows an n-3 supply
364 of about 0.4 g dose⁻¹ (15 g), considering that the sample contains 6% lipid. The Italian product
365 has a good n-6/n-3 ratio (1.4) but a low lipid content (2.5%). Thus, 15 g pollen provide only
366 0.09 g of n-3. On the contrary, Spanish sample has a high lipid content (5.9%) but a poor n-
367 6/n-3 ratio (2.1), and this brings to get an n-3 intake of 0.16 g dose⁻¹. Overall, Colombian
368 pollen provides an amount of n-3 fatty acids 4.4 and 2.5 times higher than that provided by
369 the Italian and Spanish pollen, respectively.

370 Considering that in Italy the n-3 target daily intake must be in the range 1.4-5.5 g
371 (LARN, 2014), pollens with high lipid content and low n-6/n-3 ratio could be an attractive
372 alternative to increase n-3 intake.

373 Regarding the SFA content and UFA/SFA ratio, no significant differences among the
374 pollen samples were observed. On the contrary, the monounsaturated fatty acid (MUFA)
375 content was higher in the Italian pollen than the Colombian and Spanish specimens. Twenty-
376 one fatty acids were identified in the pollen extracts. The most abundant were α -linolenic
377 (18:3n-3), linoleic (18:2n-6), palmitic (16:0) and oleic acid (18:1n-9). In particular,
378 Colombian pollen was high in α -linolenic (44.1% of fatty acids found), while Italian and
379 Spanish pollen were rich in linoleic acid, at 29.4 % and 37.5%, respectively (Table 2).

380 Free amino acid composition is reported in Table 3, with values expressed in mg g⁻¹ of
381 DBP. The sample from Spain, Italy and Colombia contained 20, 20 and 18 amino acids,
382 respectively, and the most abundant was proline. Arginine was the second most abundant AA
383 constituting on average 15.1, 12.9 and 3.8% of the total amino acids in the Spanish, Italian
384 and Colombian pollen, respectively. Thus, proline and arginine constituted about 68, 72 and

385 82% of the total AA in the Spain, Italy and Colombia pollen (Table 3). One interesting
386 finding was that arginine was the second most predominant amino acid; other bee-collected
387 pollen originating from Poland, South Korea and China showed a high content of glutamic
388 acid, aspartic acid, leucine and lysine (Szczęsna, 2006).

389 The amount of some essential amino acids such as leucine, isoleucine, lysine, and
390 phenylalanine in Spanish and Italian pollen was higher compared to Colombian sample.
391 Tryptophan (0.11-0.15 mg g⁻¹) and cysteine (0.07-0.09 mg g⁻¹) were present in low amount in
392 pollen from Spain and Italy and were not detected in Colombian one. The Spanish pollen was
393 the richest in threonine and valine. Spanish and Italian pollen presented percentages of total
394 free amino acid significantly different ($P < 0.05$) from Colombia sample. The differences in
395 the total and individual amino acid content were probably due to the different botanical
396 composition.

397 Proline (16-20 mg g⁻¹) was the main amino acid in all the sample, reaching
398 approximately a level of the 53, 59 and 78% of the total free AA content in pollen from Spain,
399 Italy and Colombia, respectively.

400 It has been reported that proline is the main free AA in the well dried and stored pollen
401 while glutamic acid is the most abundant in freshly collected pollen. When the temperature is
402 high and the drying process period excessively long, the free AA content decreases (<2 %)
403 and consequently the ratio proline/total free AA (Proline Index) increases (>80 %).
404 Conversely, if the drying process is carried out properly, the free AA content remains high
405 (>2.5 %) and the proline index is less than 80 %.

406 A minimum quantity of 2 % of free AA content is suggested to standardize the
407 commercial honeybee-collected pollen in the European market (Serra Bonvehí and Escolá
408 Jorda, 1997), and the Proline Index can be used as an “indicator” of the pollen freshness. The
409 latter must be less than 80%.

410 3.2. Identification of PA-conjugates by their fragmentation patterns

411

412 Pollen extract represents a complex mixture of variable composition. Thus, an
413 untargeted analysis was performed using an UHPLC coupled to a DAD and Orbitrap-MS.
414 High mass resolution (50 K) and high mass accuracy (2 ppm) allow obtaining the formula of
415 parent and product ions. Due to these features, together with the enhanced efficiency of the
416 UHPLC technique, this system is a powerful tool for the identification of unknown analytes in
417 the pollen extracts. Untargeted analysis, however, cannot be done based on elemental
418 composition data alone. Additional information is required, such as the UV spectrum and
419 fragmentation pattern with collision-induced dissociation (CID) of the parent ion. Given the
420 relatively high proton affinities of amine-N atoms, detection in positive ion mode is initially
421 preferred for these compounds. After this step, analyses were also conducted in the negative
422 ion mode, for the improved elucidation of the HCAs conjugated to the PAs. Examples of the
423 UHPLC profiles of the pollens examined in this study, over the 240–440 nm range, are shown
424 in Figure 1. Table 4 reports the on-line UV spectra, deprotonated ion and fragments of the
425 main flavonoids and PAs, such as spermidine and spermine conjugates to HCAs detected in
426 the DBP. Regarding PAs linked to HCAs, the fragmentation starts with a single cleavage of
427 the terminal amide bond (N^1) of the spermidine and spermine cores, resulting in the formation
428 of di- or tri-substituted fragments, respectively. On the contrary, the cleavage of the amide
429 bond at N^5 for spermidine and N^5 or N^{10} for spermine was obtained at higher collision
430 energies. Thus, the fragmentation depends on the HCA moiety substitution position in the PA
431 core. As an example, Figure 2A displays the fragmentation pattern of the N^1 -caffeoyl- $N^{5,10}$ -di-
432 *p*-coumaroyl-spermidine (m/z 598.2558 u). The main product ion, m/z 462.2036 u,
433 corresponds to the loss of caffeic acid and formation of the isocyanate group, respectively.
434 The simultaneous loss of caffeic acid and *p*-coumaric acid produces the ion with m/z 342.1459

435 u, which contains two isocyanate groups. The fragments at m/z 135.0450 and 119.0500 u have
436 been attributed to decarboxylated caffeic acid and *p*-coumaric acid, respectively. *p*-Coumaric
437 acid also undergoes loss of a water molecule, with the formation of the ion at m/z 145.0292 u.
438 Analogously, $N^{1,10}$ -di-*p*-coumaroyl- N^5 -caffeoyl-spermidine (m/z 598.2558 u) mainly
439 generated fragments at m/z 478.1980 u (Figure 2B), corresponding to the loss of *p*-coumaric
440 acid and formation of the isocyanate group. Instead, the abundance of the ions generated at
441 low-energy CID, by the loss of N^5 -caffeic acid (m/z 462.2036 u) was low. Conversely, they
442 were present at high collision energy (25 eV). Likewise, peak 2 had the formula $C_{25}H_{30}N_3O_4$,
443 $[M-H]^-$ 436.2247 u, which corresponds to $N^{1,10}$ -di-*p*-coumaroyl-spermidine or $N^{1,5}$ -di-*p*-
444 coumaroyl-spermidine. After CID at different energies, the only fragments found had m/z
445 316.1666 and 119.0502 u, corresponding to the loss of *p*-coumaric acid at N^1 (-120.0575).
446 The ions with m/z 290.1874, corresponding to the loss of *p*-coumaric acid at N^5 , were not
447 detected (Figure 3). Overall, PAs linked to HCA easily lose the moiety bound to the primary
448 N and with much more difficulty, which joined to the secondary N. Thus, loss of 120.0575,
449 136.0524, 150.0681, 180.0787 and 166.0630 corresponded to the presence of *p*-coumaric
450 acid, caffeic acid, ferulic acid, syringic acid and 4-methyl-gallic acid residue in position N^1 of
451 spermine or spermidine, respectively.

452

453 3.3. Carotenoid determination

454

455 The total carotenoid content in the saponified pollen samples was determined
456 spectrophotometrically at 450 nm, using the zeaxanthin $\epsilon^{\%}$ and by HPLC analysis by using an
457 authentic standard. The equations of calibration curve for lutein (Lut, 0.14-1.83 $\mu\text{g mL}^{-1}$),
458 zeaxanthin (Zea, 0.11-1.64 $\mu\text{g mL}^{-1}$), β -cryptoxanthin (β -Cry, 0.14-1.69 $\mu\text{g mL}^{-1}$), echinenone

459 (Ech, 0.11-1.34 $\mu\text{g mL}^{-1}$), α -carotene (α -Car, 0.11-1.58 $\mu\text{g mL}^{-1}$) and β -carotene (β -Car,
460 0.16-1.87 $\mu\text{g mL}^{-1}$) were as follows:

461 Lut: $Y = 125.7 X - 2.7$, $R^2=0.999$, $n=5$

462 Zea: $Y = 129.1 X - 1.9$, $R^2=0.999$, $n=5$

463 β -Cry: $Y = 132.2 X - 1.5$, $R^2=0.999$, $n=5$

464 Ech: $Y = 124.1 X - 2.0$, $R^2=0.997$, $n=5$

465 α -Car: $Y = 116.0 X - 2.1$, $R^2=0.998$, $n=5$

466 β -Car: $Y = 101.2 X - 1.5$, $R^2=0.998$, $n=5$

467 Where Y =peak area $\times 10^{-3}$, X = $\mu\text{g/mL}$

468

469 The total carotenoid content, determined by spectrophotometry, was 221.4, 56.9 and 24.7 μg
470 g^{-1} for DBP from Colombia, Spain and Italy, respectively. According to the Student's t -test,
471 there was no significant difference ($p=0.591$) between the methods used.

472 Characterization of the free and esterified carotenoids is necessary to obtain reliable
473 compositional data. Thus, we developed a UHPLC-DAD-ESI-HR-MS method for the
474 identification of the carotenoid composition in pollen from three different countries. Free
475 carotenoids were identified by comparing the retention time (RT), UV-vis spectra and
476 accurate mass (2 ppm) with those of authentic standards. Non-esterified carotenoids were
477 detected in small amounts (< the limit of quantification) in all pollens. Notably, the
478 Colombian pollen contained traces of lutein, zeaxanthin, β -carotene and phytoene, while only
479 β -carotene was present in the Spanish and Italian samples. In contrast, all samples contained
480 esterified carotenoids. Figure 4A and 4B are the chromatograms, integrated over the
481 wavelength range 254–600 nm, relating to non-saponified and saponified extracts from
482 Colombian pollen. The principal peaks found in all samples had m/z 932.7610 and fragment

483 ions at m/z 750.5939 and 532.4072 u, corresponding to the molecular ion $[M]^{•+}$, namely, the
484 loss of one molecule of lauric acid $[M-C_{11}CO]^{•+}$ and two residues of lauric acid and water $[M-$
485 $2C_{11}CO-2H_2O]^{•+}$, respectively. Basic hydrolysis initially produced two peaks, with different
486 intensities, at m/z 750.5939 u and after 2 h the dominant peak in the chromatogram had m/z
487 568.4275 u $[M]^{•+}$ and a lower signal at 550.4178 u, corresponding to the loss of water $[M-$
488 $H_2O]^{•+}$. Based on the RT, UV-vis and MS spectrum, the peak was identified as zeaxanthin.
489 Thus, the primary carotenoid was zeaxanthin esterified with two molecules of lauric acid
490 (C12:0). At the end of the hydrolysis, a second peak, with the same molecular ion as
491 zeaxanthin but a shorter RT, was found. This compound was identified as lutein. Thus,
492 samples contained also small amounts of the di-lauryl ester of lutein. After hydrolysis, the
493 content of carotenoids did not increase.

494

495 *3.4. Multivariate statistical analysis*

496

497 PCA was applied to check for similarity between samples according to their chemical
498 composition and geographical origin. The first two principal components (PC) were able to
499 explain up to 95% of data variability. Bee pollen samples were differentiated in PC1 (64 %)
500 based on the content of total lipid, SFA, MUFA, PUFA, 16:0, 18:1n9, 12:0, 24:1n9, 18:3n6,
501 24:0 22:0, total and single sugars, while PC2 (31 %) separated the samples according to the
502 levels of protein, ash, carotenoid, 18:2n6, 18:3n3, 20:2n6 and 20:3n6.

503 Figure 5 reports two-dimensional plot and show the response variability of the
504 experimental data obtained from the pollen from Colombia, Italy and Spain. It is noted that
505 the three samples from three different geographical regions are largely dispersed within the
506 graph, indicating a high diversity between them. In particular, the sample from Italy has a

507 greater deviation compared to Spain and Colombia on the PC1 axes (the PC with the greater
508 power of descriptiveness of the sample).

509

510 3.5. Palynological analysis

511

512 All pollen samples were found to be heterofloral, having different pollen types and
513 percentages. In the specimen from Spain, *Cistus ladanifer* and *Echium* were the PDs (>45%),
514 followed by *Achillea* and Compositae types T (*Taraxacum*) and S (*Carduus*, *Cirsium*). The
515 POs were *Vicia*, *Quercus ilex*, *Quercus r.*, *Rubus*, Pinaceae, *Filipendula*, *Trifolium*
516 *incarnatum*, *Trifolium pratense*, *Trifolium repens*, *Prunus*, *Pyrus*, *Malus* and *Oxalis*.

517 The three species of *Trifolium* have been distinguished by their size. Indeed, in decreasing
518 order *incarnatum*, *pratense* and white. Furthermore, *incarnatum* has exine with a larger mesh
519 pattern and it is more elongated than the other two. The *pratense* is also cross-linked but with
520 less evident mesh. The white clover is the smallest and with the smoother surface.

521 The PDs in the Italian sample were *Rubus ulmifolius*, *Parthenocissus quinquefolia* and
522 *Ampelopsis brevipedunculata*. The PSs were *Papaver*, *Muscari* and *Lamium*, while *Ambrosia*,
523 *Fraxinus ornus*, *Castanea sativa* and some Gramineae, such as *Cynodon dactylon*, *Phleum*
524 *pratense* and *Poa pratensis* were the POs.

525 Pollen from Colombia contained mainly *Brassica napus*, *Taraxacum officinale* and
526 *Trifolium pratense*. The POs were from Rhamnaceae (*Gouania polygama*), Rosaceae with
527 striated resin (*Prunus sp.*), Fabaceae (*Trifolium repens*, *Vicia sp.*) and *Euphorbia sp.*

528 **4. Conclusion**

529

530 In this study, bee pollen samples from three different geographical areas were analysed
531 for their nutrient contents. Moreover, the botanical origin and the phytochemical profile of
532 each pollen was evaluated. The content of carbohydrate, moisture and ash was not
533 significantly different among the pollens, while the amount of protein was higher in the
534 Colombian and Italian samples compared to the Spanish pollen. Colombian pollen also
535 contained a high amount of n-3 fatty acids relative to the Italian and Spanish pollens. Notably,
536 Colombian pollen was rich in α -linolenic, while the Italian and Spanish samples mainly
537 contained linoleic acid. The high α -linolenic acid and TL content may be attributed to the
538 presence of Brassicaceae, such as *Brassica napus*. Due to the high content of essential fatty
539 acids, pollen load could be used as a dietary supplement. The main polyphenols were
540 coumaroyl- and caffeoyl-spermidine derivatives. Proline and arginine were the main AA in all
541 the pollen samples. Low-energy CID allowed determining which phenolic acid was bound to
542 the N¹ of the PA. All pollen samples contained esterified carotenoids, and the main one was
543 zeaxanthin linked to two residues of lauric acid, while only trace amounts of non-esterified
544 carotenoids, such as lutein, zeaxanthin and β -carotene, were detected. Therefore, the
545 nutritional composition and the high content of bioactive compounds, such as flavonoids,
546 carotenoids and phenolic acids linked to PAs, could make bee pollen a valuable ingredient for
547 the food and pharmaceutical industries.

548

549 **Acknowledgments**

550

551 The authors are grateful to Dr. Gianoncelli Carla of the Fojanini Foundation (Sondrio, I)
552 for the palynological analysis, and Dr. Gargari Giorgio (DeFENS) for the statistical analysis

553 **Funding sources**

554

555 This research did not receive any specific grant from funding agencies in the public,

556 commercial, or not-for-profit sectors.

557

558 **Declaration of interest**

559

560 None.

561 **References**

- 562 Almeida-Muradian, L.B., Pamplona, L.C., Coimbra, S., Barth, O.M. (2005). Chemical
563 composition and botanical evaluation of dried bee pollen pellets. *J. Food Comp. Anal.*
564 *18*, 105–111.
- 565 Aloisi, I., Cai, G., Serafini-Fracassini, D., Del Duca, S. (2016). Polyamines in pollen: from
566 microsporogenesis to fertilization. *Frontiers in Plant Science*, *7*, 1–7.
- 567 AOAC (2005), Official Methods of Analysis of AOAC International, Association Official
568 Analytical Chemists, Rockville, USA.
- 569 Armenta, J.M., Cortes, D.F., Pisciotta, J.M., Shuman, J.L., Blakeslee, K., Rasololon, D.,
570 Ogunbiyi, O., Sullivan, D.J. Jr, Shulaev, V. (2010). Sensitive and rapid method for amino
571 acid quantitation in malaria biological samples using AccQ.Tag ultra performance liquid
572 chromatography-electrospray ionization-MS/MS with multiple reaction monitoring. *Anal.*
573 *Chem.* *82*, 548–558
- 574 Barth, O.M. (2004). Melissopalynology in Brazil: a review of pollen analysis of honeys,
575 propolis and pollen loads of bees. *Scientia Agricola*, *61*, 342–350.
- 576 Bogdanov, S. (2004). Quality and standards of pollen and beeswax. *Apiacta*, *38*, 334–341.
- 577 Bonvehi, J.S., Torrento, M.S., Lorente, E.C. (2001). Evaluation of polyphenolic and flavonoid
578 compounds in honeybee-collected pollen produced in Spain. *J. Agric. Food Chem.* *49*,
579 1848–1853.
- 580 Bunea, A., Socaciu, C., Pinte, A. (2014). Xanthophyll esters in fruits and vegetables. *Notulae*
581 *Botanicae Horti Agrobotanici (Romania)*, *42*, 310–324.
- 582 Campos, M.G.R., Bogdanov, S., de Almeida-Muradian, L.B., Szczesna, T., Mancebo, Y.,
583 Frigerio, C., Ferreira, F. (2008). Pollen composition and standardisation of analytical
584 methods. *J. Apic. Res.* *47*, 154-161.

585 Conte, G., Benelli, G., Serra, A., Signorini, F., Bientinesi, M., Nicolella, C., Mele, M.,
586 Canale, A. (2017). Lipid characterization of chestnut and willow honeybee-collected
587 pollen: Impact of freeze-drying and microwave-assisted drying. *J. Food Comp. Anal.* 55,
588 12–19

589 Conti, I., Medrzycki, P., Argenti, C., Meloni, M., Vecchione, V., Boi, M., Mariotti, M.G.
590 (2016). Sugar and protein content in different monofloral pollens - building a database.
591 *Bull. Insect.* 69,318–320.

592 Denisow, B., Denisow-Pietrzyk, M.(2016). Biological and therapeutic properties of bee
593 pollen: a review. *J. Sci. Food Agric.* 96, 4303–4309.

594 Fuenmayor, C.B., Zuluaga, C.D., Díaz, C.M., Quicazán, M.C, Cosio, M., Mannino, S. (2014).
595 Evaluation of the physicochemical and functional properties of Colombian bee pollen.
596 *Revista MVZ Córdoba (Colombia)*, 19,4003–4014.

597 Gabriele, M., Parri, E., Felicioli, A., Sagona, S., Pozzo, L., Biondi, C., Domenici, V., Pucci,
598 L. (2015). Phytochemical composition and antioxidant activity of Tuscan bee pollen of
599 different botanic origins. *It. J. Food Sci.* 27, 248–259.

600 Grienberger, E., Besseau, S., Geoffroy, P., Debayle, D., Heintz, D., Lapierre, C., Pollet, B.,
601 Heitz, T., Legrand, M. (2009). A BAHD acyltransferase is expressed in the tapetum of
602 *Arabidopsis* anthers and is involved in the synthesis of hydroxycinnamoyl spermidines.
603 *Plant J.* 58, 246–259.

604 Han, L., Liu, X., Yang, N., Li, J., Cai, B., Cheng, S. (2012). Simultaneous chromatographic
605 fingerprinting and quantitative analysis of flavonoids in Pollen Typhae by high-
606 performance capillary electrophoresis. *Acta Pharm. Sin. B* 2, 602–609.

607 Komosinska-Vassev, K., Olczyk, P., Kaźmierczak, J., Mencner, L., Olczyk, K. (2015). Bee
608 pollen: Chemical composition and therapeutic application, *Evid. Based Comp. Alter.*
609 *Med.* 2015, Article ID 297425.

610 LARN. (2014). Lipidi. In *Livelli di assunzione di riferimento di nutrienti ed energia per la*
611 *popolazione italiana* (4th ed) (pp. 108-144). Milan, I: SICS publisher.

612 Lombeida, W.O., Rubio, F., Levy, L.W. (2016). Determination of lutein and zeaxanthin esters
613 and their geometric isomers in carotenoid ester concentrates used as ingredients in
614 nutritional supplements: Validation of a combined spectrophotometric-HPLC method. *J.*
615 *AOAC Int.* 99, 1459-1469

616 Mărgăoan, R., Mărghitaș, L.A., Dezmirean, D.S., Dulf, F.V., Bunea, A., Socaci, S.A, Bobiș,
617 O. (2014). Predominant and secondary pollen botanical origins influence the carotenoid
618 and fatty acid profile in fresh honeybee-collected pollen. *J. Agric. Food Chem.* 62, 6306–
619 6316.

620 Mihajlovic, L., Radosavljevic, J., Burazer, L., Smiljanic, K., Velickovic, T.C. (2015).
621 Composition of polyphenol and polyamide compounds in common ragweed (*Ambrosia*
622 *artemisiifolia* L.) pollen and sub-pollen particles. *Phytochemistry*, 109, 125–132.

623 Milani, A., Basirnejad, M., Shahbazi, S., Bolhassani, A. (2017). Carotenoids: biochemistry,
624 pharmacology and treatment. *Br. J. Pharm.* 174, 1290–1324.

625 Nicolson, S.W. (2011). Bee food: the chemistry and nutritional value of nectar, pollen and
626 mixtures of the two. *African Zoology*, 46, 197–204.

627 Nogueira, C., Iglesias, A., Feás, X., Estevinho, L.M. (2012). Commercial bee pollen with
628 different geographical origins: a comprehensive approach. *Int. J. Mo. Sci.* 13, 11173–
629 11187.

630 Pascoal, A., Rodrigues, S., Teixeira, A., Feás, X., Estevinho, L.M. (2014). Biological
631 activities of commercial bee pollens: antimicrobial, antimutagenic, antioxidant and anti-
632 inflammatory. *Food Chem. Toxicol.* 63, 233–239.

633 Riso, P., Porrini, M. (1997). Determination of carotenoid in vegetable foods and plasma. *Int.*
634 *J. Vit. Nutr. Res.* 67, 47–54.

635 Serra Bonvehí, J., Escolá Jorda, R. (1997). Nutrient Composition and Microbiological Quality
636 of honeybee-Collected Pollen in Spain. *J. Agric. Food Chem.* 45, 725–732.

637 Stanley, R.G., Linskens, H.F. (1974). Pollen: biology, biochemistry, management. In R.G.
638 Stanley, R.G. & Linskens, H.F (Eds.), *Pollen pigments* (pp. 223–246). Berlin: Springer-
639 Verlag.

640 Szczęsna, T. (2006). Long-chain fatty acids composition of honeybee-collected pollen. *J.*
641 *Apic. Sci.* 50, 65–79.

642 Szczęsna, T., Rybak-Chmielewska, H., Chmielewski, W. (2002). Sugar composition of pollen
643 loads harvested at different periods of the beekeeping season. *J. Apic. Sci.* 46, 107–115.

644 Wang, R., Kobayashi, Y., Lin, Y., Rauwald, H.W, Fang, L., Qiao, H., Kuchta, K. (2015). A
645 phytosterol enriched refined extract of *Brassica campestris* L. pollen significantly
646 improves benign prostatic hyperplasia (BPH) in a rat model as compared to the classical
647 TCM pollen preparation Qianlie Kang Pule'an tablets. *Phytomedicine*, 22, 145–152.

648 Yamamoto, A., Nakamura, K., Furukawa, K., Konishi, Y., Ogino, T., Higashiura K., Yago,
649 H., Okamoto, K., Otsuka, M. (2002). A new nonpeptide Tachykinin NK1 receptor
650 antagonist isolated from the plants of compositae. *Chem. Pharm. Bull.* 47, 47-52

651 Yang, K., Wu, D., Ye, X., Liu, D., Chen, J., Sun, P. (2013). Characterization of chemical
652 composition of bee pollen in China. *J. Agric. Food Chem.* 61, 708–718.

653 **Figure legends**

654

655 **Figure 1.** Typical UHPLC chromatograms in the range 240–450 nm of methanolic pollen
656 extracts from Italy (A), Spain (B) and Colombia (C). See Table 4 for peak number
657 assignment.

658

659 **Figure 2.** Main fragmentation patterns of the deprotonated ion with m/z 598.2558 u,
660 corresponding to (A) N^1 -caffeoyl- $N^{5,10}$ -di-*p*-coumaroyl-spermidine and (B) $N^{1,10}$ -di-*p*-
661 coumaroyl- N^5 -caffeoyl-spermidine. The product ions were obtained by high-resolution CID at
662 20, 40 and 60 eV.

663

664 **Figure 3.** Peak 2, $[M-H]^-$ 436.2247 u, identification by fragmentation pattern evaluation.

665

666 **Figure 4.** Typical UHPLC chromatogram in the range 300–550 nm of Colombian pollen
667 extracts not saponified (A) and saponified by methanolic KOH (B).

668 Peak a: zeaxanthin-di-lauryl ester, b: β -carotene, c: phytoene, d: zeaxanthin, e: lutein.

669

670 **Figure 5.** Inter-sample diversity based on nutrient and phytochemical constituents. The PC1
671 and PC2 were responsible for 64 and 31% respectively of the total variance among the sample
672 from Colombia, Italy and Spain.

673 **Table 1** Nutritional components, moisture, ash and carotenoids of pollen load samples from
 674 Colombia, Italy and Spain

675

Analyte	Colombia	Italy	Spain
Moisture (%)	15.5±0.4 ^a	15.1±0.5 ^a	14.9±0.4 ^a
Proteins (%)	21.6±0.5 ^a	19.5±0.4 ^a	12.3±0.2 ^b
Lipids (%)	6.0±0.2 ^a	2.5±0.2 ^b	5.9±0.6 ^a
SFA	34.9±0.7 ^a	30.9±0.4 ^a	33.7±0.6 ^a
MUFA	8.9±1.8 ^a	14.3±2.2 ^b	9.6±1.1 ^a
PUFA	56.2±2.7 ^a	54.8±2.4 ^a	56.7±2.1 ^a
n-3	44.1±5.4 ^b	23.1±2.3 ^a	18.5±1.7 ^a
n-6	11.9±2.7 ^a	31.6±0.2 ^b	38.2±0.2 ^c
UFA/SFA	1.9±0.1 ^a	2.2±0.0 ^a	2.0±0.0 ^a
Ash (%)	2.1±0.2 ^a	1.8±0.2 ^a	1.6±0.2 ^a
Carbohydrate (%)	39.1±1.9 ^a	44.1±1. ^a	37.7±1.4 ^a
Fructose	18.7±1.2 ^a	23.1±1.1 ^a	17.1±1.1 ^a
Glucose	14.4±0.9 ^a	15.9±0.9 ^a	14.1±1.3 ^a
Sucrose	6.0±0.5 ^a	5.1±0.3 ^a	6.2±0.3 ^a
Carotenoids ¹ (µg Zea g ⁻¹ DW)	221.4±10.6 ^a	24.7±1.3 ^b	56.9±1.9 ^c
Carotenoids ² (µg Zea g ⁻¹ DW)	207.4±9.1 ^a	21.3±0.9 ^b	51.3±1.2 ^c

676

677

678 ^{a,b,c} Different superscript letters indicate statistical differences among the bee pollen extracts
679 ($p < 0.05$). Variables were analyzed by one-way ANOVA with pollen as dependent factors.
680 Differences were considered significant at $p \leq 0.05$; post-hoc analysis of differences between
681 treatments was assessed by the Least Significant Difference (LSD) test with $p \leq 0.05$ as the
682 level of statistical significance. Assays were carried out in triplicate and results were
683 expressed as mean values \pm standard deviation. Carotenoids¹: spectrophotometric analysis
684 after saponification; Carotenoids²: LC-DAD analysis after saponification. DW: dry weight;
685 MUFA: monounsaturated fatty acids; PUFA: polyunsaturated fatty acids; SFA: saturated fatty
686 acids; UFA: sum of unsaturated fatty acids (MUFA+PUFA); Zea: zeaxanthin.

687 **Table 2** Percentage composition (mean±S.D.) of long-chain fatty acids (FA) of honeybee-
 688 collected pollen from Colombia (PC), Italy (PI) and Spain (PS).

FA	PC	PI	PS
12:0	1.7±0.0	0.2±0.1	1.7±0.2
14:0	1.7±0.3	0.61±0.1	0.9±0.1
15:0	0.1±0.0	0.2±0.0	0.3±0.0
16:0	25.3±1.4	20.6±0.0	26.8±1.5
17:0	0.2±0.0	0.5±0.1	0.2±0.0
18:0	2.4±0.2	2.5±0.1	1.6±0.1
20:0	1.4±0.2	1.5±0.1	0.6±0.0
22:0	0.9±0.1	2.4±0.2	0.7±0.1
23:0	0.1±0.1	0.3±0.0	0.4±0.0
24:0	0.3±0.3	2.1±0.2	0.4±0.0
18:1n9	7.0±1.9	12.8±1.9	7.4±0.4
16:1n9	0.0±0.1	0.1±0.0	0.1±0.0
16:1n7	0.1±0.0	0.2±0.1	0.2±0.0
18:1n7	0.5±0.0	0.5±0.0	0.8±0.0
20:1n9	0.5±0.2	0.8±0.1	0.7±0.1
24:1n9	0.6±0.2	0.0±0.0	0.5±0.0
18:2n6	10.6±2.4	29.4±1.5	37.5±4.3
20:2n6	1.1±0.2	0.1±0.0	0.2±0.0
18:3n6	0.3±0.4	1.9±1.3	0.1±0.0
20:3n6	0.0±0.0	0.2±0.1	0.5±0.0
18:3n3	44.1±5.4	23.1±2.2	18.4±1.7

689 C16:0 palmitic acid, C18:0 stearic acid, C18:1n9 oleic acid, C18:2n6 linoleic acid, C18:3n3

690 α -linolenic acid

691 **Table 3** Free amino acid content (mg g⁻¹ DBP) in pollen sample from Colombia, Italy and
692 Spain
693

Amino acid	Colombia	Italy	Spain
His	0.68±0.1 ^a	0.49±0.10 ^a	0.59±0.03 ^a
Ser	0.23±0.02 ^a	0.33±0.04 ^b	0.47±0.01 ^c
Arg	0.96±0.08 ^a	3.81±0.55 ^b	4.64±0.05 ^c
Gly	0.08±0.01 ^a	0.14±0.04 ^b	0.25±0.01 ^c
Asp	0.21±0.03 ^a	0.28±0.07 ^b	0.46±0.01 ^b
Aspg	0.29±0.04 ^a	0.43±0.02 ^b	0.49±0.02 ^b
Glu	0.13±0.02 ^a	0.25±0.06 ^b	0.29±0.01 ^b
Glut	0.09±0.01 ^a	0.14±0.02 ^b	0.23±0.01 ^c
Thr	0.10±0.0 ^a	0.15±0.04 ^b	0.31±0.01 ^c
Ala	0.43±0.02 ^a	0.98±0.10 ^b	1.19±0.02 ^c
Pro	19.75±0.72 ^a	17.47±1.14 ^b	16.17±0.06 ^b
Lys	0.39±0.06 ^a	0.81±0.12 ^b	0.87±0.02 ^b
Tyr	0.36±0.03 ^a	0.76±0.29 ^b	0.89±0.04 ^b
Met	0.20±0.02 ^a	0.21±0.03 ^a	0.19±0.01 ^a
Val	0.29±0.03 ^a	0.30±0.08 ^a	0.70±0.01 ^b
Ile	0.18±0.02 ^a	0.53±0.10 ^b	0.64±0.04 ^b
Leu	0.47±0.05 ^a	1.26±0.07 ^b	1.34±0.03 ^b
Phe	0.44±0.04 ^a	0.86±0.09 ^b	0.87±0.02 ^b
Trp	n.d	0.15±0.02 ^a	0.11±0.01 ^a
Cys	n.d	0.09±0.01 ^a	0.07±0.01 ^a
Total AA	25.3±1.0 ^a	29.4±0.7 ^b	30.8±0.2 ^b
(Pro/Total AA)*100	78 ^a	59 ^b	53 ^c

694 Ala, alanine; Arg, arginine; Asp, aspartic acid; Aspg, asparagine; Cys, cysteine, Gly, glycine;
695 Glu, glutamic acid; Glut, glutamine; His, histidine; Ile, isoleucine; Leu, leucine; Lys, lysine;
696 Met, methionine; Phe, phenylalanine; Pro, proline; Ser, serine; Thr, threonine; Trp,
697 tryptophan; Tyr, tyrosine; Val, valine.

698 n.d: not detected (< LOD).

699 ^{a,b,c} Different superscript letters indicate statistical differences among the bee pollen extracts

700 (p<0.05). Variables were analyzed by one-way ANOVA with pollen as dependent factors.

701 Differences were considered significant at p ≤ 0.05; post-hoc analysis of differences between

702 treatments was assessed by the Least Significant Difference (LSD) test with $p \leq 0.05$ as the
703 level of statistical significance.

704 **Table 4** Polyamines linked to hydroxyl-cinnamic acids and flavonoids in pollen sample from
 705 Colombia, Italy and Spain
 706

Peak	RT	UV (nm)	[M-H] ⁻	Fragment	Compound
1	10.8	265, 350	771.1998	609.1463, 314.0434	I-(Glc)-Glc-Ara
2	11.93	296, 306	436.2247	316.1666	S-N ^{1,10} -di- <i>pC</i>
3	12.03	265, 348	609.147	446.0854, 285.0405	L-Glc-Glc
4	12.22	258, 354	625.1417	300.0275	Q-Glc-Glc
5	12.27	258, 354	639.1572	314.0434	I-Glc-Glc
6	13.5	256, 356	595.1314	300.0277, 463.0890	Q-Glc-Ara
7	13.91	265, 346	609.1469	284.0325	K-Glc-Glc
8	14.46	254, 356	609.1468	314.0433, 284.0327	I-Glc-Ara
9	14.94	255, 354	463.0891	300.0277, 271.0614	Q-Glc-Glc
10a	15.35	272, 315	477.1044	314.0434, 299.0197	I-Glc
10b	15.49	265, 346	593.1519	284.0328	K-Glc-Rha
11	15.55	253, 353	623.1623	314.0434	I-Glc-Rha
12	15.68	253, 353	609.1471	315.0514	I-Glc-Ara
13a	16.75	309	685.3249	519.2610, 399.2037, 165.0555	SP-N ¹⁴ - <i>pC</i> -N ^x - <i>pC</i> -N ¹ -4mGA
13b	16.87	265, 346	447.0938	284.0328, 255.0301	K-Glc
14a	16.89	315	715.3357	549.2717, 399.2038, 165.0555	SP-N ¹⁴ -FA-N ^x - <i>pC</i> -N ¹ -4mGA
14b	17.1	253, 353	477.1045	314.0435	I-Glc
15	18.12	245, 377	317.0305	287.0991, 151.0035	M
16	18.94	250, 354	301.9355	151.0035	Q
17	20.34	290	403.1034	271.0612, 151.0035	P-Ara
18	20.62	245, 320	630.2459	494.1932, 468.2139, 358.1409, 372.1617, 304.1668, 135.0451	S-N ^{1,5,10} -tri-CA
19	22.63	243, 309	614.251	478.1981, 452.2193, 358.1409, 161.0243, 135.0450	S-N ^{1,10} -di-CA-N ⁵ - <i>pC</i>
20	23.04	251, 346	285.0409	151.0035, 133.0293, 255.0299	K
21	23.51	290	271.0612	151.0035	P
22a	23.77	243, 320	879.3458	713.2806, 743.2929	SP-N ^{5,10,14} -tri-CA-N ¹ -4mGA
22b	23.81	243, 320	849.3351	713.2830, 577.2304	SP-N ^{1,5,10,14} -tetra-CA
23	24.42	296, 310	598.2563	462.2036, 342.1461, 135.0450, 145,0293, 478.1983	S-N ^{5,10} -di- <i>pC</i> -N ¹ -CA
24	24.71	246, 310	598.2563	478.1983, 358.1409, 161.0242	S-N ^{1,10} -di- <i>pC</i> -N ⁵ -CA
25	25.26	294, 310	582.2615	462.2036, 342.1459, 145.0293, 119.0501	S-N ^{1,5,10} -tri- <i>pC</i>
26	26.57	294, 310	582.2615	462.2036, 342.1459, 145.0293, 119.0501	S-N ^{1,5,10} -tri- <i>pC</i>
27a	26.69		612.2829	492.2140, 462.2037, 372.1564, 342.1459	S-N ^{1,10} -di- <i>pC</i> -N ⁵ -FA
27b	26.74	293, 308	642.2829	492.2139, 522.2242, 372,1563, 466.2344	S-N ^{1,10} -di-FA-N ⁵ - <i>pC</i>

27c	26.81	317,290	672.2932	522.2245, 372.1563, 175.0390, 149.0605, 135.0449	S-N ^{1,5,10} -tri-FA
28	27.49	290, 310	785.3562	665.2983, 545.2405, 145.0293	SP-N ^{1,5,10,14} -tetra-pC
29	27.91	274, 310	785.3562	665.2983, 545.2405, 145.0293	SP-N ^{1,5,10,14} -tetra-pC
30	28.17	290, 310	785.3562	665.2983, 545.2405, 145.0293	SP-N ^{1,5,10,14} -tetra-pC
31	28.44	296, 308	785.3562	665.2983, 545.2405, 145.0293	SP-N ^{1,5,10,14} -tetra-pC
32	29.21	268, 334	537.0814	151.0035, 385.0719, 443.0410	Amentoflavone
33	29.88	267, 333	537.0814	311.2234, 223.1704, 375.0511, 353.1915	bi-Apigenin

707

708 Ara: arabinose; CA: caffeic acid; FA: ferulic acid; Glc: glucose; I: isorhamnetin; K:

709 kaempferol; L: luteolin; M: myricetin; pC: *p*-coumaric acid; P: pinobanksin; Q: quercetin;

710 Rha: rhamnose; RT: retention time (min); S: spermidine; SP: spermine; x: unknown position;

711 4mGA: 4-methyl-gallic acid.

712 **Table 4** Free amino acid content (mg g⁻¹ DP) in pollen sample from Colombia, Italy and
 713 Spain
 714

Amino acid	Colombia	Italy	Spain
His	0.68±0.1 ^a	0.49±0.10 ^a	0.59±0.03 ^a
Ser	0.23±0.02 ^a	0.33±0.04 ^b	0.47±0.01 ^c
Arg	0.96±0.08 ^a	3.81±0.55 ^b	4.64±0.05 ^c
Gly	0.08±0.01 ^a	0.14±0.04 ^b	0.25±0.01 ^c
Asp	0.21±0.03 ^a	0.28±0.07 ^b	0.46±0.01 ^b
Aspg	0.29±0.04 ^a	0.43±0.02 ^b	0.49±0.02 ^b
Glu	0.13±0.02 ^a	0.25±0.06 ^b	0.29±0.01 ^b
Glut	0.09±0.01 ^a	0.14±0.02 ^b	0.23±0.01 ^c
Thr	0.10±0.0 ^a	0.15±0.04 ^b	0.31±0.01 ^c
Ala	0.43±0.02 ^a	0.98±0.10 ^b	1.19±0.02 ^c
Pro	19.75±0.72 ^a	17.47±1.14 ^b	16.17±0.06 ^b
Lys	0.39±0.06 ^a	0.81±0.12 ^b	0.87±0.02 ^b
Tyr	0.36±0.03 ^a	0.76±0.29 ^b	0.89±0.04 ^b
Met	0.20±0.02 ^a	0.21±0.03 ^a	0.19±0.01 ^a
Val	0.29±0.03 ^a	0.30±0.08 ^a	0.70±0.01 ^b
Ile	0.18±0.02 ^a	0.53±0.10 ^b	0.64±0.04 ^b
Leu	0.47±0.05 ^a	1.26±0.07 ^b	1.34±0.03 ^b
Phe	0.44±0.04 ^a	0.86±0.09 ^b	0.87±0.02 ^b
Trp	n.d	0.15±0.02 ^a	0.11±0.01 ^a
Cys	n.d	0.09±0.01 ^a	0.07±0.01 ^a
Total AA	25.3±1.0 ^a	29.4±0.7 ^b	30.8±0.2 ^b
Proline Index	78 ^a	59 ^b	53 ^c

715 Ala, alanine; Arg, arginine; Asp, aspartic acid; Aspg, asparagine; Cys, cysteine, Gly, glycine;
 716 Glu, glutamic acid; Glut, glutamine; His, histidine; Ile, isoleucine; Leu, leucine; Lys, lysine;
 717 Met, methionine; Phe, phenylalanine; Pro, proline; Ser, serine; Thr, threonine; Trp,
 718 tryptophan; Tyr, tyrosine; Val, valine.


719 n.d: not detected (< LOD). Proline Index= (Proline/Total AA) x 100.

720 ^{a,b,c} Different superscript letters indicate statistical differences among the bee pollen extracts
 721 (p<0.05). Variables were analyzed by one-way ANOVA with pollen as dependent factors.

722 Differences were considered significant at p ≤ 0.05; post-hoc analysis of differences between

723 treatments was assessed by the Least Significant Difference (LSD) test with $p \leq 0.05$ as the
724 level of statistical significance.

Figure 1




727

Figure 2

728

729



730


731

732

Figure 3

733

734




735

736

Figure 4

737




738

739

Figure 5

740



741