

1 SYNTHESIS AND BIOLOGICAL EVALUATION OF A TRISACCHARIDE REPEATING UNIT
2 DERIVATIVE OF *STREPTOCOCCUS PNEUMONIAE* 19A CAPSULAR POLYSACCHARIDE

3
4 **Laura Morelli,¹ Silvia Fallarini,² Grazia Lombardi,² Cinzia Colombo,³ Luigi Lay,³ Federica
5 Compostella^{*,1}**

6
7 ¹Dipartimento di Biotecnologie Mediche e Medicina Traslazionale, Università degli Studi di
8 Milano, Via Saldini 50, 20133 Milano, Italy

9 ²Dipartimento di Scienze del Farmaco, Università degli Studi del Piemonte Orientale, Largo
10 Donegani 2, 28100 Novara, Italy

11 ³Dipartimento di Chimica, Università degli Studi di Milano, Via Golgi 19, 20133 Milano, Italy


12 *Corresponding author: E-mail address: federica.compostella@unimi.it

13
14 **Abstract:** *Streptococcus pneumoniae* (SP) is a common human pathogen associated with a broad
15 spectrum of diseases and it is still a leading cause of mortality and morbidity worldwide, especially
16 in children. Moreover, SP is increasingly associated with drug resistance. Vaccination against the
17 pathogen may thus represent an important strategy to overcome its threats to human health. In this
18 context, revealing the molecular determinants of SP immunoreactivity may be relevant for the
19 development of novel molecules with therapeutic perspectives as vaccine components. Serogroup
20 19 comprises the immune-cross reactive types 19F, 19A, 19B and 19C and it accounts for a high
21 percentage of invasive pneumococcal diseases, mainly caused by serotypes 19F and 19A. Herein,
22 we report the synthesis and biological evaluation of an aminopropyl derivative of the trisaccharide
23 repeating unit of SP 19A. We compare two different synthetic strategies, based on different
24 disconnections between the three monosaccharides which make up the final trisaccharide, to define
25 the best approach for the preparation of the trisaccharide. Synthetic accessibility to the trisaccharide
26 repeating unit lays the basis for the development of more complex biopolymer as well as saccharide
27 conjugates. We also evaluate the binding affinity of the trisaccharide for anti-19A and anti-19F sera
28 and discuss the relationship between the chemical properties of the trisaccharide unit and biological
29 activity.

30
31 **1. Introduction**

32 *Streptococcus pneumoniae* (SP) represents a relevant cause of infections associated with high
33 mortality and morbidity: invasive pneumococcal disease (IPD) indeed still shows a high incidence
34 especially in children and in the elderly. Capsular polysaccharides (CPSs) are the primary
35 determinants of the pathogenicity of the bacterium, and account for the classification of SP in more
36 than 90 serotypes.¹ A limited subset of serotypes is responsible for the majority of pneumococcal
37 infections, and representatives of such subsets are contained in commercial licensed vaccines (for
38 example PCV7, Prevnar 7 - Wyeth Pharmaceuticals, contains serotypes 4, 6B, 9V, 14, 18C, 19F and
39 23F). Indeed, capsular polysaccharides (CPSs) are immunogenic, and the generation of type-
40 specific antibodies to CPS is protective.² The pattern of predominant IPD associated serotypes,
41 subjected to a natural fluctuation over time, contains also serotypes of low immunogenicity, such as
42 6, 14, 19 and 23, where low immunogenicity unfortunately does not equate to low virulence,
43 especially in immune-naïve hosts.³ Consequently, a lower vaccination efficacy has been observed
44 for these serotypes.⁴ This is probably not associated to the absolute antibody concentration
45 generated by the vaccine towards each single different serotype, but, more likely, to the increased
46 amount of antibodies required for killing less immunogenic serotypes. Serogroup 19, which
47 comprises the immune-cross reactive types 19F, 19A, 19B and 19C, belongs to this group, and
48 deserves particular attention since it globally accounts for a high percentage of IPD. Serogroup 19
49 IPD are mainly caused by serotypes 19F and 19A, and, in particular, type 19F is one of the most
50 common causes of IPD in children.⁵ The low immunogenicity of this serotype can be explained by
51 the thickness of the 19F capsule and increased resistance to complement deposition, which is the

1 event required to opsonize pneumococci, facilitate phagocytosis and pathogen clearance. Serogroup
 2 19 has also attracted the interest of the research community because it represents one of the most
 3 significant cases to investigate cross-protective immunity. Capsules of serotypes 19F and 19A are
 4 isopolymers, differing only in one glycosidic linkage (glucose to rhamnose, Figure 1). The high
 5 similarity of the two capsular structures suggested the inclusion of only SP 19F in the formulation
 6 of the first glycoconjugate vaccine PCV7, since antibodies to some CPS may cross-react with
 7 related types providing protection against additional types. Indeed, this is what happened for the
 8 vaccine-type 6B, included in PCV7, since 6B-induced antibodies resulted able to cross protect
 9 against the structurally similar 6A CPS, with high effectiveness against 6A disease.⁶ Unfortunately,
 10 antibodies elicited by 19F antigen present in PCV7 provided limited cross-reactive protection
 11 against 19A disease, with the consequence of increasing non-vaccine 19A serotype carriage and
 12 virulence among population in a process defined “serotype replacement”.⁷ Indeed, most of the
 13 PCV7 recipients achieved a significant concentration of antibodies for the vaccine-associated
 14 serotype, but the absence of 19A opsonophagocytic activity indicates that such antibodies are not-
 15 functional against 19A.⁸ The immunogenicity of the 19F vaccine serotype, and the level of cross-
 16 opsonophagocytic antibodies can be influenced by the conjugation method used to connect the
 17 antigenic saccharide fragment to the T-helper peptide, like reductive amination vs cyanylation.⁹ The
 18 lack of antibody-related cross-protection between serotypes 19F and 19A may be alternatively
 19 related to conformational differences between the two CPS structures.¹⁰ Of note, the problem to
 20 induce protection against 19A disease was overcome after the replacement of PCV7 with PCV13,
 21 that contains antigenic CPSs of both serotypes 19A and 19F. Remarkably, a higher level of serotype
 22 19F IgG was found in the sera of patients immunized with PCV13 with respect to PCV7 recipients,
 23 suggesting a contribution of cross-reactive 19A antibodies to the higher 19F opsonophagocytic
 24 activity titers induced by PCV13.⁸



25 **Figure 1:** Structures of serotypes 19F and 19A capsular polysaccharides
 26
 27

28 Molecular approaches investigating the structural and chemical determinants of the cross reactivity
 29 between 19F and 19A serotypes have never been reported. Nonetheless, this knowledge may be
 30 useful to elucidate the mechanism responsible for immunoreactivity. 19F and 19A CPSs are linear
 31 biopolymers made up of trisaccharide repeating units linked through phosphodiester bridges. Each
 32 trisaccharide is composed by a β -D-ManpNAc-(1→4)- α -D-Glcp disaccharide linked to C2 or C3 of
 33 an α -L-Rha unit respectively (Figure 1). In this framework, we report the synthesis of compound **1**,
 34 the trisaccharide repeating unit of SP 19A, functionalized at the reducing end with an aminopropyl
 35 linker, in turn obtained from protected trisaccharide **2** (Figure 2). Our strategy is based on the
 36 development of a new route for the synthesis of an aminopropyl functionalized rhamnosyl acceptor,
 37 compound **3** (Scheme 1). Furthermore, in search of the most straightforward approach towards 19A
 38 trisaccharide, we explored two alternative synthetic strategies, based on different disconnections
 39 between the three monosaccharides which make up the final trisaccharide. In particular,

1 trisaccharide **1** was assembled with higher yields when the α -Glc-(1 \rightarrow 3)-Rha disaccharide was
 2 glycosylated with a glucose moiety, followed by epimerization at C2.
 3 Finally, we evaluated the binding affinity of trisaccharide **1** towards anti-19A and anti-19F sera, to
 4 investigate the role of the carbohydrate portion of the repeating unit in the antibody binding affinity.
 5 Trisaccharide **1** showed a similar and moderate activity towards both sera, indicating that a limited
 6 cross recognition exists at the level of the single repeating unit.





Figure 2: Structures of the target compound **1** and its precursor **2**

10 2. Results and discussion

11 2.1 Chemistry


12 A key point in our synthetic strategy towards compound **1** has been the preparation of protected
 13 trisaccharide **2** as the direct precursor of the target derivative. Compound **2** is a very versatile
 14 molecule, which allows access to both the trisaccharide repeating unit of SP 19A (the goal of this
 15 work), and, in principle, to oligomeric and/or shifted fragments of SP 19A CPS. Elongation at the
 16 upstream residue of the trisaccharide can be performed after selective reductive opening of the
 17 benzylidene group. The functionalization at the reducing end with a 3-aminopropyl linker has been
 18 designed to allow conjugation to carrier proteins¹¹ or the preparation of multivalent systems^{12,13}
 19 appropriate for the *in-vivo* evaluation of the immunogenic activity of 19A CPS-related saccharide
 20 antigens. In this frame of thoughts, we have planned the synthesis of rhamnosyl acceptor **3**, with the
 21 aminopropyl linker already installed,¹⁴ in order to avoid the glycosylation of the aglycon acceptor at
 22 a later stage of the synthetic route.



Scheme 1: Reagents and conditions: a. *N*-Z-3-aminopropanol, $\text{BF}_3 \cdot \text{Et}_2\text{O}$, DCM, 0 °C to rt, 75%; b.
 MeONa, MeOH, 93%; c. TrCl, Py, 60 °C; BnBr, NaH, 64%; CF_3COOH , DCM/MeOH, 90%.

27 To this aim, tetraacetyl rhamnopyranoside **4**¹⁵ was glycosylated with *N*-Z-3-aminopropanol in the
 28 presence of boron trifluoride etherate to give the rhamnose aminopropyl glycoside **5** in 75% yield
 29 (Scheme 1). Zemplen deacetylation afforded deprotected rhamnoside **6** (93%), which was


- 1 regioselectively tritylated at position 3 by treatment with trityl chloride at high temperature, and
 2 then benzylated in 64% yield over two steps. Finally, the trityl group was removed by treatment
 3 with trifluoroacetic acid to give rhamnoside acceptor **3** in 90% yield.



4
5
6
7
8

Scheme 2: Reagents and conditions: a. TESOTf, DCM, - 20 °C, 93%; b. Et₃SiH, BF₃·Et₂O, DCM, 0 °C, *ms*, 60%; c. TMSOTf, DCM, - 20 °C, *ms*, 88%; d. MeONa, MeOH, DCM, 89%; e. Im₂SO₂, NaH, DMF, - 40 °C, 85%; f. NaN₃, DMF, 80 °C, 80%; g. Zn, AcOH/Ac₂O, THF, 62%; h. H₂, Pd(OH)₂, HCl, AcOEt, MeOH, quant.

1
2 Two different disconnection strategies are possible for the construction of the 19A trisaccharide
3 repeating unit (Figure 2), and all the syntheses previously reported are based on a *A-B + C*
4 approach, where a preformed β -ManNAc-(1 \rightarrow 4)-Glc (*A-B*) disaccharide is coupled with a
5 rhamnosyl acceptor (*C*).^{16,17,18,19} Based on our previous experience on the synthesis of the
6 trisaccharide related to SP 19F CPS,^{20,21} we first followed the alternative *B-C + A* pathway in which
7 an α -Glc-(1 \rightarrow 3)-Rha (*B-C*) disaccharide is initially formed in high selectivity, then β -glycosylated
8 with a glucose moiety (*A*) which is finally epimerized to *N*-acetyl-mannosamine.
9 Rhamnosyl acceptor **3** was thus glycosylated at position 3 with 2,3-*O*-benzyl-4,6-*O*-benzylidene
10 glucosyl trichloroacetimidate donor **8**²² under the catalysis of triethylsilyl triflate (Scheme 2). The
11 aminopropyl disaccharide **9** was recovered in excellent yield (93%) and complete α -selectivity.
12 Reductive opening of the benzylidene acetal to the corresponding 6-*O*-benzyl ether was next
13 accomplished by treatment of **9** with triethylsilane in the presence of boron trifluoride-diethyl
14 ether complex to give disaccharide acceptor **10** in good yield.
15 The desired trisaccharide scaffold was obtained through a high yield glycosylation between the 2-
16 *O*-acetyl-3-*O*-benzyl-4,6-*O*-benzylidene glucosyl trichloroacetimidate donor **11**²³ and
17 disaccharide acceptor **10** to give compound **12**. The β -selectivity was guaranteed by the
18 anchimeric assistance offered by the acetyl group at position 2 of glucose **11**. Trisaccharide **12**
19 was finally subjected to the synthetic sequence that allows *gluco* to *manno* epimerization. The
20 acetyl group was initially removed to give unprotected **13** through Zemplen de-acetylation.
21 Compound **13** was then reacted with sulfonyldiimidazole in the presence of sodium hydride to
22 yield saccharide **14**, which was subjected to nucleophilic displacement with sodium azide to
23 give mannoside **15**. The newly established *manno* configuration was confirmed by the broad
24 ¹H-NMR singlet for the anomeric proton of mannose. Finally, the azido group was reduced with
25 Zinc in the presence of acetic acid/acetic anhydride to give the fully protected trisaccharide **2**,
26 which upon hydrogenolysis gave the target trisaccharide **1** in quantitative yield. Overall, the
27 desired trisaccharide **1** was obtained starting from the properly protected monosaccharide
28 donors **8** and **11** and the rhamnosyl acceptor **3** in 18% overall yield over 8 steps.
29 With the goal of developing a solid protocol to trisaccharide **1**, we next planned to test the
30 feasibility of the alternative *A-B + C* disconnection strategy, which offers the advantage to reduce
31 the number of steps on the already formed trisaccharide scaffold. To this aim, we decided to exploit
32 a new synthetic strategy to obtain thio-disaccharide **16** for the glycosylation of the aminopropyl
33 rhamnosyl acceptor **3**. This approach is based on our consolidated protocols for the construction of
34 the β -mannoside linkage (Scheme 3). In this framework, disaccharide **18** was initially formed in a
35 stereoselective fashion through a high yield glycosylation (86%) between phenylthio glucoside **17**²⁴
36 and trichloroacetimidate donor **11**. Next, epimerization at the C2' of disaccharide **18**, and the
37 introduction of the acetamido group gave compound **16** in 35% overall yield over 4 steps. In detail,
38 compound **18** was initially de-acetylated to **19**, then the hydroxy group was activated in high yield
39 as imidazylate and subjected to azide displacement with sodium azide to give mannoside **21**,
40 followed by azide reduction and *N*-acetylation. Glycosylation with rhamnosyl acceptor **3** was
41 promoted using silver triflate-*N*-iodosuccinimide system as previously described,¹⁹ and gave
42 protected trisaccharide **2** in satisfactory yields but low stereoselectivity ($\alpha/\beta = 1:2$). Compound **2**
43 was finally quantitatively deprotected to the target compound **1**. The overall yield of the second
44 synthetic strategy to compound **1**, starting from the suitable building blocks **11**, **17** and **3**, is 6%
45 over 7 steps. In general, this *A-B + C* strategy shows an efficient and easy linear synthesis of the β -
46 mannosylated thioglycosyl-donor **16**, but suffers from moderate yields and low stereoselectivity in
47 the final glycosylation of rhamnose **3**.




1
2 **Scheme 3:** Reagents and conditions: a. **11**, TESOTf, DCM, - 20 °C, *ms*, 86%; b. MeONa, MeOH,
3 DCM, 75%; c. Im₂SO₂, NaH, DMF, - 40 °C, 86%; d. NaN₃, DMF, 80 °C 83%; e. Zn, AcOH/Ac₂O,
4 THF, 66%; f. **3**, AgOTf, NIS, DCM, - 35 °C to - 10 °C, α/β: 60%, α: 20%) g. H₂, Pd(OH)₂, HCl,
5 AcOEt, MeOH, quant.
6


7 2.2 Biology

8 The ability of increasing concentrations (from 10⁻⁷-10⁰ mg/mL) of the newly synthesized
9 trisaccharide to inhibit the binding between 19A polysaccharide coated onto plates (positive
10 control) and the anti 19A rabbit polyclonal antibody was evaluated by competitive ELISA. To
11 evaluate the cross-reactivity against 19F serotype, competitive ELISA was done using native 19F
12 polysaccharide and 19F reference serum. Figure 3 shows the inhibition curves obtained with
13 compound **1**, under evaluation in both systems. The relative efficacy of compound **1** was calculated
14 by measuring the maximum effect elicited in each system, while the concentration that produces
15 50% of the maximum effect (EC₅₀) was taken as indirect index of its relative potency (Figure 3).
16 As expected the natural polysaccharide exhibited higher efficacy (100% inhibition at 10⁻¹ mg/mL)
17 and affinity (EC₅₀ = 9.1x10⁻⁵ mg/mL) than synthesized compound (41% inhibition at 10⁰ mg/mL
18 and EC₅₀ = 1.3 x 10⁻³) confirming that saccharide chain length seems to be important for their
19 biological activity. The low effectiveness of the newly synthesized compound could be related to its
20 relative weak avidity, since short chain lengths saccharide antigens, like a trisaccharide, have
21 decreased strength of antibody-antigen binding. The single repeating unit of 19A polysaccharide
22 displayed inhibitory properties also in 19F system. The trisaccharide was slightly both more
23 effective and potent in 19A than in 19F system (41% and 32% of inhibition for 19A and 19F
24 respectively; EC₅₀ 1.3 x 10⁻³ and 2.7 x 10⁻² for 19A and 19F respectively). These data suggest that
25 differences in structures of the 19A and 19F trisaccharides are almost negligible at the repeating

1 unit level, and a level of cross reactivity exists. It is reasonable to speculate that saccharide
 2 fragments with chain length longer than compound **1**, resulting in more complex structures, would
 3 contain multiple epitopes leading to an increase in specificity for 19A serum and a reduction in
 4 cross-reactivity versus 19F.



Compound	EC ₅₀ ± SEM (mg/mL)	Max inhibition ^a (%)
19A PS	$(9.1 \pm 1.2) \times 10^{-5}$	100
1	$(1.3 \pm 0.02) \times 10^{-3}$	41



Compound	EC ₅₀ ± SEM (mg/mL)	Max inhibition ^a (%)
19F PS	$(1.7 \pm 0.006) \times 10^{-4}$	100
1	$(2.7 \pm 0.3) \times 10^{-2}$	32

^a The maximum inhibition elicited by each compound at 1 mg/ml.

^a The maximum inhibition elicited by each compound at 1 mg/ml.

5 **Figure 3.** Results of the Elisa experiments with compound **1**. Concentration/response curves of
 6 compound **1** on the inhibition of the binding between the 19A (on the left) or 19F (on the right)
 7 native polysaccharides, coated onto the plates, and the anti-19A or anti-19F antibodies, respectively,
 8 were evaluated by a competitive ELISA method.
 9

10 11 3. Conclusions

12 In conclusion, the synthesis of compound **1**, an aminopropyl derivative of the trisaccharide
 13 repeating unit of SP 19A, has been developed exploiting rhamnosyl acceptor **3**, already
 14 functionalized with an aminopropyl linker. We developed a new and more efficient synthetic route
 15 to the rhamnosyl acceptor, which allows to obtain compound **3** in 40% overall yield over four steps.
 16 Two different synthetic strategies were used to build trisaccharide **1**, allowing a direct comparison
 17 among the two protocols. Based on our results, we suggest that the protocol based on the *B-C + A*
 18 strategy is more effective than the *A-B + C* one. The overall yield of assembly was around 20% for
 19 the first protocol, in contrast to the more modest 6% of the second approach, which is limited by the
 20 low selectivity in the glycosylation between disaccharide *A-B* and rhamnoside **3** (*C*). The results
 21 confirmed that the stereoselectivity of the reaction of α -glucosylation is a function of the protecting
 22 groups on glucose, and the use of 4,6-*O*-benzylidene glucosyl donors, protected with no
 23 participating groups at the 2-position, are usually α -selective.²⁵ Indeed, the use of 4,6-*O*-
 24 benzylidene glucosyl donor **8** allowed the formation of the α -product in excellent yield. Overall, the
 25 first approach to trisaccharide **1** is solid and highly reproducible. Furthermore, the protected
 26 trisaccharide **2** is a valuable intermediate for the synthesis of shifted fragments of the CPS of SP
 27 19A: the elongation of the trisaccharide at the upstream residue is functional for the synthesis of
 28 oligomers functionalized at the downstream residue with the aminopropyl linker, useful for
 29 conjugation to proteins or multivalent scaffolds. We have also showed that compound **1**, which
 30 possesses moderate inhibitory activity towards anti-19A antibodies, displays a comparable activity
 31 also towards anti-19F antibodies. This data suggests that the two sera are not capable of
 32 discriminating small differences in the structure of 19F and 19A trisaccharides. Since differences in

1 conformational preferences have been described for the repeating units of SP 19A and 19F,¹⁰ it is
2 reasonable to assume that longer and structured fragments are needed to significantly affect the
3 binding specificity of the antibodies to the saccharide antigens.

4. Experimental Section:

4.1 Synthetic procedures

Standard laboratory procedures were followed to carry out the reactions and to prepare dry solvents.²⁶ Optical rotations were measured with a Perkin-Elmer 241 polarimeter at 20 °C. ¹H and ¹³C NMR spectra were recorded with a Bruker AVANCE-500 spectrometer at a sample temperature of 298 K.²⁷ Mass spectrometric analyses were performed on a Thermo Quest Finnigan LCQ™DECA ion trap mass spectrometer; equipped with a Finnigan ESI interface. High-resolution mass spectra were collected by electrospray ionization (ESI) spectroscopy on a QToF SYNAPT G2Si Mass Spectrometer. NaH was washed with hexane three times prior to use.

4.1.1. Synthesis of *N*-(benzyloxycarbonyl)aminopropyl 2,3,4-tri-*O*-acetyl- α -L-rhamnopyranoside (**5**)
BF₃ · Et₂O (5.0 mL, 39.45 mmol) was slowly added through a dropping funnel to a solution at 0 °C under argon of compound **4** (2.28 g, 6.86 mmol) and N-CBz-aminopropanol (3.59 g, 17.15 mmol) in dry CH₂Cl₂ (70 mL). The reaction was stirred at room temperature, monitored by TLC (hexane/ethyl acetate, 1:1) and appeared to be complete after 12 h. The reaction was washed with saturated NaHCO₃ solution (2 x 100 mL), and the combined aqueous phases extracted with AcOEt (2 x 100 mL). The combined organic layers were dried over Na₂SO₄ and concentrated. Purification by flash chromatography (hexane/AcOEt, 6:4) gave pure **5** (2.48 g, 75 %) as a colorless oil. $[\alpha]_D^{20} = -43.6$ (*c* = 0.5 in chloroform); ¹H NMR (CDCl₃): $\delta = 7.40$ -7.30 (m, 5H, arom.), 5.29 (dd, 1 H, *J*_{2,3} = 3.5 Hz, *J*_{3,4} = 10.0 Hz, H-3), 5.25 (dd, 1 H, *J*_{1,2} = 1.7 Hz, *J*_{2,3} = 3.5 Hz, H-2), 5.13 (s, 2H, CH₂Ph), 5.08 (t, 1 H, *J*_{3,4} = *J*_{4,5} = 10.0 Hz, H-4), 4.95-4.88 (m, 1 H, NH), 4.73 (*br s*, 1 H, H-1), 3.91 – 3.83 (m, 1 H, H-5), 3.81-3.74 (m, 1 H, H-a), 3.54-3.47 (m, 1H, H-a'), 3.37-3.29 (m, 2 H, 2 H-c), 2.17 (s, 3 H, CH₃CO), 2.06 (s, 3 H, CH₃CO), 2.01(s, 3 H, CH₃CO), 1.92-1.80 (m, 2 H, 2 H-b), 1.24 (d, 3 H, *J*_{5,6} = 6.3Hz, 3 H-6); ¹³C NMR (CDCl₃): $\delta = 170.2$ (C=O), 170.0 (C=O), 169.9 (C=O), 156.4 (C=O, Cbz), 136.6 (arom), 128.5-128.1 (5 C arom), 97.5 (C-1), 71.1 (C-4), 69.8 (C-2), 69.1 (C-3), 66.7 (CH₂Ph), 66.5 (C-5), 65.8 (C-a), 38.4 (C-c), 29.6 (C-b), 20.9 (CH₃), 20.8 (CH₃), 20.7 (CH₃), 17.4(C-6). MS (ESI) *m/z* (%): 504.1 (100) [M+Na]⁺. HRMS (ESI): *m/z* calcd for C₂₃H₃₁NO₁₀Na 504.1846 [M+Na]⁺, found 504.1836.

4.1.2. Synthesis of *N*-(benzyloxycarbonyl)aminopropyl α -L-rhamnopyranoside (**6**)

Compound **5** (2.40 g, 4.98 mmol) was dissolved in dry dichloromethane (50 mL) and sodium methoxide in dry methanol (0.2 M solution, 12 mL) was added. The reaction was stirred for 3h at room temperature, then it was neutralized with an ion exchange resin (Dowex 50 × 8, H⁺ form), filtered and concentrated. The crude was subjected to flash chromatography (CH₂Cl₂/MeOH, 9:1) to give compound **6** (1.64 g, 93 %) as a colorless oil. $[\alpha]_D^{20} = -38.5$ (*c* = 0.5 in chloroform) ¹H NMR (MeOD): $\delta = 7.40$ -7.28 (m, 5H, arom.), 5.09 (*br s*, 2H, CH₂Ph), 4.67 (*br s*, 1H, H-1), 3.83-3.80 (m, 1H, H-2), 3.77-3.70 (m, 1H, H-a), 3.66 (dd, 1 H, *J*_{2,3} = 3.3 Hz, *J*_{3,4} = 9.5 Hz, H-3), 3.61-3.55 (m, 1H, H-5), 3.47-3.41 (m, 1H, H-a), 3.38 (t, 1 H, *J*_{3,4} = *J*_{4,5} = 9.5 Hz, H-4), 3.28-3.18 (m, 2H, 2 H-c), 1.83-1.75 (m, 2H, 2 H-b), 1.27 (d, 3 H, *J*_{5,6} = 6.4Hz, 3 H-6); ¹³C NMR (MeOD): $\delta = 157.5$ (C=O), 137.0 (arom), 128.1-127.4 (5 C arom), 100.3 (C-1), 72.6 (C-4), 71.0 (C-3), 70.9 (C-2), 68.4 (C-5), 66.0 (CH₂Ph), 64.5 (C-a), 37.6 (C-c), 29.4 (C-b), 16.6 (C-6). MS (ESI) *m/z* (%): 378.1 (100) [M+Na]⁺, 732.8 (12) [2M+Na]⁺. HRMS (ESI): *m/z* calcd for C₁₇H₂₅NO₇Na 378.1529 [M+Na]⁺, found 378.1526.

1 4.1.3. Synthesis of *N*-benzyl-*N*-benzyloxycarbonyl-3-aminopropyl 2,4-di-*O*-benzyl-3-*O*-trityl- α -*L*-
2 rhamnopyranoside (**7**)

3 A mixture of **6** (1.60 g, 4.50 mmol), trityl chloride (2.51 g, 9.00 mmol) and dry pyridine (15 mL)
4 was stirred at 60 °C for 20 h. After the addition of Et₃N (2 mL), the reaction was diluted with
5 EtOAc (50 mL) and washed with HCl 1N (2 x 50 mL). The combined aqueous phases were
6 extracted with AcOEt (3 x 40 mL), and then the combined organics were washed with satd.
7 NaHCO₃ soln. (1 x 60 mL), dried over Na₂SO₄ and evaporated under reduced pressure. To a
8 solution of the crude and benzyl bromide (3.2 mL, 27 mmol) in dry DMF (50 mL), NaH (60 % in
9 oil, 1.21 g, 31.5 mmol) was added portionwise at 0 °C. The reaction was warmed to room
10 temperature. After 5 h, an additional amount of NaH (60 % in oil, 0.34 g, 9.00 mmol) was added
11 and the reaction stirred for 12 h. The mixture was quenched by carefully addition of MeOH (5 mL),
12 then diluted with HCl 1N (100 mL), and extracted with AcOEt (3 x 100 mL). The combined
13 organics were washed with brine (2 x 150 mL), dried over Na₂SO₄ and evaporated. The crude was
14 purified through flash chromatography (hexane/ AcOEt, 82:25) to give product **7** (2.5 g, 64 %) as a
15 light yellow viscous oil. $[\alpha]_D^{20} = -8.7$ ($c = 0.5$ in chloroform)

16 ¹H NMR (CDCl₃): $\delta = 7.65$ -7.10 (m, 35H, arom.), 5.25-5.00 (m, 3H, CH₂Ph), 4.80-4.65 (m, 1H,
17 CH₂Ph), 4.55-4.35 (m, 3H, H-1 and CH₂Ph), 4.30-4.15 (m, 2H, CH₂Ph), 4.10 (dd, 1H, $J_{2,3} = 2.6$ Hz,
18 $J_{3,4} = 9.2$ Hz, H-3), 3.90-3.70 (m, 1H, H-4), 3.65-3.35 (m, 2H, H-5 and H-a), 3.35-3.05 (m, 3H, H-a
19 and 2 H-c), 2.45-2.25 (m, 1H, H-2), 1.75-1.55 (m, 2H, 2 H-b), 1.33(d, 3H, $J_{5,6} = 6.2$ Hz, 3 H-6); ¹³C
20 NMR (CDCl₃): $\delta = 156.8$ (C=O), 145.1-127.0 (42 C, arom), 97.2 (C-1), 87.4 (C trityl), 80.5 (C-4),
21 77.9 (C-2), 75.3 (CH₂Ph), 73.8 (C-3), 71.9 (CH₂Ph), 69.1 (C-5), 67.2 (CH₂Ph), 65.0 (C-a), 51.0
22 (CH₂Ph), 45.1-44.1 (m, C-c), 28.6-28.0 (m, C-b), 18.4 (C-6); MS (ESI) m/z (%): 890.5 (100)
23 $[M+Na]^+$. HRMS (ESI): m/z calcd for C₅₇H₅₇NO₇Na 890.4033 $[M+Na]^+$, found 890.4029.

24

25 4.1.4. Synthesis of *N*-benzyl-*N*-benzyloxycarbonyl-3-aminopropyl 2,4-di-*O*-benzyl- α -*L*-
26 rhamnopyranoside (**3**)

27 To a solution of compound **7** (0.90 g, 1.04 mmol) in 21 mL of DCM/MeOH (6:1, v/v),
28 trifluoroacetic acid (0.60 mL, 7.88 mmol) was added dropwise. The reaction was stirred at room
29 temperature for 5 h, then quenched to neutrality through addition of TEA. The solvent was
30 evaporated under reduced pressure, and the crude purified by flash chromatography (hexane/
31 AcOEt, 82:25) to give rhamnoside **3** (0.58 g, 90 %) as a colorless oil. $[\alpha]_D^{20} = -13.8$ ($c = 0.1$ in
32 chloroform)

33 ¹H NMR (CDCl₃): $\delta = 7.43$ -7.13 (m, 20 H, arom), 5.23-5.14 (m, 2H, CH₂Ph), 4.92 (d, 1H, $J = 11.1$
34 Hz, CH₂Ph), 4.81-4.70 (m, 2H, H-1 and CH₂Ph), 4.67 (d, 1H, $J = 11.1$ Hz, CH₂Ph), 4.63-4.43 (m,
35 3H, CH₂Ph), 3.97-3.87 (m, 1H, H-3), 3.74-3.17 (m, 3H, H-2,5 and H-a), 3.48-3.25 (m, 4H, H-4, H-a
36 and 2 H-c), 1.87-1.70 (m, 2H, 2 H-b), 1.33 (d, 3H, $J_{5,6} = 6.2$ Hz, 3 H-6); ¹³C NMR (CDCl₃): $\delta =$
37 156.2 (C=O), 138.6-127.3 (24 C, arom), 97.0 (C-1), 82.3 (C-4), 78.6 (C-2), 75.1 (CH₂Ph), 73.0
38 (CH₂Ph), 71.7 (C-3), 67.2 (2C, C-5 and CH₂Ph), 65.0 (C-a), 50.5 and 50.7 (d, NCH₂Ph), 44.5 and
39 43.7 (d, C-c), 28.3 and 27.8 (C-b), 18.0 (C-6); MS (ESI) m/z (%): 684.4 (100) $[M+Na]^+$. HRMS
40 (ESI): m/z calcd for C₃₈H₄₃NO₇Na $[M+Na]^+$ 648.2937, found 648.2936.

41

42 4.1.5. Synthesis of *N*-benzyl-*N*-benzyloxycarbonyl-3-aminopropyl 2,3-di-*O*-benzyl-4,6-*O*-
43 benzylidene- α -*D*-glucopyranosyl-(1→3)-2,4-di-*O*-benzyl- α -*L*-rhamnopyranoside (**9**)

44 A solution of glucosyl trichloroacetimidate **8** (0.70 g, 1.20 mmol) and rhamnoside **3** (0.30 g, 0.48
45 mmol) in DCM (16 mL) was cooled at -20 °C, then triethylsilyl trifluoromethanesulfonate (0.1 M
46 solution in DCM, 0.95 mL) was added dropwise. After 1.5 h the reaction was quenched by the
47 addition of TEA, and allowed to warm to room temperature. The reaction was concentrated, then
48 purified by flash chromatography (hexane/AcOEt, 8:2) to give disaccharide **9** (0.47 g, 93 %) as an
49 oil. $[\alpha]_D^{20} = +3.8$ ($c = 1$ in chloroform). ¹H NMR (CDCl₃): $\delta = 7.50$ -7.14 (m, 35H, arom.), 5.56 (s,
50 1H, CHPh), 5.18 (*br s*, 2H, CH₂Ph), 5.14 (d, 1H, $J_{1',2'} = 3,4$ Hz, H-1'), 4.98-4.93 (m, 2H, CH₂Ph),

1 4.85-4.41 (m, 9H, H-1 and CH₂Ph), 4.21-4.04 (m, 4H, H-3, 6' and 2H_s), 3.90-3.81 (m, 1H, H-2),
2 3.69-3.55 (m, 6H, H-a, 4, 5, 2', 6' and 1H), 3.44-3.22 (m, 3H, 1 H-a and 2 H-c), 1.84-1.69 (m, 2H,
3 2 H-b), 1.31 (*br d*, 3H, 3 H-6). ¹³C NMR (CDCl₃): δ = 163.3 (C=O), 138.6-126.2 (42C, arom.),
4 101.3 (PhCH), 98.1 (C-1), 96.3 (C-1'), 82.6, 80.2, 79.2, 78.5, 76.7(C-3), 75.5 (2C, C-2 and CH₂Ph),
5 75.1 (CH₂Ph), 73.7 (CH₂Ph), 73.2 (CH₂Ph), 69.0 (C-6'), 68.4, 67.2 (CH₂Ph), 65.1 (C-a), 63.0, 50.52
6 and 50.75 (NCH₂Ph), 43.73 and 44.50 (C-c), 27.81 and 28.30 (C-b), 18.0 (C-6). MS (ESI) *m/z* (%):
7 1079.1 (100) [M + 1 + Na]⁺. HRMS (ESI): *m/z* calcd for C₆₅H₆₉NO₁₂Na 1078.4717 [M+Na]⁺, found
8 1078.4712.

9 10 4.1.6. Synthesis of *N*-benzyl-*N*-benzyloxycarbonyl-3-aminopropyl 2,3,6-tri-*O*-benzyl- α -*D*- 11 glucopyranosyl-(1 \rightarrow 3)-2,4-di-*O*-benzyl- α -*L*-rhamnopyranoside (**10**)

12 Compound **9** (0.45 g, 0.43 mmol) and 4 Å m.s. (0.45 g) were dissolved in DCM (10 mL), stirred at
13 room temperature for 15 minutes, then the suspension was cooled at 0 °C. Triethylsilane (0.63 mL,
14 4.30 mmol) was added, followed by the slow dropwise addition of BF₃·Et₂O (0.27 mL, 2.15 mmol).
15 The reaction was stirred at 0 °C for 2 h, then quenched with triethylamine, diluted with DCM,
16 filtered over celite, and concentrated *in vacuo*. The residue was purified by flash chromatography
17 (Hexane/AcOEt, 8:2) to afford compound **10** (0.27 g, 60%) as an oil. [α]_D²⁰ = +10.8 (*c* = 1 in
18 chloroform) ¹H NMR (CDCl₃): δ = 7.40-7.20 (m, 35H, arom), 5.21-5.13 (m, 3H, H-1' and CH₂Ph),
19 5.00-4.80 (m, 2H, CH₂Ph), 4.84-4.83 (m, 11H, H-1 and CH₂Ph), 4.14-4.06 (m, 1H, H-3), 4.04-3.97
20 (m, 1H, H-5'), 3.96-3.80 (m, 2H, H-2, 3'), 3.74-3.45 (m, 7H, H-a, 4, 5, 2', 4', 6'a, 6'b), 3.42-3.19
21 (m, 3H, 2 H-c and 1 H-a), 2.25-2.05 (*br s*, 1H, OH), 1.81-1.65 (m, 2H, 2 H-b), 1.33 (*br s*, 3H, 3 H-
22 6). ¹³C NMR (CDCl₃): δ = 156.4 (C=O), 138.8-127.3 (42C, arom.), 98.2 (C-1), 95.0 (C-1'), 81.3
23 (C-3'), 80.1, 79.4, 76.0 (C-3), 75.5 (C-2), 75.2 (2C, CH₂Ph), 73.4 (CH₂Ph), 73.2 (CH₂Ph), 73.0
24 (CH₂Ph), 71.2, 70.2 (C-5'), 69.5 (C-6'), 68.4, 67.2 (CH₂Ph of Cbz), 65.1 (C-a), 50.5 and 50.8
25 (NCH₂Ph), 43.7 and 44.5 (C-c), 27.8 and 28.3 (C-b), 18.1 (C-6). MS (ESI) *m/z* (%): 1080.1 (100)
26 [M+Na]⁺. HRMS (ESI): *m/z* calcd for C₆₅H₇₁NO₁₂Na 1080.4874 [M+Na]⁺, found 1080.4883.

27 28 4.1.7. Synthesis of *N*-benzyl-*N*-benzyloxycarbonyl-3-aminopropyl 2-*O*-acetyl-3-*O*-benzyl-4,6-*O*- 29 benzylidene- β -*D*-glucopyranosyl-(1 \rightarrow 4)-2,3,6-tri-*O*-benzyl- α -*D*-glucopyranosyl-(1 \rightarrow 3)-2,4-di-*O*- 30 benzyl- α -*L*-rhamnopyranoside (**12**)

31 A suspension of 2-*O*-acetyl-glucosyl trichloroacetimidate **11** (0.42 g, 0.78 mmol), disaccharide **10**
32 (0.23 g, 0.22 mmol) and 4 Å m.s. (0.23 g) in DCM (7 mL) was stirred for 0.15 min at room
33 temperature, then cooled at -20 °C. Triethylsilyl trifluoromethanesulfonate (0.1 M solution in
34 DCM, 0.44 mL) was added dropwise and the disappearance of the starting material was followed by
35 TLC (Toluene/Acetone, 7:3; hexane/AcOEt, 7:3). After 1.5 h, the reaction was quenched with
36 triethylamine, diluted with DCM, filtered over Celite, and the solvent evaporated. The crude
37 product was purified by flash chromatography (hexane/AcOEt, 8:2) to give **12** (0.28 g, 88%) as an
38 amorphous solid. [α]_D²⁰ = +5.9 (*c* = 1 in chloroform). ¹H NMR (CDCl₃): δ = 7.53-7.10 (m, 45H,
39 arom.), 5.48 (s, 1H, PhCH), 5.23-5.13 (m, 3H, H-1' and CH₂Ph), 4.97-4.82 (m, 4H, H-2'' and
40 CH₂Ph), 4.76 (d, 2H, CH₂Ph), 4.72-4.53 (m, 7H, H-1 and CH₂Ph), 4.52-4.44 (m, 3H, H-1'' and
41 NCH₂Ph), 4.29-4.22 (m, 1H, CH₂Ph), 4.17-4.11 (m, 1H, H-6a''), 4.08-3.99 (m, 1H, H-3), 3.99-3.91
42 (m, 3H, H-3', 4', 5'), 3.91-3.82 (m, 1H, H-2), 3.74-3.53 (m, 6H, H-a, 4, 5, 2', 6a', 4''), 3.51-3.36
43 (m, 3H, H-6b', 3'', 6b''), 3.35-3.21 (m, 3H, 1 H-a and 2 H-c), 3.18-3.10 (m, 1H, H-5''), 1.82 (s,
44 3H, COCH₃), 1.81-1.67 (m, 2H, 2 H-b), 1.22 (*br d*, 3H, 3 H-6). ¹³C NMR (CDCl₃): δ = 168.9
45 (C=O), 139.3-126.0 (54C, arom.), 101.1 (CHPh), 100.8 (C-1''), 98.2 (C-1), 96.5 and 96.3 (C-1'),
46 81.6 (C-4''), 80.1, 79.9, 79.2, 78.7, 77.4 (C-3), 76.7, 76.1 (C-2), 75.0 (CH₂Ph), 74.9 (CH₂Ph), 74.0
47 (CH₂Ph), 73.6 (CH₂Ph), 73.3 (2C, C-2'' and CH₂Ph), 73.2 (CH₂Ph), 70.7, 68.6 (C-6''), 68.2, 67.6
48 (C-6'), 67.2 (CH₂Ph), 65.9 (C-5''), 65.1 (C-a), 50.8 and 50.5 (NCH₂Ph), 44.6 and 43.7 (C-c), 28.3
49 and 27.9 (C-b), 20.8 (CH₃CO), 18.0 (C-6). MS (ESI) *m/z* (%): 1463.5 (100) [M + 1 + Na]⁺. HRMS
50 (ESI): *m/z* calcd for C₈₇H₉₃NO₁₈Na 1462.6290 [M+Na]⁺, found 1462.6276.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

4.1.8. Synthesis of *N*-benzyl-*N*-benzyloxycarbonyl-3-aminopropyl 3-*O*-benzyl-4,6-*O*-benzylidene- β -*D*-glucopyranosyl-(1 \rightarrow 4)-2,3,6-tri-*O*-benzyl- α -*D*-glucopyranosyl-(1 \rightarrow 3)-2,4-di-*O*-benzyl- α -*L*-rhamnopyranoside (**13**)

To a stirred solution of **12** (0.27 g, 0.19 mmol) in DCM/MeOH 1:1 (6 mL) sodium methoxide in methanol (1 M solution, 0.19 mL) was added. The reaction was stirred for 48 h at room temperature, then it was neutralized with an ion exchange resin (Dowex 50 \times 8, H⁺ form), filtered and concentrated. The crude product was subjected to flash chromatography (hexane/AcOEt, 7:3) to give pure **13** (0.24 g, 89%) as an amorphous solid. $[\alpha]_D^{20} = +15.3$ ($c = 1$ in chloroform). ¹H NMR (CDCl₃): $\delta = 7.52$ -7.16 (m, 45H, arom.), 5.48 (s, 1H, PhCH), 5.24-5.11 (m, 2H, H-1' and CH₂Ph), 5.00-4.61 (m, 10H, H-1 and CH₂Ph), 4.61-4.41 (m, 4H, CH₂Ph), 4.37 (d, 1H, J_{1'',2''} = 7.5 Hz, H-1''), 4.33-4.28 (m, 1H, CH₂Ph), 4.12-3.95 (m, 5H, H-3, 3', 4', 5', 6a''), 3.89-3.80 (m, 1H, H-2), 3.80-3.74 (m, 1H, H-6a'), 3.74-3.58 (m, 5H, H-a, 4, 5, 2'), 3.58-3.51 (t, 1H, J_{3'',4''} = J_{4'',5''} = 9.3 Hz, H-4''), 3.51-3.37 (m, 3H, H-6b', 3'', 6b''), 3.37-3.21 (m, 4H, H-a, 2'' and 2 H-c), 3.11-3.04 (m, 1H, H-5''), 1.85-1.60 (m, 3H, 2 H-b and OH), 1.35 (d, 3H, J = 5.7 Hz, 3 H-6). ¹³C NMR (CDCl₃): $\delta = 157.7$ (C=O), 128.4-126.0 (54C, arom), 103.4 (C-1''), 101.2 (CHPh), 98.3 (C-1), 94.5 (*br s*, C-1'), 81.2 (C-4''), 80.6, 80.3 (C-3''), 80.0, 79.1 (C-2'), 77.4, 76.1 (C-3), 75.2 (2C, C-2, 2''), 75.1-73.3 (6C, CH₂Ph), 70.0 (C-5'), 68.7 (C-6''), 68.4, 68.2 (C-6'), 67.2 (CH₂Ph), 66.1 (C-5''), 65.2 (C-a), 50.6 (*br s*, NCH₂Ph), 44.5 and 44.4 (C-c), 27.9 and 27.6 (C-b), 18.0 (C-6). MS (ESI) m/z (%): 1420.7 (100) [M+NaM+Na]⁺. HRMS (ESI): m/z calcd for C₈₅H₉₁NO₁₇Na 1420.6185 [M+Na]⁺, found 1420.6194.

4.1.9. Synthesis of *N*-benzyl-*N*-benzyloxycarbonyl-3-aminopropyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-*O*-(*N*-imidazole-1-sulfonyl)- β -*D*-glucopyranosyl-(1 \rightarrow 4)-2,3,6-tri-*O*-benzyl- α -*D*-glucopyranosyl-(1 \rightarrow 3)-2,4-di-*O*-benzyl- α -*L*-rhamnopyranoside (**14**)

NaH (60 % in oil, 0.070 g, 1.76 mmol) was added to a stirred solution of compound **13** (0.23 g, 0.16 mmol) in dry DMF (3.5 mL) at room temperature. After 1 h, the suspension was cooled at -40 °C and 1,1'-sulfonyl-diimidazole (0.22 g, 1.12 mmol) in dry DMF (1.5 mL) was added. After 24 h the reaction mixture was quenched with MeOH and allowed to warm to room temperature, then diluted with water (40 mL). The mixture was extracted with AcOEt (3 x 50 mL). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and evaporated. Flash chromatography (hexane/AcOEt, 7:3) of the crude product gave trisaccharide **14** (0.21 g, 85%) as an amorphous solid. $[\alpha]_D^{20} = +1.8$ ($c = 1$ in chloroform). ¹H NMR (CDCl₃): $\delta = 7.94$ (*br s*, 1H, Im), 7.53-7.10 (m, 45H, arom), 7.00 (m, 2H, Im), 5.46 (s, 1H, PhCH), 5.25-5.12 (m, 3H, H-1' and CH₂Ph), 4.97 (d, 1H, J = 11.1 Hz, CH₂Ph), 4.85-4.58 (m, 10H, H-1 and CH₂Ph), 4.58-4.44 (m, 4H, H-2'' and CH₂Ph), 4.32 (d, 1H, J_{1'',2''} = 7.9 Hz, H-1''), 4.23 (dd, 1H, J_{5'',6''} = 4.9 Hz, J_{6a'',6b''} = 10.6 Hz, H-6a''), 4.16-4.09 (m, 1H, CH₂Ph), 4.09-3.80 (m, 5H, H-2, 3, 3', 4', 5'), 3.75-3.52 (m, 5H, H-a, 4, 5, 2', 4''), 3.52-3.20 (m, 7H, H-a', c, c', 6a', 6b', 3'', 6b''), 3.10-2.98 (m, 1H, H-5''), 1.91-1.69 (m, 2H, 2 H-b), 1.29-1.21 (*br d*, 3H, 3 H-6). ¹³C NMR (CDCl₃): $\delta = 155.6$ (C=O), 139.1-136.5 (9C, arom), 136.8 (C Im), 129.2-126.0 (46C, arom.), 118.6 (C Im), 101.4 (CHPh), 98.6 (C-1''), 98.2 (C-1), 97.1 (*br s*, C-1'), 85.7 (C-2''), 81.9 (C-4''), 80.3 (*br s*, C-4), 79.5 (C-3'), 79.2 (C-2'), 78.4 (*br s*, C-3), 76.7 (C-3''), 76.5 (*br s*, C-2), 76.2 (C-4'), 75.2 (2C, CH₂Ph), 74.6 (*br s*, CH₂Ph), 74.3 (CH₂Ph), 73.6 (CH₂Ph), 73.1 (*br s*, CH₂Ph), 70.2 (C-5'), 68.4 (C-6''), 68.3 (C-5), 67.3 (C-6'), 67.2 (CH₂Ph), 65.7 (C-5''), 65.2 (C-a), 50.8 and 50.5 (NCH₂Ph), 44.6 and 43.7 (C-c), 28.4 and 27.9 (C-b), 18.0 (C-6). MS (ESI) m/z (%): 1550.3 (100) [M+NaM+Na]⁺. HRMS (ESI): m/z calcd for C₈₈H₉₃N₃O₁₉NaS 1550.6022 [M+Na]⁺, found 1550.6055.

4.1.10. Synthesis of *N*-benzyl-*N*-benzyloxycarbonyl-3-aminopropyl 2-azido-3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- β -*D*-mannopyranosyl-(1 \rightarrow 4)-2,3,6-tri-*O*-benzyl- α -*D*-glucopyranosyl-(1 \rightarrow 3)-2,4-di-*O*-benzyl- α -*L*-rhamnopyranoside (**15**)

1 To a stirred solution of **14** (0.20 g, 0.13 mmol) in dry DMF (4 mL), sodium azide (0.085g, 1.30
2 mmol) was added and the resulting solution was heated at 80 °C. After 5 h, the reaction was cooled
3 to room temperature, diluted with H₂O and extracted with AcOEt (3 x 40 mL). The combined
4 organic layers were dried over Na₂SO₄, filtered and evaporated. The crude product was purified by
5 flash chromatography (hexane/AcOEt, 75:25) to give compound **15** (0.15g, 80%) as colourless oil
6 $[\alpha]_D^{20} = -4.1$ ($c = 1$ in chloroform). ¹H NMR (CDCl₃): $\delta = 7.63$ -7.08 (m, 45H, arom.), 5.52 (s, 1H,
7 PhCH), 5.25-5.16 (m, 2H, CH₂Ph), 5.13 (d, 1H, $J_{1',2'} = 3.4$ Hz, H-1'), 5.06 (d, 1H, $J = 10.5$ Hz,
8 CH₂Ph), 4.92 (d, 1H, $J = 11.5$ Hz, CH₂Ph), 4.88-4.82 (m, 2H, CH₂Ph), 4.82-4.75 (m, 2H, CH₂Ph),
9 4.74-4.45 (m, 8H, H-1 and CH₂Ph), 4.34 (*br s*, 1H, H-1''), 4.19-3.95 (m, 6H, H-3, 3', 4', 5', 6a''
10 and 1H x CH₂Ph), 3.88 (t, 1H, $J_{3'',4''} = J_{4'',5''} = 9.4$ Hz, H-4''), 3.85-3.76 (m, 1H, H-2), 3.75-3.58 (m,
11 4H, H-a, 4, 5, 2'), 3.56-3.48 (m, 2H, H-2'', 6b''), 3.48-3.20 (m, 6H, H-a', 6a', 6b', 3'' and 2 H-c),
12 3.00-2.92 (m, 1H, H-5''), 1.88-1.72 (m, 2H, 2 H-b), 1.33 (d, 3H, $J_{5,6} = 5.8$ Hz, 3 H-6). ¹³C NMR
13 (CDCl₃): $\delta = 156.6$ and 156.1 (C=O), 137.9-126.0 (54C, arom.), 101.5 (CHPh), 99.7 (C-1''), 98.4
14 (C-1), 95.5 (*br s*, C-1'), 80.5 (C-3'), 79.8 (C-4), 79.0 (C-2'), 78.5 (C-4''), 76.9 (C-3), 76.7 (C-4'),
15 76.4 (C-3''), 75.9 (C-2), 75.1 (CH₂Ph), 74.7 (CH₂Ph), 73.7 (CH₂Ph), 73.6 (CH₂Ph), 73.5 (CH₂Ph),
16 72.5 (CH₂Ph), 69.7 (C-5'), 68.6 (C-5), 68.4 (C-6''), 68.2 (C-6'), 67.2 (CH₂Ph), 67.1 (C-5''), 65.2
17 (C-a), 63.2 (C-2''), 50.8 and 50.6 (NCH₂Ph), 44.6 and 43.7 (C-c), 28.3 and 27.9 (C-b), 18.0 (C-6).
18 MS (ESI) m/z (%): 1446.4 (100) [M + 1 + Na]⁺. HRMS (ESI): m/z calcd for C₈₅H₉₀N₄O₁₆Na
19 1445.6250 [M+Na]⁺, found 1445.6246.

20

21 *4.1.11. Synthesis of phenyl 2-O-acetyl-3-O-benzyl-4,6-O-benzylidene-β-D-glucopyranosyl-(1→4)-*
22 *2,3,6-tri-O-benzyl-1-thio-β-D-glucopyranoside (18)*

23 Glucosyl trichloroacetimidate **11** (0.31 g, 0.57 mmol), phenylthio glucoside **17** (0.20 g, 0.37 mmol)
24 and 4Å molecular sieves (0.20 g) were diluted in DCM (4 mL). The suspension was cooled at -20
25 °C, then triethylsilyl trifluoromethanesulfonate (0.1 M solution in DCM, 0.74 mL) was added
26 dropwise. The reaction was monitored by TLC (toluene/acetone, 9:1). After 1 h the reaction was
27 quenched by the addition of TEA, diluted with AcOEt, and filtered over a Celite pad. After
28 evaporation of the solvent, then crude was purified by flash chromatography (hexane/AcOEt, from
29 8:2 to 7:3) to give disaccharide **18** (0.29 g, 86 %) as an amorphous white solid. $[\alpha]_D^{20} = +20.6$ ($c = 1$
30 in chloroform). ¹H NMR (CDCl₃): $\delta = 7.65$ -7.20 (m, 30H, arom), 5.50 (s, 1H, PhCH), 5.02-4.95 (m,
31 2H, H- 2' and CH₂Ph), 4.92-4.70 (m, 5H, CH₂Ph), 4.68-4.62 (m, 3H, H-1,1' and CH₂Ph), 4.52 (d,
32 1H, $J = 12.0$ Hz, CH₂Ph), 4.15 (dd, 1H, $J_{5',6'a} = 5.0$ Hz, $J_{6'a,6'b} = 10.5$ Hz, H-6'a), 3.98 (t, 1H, $J_{3,4} =$
33 $J_{4,5} = 9.4$ Hz, H-4), 3.79-3.77 (m, 2H, 2 H-6), 3.68 (t, 1H, $J_{3',4'} = J_{4',5'} = 9.3$ Hz, H-4'), 3.65-3.56 (m,
34 2H, H-3, 3'), 3.52-3.43 (m, 2H, H-2, 6'b), 3.38(dt, 1H, $J_{4,5} = 9.4$ Hz, $J_{5,6} = 2.6$ Hz, H-5), 3.21 (dt,
35 1H, $J_{4',5'} = 9.3$ Hz, $J_{5',6'a} = 5.0$ Hz, $J_{5',6'b} = 9.8$ Hz, H-5'), 1.97 (s, 3H, CH₃). ¹³C NMR (CDCl₃): $\delta =$
36 169.1 (C=O), 138.2-126.0 (36C, arom.), 101.2 (CHPh), 100.8 (C-1'), 87.4 (C-1), 84.7 (C-3), 81.7
37 (C-4'), 80.2 (C-2), 79.0 (C-5), 78.6 (C-3'), 76.7 (C-4), 75.5 (CH₂Ph), 75.4 (CH₂Ph), 74.1 (CH₂Ph),
38 73.6 (CH₂Ph), 73.3 (C-2'), 68.6 (C-6'), 67.9 (C-6), 66.1 (C-5'), 20.9(CH₃). MS (ESI) m/z (%): 947.6
39 (100) [M+Na]⁺. HRMS (ESI): m/z calcd for C₅₅H₅₆O₁₁NaS 947.3441 [M+Na]⁺, found 947.3439.

40

41 *4.1.12. Synthesis of phenyl 3-O-benzyl-4,6-O-benzylidene-β-D-glucopyranosyl-(1→4)-2,3,6-tri-O-*
42 *benzyl-1-thio-β-D-glucopyranoside (19)*

43 To a stirred solution of **18** (0.28 g, 0.30 mmol) in DCM (6 mL) sodium methoxide in methanol (0.1
44 M solution, 0.60 mL) was added. The reaction was stirred for 17 h at room temperature, then it was
45 neutralized with an ion exchange resin (Dowex 50 × 8, H⁺ form), filtered and concentrated. The
46 crude product was subjected to flash chromatography (hexane/AcOEt, 8:2) to give pure **19** (0.20 g,
47 75%) as an amorphous white solid. $[\alpha]_D^{20} = +0.20$ ($c = 1$ in chloroform). ¹H NMR (CDCl₃): $\delta =$
48 7.60 -7.27 (m, 30H, arom.), 5.49 (s, 1H, CHPh), 5.33 (s, 1H, OH), 4.98-4.94 (m, 2H, CH₂Ph), 4.87-
49 4.78 (m, 3H, CH₂Ph), 4.74-4.60 (m, 5H, H-1,1' and CH₂Ph), 4.10-4.01 (m, 2H, H-3, 6a), 3.97 (dd,
50 1H, $J_{5',6'a} = 5$ Hz, $J_{6'a,6'b} = 10.4$ Hz, H-6'a), 3.87-3.84 (m, 1H, H-6b), 3.68 (dd, 1H, $J_{3,4} = J_{4,5} = 8.8$ Hz,

1 H-4), 3.63-3.48 (m, 6H, H-2, 4, 5, 2', 3', 6'), 3.18-3.12 (m, 1H, H-5'). ¹³C NMR (CDCl₃): δ =
2 138.8-126.0 (36 C, arom.), 103.6 (C-1'), 101.2 (CHPh), 87.5 (C-1), 85.5 (C-4), 81.3, 80.5, 80.4,
3 78.8, 77.0 (C-3), 75.5, 75.3 (2 C, CH₂Ph), 74.6 (CH₂Ph), 73.6 (CH₂Ph), 68.6 (C-6'), 68.5 (C-6),
4 66.4 (C-5'). MS (ESI) 905.3 (100) [M+Na]⁺, 1787.9 (40) [2M+Na]⁺. HRMS (ESI): *m/z* calcd for
5 C₅₃H₅₄O₁₀NaS 905.3335 [M+Na]⁺, found 905.3331.

6

7 *4.1.13. Synthesis of phenyl 3-O-benzyl-4,6-O-benzylidene-2-O-(N-imidazole-1-sulfonyl)-β-D-*
8 *glucopyranosyl-(1→4)-2,3,6-tri-O-benzyl-1-thio-β-D-glucopyranoside (20)*

9 NaH (60 % in oil, 0.13 g, 3.30 mmol) was added to a stirred solution of compound **19** (0.19 g, 0.22
10 mmol) in dry DMF (6 mL) at room temperature. After 1 h, the suspension was cooled at -40 °C and
11 1,1'-sulfonyl-diimidazole (0.44 g, 2.20 mmol) in dry DMF (3 mL) was added. After 2 h the reaction
12 mixture was quenched with MeOH and allowed to warm to room temperature, then diluted with
13 AcOEt (40 mL) and washed with brine (2 x 30 mL). The organic layers were dried over Na₂SO₄,
14 filtered and evaporated. Flash chromatography (hexane/AcOEt, 8:2) of the crude product gave
15 compound **20** (0.19 g, 86%) as a foamy solid. [α]_D²⁰ = -10.6 (*c* = 1 in chloroform). ¹H NMR
16 (CDCl₃): δ = 7.89 (s, 1H, H imidazole), 7.59-7.22 (m, 31H, arom.), 7.03 (s, 1H, H imidazole), 5.49
17 (s, 1H, CHPh), 4.89-4.75 (m, 6H, CH₂Ph), 4.64-4.58 (m, 3H, H-1, 1' and 1H of CH₂Ph), 4.52 (t, 1H,
18 J_{1',2'} = J_{2',3'} = 8.7 Hz, H-2'), 4.42 (d, 1H, J = 11.8 Hz, 1H of CH₂Ph), 4.26 (dd, 1H, J_{5,6'a} = 5.0 Hz, J_{6'a,6'b}
19 = 10.5 Hz, H-6'a), 4.06 (t, 1H, J_{3,4} = J_{4,5} = 9.5 Hz, H-4), 3.70-3.44 (m, 7H, H-2, 3, 6a, 6b, 3', 4',
20 6'b), 3.18-3.07 (m, 2H, H-5, 5'). ¹³C NMR (CDCl₃): δ = 138.7-126.0 (38 C, 2 C imidazole and 36 C
21 arom), 118.5 (C imidazole), 101.5 (CHPh), 98.4 (C-1'), 87.5 (C-1), 85.5 (C-2'), 84.1, 81.9, 80.3,
22 78.1 (C-5), 77.0, 75.6, 75.5 (CH₂Ph), 75.4 (CH₂Ph), 74.5 (CH₂Ph), 73.6 (CH₂Ph), 68.4 (C-6'), 67.6
23 (C-6), 65.9 (C-5'). MS (ESI) *m/z* (%): 1035.2 (100) [M+Na]⁺. HRMS (ESI): *m/z* calcd for
24 C₅₆H₅₆N₂O₁₂NaS₂ 1035.3172 [M+Na]⁺, found 1035.3165.

25

26 *4.1.14. Synthesis of phenyl 2-azido-3-O-benzyl-4,6-O-benzylidene-2-deoxy-β-D-glucopyranosyl-*
27 *(1→4)-2,3,6-tri-O-benzyl-1-thio-β-D-glucopyranoside (21)*

28 To a stirred solution of **20** (0.18 g, 0.18 mmol) in dry DMF (3.5 mL), sodium azide (0.12 g, 1.80
29 mmol) was added and the resulting solution was heated at 85 °C. After 4 h, the reaction was cooled
30 to room temperature, diluted with brine (30 mL) and extracted with AcOEt (3 x 20 mL). The
31 combined organic layers were dried over Na₂SO₄, filtered and evaporated. The crude product was
32 purified by flash chromatography (hexane/AcOEt, 75:25) to give compound **21** (0.13 g, 83%) as a
33 foamy white solid. [α]_D²⁰ = -23.9 (*c* = 1 in chloroform). ¹H NMR (CDCl₃): δ = 7.60-7.28 (m, 30H,
34 arom.), 5.53 (s, 1H, CHPh), 5.01 (d, 1H, J = 10.5 Hz, 1H of CH₂Ph), 4.88-4.76 (m, 4H, CH₂Ph),
35 4.73-4.70 (m, 2H, H-1' and 1H of CH₂Ph), 4.68-4.64 (m, 2H, H-1 and 1H of CH₂Ph), 4.50 (d, 1H, J
36 = 10.5 Hz, 1H of CH₂Ph), 4.05-3.99 (m, 2H, H-3, 6'a), 3.94 (t, 1H, J_{3',4'} = J_{4',5'} = 9.5 Hz, H-4'), 3.86
37 (dd, 1H, J_{1',2'} = 1.1 Hz, J_{2',3'} = 3.6 Hz, H-2'), 3.81-3.78 (m, 2H, 2 H-6), 3.72 (t, 1H, J_{3,4} = J_{4,5} = 8.9 Hz,
38 H-3), 3.58-3.49 (m, 4H, H-2, 5, 3', 6'b), 3.09-3.04 (m, 1H, H-5'). ¹³C NMR (CDCl₃): δ = 138.8-
39 126.0 (36C, arom.), 101.5 (CHPh), 100.3 (C-1'), 87.5 (C-1), 85.0 (C-4), 80.3, 78.5, 76.5 (C-4'),
40 77.4 (C-3), 76.7, 75.5 (2C, CH₂Ph), 73.7 (CH₂Ph), 72.8 (CH₂Ph), 68.8 (C-6), 68.3 (C-6'), 67.3 (C-
41 5'), 63.6 (C-2'). MS (ESI) *m/z* (%): 930.3 (100) [M+Na]⁺, 1837.5 (20) [2M+Na]⁺. HRMS (ESI): *m/z*
42 calcd for C₅₃H₅₃N₃O₉NaS 930.3400 [M+Na]⁺, found 930.3403.

43

44 *4.1.15. Synthesis of phenyl 2-acetamido-3-O-benzyl-4,6-O-benzylidene-2-deoxy-β-D-*
45 *glucopyranosyl-(1→4)-2,3,6-tri-O-benzyl-1-thio-β-D-glucopyranoside (16)*

46 A mixture of **21** (0.12 g, 0.13 mmol) and Zinc (0.43 g, activated with aq. 2% CuSO₄) in
47 THF/Ac₂O/AcOH 3:2:1 (5 mL) was stirred for 1h at room temperature. The reaction was diluted
48 with AcOEt and filtered over a Celite pad. Satd. aq. NaHCO₃ was added (30 mL) and, after
49 separation, the aqueous phases were extracted with AcOEt (2 x 20mL). The combined organics

1 were dried over NaSO₄, filtered and concentrated. Flash chromatography (Hexane/AcOEt, 6:4) of
2 the crude product gave pure **16** (0.080 g, 66%) as a foam. [α]_D²⁰ = -34.0 (*c* = 1 in chloroform)
3 ¹H NMR (CDCl₃): δ = 7.60-7.24 (m, 30H, arom.), 5.58 (*br d*, 1H, *J* = 9.1 Hz, NH), 5.50 (s, 1H,
4 CHPh), 4.89-4.86 (m, 3H, CH₂Ph), 4.77-4.67 (m, 5H, H-1', 2' and CH₂Ph), 4.65 (d, 1H, *J*_{1,2} = 9.7
5 Hz, H-1), 4.57-4.53 (m, 2H, CH₂Ph), 4.18-4.06 (m, 2H, H-4,6'a), 3.84-3.77 (m, 2H, 2 H-6), 3.66-
6 3.58 (m, 3H, H-3, 4', 6'b), 3.54-3.51 (m, 2H, H-2, 3'), 3.46-3.44 (m, 1H, H-5), 3.22-3.15 (m, 1H,
7 H-5'), 1.87 (s, 3H, CH₃). ¹³C NMR (CDCl₃): δ = 170.4 (C=O), 139.0-126.1 (36C, arom.), 101.6
8 (CHPh), 100.1 (C-1'), 87.5 (C-1), 85.3, 80.6, 78.7 (C-5), 78.6, 76.5 (C-4), 75.8, 75.4 (CH₂Ph), 75.2
9 (CH₂Ph), 73.5 (CH₂Ph), 71.5 (CH₂Ph), 68.7 (C-6), 68.6 (C-6'), 67.1 (C-5'), 50.4 (C-2'), 23.2 (CH₃).
10 MS (ESI) *m/z* (%): 946.4 (100) [M+Na]⁺, 1869.7 (75) [2M+Na]⁺. HRMS (ESI): *m/z* calcd for
11 C₅₅H₅₇NO₁₀NaS 946.3601 [M+Na]⁺, found 946.3605.

12
13 *4.1.16. Synthesis of N-benzyl-N-benzoyloxycarbonyl-3-aminopropyl 2-acetamido-3-O-benzyl-4,6-O-*
14 *benzylidene-2-deoxy- β -D-mannopyranosyl-(1 \rightarrow 4)-2,3,6-tri-O-benzyl- α -D-glucopyranosyl-(1 \rightarrow 3)-*
15 *2,4-di-O-benzyl- α -L-rhamnopyranoside (2)*

16 *From compound 15:* A mixture of **15** (0.14 g, 0.10 mmol) and Zinc (0.44 g, activated with aq. 2%
17 CuSO₄) in THF/Ac₂O/AcOH 3:2:1 (5 mL) was stirred for 3h at room temperature. The reaction was
18 diluted with AcOEt and filtered over a Celite pad. Satd. aq. NaHCO₃ was added (30 mL) and, after
19 separation, the aqueous phases were extracted with AcOEt (2 x 20mL). The combined organics
20 were washed with brine, dried over NaSO₄, filtered and concentrated. Flash chromatography
21 (Hexane/AcOEt, 7:3) of the crude product gave pure **16** (0.091 g, 62%) as an amorphous glassy
22 solid.

23 *From compound 16:* A solution of **16** (0.06 g, 0.065 mmol) and **3** (0.08 g, 0.13 mmol) in dry DCM
24 (2 mL) containing 4Å molecular sieves (0.15 g) was stirred at room temperature for 0.5 h. The
25 suspension was cooled to -35 °C, and then NIS (0.022 g, 0.097 mmol) followed by AgOTf (8 mg,
26 0.033 mmol) were added. After the addition, the reaction was allowed to warm to -10 °C and was
27 stirred at that temperature. After 0.45 h, TLC (Hexane/AcOEt, 6:4) showed the disappearances of
28 the donor. The reaction was diluted with DCM (30 mL) and filtered over a Celite pad. The organic
29 solution was then washed with 10 % aq. Na₂S₂O₃ (30 mL) and satd aq. NaHCO₃ (30 mL). The
30 organics were then dried over NaSO₄, filtered and concentrated. The residue was purified by flash
31 chromatography (Hexane/AcOEt, 7:3) to give first α -**2** (0.018 g), followed by β -**2** (0.037 g) with an
32 overall glycosylation yield of 60%. [α]_D²⁰ = +0.77 (*c* = 0.5 in chloroform). ¹H NMR (CDCl₃): δ =
33 7.59-7.01 (m, 45H, arom.), 5.53-5.43 (m, 2H, 1H x CH₂Ph and NH), 5.23-5.17 (m, 2H, CH₂Ph),
34 5.15 (d, 1H, *J*_{1,2} = 2.9 Hz, H-1), 4.95 (d, 1H, *J* = 11.7 Hz, 1 x CH₂Ph), 4.89-4.74 (m, 4H, CH₂Ph),
35 4.74-4.46 (m, 10H, H-1, 2" and CH₂Ph), 4.43 (*br s*, 1H, H-1"), 4.25-4.11 (m, 2H, H-6a'' and 1 x
36 CH₂Ph), 4.07-4.01 (m, 2H, H-3, 3'), 3.99-3.92 (m, 2H, H-4', 5'), 3.84 (*br d*, 1H, H-2), 3.75-3.58 (m,
37 5H, H-a, 4, 5, 2', 6b''), 3.55 (t, 1H, *J*_{3,4''} = *J*_{4,5''} = 9.6 Hz, H-4''), 3.47-3.20 (m, 6H, H-a', 6a', 6b',
38 3'' and 2 H-c), 3.13-3.03 (m, 1H, H-5''), 1.87-1.73 (m, 5H, 2 H-b and CH₃CO), 1.32 (d, 3H, *J*_{5,6} =
39 6.0 Hz, 3 H-6). ¹³C NMR (CDCl₃): δ = 170.25 (C=O), 156.6 and 156.1 (C=O), 139.4-126.1 (54C,
40 arom.), 101.6 (CHPh), 99.4 (C-1''), 98.3 (C-1), 96.3 (C-1'), 80.8 (C-4'), 79.9 (C-4), 79.4 (C-2'),
41 78.7 (C-4''). 77.6 (*br s*, C-3'), 76.3 (C-2), 75.8 (C-3''), 75.7 (C-3), 74.9 (CH₂Ph), 74.8 (CH₂Ph),
42 73.6 (CH₂Ph), 73.4 (CH₂Ph), 73.3 (CH₂Ph), 71.3 (CH₂Ph), 70.2 (C-5'), 68.7 (C-6''), 68.5 (C-5),
43 68.0 (C-6'), 67.2 (CH₂Ph), 67.0 (C-5''), 65.1 (C-a), 50.8 and 50.6 (2C, C-2'' and NCH₂Ph), 44.6
44 and 43.7 (C-c), 28.3 and 27.9 (C-b), 23.1 (CH₃CO), 18.1 (C-6). MS (ESI) *m/z* (%): 1461.9 (100)
45 [M+Na]⁺. HRMS (ESI): *m/z* calcd for C₈₇H₉₄N₂O₁₇Na 1461.6450 [M+Na]⁺, found 1461.6449.

46
47 *4.1.17. Synthesis of 3-aminopropyl 2-acetamido-2-deoxy- β -D-mannopyranosyl-(1 \rightarrow 4)- α -D-*
48 *glucopyranosyl-(1 \rightarrow 3)- α -L-rhamnopyranoside hydrochloride salt (1)*

49 Compound **2** (0.080 g, 0.056 mmol) in AcOEt/MeOH/0.02M HCl, 1:1:1 (9 mL) was
50 hydrogenolyzed over Pd(OH)₂ (0.070 g) for 4 days. The mixture was filtered over pleated filter

1 paper, the filtrate was concentrated to 1 mL, and then lyophilized to give trisaccharide **1** (0.034 g,
2 97%) as an amorphous white solid. $[\alpha]_D^{20} = -19.6$ ($c = 0.5$ in water). $^1\text{H-NMR}$ (D_2O): $\delta = 4.98$ (d,
3 1H, $J_{1,2} = 3.7$ Hz, H-1'), 4.81 (*br* d, 1H, $J_{1,2} = 1.3$ Hz, H-1''), 4.77 (*br* d, 1H, $J_{1,2} = 1.7$ Hz, H-1),
4 4.47 (dd, 1H, $J_{2,3} = 1.4$ Hz, $J_{2,3} = 4.4$ Hz, H-2''), 4.09-4.05 (m, 1H, H-2), 3.98-3.92 (m, 1H, H-5'),
5 3.89-3.34 (m, 15H), 3.10-2.95 (m, 2H, 2 H-c), 1.99 (s, 3H, CH_3CO), 1.95-1.87 (m, 2H, H-b), 1.23
6 (d, 3H, 3 H-6). $^{13}\text{C NMR}$ (D_2O): $\delta = 175.4$ (C=O), 99.4 (2C, C-1, 1''), 95.5 (C-1'), 78.6, 76.5 (C-
7 5''), 75.9 (C-3), 71.9 (C-3''), 71.5 (C-3'), 71.2 (C-2'), 70.2 (2C), 68.8, 66.8 (C-2), 66.7, 64.9 (C-a),
8 60.4 (C-6''), 59.7 (C-6'), 53.3 (C-2''), 37.4 (C-c), 26.7 (C-b), 22.0 (CH_3CO), 16.7 (C-6). MS (ESI)
9 m/z (%): 609.3 (100) $[\text{M}+\text{Na}]^+$. HRMS (ESI): m/z calcd for $\text{C}_{23}\text{H}_{43}\text{N}_2\text{O}_{15}$ 587.2663 $[\text{M}+\text{H}]^+$, found
10 587.2664.

11 4.2. Biological test

12 **Competitive ELISA assay:** 96-well flat-bottomed plates were incubated overnight at 4-8°C with a
13 mixture of *S. pneumoniae* CPS 19A (1 mg/mL, Statens Serum Institut, Artillerivej, Denmark) or
14 19F (1 mg/mL, Sanofi-Aventis, France) and methylated human serum albumin (1 mg/mL). A
15 solution of foetal calf serum (5%) in phosphate-buffered saline supplemented with Brij-35 (0.1%)
16 and sodium azide (0.05%) was applied to the plates for blocking of nonspecific binding sites. The
17 plates were incubated overnight at 4-8°C with a solution (1:200) of rabbit anti-19A or 19F, used as
18 reference serum (Statens Serum Institut, Artillerivej, Denmark). When trisaccharide was tested, it
19 was added to each well immediately before the addition of the reference serum. The plates were
20 then incubated with alkaline phosphatase conjugate goat anti-rabbit IgG (Sigma-Aldrich, Milan,
21 Italy), stained with *p*-nitrophenylphosphate, and the absorbance was measured at 405 nm with an
22 Ultramark microplate reader (Bio-Rad Laboratories S.r.l., Milan, Italy).

23 **Acknowledgements:** This work was supported by the Italian Ministry of University and Research
24 (PRIN 2015 grant, prot. 2015RNWJAM, Nanoplatforms for enhanced immune response).

25 References

- 26 1. Geno KA, Gilbert GL, Song JY, Skovsted IC, Klugman KP, Jones C, Konradsen HB, Nahm MH. Pneumococcal
27 Capsules and Their Types: Past, Present, and Future. *Clinical microbiology reviews* 2015; 28: 871-899.
- 28 2. Mehr S, Wood N. Streptococcus pneumoniae--a review of carriage, infection, serotype replacement and
29 vaccination. *Paediatric respiratory reviews* 2012; 13: 258-264.
- 30 3. Pomat WS, Lehmann D, Sanders RC, Lewis DJ, Wilson J, Rogers S, Dyke T, Alpers MP. Immunoglobulin G
31 antibody responses to polyvalent pneumococcal vaccine in children in the highlands of Papua New Guinea.
32 *Infection and immunity* 1994; 62: 1848-1853.
- 33 4. Whitney CG, Pilishvili T, Farley MM, Schaffner W, Craig AS, Lynfield R, Nyquist AC, Gershman KA, Vazquez
34 M, Bennett NM, Reingold A, Thomas A, Glode MP, Zell ER, Jorgensen JH, Beall B, Schuchat A. Effectiveness
35 of seven-valent pneumococcal conjugate vaccine against invasive pneumococcal disease: a matched case-
36 control study. *Lancet* 2006; 368: 1495-1502.
- 37 5. Morona JK, Morona R, Paton JC. Comparative genetics of capsular polysaccharide biosynthesis in
38 *Streptococcus pneumoniae* types belonging to serogroup 19. *Journal of bacteriology* 1999; 181: 5355-5364.
- 39 6. Millar EV, Pimenta FC, Roundtree A, Jackson D, Carvalho Mda G, Perilla MJ, Reid R, Santosham M,
40 Whitney CG, Beall BW, O'Brien KL. Pre- and post-conjugate vaccine epidemiology of pneumococcal
41 serotype 6C invasive disease and carriage within Navajo and White Mountain Apache communities. *Clinical
42 infectious diseases : an official publication of the Infectious Diseases Society of America* 2010; 51: 1258-
43 1265.
- 44 7. Hausdorff WP, Hoet B, Schuerman L. Do pneumococcal conjugate vaccines provide any cross-protection
45 against serotype 19A? *BMC pediatrics* 2010; 10: 4.
- 46 8. Grant LR, O'Brien SE, Burbidge P, Haston M, Zancolli M, Cowell L, Johnson M, Weatherholtz RC, Reid R,
47 Santosham M, O'Brien KL, Goldblatt D. Comparative immunogenicity of 7 and 13-valent pneumococcal
48
49
50

1 conjugate vaccines and the development of functional antibodies to cross-reactive serotypes. *PLoS one*
2 2013; 8: e74906.

3 9. Poolman J, Frasch C, Nurkka A, Kayhty H, Biemans R, Schuerman L. Impact of the conjugation method on
4 the immunogenicity of Streptococcus pneumoniae serotype 19F polysaccharide in conjugate vaccines.
5 *Clinical and vaccine immunology : CVI* 2011; 18: 327-336.

6 10. Kuttel MM, Jackson GE, Mafata M, Ravenscroft N. Capsular polysaccharide conformations in
7 pneumococcal serotypes 19F and 19A. *Carbohydrate research* 2015; 406: 27-33.

8 11. Morelli L, Cancogni D, Tontini M, Nilo A, Filippini S, Costantino P, Romano MR, Berti F, Adamo R, Lay L.
9 Synthesis and immunological evaluation of protein conjugates of Neisseria meningitidis X capsular
10 polysaccharide fragments. *Beilstein journal of organic chemistry* 2014; 10: 2367-2376.

11 12. Vetro M, Safari D, Fallarini S, Salsabila K, Lahmann M, Penades S, Lay L, Marradi M, Compostella F.
12 Preparation and immunogenicity of gold glyco-nanoparticles as antipneumococcal vaccine model.
13 *Nanomedicine* 2017; 12: 13-23.

14 13. Compostella F, Pitirollo O, Silvestri A, Polito L. Glyco-gold nanoparticles: synthesis and applications.
15 *Beilstein journal of organic chemistry* 2017; 13: 1008-1021.

16 14. Vansteijn AMP, Kamerling JP, Vliegthart JFG. Synthesis of 4 Spacer-Containing Trisaccharides with the
17 4-O-(Beta-L-Rhamnopyranosyl)-D-Glucopyranose Unit in Common, Representing Fragments of Capsular
18 Polysaccharides from Streptococcus-Pneumoniae Type-2, Type-7f, Type-22f, and Type-23f. *J Carbohydr*
19 *Chem* 1992; 11: 665-689.

20 15. Su Y, Xie J, Wang Y, Hu X, Lin X. Synthesis and antitumor activity of new shikonin glycosides. *European*
21 *journal of medicinal chemistry* 2010; 45: 2713-2718.

22 16. Paulsen H, Helpap B, Lorentzen JP. [Synthesis of trisaccharide units from capsular polysaccharides of
23 Streptococcus pneumoniae]. *Carbohydrate research* 1988; 179: 173-197.

24 17. Panza L, Ronchetti F, Toma L. Streptococcus-Pneumoniae Type-19a Polysaccharide - Synthesis of the
25 Trisaccharide Component of the Repeating Unit. *Carbohydrate research* 1988; 181: 242-245.

26 18. Kaji E, Osa Y, Tanaike M, Hosokawa Y, Takayanagi H, Takada A. An alternative access to a trisaccharide
27 repeating unit of the capsular polysaccharide of Streptococcus pneumoniae serotype 19A. *Chemical &*
28 *pharmaceutical bulletin* 1996; 44: 437-440.

29 19. Bonaccorsi F, Catelani G, Oscarson S. A new route for the synthesis of Streptococcus pneumoniae 19F
30 and 19A capsular polysaccharide fragments avoiding the beta-mannosamine glycosylation step.
31 *Carbohydrate research* 2009; 344: 1442-1448.

32 20. Bousquet E, Khitri M, Lay L, Nicotra F, Panza L, Russo G. Capsular polysaccharide of Streptococcus
33 pneumoniae type 19F: synthesis of the repeating unit. *Carbohydrate research* 1998; 311: 171-181.

34 21. Legnani L, Ronchi S, Fallarini S, Lombardi G, Campo F, Panza L, Lay L, Poletti L, Toma L, Ronchetti F,
35 Compostella F. Synthesis, molecular dynamics simulations, and biology of a carba-analogue of the
36 trisaccharide repeating unit of Streptococcus pneumoniae 19F capsular polysaccharide. *Org Biomol Chem*
37 2009; 7: 4428-4436.

38 22. Liotta LJ, Capotosto RD, Garbitt RA, Horan BM, Kelly PJ, Koleros AP, Brouillette LM, Kuhn AM,
39 Targonsidis S. Synthesis of methyl alpha-D-glucopyranosyl-(1 -> 4)-alpha-D-galactopyranoside and methyl
40 alpha-D-xylo-hex-4-ulopyranosyl-(1 -> 4)-alpha-D-galactopyranoside. *Carbohydrate research* 2001; 331:
41 247-253.

42 23. Nitz M, Bundle DR. Synthesis of di- to hexasaccharide 1,2-linked beta-mannopyranan oligomers, a
43 terminal S-linked tetrasaccharide congener and the corresponding BSA glycoconjugates. *J Org Chem* 2001;
44 66: 8411-8423.

45 24. Baek JY, Kwon HW, Myung SJ, Park JJ, Kim MY, Rathwell DCK, Jeon HB, Seeberger PH, Kim KS. Directing
46 effect by remote electron-withdrawing protecting groups at O-3 or O-4 position of donors in glycosylations
47 and galactosylations. *Tetrahedron* 2015; 71: 5315-5320.

48 25. Mydock LK, Demchenko AV. Mechanism of chemical O-glycosylation: from early studies to recent
49 discoveries. *Org Biomol Chem* 2010; 8: 497-510.

50 26. Franchini L, Compostella F, Colombo D, Panza L, Ronchetti F. Synthesis of the Sulfonate Analogue of
51 Seminolipid via Horner-Wadsworth-Emmons Olefination. *J Org Chem* 2010; 75: 5363-5366.

- 1 27. Vetro M, Costa B, Donvito G, Arrighetti N, Cipolla L, Perego P, Compostella F, Ronchetti F, Colombo D.
- 2 Anionic glycolipids related to glucuronosyldiacylglycerol inhibit protein kinase Akt. *Org Biomol Chem* 2015;
- 3 13: 1091-1099.
- 4