


Accepted: 27 May 2018

DOI: 10.1111/clar.13311

SUPPLEMENT ARTICLE

WILEY  CLINICAL ORAL IMPLANTS RESEARCH

Implant supported cantilevered fixed dental rehabilitations in partially edentulous patients: Systematic review of the literature. Part I.

Stefano Storelli¹  | Massimo Del Fabbro² | Massimo Scanferla¹ | Giulia Palandrani¹ | Eugenio Romeo¹

¹Department of Biomedical, Surgical and Dental Sciences, Clinica Odontoiatrica ASST Santi Paolo e Carlo, University of Milan, Milan, Italy

²Department of Biomedical, Surgical and Dental Sciences, IRCCS Orthopedic Institute Galeazzi, University of Milan, Milan, Italy

Correspondence

Stefano Storelli, Department of Biomedical, Surgical and Dental Sciences, University of Milan, Milan, Italy.

Email: stefano.storelli@studioplinio.it

Abstract

Objectives: To investigate in which clinical situations a cantilever fixed implant supported restorations can be a treatment alternative and which complications are reported.

Materials and Methods: Two operators screened the literature (MEDLINE, EMBASE) and performed a hand search on the main journals dealing with implantology and prosthetics until 31 December 2017. Only articles that considered cantilever implant fixed restorations with at least 10 patients and with a mean follow-up of at least 5 year were selected. The outcome variables were survival of implants and prosthesis, mechanical, technical and biological complications, marginal bone loss. The review was performed according to the PRISMA statements. Risk of bias assessment was evaluated. Failure and complication rates were analysed using random effect Poisson regression models to obtain summary estimate of 5- and 10-year survival and complication rates.

Results: A total of nine papers were selected for partially edentulous patients and reported high survival rate of the prosthesis. The estimated survival rate for 5–10 years was calculated to be 98.4% for the implants and 99.2% for the rehabilitations. Mechanical, technical and biological complications were reported with a cumulative 5–10 years complication rate of 28.66% and 26.57% for the patients and for the prosthesis, respectively. Two papers for single implant supporting 2-unit cantilever were not sufficient to draw conclusions.

Conclusions: There is evidence that cantilever can be successful treatment in partially edentulous patients. In two adjacent edentulous sites, data are not yet sufficient.

1 | INTRODUCTION

Implant placement can be limited by anatomical conditions that may be overcome with different solutions: reduced dimension implants, surgical bone augmentation procedures or different prosthetic designs. Other systematic reviews have pointed out how major reconstructions can be effective but need to be carefully applied in cases

with ideal conditions (Chiapasco, Zaniboni, & Boisco, 2006; Esposito, Grusovin, Worthington, & Coulthard, 2006). On the other hand, short or tilted implants can be a less invasive and effective procedure, provided the bone is sufficient for their placement (Del Fabbro, Bellini, Romeo, & Francetti, 2012; Zinsli, Sägesser, & Mericske, 2004).

This concept was borrowed from the prosthetic rehabilitation of periodontally treated patients, where cantilevered

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2018 The Authors. *Clinical Oral Implants Research* Published by John Wiley & Sons Ltd.




FIGURE 1 Flow chart

prosthesis supported by natural teeth was used. Tooth supported cantilever FPD were reported to have statistically higher incidence of failure than non-cantilevered tooth

supported FPDs (Pjetursson, Bragger, Lang, & Zwahlen, 2007).

One of the prosthetic alternatives is the use of cantilevered prostheses (Implant cantilevered fixed dental prostheses, ICFDP). This is an option in anatomical compromised locations, or in patients that have limited financial means to afford complex treatments. In such design, nor implants nor biomaterials are placed in resorbed areas, thus reducing the risk for failures and lowering the invasiveness of the treatment. The biomechanical risk of cantilever may be that of overloading the rehabilitations, leading to implant and/or prosthetic failure. In vitro studies have revealed that higher stress to the implant closest to the cantilever extension may be concentrated at the marginal bone level and may pose a risk to marginal bone loss. (Sertgoz & Guvener, 1996; Stegaroiu, Sato,

TABLE 1 Excluded studies

Cantilever data not retrievable	Koller, Pereira-Cenci and Boscatto (2016); Ozgur, Kazancioglu, Demirtas, Deger and Ak (2016); Mangano et al. (2014); Ekfeldt, Zellmer and Carlsson (2013); Degidi, Nardi and Piattelli (2013); Wittneben et al. (2014); Heschl et al. (2012); Ortorp and Jemt (2012); Malo, de Araújo Nobre, Lopes, Moss and Molina (2011); Krennmair, Seemann, Schmidinger, Ewers and Piehlslinger (2010); Eliasson et al. (2010); Davó (2009); Isaksson, Becktor, Brown, Laurizohn and Isaksson (2009); Degidi, Iezzi, Perrotti and Piattelli (2009); Ortorp and Jemt (2009); Gualini, Gualini, Cominelli and Lekholm (2009); Blanes, Bernard, Blanes and Belser (2007); Rasmusson, Roos and Bystedt (2005); Hartman and Cochran (2004); Åstrand et al. (2004); Attard and Zarb (2004); Ekelund, Lindquist, Carlsson and Jemt (2003); Murphy, Absi, Gregory and Williams (2002); Raghoobar, Timmenga, Reintsema, Stegenga and Vissink (2001); Tinsley, Watson and Russell (2001); Bragger, Aeschlimann, Bürgin, Hämmerle and Lang (2001); Friberg, Gröndahl, Lekholm and Brånemark (2000); Becker and Kaiser (2000); Schwartz-Arad, Gulayev and Chaushu (2000); Arvidson, Bystedt, Frykholm, von Konow and Lothigius (1998); Schwartz-Arad and Chaushu (1998); Keller, Tolman and Eckert (1998); Parein, Eckert, Wollan and Keller (1997); Schnitman, Wöhrle, Rubenstein, DaSilva and Wang (1997); Jemt and Lekholm (1995); Brånemark, Svensson and van Steenberghe (1995); Hemmings, Schmitt and Zarb (1994); Naert, Quirynen, van Steenberghe and Darius (1992); Zarb and Schmitt (1991)
Non-human study	Costa, Santos, Nary and Brånemark (2015); Kupeyan and Clayton (2004); McAlarney and Stavropoulos (2000)
Mean follow-up <5 years	Correia, Gouveia, Felino, Costa and Almeida (2017); Wang, Judge and Bailey (2016); Tartaglia, Maiorana, Gallo, Codari and Sforza (2016); Francetti et al. (2015); Mundt, Heinemann, Schwahn and Biffar (2012); Lee et al. (2011); Francetti, Romeo, Corbella, Taschieri and Del Fabbro (2012); Mangano et al. (2011); Nedir, Bischof, Szmulker-Moncler, Belser and Samson (2006); Ibañez et al. (2005); Balshi, Wolfinger and Balshi (2005); Becker (2004); Romeo et al. (2003); Engstrand et al. (2003); Ahrén and Kahnberg (2001); Brocard et al. (2000); Eliasson, Palmqvist, Svensson and Sondell (2000); Haas, Mendorff-Pouilly, Mailath and Bernhart (1998); Kucey (1997); Gotfredsen (1997); Carlson and Carlsson (1994)
Number of patients < 10	Deporter, Ogiso, Sohn, Ruljancich and Pharoah (2008); Van Nimwegen et al. (2017); Fischer and Stenberg (2013)
On natural teeth	Lam, Botelho and McGrath (2013); Cordaro, Ercoli, Rossini, Torsello and Feng (2005)
Out of topic (no cantilever)	Agliardi, Romeo, Panigatti, de Araújo Nobre and Maló (2017); Malo, de Araujo Nobre, Guedes and Almeida (2017); Niedermaier et al. (2017); Zanolta et al. (2016); Lee, Kweon, Choi and Kim (2016); Esposito et al. (2016); Cavalli et al. (2016); Zhang, Shi, Gu and Lai (2016); Imburgia and Del Fabbro (2015); Ata-Ali et al. (2015); Tealdo et al. (2014); Pettersson and Sennerby (2015); Raval, Dahlgren, Teiwik and Gröndahl (2013); Kim et al. (2013); Al-Nawas et al. (2012); Ozkan, Akoğlu and Kulak-Ozkan (2011); Browaeys et al. (2011); Lethaus, Kälber, Petrin, Brandstätter and Weingart (2011); Mura (2012); Schrott, Jimenez, Hwang, Fiorellini and Weber (2009); Botticelli, Renzi, Lindhe and Berglundh (2008); Friberg, Raghoobar, Grunert, Hobkirk and Tepper (2008); Astrand, Ahlqvist, Gunne and Nilson (2008); Glauser, Zembic, Ruhstaller and Windisch (2007); Jaffin, Kolesar, Kumar, Ishikawa and Fiorellini (2007); Romeo, Ghisolfi, Rozza, Chiapasco and Lops (2006); Romeo, Lops, et al. (2006); Sullivan, Vincenzi and Feldman (2005); Quirynen et al. (2005); Degidi and Piattelli (2005); Vigolo, Givani, Majzoub and Cordioli (2004); Zinsli et al. (2004); Lambrecht, Filippi, Künzel and Schiel (2003); Davis, Packer and Watson (2003); Weng et al. (2003); Attard and Zarb (2003); Brosky, Koriath and Hodges (2003); Naert et al. (2002); Fortin, Sullivan and Rangert (2002); Wyatt and Zarb (2002); Attard and Zarb (2002); Zarb and Zarb (2002); Ferrigno, Laureti, Fanali and Grippaudo (2002); Sullivan, Sherwood and Porter (2001); Ekfeldt et al. (2001); Hellem et al. (2001); Merickse-Stern, Aerni, Geering and Buser (2001); Allen, McMillan and Walshaw (2001); Vajdovich and Fazekas (1999); Noack, Willer and Hoffmann (1999); Schliephake, Schmelzeisen, Husstedt and Schmidt-Wondera (1999); Chaushu and Schwartz-Arad (1999); Makkonen et al. (1997); Zarb and Schmitt (1993)
Same pool of patients of other article	Cavalli, Corbella, Taschieri and Francetti (2015); Fischer, Stenberg, Hedin and Sennerby (2008)

TABLE 2 Study and patient characteristics of the included studies (partial prostheses)

Study	Implant system	Study design	Mean Follow-up, years (range)	No. implants placed (ICFDPs/patients)	No. implants available for analysis (ICFDPs/patients)	Age range (mean), years	Setting	Type of extension	Location of reconstruction
Romeo et al. (2009)	Straumann Dental Implant System®	Prospective	8	148 (75/59)	116 (59/45)	42–100 (63)	Hospital	Mixed	33 maxilla, 26 mandible
Kreissl et al. (2007)	3i Ossotite®	Prospective	5	61 (23/20)	61 (23/20)	NR (NR)	NR	Monolateral	Maxilla and mandible
Brägger et al. (2005)	Straumann Dental Implant System®	Prospective	9.4	33 (18/14)	33 (18/14)	20–78 (42.9)	University	Mixed	11 maxilla, 7 mandible
Wennström et al. (2004)	Astra Tech® Dental Implant System	Prospective	5	NR (28/28)	66 (24/24)	NR (57)	University	Distal	16 maxilla, 8 mandible
Jokstad et al. (2017)	Brånemark System®	Retrospective	17.5	NR (NR/NR)	58 (24/24)	17–73 (45)	University	Monolateral	Maxilla and mandible
Malo et al. (2013)	Brånemark System®	Retrospective	5	185 (150/137)	153 (119/113)	17–84 (49)	Private clinic	Monolateral	Maxilla and mandible
Aglietta et al. (2012)	Straumann Dental Implant System®	Retrospective	5.5	42 (21/21)	42 (21/21)	NR (54.6)	University	Monolateral (9 mesial/2distal)	14 maxilla, 7 mandible
Hälg et al. (2008)	Straumann Dental Implant System®	Retrospective	5	38 (19/19)	34 (17/17)	44–83 (61.9)	Private clinic	Monolateral	9 maxilla, 10 mandible
Eliasson et al. (2006)	Brånemark System®	Retrospective	9.8	209 (84/84)	176 (71/71)	NR (NR)	University	Mixed	NR

ICFDPs: implant-supported, cantilever-fixed dental prostheses; NR: not reported.

TABLE 3 Implant and prosthesis characteristics of the included studies (Partial prostheses)

Study Prospective/Retrospective	No. implants available for analysis (ICFDPs/patients)	Implant system	Surface	Implant diameter (mm)	Implant length (mm)	Loading	Type of extension	Location of reconstruction	Fixation	Type of material
Romeo et al. (2009) Prospective	116 (59/45)	Straumann Dental Implant System®	N/A	3.3–4.1–4.8	8.0–14.0	Conventional >3 months	Mixed	33 maxilla, 26 mandible	Cemented and screw-retained	Metal-ceramic
Kreissl et al. (2007) Prospective	61 (23/20)	3i Osseotite®	Smooth	NR	NR	NR	Monolateral	Maxilla and mandible	Screw-retained	Metal-ceramic
Brägger et al. (2005) Prospective	33 (18/14)	Straumann Dental Implant System®	Rough	NR	NA	Conventional >3 months	Mixed	11 maxilla, 7 mandible	Cemented and screw-retained	Metal-ceramic
Wennström et al. (2004) Prospective	66 (24/24)	Astra Tech® Dental Implant System	Rough/Smooth	3.5	8–19	Conventional >3 months	Distal	16 maxilla, 8 mandible	Screw-retained	Metal-ceramic
Jokstad et al. (2017) Retrospective	58 (24/24)	Brånemark System®	Smooth	≤4.5	≤14	NR	Monolateral	Maxilla and mandible	Screw-retained	Resin
Malo et al. (2013) Retrospective	153 (119/113)	Brånemark System®	Rough/Smooth	3.3–3.75–4.0	8.5–18.0	Immediate/Early/Conventional	Monolateral	Maxilla and mandible	Cemented and screw-retained	Ceramic; Metal ceramic; Metal-resin
Aglietta et al. (2012) Retrospective	42 (21/21)	Straumann Dental Implant System®	Rough	4.1–4.8	NR	NR	Monolateral mesial/distal	14 maxilla, 7 mandible	Cemented and screw-retained	NA
Hälg et al. (2008) Retrospective	34 (17/17)	Straumann Dental Implant System®	Smooth	<4.5	6.0–12.0	Conventional >3 months	Monolateral	9 maxilla, 10 mandible	Cemented	Metal-ceramic
Eliasson et al. (2006) Retrospective	176 (71/71)	Brånemark System®	Smooth	3.75	6–18	NR	Mixed	NR	Screw-retained	Metal-resin/ Metal-ceramic

ICFDPs: implant-supported, cantilever-fixed dental prostheses; NR: not reported.

TABLE 4 Study and patient characteristics of the included studies (single implant)

Study	Implant system	Study design	Mean		No. implants placed (ICFDPs/patients)	No. implants available for analysis (ICFDPs/patients)	Age range (mean), years	Setting	Type of extension	Location of reconstruction
			Follow-up, years (range)	Follow-up, years (range)						
De Angelis et al. (2017)	Winsix	Retrospective	13.6 (10–18)	28 (28/28)	25 (25/25)	NR (NR)	University	Monolateral	15 maxilla, 10 mandible	
Aglietta et al. (2012)	Straumann dental implant system	Retrospective	6.5 (NR)	19 (19/17)	19 (19/17)	NR (54.6)	University	Monolateral, 9 mesial, 10 distal	5 maxilla 14 mandible	

ICFDPs: implant-supported, cantilever-fixed dental prostheses; NR: not reported.

TABLE 5 Risk of bias summary (partial prostheses)

	Blinding of outcome assessment	Comparability of control and treatment groups at entry	Clear definition of inclusion and exclusion criteria	Outcome measurement description	Completeness of the outcome data reported	Recall rate	Sample size	Number of surgeon involved
Hälg et al. (2008)	-	+	?	+	+	?	+	+
Wennström et al. (2004)	-	+	+	+	+	+	-	+
Romeo et al. (2009)			+	+	+	-	+	?
Kreissl et al. (2007)			-	+	+	+	-	?
Brägger et al. (2005)			-	+	+	?	-	-
Eliasson et al. (2006)			-	-	+	+	+	?
Jokstad et al. (2017)			-	+	+	+	+	?
Malo et al. (2013)			+	+	+	+	+	?
Aglietta et al. (2012)			+	+	+	+	-	?

Note. +, Low risk of bias; -, High risk of bias; ?, moderate risk of bias.




FIGURE 2 Risk of bias graph (partial prostheses)

Kusakari, & Miyakawa, 1998; Zampelis, Rangert, & Heijl, 2007). By contrast, in humans the results of higher stresses on implants remain unclear.

Few systematic reviews have been published on the topic in the past years, two in 2009 (Aglietta et al., 2009; Zurdo, Romão, & Wennström, 2009) and one in 2012 (Romeo & Storelli, 2012). All of them considered the outcome of cantilevered prostheses used in fixed partial dentures (Partial implant cantilevered fixed dental prosthesis, PICFDP): the analysis was carried out on papers treating the cantilever solution in partial edentulism and mostly in posterior areas.

In literature, by contrast, other use of cantilever can be found also in fully edentulous cases (full arch implant cantilevered fixed dental prosthesis, FAICFDP) and in cases where one implant supports two teeth (single implant cantilevered fixed dental prosthesis, SICFDP) (Aglietta et al., 2012).

The main objective of this systematic review was to assess the survival and complication rate of implant supported cantilever fixed dental prosthesis (ICFDP) in different clinical situations.

2 | MATERIAL AND METHODS

The present systematic review was designed to report data on full arch and partial fixed reconstructions with cantilever. The present review is reported according to the PRISMA (Preferred Reporting Items Systematic review and Meta-Analyses) statement (Liberati et al., 2009; Moher, Liberati, Tetzlaff, & Altman, 2009).

The focused question was: "In what clinical situations and with what implant systems are cantilevers a successful treatment modality?" The preliminary PICO assessment was used to define the search strategy with the following criteria.

TABLE 6 Risk of bias summary (single implant)

	Clear definition of inclusion and exclusion criteria	Outcome measurement method description	Completeness of the outcome data reported	Recall rate	Sample size	Number of surgeon involved
De Angelis et al. (2017)	+	-	+	+	+	?
Aglietta et al. (2012)	+	+	+	+	-	?

Note. +, Low risk of bias; -, High risk of bias; ?, moderate risk of bias.

2.1 | Types of participants

Patients who received cantilevered implant supported rehabilitations.

2.2 | Types of interventions

Any rehabilitations that was produced with cantilevered teeth. Three different kinds of restorations were investigated: full-arch fixed restorations, fixed partial restorations and single implants supporting two-crown restorations.

2.3 | Types of outcome measures

Several variables were considered for analysis:

- Implant survival rate
- Prosthetic survival rate
- Biological complications
- Prosthetic complications (Mechanical and Technical)
- Marginal bone loss

Other variables were searched and described when present: loading time of the rehabilitations, reconstruction material, implant system used.

2.4 | Types of studies

The present systematic review considered both prospective and retrospective studies, randomized and controlled clinical trials as well as cohort studies and case series. Studies had to report data on minimum 10 participants and have a minimum of 5-year follow-up.

2.5 | Search strategy

The English literature was first searched up to July 2017 and a second search was carried out up to December 2017. Two electronic databases were searched: The National Library of Medicine (MEDLINE by PubMed) and EMBASE. The following terms were searched in combination: dental implant AND (cantilever or extension or "fixed dental prosthesis" or "fixed partial denture" or "full arch" or "fixed complete restoration" or "fixed complete prostheses" or "single implant" or "single tooth"). Moreover, the issues from 2015 to July 2017 of the following journals were hand searched: Clinical




FIGURE 3 Risk of bias graph (single implant)

Oral Implants Research, International Journal of Periodontics and Restorative Dentistry, Journal of Periodontology, Journal of Clinical Periodontology, International Journal of Oral and Maxillofacial Implants, Journal of Prosthetic Dentistry, Journal of Prosthodontics, Journal of Oral Rehabilitations. Moreover, the bibliographies of previous systematic reviews on the topic as well as selected articles were thoroughly screened. The aim of the review was to screen the literature for papers reporting at least a mean of 5-year follow-up data on cantilevered rehabilitations, both in fully edentulous and partially edentulous cases.

2.6 | Inclusion criteria

Both retrospective and prospective studies were selected with a mean follow-up of a minimum of 5 years and at least 10 rehabilitations. RCTs, Cohort and Case-Control studies on implant supported cantilever restorations were considered. The primary outcome was prosthetic and implant survival. Secondary outcome was complication rates (mechanical, technical and biological) and marginal bone loss. Moreover, information regarding implant manufacturer and abutment characteristics as well as influence of retention (cemented or screw retained) was assessed.

2.7 | Exclusion criteria

Papers were not meeting all inclusion criteria. Papers with a less than 5-year follow-up and/or with less than 10 patients were excluded. Letters, narrative reviews, questionnaires and charts were also excluded. Studies from which data on selected outcome variables could not at all be retrieved or calculated were not considered. Also, papers reporting data from the same cohort were excluded, except for the one with the longest follow-up.

2.8 | Study selection

The pool of retrieved articles was screened for duplicates by undergraduate students of the department (Stefano Corti and Elisabetta Morfini). All identified titles and abstract were then independently screened by two review authors (SS and GP). Full text was obtained either for articles meeting the inclusion criteria or for those whose abstract presented unclear data. The full texts were then assessed

by two authors (SS and GP) that defined if the articles were to be included or not. Any disagreement was resolved by discussion with the other reviewers (ER and MDF).

2.9 | Data extraction

Data were extracted by two review authors (SS and GP) using data collection forms. Study setting and design, implant manufacturer and data on restorations were extracted. Survival rate of implant and prosthesis were extracted or calculated from the original articles. Implant survival was considered if the implant was present at the follow-up examination; prosthesis survival was considered if the restoration was present at the follow-up visit without any modifications. Prosthesis complications were considered all the events affecting the abutment and/or the meso- and/or the supra-structures' integrity and were divided into mechanical and technical complications.

Implant/abutment related technical complications were considered those affecting the integrity of the implant and the abutment and were reported in tables as fracture of the implant, abutment and screws and abutment loosening. Restoration-related technical complications were considered to be those affecting the prosthetic rehabilitation: loss of retention (i.e., unscrewing of occlusal screws for screw-retained rehabilitations and decementations for cemented restorations), veneer chipping, fracture of framework. Biological complications comprised peri-implantitis and mucositis. Moreover, when reported, data on marginal bone loss were also extracted.

When the reported data were unclear, authors contacted by emails the corresponding authors and asked for more informations.

2.10 | Risk of bias assessment

The risk of bias assessment for the included trials was performed independently by two reviewers (SS and GP), using a purposely designed risk of bias assessment tool with the following domains:

Randomized Studies:

- Random sequence generation method
- Allocation concealment

Comparative Studies:

- Blinding of outcome assessment
- Comparability of control and treatment groups at entry

All Studies:

- Definition of inclusion and exclusion criteria
- Outcome measurement method description
- Completeness of the outcome data reported

Randomized studies were not considered as such if the randomization purpose was not the use of cantilever restorations. In that case, the study was considered only a prospective study.

Recall rate (it was assumed as low risk if the dropout rate was <10%, unclear if it was between 10% and 20%, high risk if it was >20%).

TABLE 7 Annual failure rates and survival of implants/ICFDPs (partial prostheses)

Study	No. implants/ ICFDPs available for the analysis (patients)	Mean follow-up time (years)	No. failures implants/ICFDPs	Total implants/ ICFDPs exposure time	Estimated failure rate (per 100 implants/ICFDPs years)	Estimated implant/ ICFDP survival rate after 5–10 years	Mean MBL, mm	Estimated MBL after 5 years
Romeo et al. (2009)	116/59 (45)	8	0/0	928/472	0.00/0.00	100%/100%	1.1	0.69
Kreissl et al. (2007)	61/23 (20)	5	1/1	305/115	0.33/0.87	98.4%/95.7%	NR	NE
Brägger et al. (2005)	33/18 (14)	9.4	1/3	310.2/169.2	0.32/1.77	98.4%/91.1%	NR	NE
Wennström et al. (2004)	66/24 (24)	5	2/2	330/130	0.61/1.54	97.0%/92.3%	0.39 ± 1.04	0.39
Summary estimate prospective studies								0.54 (−1.36, 2.45)
Jokstad et al. (2017)	58/24 (24)	17.5	1/0	1,015/420	0.10/0.00	99.5%/100%	NR	NE
Malo et al. (2013)	153/119 (113)	5	3/1	765/595	0.39/0.17	98.0%/99.2%	1.84	1.84
Aglietta et al. (2012)	42/21 (21)	5.5	0/0	231/115.5	0.00/0.00	100%/100%	0.25 ± 0.55	0.23
Hälg et al. (2008)	34/17 (17)	5	2/3	170/85	1.18/3.53	94.1%/82.4%	0.27	0.27
Eliasson et al. (2006)	176/71 (71)	9.8	3/0	1,724.8/695.8	0.17/0.00	99.1%/100%	NR	NE
Summary estimate retrospective studies								0.78 (−1.5, 3.06)
Total Summary estimate (95% CI)*						98.85 (88.4, 101.1) 99.0 (86.65, 106.0)		0.68 (−0.15, 1.52)

ICFDPs: implant-supported, cantilever-fixed dental prostheses; MBL: marginal bone loss; NR: not reported.

*Based on random effects Poisson regression, test for heterogeneity, $p = 0.34/p = 0.11$.

TABLE 8 Overall complications (partial prostheses)

Study	No. implants/CFDPs available for the analysis (patients)	Mean follow-up time (years)	Total patients exposure time	Total prostheses exposure time	Total No. complications	Estimated complication rate (per 100 patients years)	Estimated complication rate (per 100 prostheses years)
Romeo et al. (2009)	116/59 (45)	8	360	472	22	6.11	4.66
Kreissl et al. (2007)	61/23 (20)	5	100	115	14	14.00	12.17
Brägger et al. (2005)	33/18 (14)	9.4	131.6	169.2	14	10.64	8.27
Wennström et al. (2004)	66/24 (24)	5	120	120	6	5.00	5.00
Cumulative 5-year complication rate prospective studies						39.35% (11.56, 77.81)	31.96% (9.79, 65.48)
Jokstad et al. (2017)	58/24 (24)	17.5	420	420	8	1.90	1.90
Malo et al. (2013)	153/119 (113)	5	565	595	51	9.03	8.57
Aglietta et al. (2012)	42/21 (21)	5.5	115.5	115.5	NR	NR	NR
Hälg et al. (2008)	34/17 (17)	5	85	85	9	10.59	10.59
Eliasson et al. (2006)	176/71 (71)	9.8	695.8	695.8	18	2.59	2.59
Cumulative 5-year complication rate retrospective studies						24.35% (-5.04, 65.30)	23.94% (-4.83, 63.9)
Cumulative 5–10 year complication rate (95% CI)*						28.66% (19.56, 55.26)	26.57% (17.96, 49.24)

CFDPs: implant-supported, cantilever-fixed dental prostheses; NE: not estimable; NR: not reported.

*Based on random effects Poisson regression, test for heterogeneity for complications (per patients/prostheses), $p = 0.18/p = 0.08$.

TABLE 9 Mechanical complications: implant/abutment-related complications (partial prostheses)

Study	No. implants available for the analysis (patients)	Mean follow-up time (years)	Total implants exposure time	No. of implant fracture	Estimated rate of implant fractures (per 100 implants years)	No of abutment or screw fracture	Estimated rate of abutment or screw fractures (per 100 patients/year)
Romeo et al. (2009)	116 (45)	8	928	0	0	0	0.00
Kreissl et al. (2007)	61 (20)	5	305	0	0	0	0.00
Brägger et al. (2005)	33 (14)	9.4	310.2	1	0.32	1	0.76
Wennström et al. (2004)	66 (24)	5	330	0	0	1	0.83
Cumulative 5-year complication rate prospective studies					0.27% (−0.88, 1.68)		1.41% (−1.68, 5.66)
Jokstad et al. (2017)	58 (24)	17.5	1,015	0	0	1	0.24
Malo et al. (2013)	153 (113)	5	765	NR	NE	NR	NE
Aglietta et al. (2012)	42 (21)	5.5	231	NR	NE	NR	NE
Hälg et al. (2008)	34 (17)	5	170	2	1.18	0	0.00
Eliasson et al. (2006)	176 (71)	9.8	1,724.8	0	0	3	0.43
Cumulative 5-year complication rate retrospective studies					0.34% (−6.47, 10.39)		1.67% (−1.57, 3.80)
Cumulative 5–10 year complication rates (95% CI)*					0.31% (−0.97, 3.11)		1.57% (−0.05, 3.29)

*Based on random effects Poisson regression, test for heterogeneity $p = 0.10$ for implant fractures. Based on Poisson regression, test for heterogeneity $p = 0.08$ for abutment/screw fractures.

Sample size (it was considered low risk if >30 patients were treated, high risk if <30 patients were treated).

Number of surgeons involved (it was considered low risk if the same surgeon performed all operations, high risk if more than one surgeon performed all operations).

Each domain was judged as at low, unclear or high risk of bias according to the evaluation criteria as described in the Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0. A domain was evaluated as unclear when it was doubtful or not specified in the article. Cases of disagreement were resolved by discussion.

After judgement was given for each of the above-mentioned domains, studies were grouped into the following categories:

- Low risk of bias (plausible bias unlikely to seriously alter the results) if all criteria were met
- Moderate risk of bias (plausible bias that raises some doubt about the results) if one or more criteria were partly met or were assessed as unclear
- High risk of bias (plausible bias that seriously weakens confidence in the results) if one or more criteria were not met.

2.11 | Statistical analysis

Failure and complication rates were calculated by dividing the number of events (failures or complications) in the numerator by the total exposure time (implant, patient or prosthesis-time) in the denominator, similar to previous systematic reviews (Romeo & Storelli, 2012). Failures and complications were directly extracted from the publications, as well as the mean follow-up time. Exposure time was calculated by multiplying the mean follow-up time by the number of implants or ICFDPs available. The mean follow-up duration was directly extracted by the articles, provided by adjunctive information by the authors or estimated from the original data. For further analysis, the total number of events was considered to be Poisson distributed for a given sum of implant exposure years, and Poisson regression with a logarithmic link function and total exposure time per study as an offset variable was used (Kirkwood & Sterne, 2003). Event rates for implants and prostheses were calculated by dividing the total number of events by the respective total exposure time in years.

Robust standard errors were calculated to obtain 95% confidence intervals of the summary estimates of the event rates. To assess heterogeneity of the study-specific event rates, the Spearman goodness-of-fit statistics and associated *p*-value were calculated. If the goodness-of-fit *p*-value was below 0.05, indicating heterogeneity, random effects Poisson regression (with Gamma-distributed random effects) was used to obtain a summary estimate of the event rates. Five- and 10-year survival and complication proportions were estimated through the relationship between event rate and survival function S , $S(T) = \exp(-T \times \text{event rate})$, assuming constant event rates (Kirkwood & Sterne, 2003).

Ninety-five per cent confidence intervals (CIs) of the summary estimates of the event rates obtained from the Poisson regression were reported. The 95% CIs for survival probabilities were obtained using the 95% confidence limits from the summary event rates. All analyses were performed using IBM SPSS Statistics version 24 (IBM Corporation, Armonk, NY).

Regarding the reported radiographic bone loss, the mean difference between implants close to and distant from cantilevers, or belonging to non-cantilevered FPDs, and its standard error was calculated for each study. Such study-specific differences were then meta-analysed using the inverse-variance weighting method. Random effects model was used if no heterogeneity among studies was detected, otherwise a fixed effects model was chosen, following the directions of the Cochrane Handbook for Systematic Reviews of Interventions (Version 5.1.0, March 2011). Heterogeneity among studies was assessed with Cochran's test for heterogeneity, with a significance threshold of $p < 0.1$. The quantification of the heterogeneity was calculated with I^2 statistics. Review Manager 5.3 (The Nordic Cochrane Center, The Cochrane Collaboration, Copenhagen, Denmark) was used for meta-analysis calculations and plots concerning radiographic peri-implant bone loss.

3 | RESULTS

The electronic search identified a total of 6,926 titles (4386 MEDLINE, 2540 EMBASE). Another 23 titles were included after manual search. After de-duplication a total of 5,336 studies were screened. A total of 149 papers underwent full-text analysis (Figure 1). After full-text reading, 125 papers were excluded. Reasons for excluding papers were mainly follow-up less than 5 years, papers on natural teeth, in vitro or non-clinical studies. Also, papers non-clearly reporting data on cantilever were excluded. When, after discussion, there was still a doubt, authors were contacted by email and asked for better explanations. Reason for exclusion can be found in Table 1. Any disagreement was resolved by discussion. Finally, 24 papers were selected and included in the review: 10 papers were selected for the partially edentulous and 14 for the fully edentulous cantilevered restorations. In the present review only those concerning partially edentulous cantilevered restorations were considered (Tables 2, 3 and 4).

3.1 | Excluded studies

The main reason for exclusion of the full text is reported in Table 1. Out of 125 excluded papers, 54 examined prostheses without cantilevers, 39 did not report data about cantilever, 22 had a follow-up less than 5 years, 3 were non-human studies, 3 had number of patients less than 10, 2 were about rehabilitations on natural teeth, 2 had the same pool of patients as other articles with longer follow-up already included in the study. Additional 14 studies were not considered in the present review because they were included in part II (Storelli, Scanferla, Palandrani, Mosca, & Romeo, 2017).

TABLE 10 Technical complications: veneer fractures and decementation/screw loosening (partial prostheses)

Study	No. of ICFDPs available for the analysis (patients)	Mean follow-up time (years)	Total ICFDPs exposure time	No. of veneer fractures	Estimated rate of veneer fractures (per 100 patients/year)	No. of cemented ICFDPs available for loss of retention analysis
Romeo et al. (2009)	59 (45)	8	472	17	4.72	46
Kreissl et al. (2007)	23 (20)	5	115	8	8.00	0
Brägger et al. (2005)	18 (14)	9.4	169.2	1	0.76	13
Wennström et al. (2004)	24 (24)	5	120	1	0.83	0
Cumulative 5-year complication rate prospective studies					18.97% (−9.79, 45.58)	
Jokstad et al. (2017)	24 (24)	17.5	420	4	0.95	0
Malo et al. (2013)	119 (113)	5	595	29	5.13	NR
Aglietta et al. (2012)	21 (21)	5.5	115.5	NR	NE	NR
Hälg et al. (2008)	17 (17)	5	85	4	4.71	17
Eliasson et al. (2006)	71 (71)	9.8	695.8	5	0.72	0
Cumulative 5-year complication rate retrospective studies					11.99% (−4.44, 33.21)	
Total 5-year complication rate Summary estimate (95% CI)*					13.93% (4.52, 27.76)	

Test for heterogeneity $p < 0.75$ for veneers fractures, $p = 0.01$ for loss of retention, $p = 0.06$ for screw loosening.

CI: confidence interval; ICFDPs: implant-supported, cantilever-fixed dental prostheses; NA: not applicable; NE: not estimable; NR: not reported.

3.2 | Study characteristics

Studies were divided into Fixed Partial Denture (Aglietta et al., 2012; Brägger et al., 2005; De Angelis et al., 2017; Eliasson, Eriksson, Johansson, & Wennerberg, 2006; Hälg, Schmid, & Hämmerle, 2008; Jokstad et al., 2017; Kreissl, Gerds, Muche, Heydecke, & Strub, 2007; Malo, de Araujo Nobre, & Lopes, 2013; Romeo, Tomasi, Finini, Casentini, & Lops, 2009; Wennström et al., 2004) and Single Implant supporting two crown (Aglietta et al., 2012; De Angelis et al., 2017). Descriptive data regarding the characteristics of included studied were reported in Tables 2, 3 and 4.

3.3 | Risk of bias

The risk of bias summary is presented in Table 5 and Figure 2 for studies about PCFDP and in Table 6 and Figure 3 for the studies about SICFDP.

Among the studies about PCFDP nine were classified as high risk of bias (Aglietta et al., 2012; Brägger et al., 2005; De Angelis et al., 2017; Eliasson et al., 2006; Hälg et al., 2008; Jokstad et al., 2017; Kreissl et al., 2007; Romeo et al., 2009; Wennström et al., 2004) and one was classified as unclear risk of bias (Malo et al., 2013).

Both studies about SICFDP were classified as high risk of bias (Aglietta et al., 2012; De Angelis et al., 2017).

3.4 | Fixed partial rehabilitations

Nine papers were found to be suitable for fixed partial denture analysis (Aglietta et al., 2012; Brägger et al., 2005; Eliasson et al., 2006;

Hälg et al., 2008; Jokstad et al., 2017; Kreissl et al., 2007; Malo et al., 2013; Romeo et al., 2009; Wennström et al., 2004). One additional study (De Angelis et al., 2017) was included in the single implant analysis but was excluded from the fixed partial denture since less than 10 patients with ICFDP were treated. Five retrospective and four prospective studies were selected: A total of 739 implants supporting 376 rehabilitations in 349 patients were followed for at least 5 years (range 5–17.5). Thirteen implants failed as leading to 10 failed rehabilitations. The estimated survival rate for 5–10 years was calculated to be 98.9% for the implants and 98.2% for the rehabilitations (Table 7). Prospective studies reported a 5–10 years survival rate of 98.93% (96.5, 100, 4% CI) and 96.6% (95.19; 101.1, 95% CI) at implant and prosthesis level, respectively. Retrospective studies reported 5–10 years survival rate of 98.85% (88.4; 101.1, 95% CI) and 99.0% (86.65, 106.0, 95% CI) at implant and prosthesis level, respectively.

A total of 142 complications (mechanical, technical and biological) were reported with a cumulative 5–10 years complication rate of 28.66% (19.56, 55.26, 95% CI) and 26.57% (17.96, 49.24, 95% CI) for the patients and for the prosthesis, respectively (Table 8).

Mechanical complications were reported in 7 studies with a total of 544 implants and 215 rehabilitations (Table 9). Three implant fractures were documented with a cumulative 5–10 years complications rate of 0.31% (−0.97; 3.11). Five cases of abutment screw fracture were documented with a 5–10 years complications rate of 1.57% (−0.05; 3.29).

Technical complications were reported in eight studies (Table 10). Six studies reported on screw retained restorations (160 restorations followed for 5–17.5 years with 16 cases of

Total cemented ICFDPs exposure time	No. of cases of loss of retention	Estimated rate of loss of retention (per 100 ICFDP year)	No. of screw-retained ICFDPs available for screw loosening analysis	Total screw-retained ICFDPs exposure time	No. of cases of screw loosening	Estimated rate of screw loosening (per 100 screw-retained ICFDP year)
368	3	0.83	13	104	0	0
NA	NA	NA	23	115	5	4.35
122.2	2	1.64	5	47	0	0
NA	NA	NA	24	120	2	1.67
		5.10% (-19.92, 32.18)				9.07% (-8.81, 23.85)
NA	NA	NA	24	420	2	0.48
NE	NR	NE	NR	NE	13	NE
NE	NR	NE	NR	NE	NR	NE
85	1	1.18	0	NA	NA	NA
NA	NA	NA	71	695.8	7	1.01
		5.88%				4.03% (-13.13, 20.54)
		5.22% (0.94, 11.15)				5.33% (-2.39, 14.89)

screw loosening), with a cumulative 5–10 years complications rate of 5.33% (-2.39; 14.89). Three studies reported on cemented restorations (76 restorations followed for 5–9.4 years with 6 cases of decementation) with a cumulative 5–10 years complications rate of 5.22% (0.94; 11.15). Eight studies (328 rehabilitations followed for 5–17.5 years) reported 69 cases of veneer fractures with a cumulative 5–10 years complications rate of 13.93% (4.52; 27.76). Six studies reported 0 framework fractures (201 rehabilitations followed for 5–17.5 years) with a cumulative 5–10 years complications rate of 0%.

Biological complications were reported in four studies (Table 11). No study reported on mucositis, instead the data retrieved from four studies showed that peri-implantitis has a cumulative 5–10 years complication rate of 3.68% (-4.84, 13.78) for the implants and 6.06% (-9.53, 24.93) for the prosthesis (95% CI).

MBL was reported in 5 studies with a range from 0.25 to 1.84 mm and an estimated MBL after 5 years of 0.68 mm (-0.15, 1.52, CI 95%) (Table 7).

Three studies (Aglietta et al., 2012; Hälg et al., 2008; Wennström et al., 2004) reported on MBL of implants near to the cantilever and distant from the cantilever. The Forest Plot (Figure 4) reported a summary estimated mean difference in bone loss per year of -0.03 (-0.24, 0.17, CI 95%). The result was not statistically significant ($p = 0.75$).

In three studies rehabilitations were supported by Branemark system implants (387 implants and 214 rehabilitations), in four studies rehabilitations were supported by Straumann dental implants system (225 implants and 115 rehabilitations) and the remaining two studies by 3i Osseotite (61 implants and 23 rehabilitations), Astra

Tech Dental Implant System (66 implants and 24 rehabilitations) (Table 3).

3.5 | Single implant supporting two crowns

Two papers were selected and reported in Table 4, follow-up ranged from 6.5 to 13.6 years (Aglietta et al., 2012; De Angelis et al., 2017). Both studies were retrospective. Three additional studies already included in this review in PICFDP were excluded from the SICFDP analysis. Romeo et al. (2009) had less than 10 patients treated with single implants, Hälg et al. (2008) and Malo et al. (2013) had a mean follow-up for SICFDP < 5 years.

A total of 44 prosthesis supported by 44 implants in 42 patients were analysed. All the rehabilitations supported monolateral cantilevers, 10 distal and 34 mesial. Both studies included either maxillary or mandibular rehabilitations.

Both studies reported on implants and prosthetic failure. Three implants out of 44 and 4 prosthesis out of 44 failed. Two implants were lost due to severe peri-implantitis, one due to implant fracture. Two prostheses failed due to screw fracture, two due to abutment fracture. The estimated 5–10 years survival rate was calculated to be 97.80% (69.85–125.8) and 97.05% (59.57–134.5) for the implants and the prosthesis, respectively (Table 12).

In the paper by Aglietta et al. (2012) data regarding mechanical, technical and biological complications were not reported.

In the study by De Angelis et al. (2017) mechanical, technical and biological complications were reported: four prostheses failed due to abutment or screw fractures, two and two, respectively. The

TABLE 11 Biological complications (partial prostheses)

Study	No. implants/ICFDPs available for the analysis (patients)	Mean follow-up time (years)	Total implants/ICFDPs exposure time	No. of biological complications (peri-implantitis)	Estimated implant/ICFDP complication rate (per 100 implants/prostheses years)
Romeo et al. (2009)	116/59 (45)	8	928/472	2 (2)	0.22/0.42
Kreissl et al. (2007)	61/23 (20)	5	305/115	NR	NE
Brägger et al. (2005)	33/18 (14)	9.4	310.2/169.2	8 (8)	2.58/4.73
Wennström et al. (2004)	66/24 (24)	5	330/130	NR	NR
Cumulative 5-year complication rate prospective studies					4.04% (-68.05, 82.02) 7.80% (-123.8, 149.6)
Jokstad et al. (2017)	58/24 (24)	17.5	1,015/420	NR	NE
Malo et al. (2013)	153/119 (113)	5	765/595	6 (6)	0.78/1.01
Aglietta et al. (2012)	42/21 (21)	5.5	231/115.5	NR	NE
Hälg et al. (2008)	34/17 (17)	5	170/85	0	0.00/0.00
Eliasson et al. (2006)	176/71 (71)	9.8	1,724.8/695.8	NR	NE
Cumulative 5-year complication rate retrospective studies					3.21% (-22.94, 26.06) 4.41 (-29.5, 34.54)
Cumulative 5–10 year complication rate (95% CI)*					3.68% (-4.84, 13.78) 6.06% (-9.53, 24.93)

ICFDPs: implant-supported, cantilever-fixed dental prostheses; NE: not estimable; NR: not reported.

*Based on random effects Poisson regression, test for heterogeneity, $p = 0.80/p = 0.60$.




FIGURE 4 Marginal bone loss for PICFDP

study reported on 16 cemented and 9 screw retained restorations, followed for 10–18 years, with 5 veneer fractures, 2 unscrewed prostheses, 6 decemented prostheses. Two cases of periimplantitis and eight cases of mucositis were reported.

Marginal bone loss was reported in both studies with a range from 0.1 to 2.5 mm.

In one study rehabilitations were supported by Winsix Implants (25 implants and 25 rehabilitations), in one study rehabilitations were supported by Straumann Dental Implants System (19 implants and 19 rehabilitations).

4 | DISCUSSION

The focused question of the present review was “In what clinical situations and with what implant systems are cantilever a successful treatment modality?” Both retrospective and prospective studies were selected, with a minimum follow-up of 5 years and at least 10 patients. Fully edentulous situations treated with implant supported fixed reconstructions with cantilever (FAICFDP) were considered and reported in part II (Storelli, 2018). Partially edentulous patients were divided into cases where one implant was supporting two teeth (SICFDP) and cases of two or more implants were supporting cantilevered prosthesis (PICFDP). A total of 25 papers were selected, of which 14 for the fully edentulous and 11 for the partially edentulous (9 PICFDP and 2 SICFDP). The screening phase was quite complicated by the fact that several papers did not specifically report on cantilever but were showing images and radiographs of cantilevered rehabilitations. Several emails were sent to the authors but the answer was quite scarce and very few authors were able to help in retrieving additional data for this review.

4.1 | Fixed partial rehabilitations

In this systematic review, five retrospective and four prospective studies were selected: a total of 739 implants supporting 376 rehabilitations in 349 patients were followed for at least 5 years (range 5–17.5). The estimated survival rate after 5–10 years was calculated to be 98.4% for the implants and 99.2% for the rehabilitations. A previous systematic review focused on PICFDP (Romeo & Storelli, 2012) reported an estimated survival rate of prospective and retrospective studies of 95.4% and 98.2% for the rehabilitations. Another systematic review (Pjetursson, Thoma, Jung, Zwahlen, & Zembic, 2012) assessed the survival rate of cantilevered

and non-cantilevered partial rehabilitations: the survival rate of the prostheses was calculated to be 95.4% at 5 years. The survival rate of PICFDP rehabilitations appears to be similar to that of non-cantilevered restorations.

Complications of PICFDP were calculated to be in the present review at 5–10 years to be 26.6% for the rehabilitations. This is in agreement with the review of Pjetursson et al. (2012) which assessed that the success rate (i.e., the complications free patients) were 66.4% at 5 years. Although many complications can be considered as minor, it must be stressed the fact that these complications indeed occur and must be accounted for. Among the complications, implant fracture and peri-implantitis can be considered two major ones. In the current review, three implant fractures were documented with a cumulative 5–10 years complications rate of 0.31% (–0.97; 3.11). Pjetursson et al. (2012) calculated that the cumulative incidence of implant fractures was 0.5% at 5 years. Although a small figure, this incident needs to be addressed by clinician and manufacturers. In partially edentulous sites ceramic was the most used veering material. This choice is probably due to the aesthetic results that dental technician can obtain with ceramics. The chipping rate evaluated in the present paper was 18.9% in prospective studies and 11.9% in retrospective studies with follow-up ranging from 5 to 17 years. Resin veneering was reported only in two included studies. In a previous review, Ceramic chipping in implant supported fixed partial denture was calculated to be 8.8% at 5 years, while resin fractures were up to 15.7% at 5 years (Pjetursson et al., 2007).

In the current review, no studies reported on mucositis, while the data retrieved from four studies showed that peri-implantitis has a cumulative 5–10 years complication rate of 6.06% for the prosthesis. In Pjetursson review (2012), the cumulative rate of biological complications after 5 years for implant supported fixed partial dentures was 8.5%.

In the current review, MBL was reported in 5 studies with a range from 0.25 to 1.84 mm and a calculated estimated MBL after 5 years of 0.54 mm (–0.15, 1.52, CI 95%).

The summary estimated mean difference in bone loss between implants close to and distant from cantilevers (reported in three studies only) is –0.03 mm per year (–0.24, 0.17, CI 95%, *p*: 0.75). Similar results were obtained from Aglietta et al. (2009) and Romeo and Storelli (2012), their review reported a summary estimate mean difference in bone loss per year of 0.033 (0.02–0.087, CI 95%, *p*: 0.14). All the authors reported that there is no statistically significant difference in bone loss between implants close to and distant from cantilevers.

TABLE 12 Annual failure rates and survival of implants/ICFDPs (single implant)

Study	No. implants/ICFDPs available for the analysis (patients)	Mean follow-up time (years)	No. failures implants/ICFDPs	Total implants/ICFDPs exposure time	Estimated failure rate (per 100 implants/ICFDPs years)	Estimated implant/ICFDP survival rate after 5–10 years
De Angelis et al. (2017)	25/25 (25)	13.6	3/4	340	0.88/1.18	95.6%/94.1%
Aglietta et al. (2012)	19/19 (17)	6.5	0/0	123.5	0.00/0.00	100%/100%
Total Summary estimate (95% CI)*						97.80 (69.85;125.8) /97.05 (59.57; 134.5)

ICFDPs: implant-supported, cantilever-fixed dental prostheses.

*Based on random effects Poisson regression; test for heterogeneity, not calculable.

4.2 | Single implant supporting two crowns

In this systematic review, two retrospective studies were selected: a total of 44 implants supporting 44 rehabilitations in 42 patients were followed for at least 5 years (range 5–17.5). Estimated survival rate after 5–10 years was 97.80%(69.85;125.8) for implants and 97.05 (59.57; 134.5) for the rehabilitations. Romeo et al. (2009) analysed less than 10 cases of single implant supporting cantilevered prostheses and was therefore excluded, as well as Hälg et al. (2008) and Malo et al. (2013), whose mean follow-ups were less than 5 years.

In a recent systematic review (Van Nimwegen, Raghoobar, Tymstra, Vissink, & Meijer, 2017), single implant supporting two crowns were analysed. The review included five studies with a mean follow-up <5 years, none of which met the inclusion criteria in the present review. Survival rate ranged from 96.6% to 100% up to 3 years. In the present review, not enough data were retrieved about prosthetic and biological complications. Therefore, no conclusion can be drawn at the moment concerning complications in these clinical situations.

One limitation of the present review was that studies with different designs (both retrospective and prospective studies) were selected and analysed together. This was done in order to consider the widest possible amount of data available for analysis but might have contributed to increase heterogeneity of the datasets.

5 | CONCLUSIONS

On the basis of the present review, it is possible to acknowledge the use of cantilevered rehabilitations in partially edentulous patients. Implant-supported restorations with cantilever appear to be able to provide a high survival rate of the restorations in partially edentulous patients. Complications single implant supporting 2-unit cantilever appear to have scarce evidence concerning the survival and rate of complications.

ORCID

Stefano Storelli  <http://orcid.org/0000-0002-7412-0596>

REFERENCES

- Agliardi, E. L., Romeo, D., Panigatti, S., de Araújo Nobre, M., & Maló, P. (2017). Immediate full-arch rehabilitation of the severely atrophic maxilla supported by zygomatic implants: A prospective clinical study with minimum follow-up of 6 years. *International Journal of Oral and Maxillofacial Surgery*, 46(12), 1592–1599. <https://doi.org/10.1016/j.ijom.2017.05.023>
- Aglietta, M., Iorio Siciliano, V., Blasi, A., Sculean, A., Brägger, U., Lang, N. P., & Salvi, G. E. (2012). Clinical and radiographic changes at implants supporting single-unit crowns (SCs) and fixed dental prostheses (FDPs) with one cantilever extension. A retrospective study. *Clinical Oral Implants Research*, 23(5), 550–555. <https://doi.org/10.1111/j.1600-0501.2011.02391.x>
- Aglietta, M., Siciliano, V. I., Zwahlen, M., Brägger, U., Pjetursson, B. E., Lang, N. P., & Salvi, G. E. (2009). A systematic review of the survival

- and complication rates of implant supported fixed dental prostheses with cantilever extensions after an observation period of at least 5 years. *Clinical Oral Implants Research*, 20, 441–451. <https://doi.org/10.1111/j.1600-0501.2009.01706.x>
- Ahrén, S., & Kahnberg, K. E. (2001). The adaptation of implant-supported superstructures to the alveolar crest: A follow-up of 49 cases. *Implant Dentistry*, 10(3), 172–177. <https://doi.org/10.1097/00008505-200107000-00006>
- Allen, P. F., McMillan, A. S., & Walshaw, D. (2001). A patient-based assessment of implant-stabilized and conventional complete dentures. *The Journal of Prosthetic Dentistry*, 85(2), 141–147. <https://doi.org/10.1067/mpr.2001.113214>
- Al-Nawas, B., Kämmerer, P. W., Morbach, T., Ladwein, C., Wegener, J., & Wagner, W. (2012). Ten-year retrospective follow-up study of the TiOblast dental implant. *Clinical Implant Dentistry and Related Research*, 14(1), 127–134. <https://doi.org/10.1111/j.1708-8208.2009.00237.x>
- Arvidson, K., Bystedt, H., Frykholm, A., von Konow, L., & Lothigius, E. (1998). Five-year prospective follow-up report of the Astra Tech Dental Implant System in the treatment of edentulous mandibles. *Clinical Oral Implants Research*, 9(4), 225–234. <https://doi.org/10.1034/j.1600-0501.1998.090403.x>
- Astrand, P., Ahlqvist, J., Gunne, J., & Nilson, H. (2008). Implant treatment of patients with edentulous jaws: A 20-year follow-up. *Clinical Implant Dentistry and Related Research*, 10(4), 207–217.
- Åstrand, P., Engquist, B., Dahlgren, S., Gröndahl, K., Engquist, E., & Feldmann, H. (2004). Astra Tech and Brånemark system implants: A 5-year prospective study of marginal bone reactions. *Clinical Oral Implants Research*, 15, 413–420. <https://doi.org/10.1111/j.1600-0501.2004.01028.x>
- Ata-Ali, J., Flichy-Fernández, A. J., Alegre-Domingo, T., Ata-Ali, F., Palacio, J., & Peñarocha-Diago, M. (2015). Clinical, microbiological, and immunological aspects of healthy versus peri-implantitis tissue in full arch reconstruction patients: A prospective cross-sectional study. *BMC Oral Health*, 15, 43. <https://doi.org/10.1186/s12903-015-0031-9>
- Attard, N., & Zarb, G. A. (2002). Implant prosthodontic management of posterior partial edentulism: Long-term follow-up of a prospective study. *The Journal of the Canadian Dental Association*, 68(2), 118–124.
- Attard, N. J., & Zarb, G. A. (2003). Implant prosthodontic management of partially edentulous patients missing posterior teeth: The Toronto experience. *The Journal of Prosthetic Dentistry*, 89(4), 352–359. <https://doi.org/10.1067/mpr.2003.91>
- Attard, N. J., & Zarb, G. A. (2004). Long-term treatment outcomes in edentulous patients with implant-fixed prostheses: The Toronto study. *The International Journal of Prosthodontics*, 17(4), 425–433.
- Balshi, S. F., Wolfinger, G. J., & Balshi, T. J. (2005). A prospective study of immediate functional loading, following the Teeth in a Day protocol: A case series of 55 consecutive edentulous maxillas. *Clinical Implant Dentistry and Related Research*, 7(1), 24–31. <https://doi.org/10.1111/j.1708-8208.2005.tb00043.x>
- Becker, C. M. (2004). Cantilever fixed prostheses utilizing dental implants: A 10-year retrospective analysis. *Quintessence International*, 35(6), 437–441.
- Becker, C. M., & Kaiser, D. A. (2000). Implant-retained cantilever fixed prosthesis: Where and when. *The Journal of Prosthetic Dentistry*, 84(4), 432–435. <https://doi.org/10.1067/mpr.2000.110259>
- Blanes, R. J., Bernard, J. P., Blanes, Z. M., & Belser, U. C. (2007). A 10-year prospective study of ITI dental implants placed in the posterior region. I: Clinical and radiographic results. *Clinical Oral Implants Research*, 18(6), 699–706. <https://doi.org/10.1111/j.1600-0501.2006.01306.x>
- Botticelli, D., Renzi, A., Lindhe, J., & Berglundh, T. (2008). Implants in fresh extraction sockets: A prospective 5-year follow-up clinical study. *Clinical Oral Implants Research*, 19(12), 1226–1232. <https://doi.org/10.1111/j.1600-0501.2008.01620.x>
- Brägger, U., Aeschlimann, S., Bürgin, W., Hämmerle, C. H., & Lang, N. P. (2001). Biological and technical complications and failures with fixed partial dentures (FPD) on implants and teeth after four to five years of function. *Clinical Oral Implants Research*, 12(1), 26–34. <https://doi.org/10.1034/j.1600-0501.2001.012001026.x>
- Brägger, U., Karoussis, I., Persson, R., Pjetursson, B., Salvi, G., & Lang, N. (2005). Technical and biological complications/failures with single crowns and fixed partial dentures on implants: A 10-year prospective cohort study. *Clinical Oral Implants Research*, 16(3), 326–334. <https://doi.org/10.1111/j.1600-0501.2005.01105.x>
- Brånemark, P. I., Svensson, B., & van Steenberghe, D. (1995). Ten-year survival rates of fixed prostheses on four or six implants ad modum Brånemark in full edentulism. *Clinical Oral Implants Research*, 6(4), 227–231. <https://doi.org/10.1034/j.1600-0501.1995.060405.x>
- Brocard, D., Barthet, P., Baysse, E., Duffort, J. F., Eller, P., Justumus, P., ... Brunel, G. (2000). A multicenter report on 1,022 consecutively placed ITI implants: A 7-year longitudinal study. *International Journal of Oral and Maxillofacial Implants*, 15(5), 691–700.
- Brosky, M. E., Koriath, T. W., & Hodges, J. (2003). The anterior cantilever in the implant-supported screw-retained mandibular prosthesis. *Journal of Prosthetic Dentistry*, 89(3), 244–249. <https://doi.org/10.1067/mpr.2003.43>
- Browaeys, H., Defrancq, J., Dierens, M. C., Miremadi, R., Vandeweghe, S., Van de Velde, T., & De Bruyn, H. (2011). A retrospective analysis of early and immediately loaded osseointegrated implants in cross-arch rehabilitations in edentulous maxillas and mandibles up to 7 years. *Clinical Implant Dentistry and Related Research*, 13(3), 380–389.
- Carlson, B., & Carlsson, G. E. (1994). Prosthodontic complications in osseointegrated dental implant treatment. *International Journal of Oral and Maxillofacial Implants*, 9(1), 90–94. <https://doi.org/10.1097/00008505-199412000-00012>
- Cavalli, N., Austoni, C., Corbella, S., Taschieri, S., Barbaro, B., Azzola, F., & Francetti, L. (2016). Retrospective analysis of the prevalence of peri-implant diseases in non-smoking patients rehabilitated with a fixed full-arch restoration, supported by two mesial axial and two distal tilted implants. *Minerva Stomatologica*, 65(3), 164–175.
- Cavalli, N., Corbella, S., Taschieri, S., & Francetti, L. (2015). Prevalence of Peri-Implant Mucositis and Peri-Implantitis in Patients Treated with a Combination of Axial and Tilted Implants Supporting a Complete Fixed Denture. *ScientificWorldJournal*, 2015, 874842. <https://doi.org/10.1155/2015/874842>
- Chausu, G., & Schwartz-Arad, D. (1999). Full-arch restoration of the jaw with fixed ceramo-metal prosthesis: Late implant placement. *Journal of Periodontology*, 70(1), 90–94. <https://doi.org/10.1902/jop.1999.70.1.90>
- Chiapasco, M., Zaniboni, M., & Boisco, M. (2006). Augmentation procedures for the rehabilitation of deficient edentulous ridges with oral implants. *Clinical Oral Implants Research*, 17(S2), 136–159. <https://doi.org/10.1111/j.1600-0501.2006.01357.x>
- Cordaro, L., Ercoli, C., Rossini, C., Torsello, F., & Feng, C. (2005). Retrospective evaluation of complete-arch fixed partial dentures connecting teeth and implant abutments in patients with normal and reduced periodontal support. *Journal of Prosthetic Dentistry*, 94(4), 313–320. <https://doi.org/10.1016/j.prosdent.2005.08.007>
- Correia, F., Gouveia, S., Felino, A. C., Costa, A. L., & Almeida, R. F. (2017). Survival Rate of Dental Implants in Patients with History of Periodontal Disease: A Retrospective Cohort Study. *International Journal of Oral and Maxillofacial Implants*, 32(4), 927–934. <https://doi.org/10.11607/jomi.3732>
- Costa, R. S., Santos, P. A., Nary, H. F., & Brånemark, P. I. (2015). Key biomechanical characteristics of complete-arch fixed mandibular prostheses supported by three implants developed at P-I Brånemark Institute, Bauru. *The International Journal of Oral & Maxillofacial Implants*, 30(6), 1400–1404. <https://doi.org/10.11607/jomi.3944>

- Davis, D. M., Packer, M. E., & Watson, R. M. (2003). Maintenance requirements of implant-supported fixed prostheses opposed by implant-supported fixed prostheses, natural teeth, or complete dentures: A 5-year retrospective study. *The International Journal of Prosthodontics*, 16(5), 521–523.
- Davó, R. (2009). Zygomatic implants placed with a two-stage procedure: A 5-year retrospective study. *European Journal of Oral Implantology*, 2(2), 115–124.
- De Angelis, F., Papi, P., Mencio, F., Rosella, D., Di Carlo, S., & Pompa, G. (2017). Implant survival and success rates in patients with risk factors: Results from a long-term retrospective study with a 10 to 18 years follow-up. *European Review for Medical and Pharmacological Sciences*, 21(3), 433–437.
- Degidi, M., Iezzi, G., Perrotti, V., & Piattelli, A. (2009). Comparative analysis of immediate functional loading and immediate nonfunctional loading to traditional healing periods: A 5-year follow-up of 550 dental implants. *Clinical Implant Dentistry and Related Research*, 11(4), 257–266. <https://doi.org/10.1111/j.1708-8208.2008.00117.x>
- Degidi, M., Nardi, D., & Piattelli, A. (2013). A six-year follow-up of full-arch immediate restorations fabricated with an intraoral welding technique. *Implant Dentistry*, 22(3), 224–231. <https://doi.org/10.1097/ID.0b013e31829261ed>
- Degidi, M., & Piattelli, A. (2005). 7-year follow-up of 93 immediately loaded titanium dental implants. *Journal of Oral Implantology*, 31(1), 25–31. <https://doi.org/10.1563/0-730.1>
- Del Fabbro, M., Bellini, C. M., Romeo, D., & Francetti, L. (2012). Tilted Implants for the Rehabilitation of Edentulous Jaws: A Systematic Review. *Clinical Implant Dentistry and Related Research*, 14(4), 612–621. <https://doi.org/10.1111/j.1708-8208.2010.00288.x>
- Deporter, D., Ogiso, B., Sohn, D. S., Ruljancich, K., & Pharoah, M. (2008). Ultrashort sintered porous-surfaced dental implants used to replace posterior teeth. *Journal of Periodontology*, 79(7), 1280–1286. <https://doi.org/10.1902/jop.2008.070496>
- Ekelund, J. A., Lindquist, L. W., Carlsson, G. E., & Jemt, T. (2003). Implant treatment in the edentulous mandible: A prospective study on Brånemark system implants over more than 20 years. *The International Journal of Prosthodontics*, 16(6), 602–608.
- Eckfeldt, A., Christiansson, U., Eriksson, T., Lindén, U., Lundqvist, S., Rundcrantz, T., ... Billström, C. (2001). A retrospective analysis of factors associated with multiple implant failures in maxillae. *Clinical Oral Implants Research*, 12(5), 462–467. <https://doi.org/10.1034/j.1600-0501.2001.120505.x>
- Eckfeldt, A., Zellmer, M., & Carlsson, G. E. (2013). Treatment with implant-supported fixed dental prostheses in patients with congenital and acquired neurologic disabilities: A prospective study. *The International Journal of Prosthodontics*, 26(6), 517–524. <https://doi.org/10.11607/jip.3511>
- Eliasson, A., Eriksson, T., Johansson, A., & Wennerberg, A. (2006). Fixed partial prostheses supported by 2 or 3 implants: A retrospective study up to 18 years. *International Journal of Oral and Maxillofacial Implants*, 21(4), 567–574.
- Eliasson, A., Narby, B., Ekstrand, K., Hirsch, J., Johansson, A., & Wennerberg, A. (2010). A 5-year prospective clinical study of submerged and nonsubmerged Paragon system implants in the edentulous mandible. *The International Journal of Prosthodontics*, 23(3), 231–238.
- Eliasson, A., Palmqvist, S., Svenson, B., & Sondell, K. (2000). Five-year results with fixed complete-arch mandibular prostheses supported by 4 implants. *International Journal of Oral and Maxillofacial Implants*, 15(4), 505–510.
- Engstrand, P., Gröndahl, K., Ohnell, L. O., Nilsson, P., Nannmark, U., & Brånemark, P. I. (2003). Prospective follow-up study of 95 patients with edentulous mandibles treated according to the Brånemark Novum concept. *Clinical Implant Dentistry and Related Research*, 5(1), 3–10. <https://doi.org/10.1111/j.1708-8208.2003.tb00176.x>
- Esposito, M., Grusovin, M. G., Worthington, H. V., & Coulthard, P. (2006). Interventions for replacing missing teeth: Bone augmentation techniques for dental implant treatment (Review). *The Cochrane Database of Systematic Reviews*, 25, CD003607.
- Esposito, M., Maghaireh, H., Pistilli, R., Grusovin, M.G., Lee, S.T., Trullenque-Eriksson, A., & Gualini, F. (2016). Dental implants with internal versus external connections: 5-year post-loading results from a pragmatic multicenter randomised controlled trial. *European Journal of Oral Implantology*, 9Suppl 1(2):129–141.
- Ferrigno, N., Laureti, M., Fanali, S., & Grippaudo, G. (2002). A long-term follow-up study of non-submerged ITI implants in the treatment of totally edentulous jaws. Part I: Ten-year life table analysis of a prospective multicenter study with 1286 implants. *Clinical Oral Implants Research*, 13(3), 260–273. <https://doi.org/10.1034/j.1600-0501.2002.130305.x>
- Fischer, K., & Stenberg, T. (2013). Prospective 10-year cohort study based on a randomized, controlled trial (RCT) on implant-supported full-arch maxillary prostheses. part II: Prosthetic outcomes and maintenance. *Clinical Implant Dentistry and Related Research*, 15(4), 498–508. <https://doi.org/10.1111/j.1708-8208.2011.00383.x>
- Fischer, K., Stenberg, T., Hedin, M., & Sennerby, L. (2008). Five-year results from a randomized, controlled trial on early and delayed loading of implants supporting full-arch prosthesis in the edentulous maxilla. *Clinical Oral Implants Research*, 19(5), 433–441. <https://doi.org/10.1111/j.1600-0501.2007.01510.x>
- Fortin, Y., Sullivan, R. M., & Rangert, B. R. (2002). The Marius implant bridge: Surgical and prosthetic rehabilitation for the completely edentulous upper jaw with moderate to severe resorption: A 5-year retrospective clinical study. *Clinical Implant Dentistry and Related Research*, 4(2), 69–77. <https://doi.org/10.1111/j.1708-8208.2002.tb00155.x>
- Francetti, L., Rodolfi, A., Barbaro, B., Taschieri, S., Cavalli, N., & Corbella, S. (2015). Implant success rates in full-arch rehabilitations supported by upright and tilted implants: A retrospective investigation with up to five years of follow-up. *Journal of Periodontal and Implant Science*, 45(6), 210–215. <https://doi.org/10.5051/jpis.2015.45.6.210>
- Francetti, L., Romeo, D., Corbella, S., Taschieri, S., & Del Fabbro, M. (2012). Bone level changes around axial and tilted implants in full-arch fixed immediate restorations. Interim results of a prospective study. *Clinical Implant Dentistry and Related Research*, 14(5), 646–654. <https://doi.org/10.1111/j.1708-8208.2010.00304.x>
- Friberg, B., Gröndahl, K., Lekholm, U., & Brånemark, P. I. (2000). Long-term follow-up of severely atrophic edentulous mandibles reconstructed with short Brånemark implants. *Clinical Implant Dentistry and Related Research*, 2(4), 184–189. <https://doi.org/10.1111/j.1708-8208.2000.tb00116.x>
- Friberg, B., Raghoebar, G. M., Grunert, I., Hobkirk, J. A., & Tepper, G. (2008). A 5-year prospective multicenter study on 1-stage smooth-surface Brånemark System implants with early loading in edentulous mandibles. *International Journal of Oral and Maxillofacial Implants*, 23(3), 481–486.
- Glauser, R., Zembic, A., Ruhstaller, P., & Windisch, S. (2007). Five-year results of implants with an oxidized surface placed predominantly in soft quality bone and subjected to immediate occlusal loading. *Journal of Prosthetic Dentistry*, 97(6 Suppl), S59–S68. [https://doi.org/10.1016/S0022-3913\(07\)60009-2](https://doi.org/10.1016/S0022-3913(07)60009-2)
- Gotfredsen, K. (1997). Implant supported overdentures—the Copenhagen experience. *Journal of Dentistry*, 25(Suppl 1), S39–S42. [https://doi.org/10.1016/S0300-5712\(97\)87699-3](https://doi.org/10.1016/S0300-5712(97)87699-3)
- Gualini, F., Gualini, G., Cominelli, R., & Lekholm, U. (2009). Outcome of Brånemark Novum implant treatment in edentulous mandibles: A retrospective 5-year follow-up study. *Clinical Implant Dentistry and Related Research*, 11(4), 330–337. <https://doi.org/10.1111/j.1708-8208.2008.00118.x>

- Haas, R., Mendorff-Pouilly, N., Mailath, G., & Bernhart, T. (1998). Five-year results of maxillary intramobile Zylinder implants. *British Journal of Oral and Maxillofacial Surgery*, 36(2), 123–128. [https://doi.org/10.1016/S0266-4356\(98\)90180-2](https://doi.org/10.1016/S0266-4356(98)90180-2)
- Hälgl, G. A., Schmid, J., & Hämmerle, C. H. (2008). Bone level changes at implants supporting crowns or fixed partial dentures with or without cantilevers. *Clinical Oral Implants Research*, 19(10), 983–990. <https://doi.org/10.1111/j.1600-0501.2008.01556.x>
- Hartman, G. A., & Cochran, D. L. (2004). Initial implant position determines the magnitude of crestal bone remodeling. *Journal of Periodontology*, 75(4), 572–577. <https://doi.org/10.1902/jop.2004.75.4.572>
- Hellem, S., Karlsson, U., Almfeldt, I., Brunell, G., Hamp, S. E., & Astrand, P. (2001). Nonsubmerged implants in the treatment of the edentulous lower jaw: A 5-year prospective longitudinal study of ITI hollow screws. *Clinical Implant Dentistry and Related Research*, 3(1), 20–29. <https://doi.org/10.1111/j.1708-8208.2001.tb00125.x>
- Hemmings, K. W., Schmitt, A., & Zarb, G. A. (1994). Complications and maintenance requirements for fixed prostheses and overdentures in the edentulous mandible: A 5-year report. *International Journal of Oral and Maxillofacial Implants*, 9(2), 191–196.
- Heschl, A., Payer, M., Platzer, S., Wegscheider, W., Pertl, C., & Lorenzoni, M. (2012). Immediate rehabilitation of the edentulous mandible with screw type implants: Results after up to 10 years of clinical function. *Clinical Oral Implants Research*, 23(10), 1217–1223. <https://doi.org/10.1111/j.1600-0501.2011.02292.x>
- Ibañez, J. C., Tahhan, M. J., Zamar, J. A., Menendez, A. B., Juaneda, A. M., Zamar, N. J., & Monqaut, J. L. (2005). Immediate occlusal loading of double acid-etched surface titanium implants in 41 consecutive full-arch cases in the mandible and maxilla: 6- to 74-month results. *Journal of Periodontology*, 76(11), 1972–1981. <https://doi.org/10.1902/jop.2005.76.11.1972>
- Imburgia, M., & Del Fabbro, M. (2015). Long-Term Retrospective Clinical and Radiographic Follow-up of 205 Brånemark System Mk III TiUnite Implants Submitted to Either Immediate or Delayed Loading. *Implant Dentistry*, 24(5), 533–540. <https://doi.org/10.1097/ID.0000000000000265>
- Isaksson, R., Becktor, J. P., Brown, A., Laurizohn, C., & Isaksson, S. (2009). Oral health and oral implant status in edentulous patients with implant-supported dental prostheses who are receiving long-term nursing care. *Gerodontology*, 26(4), 245–249. <https://doi.org/10.1111/j.1741-2358.2009.00275.x>
- Jaffin, R., Kolesar, M., Kumar, A., Ishikawa, S., & Fiorellini, J. (2007). The radiographic bone loss pattern adjacent to immediately placed, immediately loaded implants. *International Journal of Oral and Maxillofacial Implants*, 22(2), 187–194.
- Jemt, T., & Lekholm, U. (1995). Implant treatment in edentulous maxillae: A 5-year follow-up report on patients with different degrees of jaw resorption. *International Journal of Oral and Maxillofacial Implants*, 10(3), 303–311.
- Jokstad, A., Gussgard, A. M., Fava, J., Lin, M., Shokati, B., Somogyi-Ganss, E., ... Zahran, M. (2017). Benchmarking Outcomes in Implant Prosthodontics: Partial Fixed Dental Prostheses and Crowns Supported by Implants with a Turned Surface over 10 to 28 Years at the University of Toronto. *The International Journal of Oral & Maxillofacial Implants*, 32(4), 880–892. <https://doi.org/10.11607/jomi.5454>
- Keller, E. E., Tolman, D., & Eckert, S. (1998). Endosseous implant and autogenous bone graft reconstruction of mandibular discontinuity: A 12-year longitudinal study of 31 patients. *International Journal of Oral and Maxillofacial Implants*, 13(6), 767–780.
- Kim, S. S., Yeo, I. S., Lee, S. J., Kim, D. J., Jang, B. M., Kim, S. H., & Han, J. S. (2013). Clinical use of alumina-toughened zirconia abutments for implant-supported restoration: Prospective cohort study of survival analysis. *Clinical Oral Implants Research*, 24(5), 517–522. <https://doi.org/10.1111/j.1600-0501.2011.02413.x>
- Kirkwood, B. R., & Sterne, J. A. C. (2003). *Essential Medical Statistics*. Oxford, UK: Blackwell Science.
- Koller, C. D., Pereira-Cenci, T., & Boscatto, N. (2016). Parameters associated with marginal bone loss around implant after prosthetic loading. *Brazilian Dental Journal*, 27(3), 292–297. <https://doi.org/10.1590/0103-6440201600874>
- Kreissl, M. E., Gerdts, T., Muche, R., Heydecke, G., & Strub, J. R. (2007). Technical complications of implant-supported fixed partial dentures in partially edentulous cases after an average observation period of 5 years. *Clinical Oral Implants Research*, 18(6), 720–726. <https://doi.org/10.1111/j.1600-0501.2007.01414.x>
- Krennmair, G., Seemann, R., Schmidinger, S., Ewers, R., & Piehslinger, E. (2010). Clinical outcome of root-shaped dental implants of various diameters: 5-year results. *International Journal of Oral and Maxillofacial Implants*, 25(2), 357–366.
- Kucey, B. K. (1997). Implant placement in prosthodontics practice: A five-year retrospective study. *Journal of Prosthetic Dentistry*, 77(2), 171–176. [https://doi.org/10.1016/S0022-3913\(97\)70231-2](https://doi.org/10.1016/S0022-3913(97)70231-2)
- Kupeyan, H. K., & Clayton, J. A. (2004). The lingual arm cantilever. *Journal of Prosthetic Dentistry*, 92(1), 90–92. <https://doi.org/10.1016/j.prosdent.2004.03.012>
- Lai, H. C., Zhang, Z. Y., Zhuang, L. F., Wang, F., Liu, X., & Pu, Y. P. (2008). Early loading of ITI implants supporting maxillary fixed full-arch prostheses. *Clinical Oral Implants Research*, 19(11), 1129–1134. <https://doi.org/10.1111/j.1600-0501.2008.01563.x>
- Lam, W. Y., Botelho, M. G., & McGrath, C. P. (2013). Longevity of implant crowns and 2-unit cantilevered resin-bonded bridges. *Clinical Oral Implants Research*, 24(12), 1369–1374. <https://doi.org/10.1111/clr.12034>
- Lambrecht, J. T., Filippi, A., Künzel, A. R., & Schiel, H. J. (2003). Long-term evaluation of submerged and nonsubmerged ITI solid-screw titanium implants: A 10-year life table analysis of 468 implants. *The International Journal of Oral & Maxillofacial Implants*, 18(6), 826–834.
- Lee, J. H., Kweon, H. H., Choi, S. H., & Kim, Y. T. (2016). Association between dental implants in the posterior region and traumatic occlusion in the adjacent premolars: A long-term follow-up clinical and radiographic analysis. *Journal of Periodontal and Implant Science*, 46(6), 396–404. <https://doi.org/10.5051/jpis.2016.46.6.396>
- Lee, J. Y., Park, H. J., Kim, J. E., Choi, Y. G., Kim, Y. S., Huh, J. B., & Shin, S. W. (2011). A 5-year retrospective clinical study of the Dentium implants. *The Journal of Advanced Prosthodontics*, 3(4), 229–235. <https://doi.org/10.4047/jap.2011.3.4.229>
- Lethaus, B., Kälber, J., Petrin, G., Brandstätter, A., & Weingart, D. (2011). Early loading of sandblasted and acid-etched titanium implants in the edentulous mandible: A prospective 5-year study. *The International Journal of Oral & Maxillofacial Implants*, 26(4), 887–892.
- Liberati, A., Altman, D. G., Tetzlaff, J., Mulrow, C., Gøtzsche, P. C., Ioannidis, J. P., ... Moher, D. (2009). The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: Explanation and elaboration. *Annals of Internal Medicine*, 151, W65–W94. <https://doi.org/10.7326/0003-4819-151-4-200908180-00136>
- Makkonen, T. A., Holmberg, S., Niemi, L., Olsson, C., Tammisalo, T., & Peltola, J. (1997). A 5-year prospective clinical study of Astra Tech dental implants supporting fixed bridges or overdentures in the edentulous mandible. *Clinical Oral Implants Research*, 8(6), 469–475. <https://doi.org/10.1034/j.1600-0501.1997.080605.x>
- Malo, P., de Araujo Nobre, M. A., Guedes, C. M., & Almeida, R. (2017). Outcomes of immediate function implant prosthetic restorations with mechanical complications: A retrospective clinical study with 5 years of follow-up. *The European Journal of Prosthodontics and Restorative Dentistry*, 25(1), 26–34.
- Malo, P., de Araujo Nobre, M., & Lopes, A. (2013). The prognosis of partial implant-supported fixed dental prostheses with cantilevers. A 5-year

- retrospective cohort study. *European Journal of Oral Implantology*, 6(1), 51–59.
- Malo, P., de Araújo Nobre, M., Lopes, A., Moss, S. M., & Molina, G. J. (2011). A longitudinal study of the survival of All-on-4 implants in the mandible with up to 10 years of follow-up. *Journal of the American Dental Association*, 142(3), 310–320. <https://doi.org/10.14219/jada.archive.2011.0170>
- Mangano, F., Macchi, A., Caprioglio, A., Sammons, R. L., Piattelli, A., & Mangano, C. (2014). Survival and complication rates of fixed restorations supported by locking-taper implants: A prospective study with 1 to 10 years of follow-up. *Journal of Prosthodontics*, 23(6), 434–444. <https://doi.org/10.1111/jopr.12152>
- Mangano, C., Mangano, F., Shibli, J. A., Tettamanti, L., Figliuzzi, M., d'Avila, S., ... Piattelli, A. (2011). Prospective evaluation of 2,549 Morse taper connection implants: 1- to 6-year data. *Journal of Periodontology*, 82(1), 52–61. <https://doi.org/10.1902/jop.2010.100243>
- McAlarney, M. E., & Stavropoulos, D. N. (2000). Theoretical cantilever lengths versus clinical variables in fifty-five clinical cases. *Journal of Prosthetic Dentistry*, 83(3), 332–343. [https://doi.org/10.1016/S0022-3913\(00\)70137-5](https://doi.org/10.1016/S0022-3913(00)70137-5)
- Merckse-Stern, R., Aerni, D., Geering, A. H., & Buser, D. (2001). Long-term evaluation of non-submerged hollow cylinder implants. Clinical and radiographic results. *Clinical Oral Implants Research*, 12(3), 252–259. <https://doi.org/10.1034/j.1600-0501.2001.012003252.x>
- Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *Journal of Clinical Epidemiology*, 62, 1006–1012. <https://doi.org/10.1016/j.jclinepi.2009.06.005>
- Mundt, T., Heinemann, F., Schwahn, C., & Biffar, R. (2012). Retrievable, tooth-implant-supported, complete-arch fixed restorations in the maxilla: A 6-year retrospective study. *Biomedizinische Technik/Biomedical Engineering*, 57(1), 39–43. <https://doi.org/10.1515/bmt-2011-0033>
- Mura, P. (2012). Immediate loading of tapered implants placed in postextraction sockets: Retrospective analysis of the 5-year clinical outcome. *Clinical Implant Dentistry and Related Research*, 14(4), 565–574. <https://doi.org/10.1111/j.1708-8208.2010.00297.x>
- Murphy, W. M., Absi, E. G., Gregory, M. C., & Williams, K. R. (2002). A prospective 5-year study of two cast framework alloys for fixed implant-supported mandibular prostheses. *The International Journal of Prosthodontics*, 15(2), 133–138.
- Naert, I., Koutsikakis, G., Quirynen, M., Duyck, J., van Steenberghe, D., & Jacobs, R. (2002). Biologic outcome of implant-supported restorations in the treatment of partial edentulism. Part 2: A longitudinal radiographic study. *Clinical Oral Implants Research*, 13(4), 390–395. <https://doi.org/10.1034/j.1600-0501.2002.130407.x>
- Naert, I., Quirynen, M., van Steenberghe, D., & Darius, P. (1992). A study of 589 consecutive implants supporting complete fixed prostheses. Part II: Prosthetic aspects. *Journal of Prosthetic Dentistry*, 68(6), 949–956. [https://doi.org/10.1016/0022-3913\(92\)90557-Q](https://doi.org/10.1016/0022-3913(92)90557-Q)
- Nedir, R., Bischof, M., Szmukler-Moncler, S., Belser, U. C., & Samson, J. (2006). Prosthetic complications with dental implants: From an up-to-8-year experience in private practice. *The International Journal of Oral & Maxillofacial Implants*, 21(6), 919–928.
- Niedermaier, R., Stelzle, F., Riemann, M., Bolz, W., Schuh, P., & Wachtel, H. (2017). Implant-Supported Immediately Loaded Fixed Full-Arch Dentures: Evaluation of Implant Survival Rates in a Case Cohort of up to 7 Years. *Clinical Implant Dentistry and Related Research*, 19(1), 4–19. <https://doi.org/10.1111/cid.12421>
- Noack, N., Willer, J., & Hoffmann, J. (1999). Long-term results after placement of dental implants: Longitudinal study of 1,964 implants over 16 years. *The International Journal of Oral & Maxillofacial Implants*, 14(5), 748–755.
- Ortorp, A., & Jemt, T. (2009). Early laser-welded titanium frameworks supported by implants in the edentulous mandible: A 15-year comparative follow-up study. *Clinical Implant Dentistry and Related Research*, 11(4), 311–322. <https://doi.org/10.1111/j.1708-8208.2008.00119.x>
- Ortorp, A., & Jemt, T. (2012). CNC-milled titanium frameworks supported by implants in the edentulous jaw: A 10-year comparative clinical study. *Clinical Implant Dentistry and Related Research*, 14(1), 88–99. <https://doi.org/10.1111/j.1708-8208.2009.00232.x>
- Ozgun, G. O., Kazancioglu, H. O., Demirtas, N., Deger, S., & Ak, G. (2016). Risk Factors Associated With Implant Marginal Bone Loss: A Retrospective 6-Year Follow-Up Study. *Implant Dentistry*, 25(1), 122–127. <https://doi.org/10.1097/ID.0000000000000366>
- Ozkan, Y., Akoğlu, B., & Kulak-Ozkan, Y. (2011). Five-year treatment outcomes with four types of implants in the posterior maxilla and mandible in partially edentulous patients: A retrospective study. *The International Journal of Oral & Maxillofacial Implants*, 26(3), 639–647.
- Parein, A. M., Eckert, S. E., Wollan, P. C., & Keller, E. E. (1997). Implant reconstruction in the posterior mandible: A long-term retrospective study. *Journal of Prosthetic Dentistry*, 78(1), 34–42. [https://doi.org/10.1016/S0022-3913\(97\)70085-4](https://doi.org/10.1016/S0022-3913(97)70085-4)
- Pettersson, P., & Sennerby, L. (2015). A 5-year retrospective study on Replace Select Tapered dental implants. *Clinical Implant Dentistry and Related Research*, 17(2), 286–295. <https://doi.org/10.1111/cid.12105>
- Pjetursson, B. E., Bragger, U., Lang, N. P., & Zwahlen, M. (2007). Comparison of survival and complication rates of tooth-supported fixed dental prostheses (FDPs) and implant-supported FDPs and single crowns (SCs). *Clinical Oral Implants Research*, 18(Suppl 3), 97–113. <https://doi.org/10.1111/j.1600-0501.2007.01439.x>
- Pjetursson, B. E., Thoma, D., Jung, R., Zwahlen, M., & Zembic, A. (2012). A systematic review of the survival and complication rates of implant-supported fixed dental prostheses (FDPs) after a mean observation period of at least 5 years. *Clinical Oral Implants Research*, 23, 22–38. <https://doi.org/10.1111/j.1600-0501.2012.02546.x>
- Quirynen, M., Alsaadi, G., Pauwels, M., Haffajee, A., van Steenberghe, D., & Naert, I. (2005). Microbiological and clinical outcomes and patient satisfaction for two treatment options in the edentulous lower jaw after 10 years of function. *Clinical Oral Implants Research*, 16(3), 277–287. <https://doi.org/10.1111/j.1600-0501.2005.01127.x>
- Raghoebar, G. M., Timmenga, N. M., Reintsema, H., Stegenga, B., & Vissink, A. (2001). Maxillary bone grafting for insertion of endosseous implants: Results after 12–124 months. *Clinical Oral Implants Research*, 12(3), 279–286. <https://doi.org/10.1034/j.1600-0501.2001.012003279.x>
- Rasmusson, L., Roos, J., & Bystedt, H. A. (2005). 10-year follow-up study of titanium dioxide-blasted implants. *Clinical Implant Dentistry and Related Research*, 7, 36–42. <https://doi.org/10.1111/j.1708-8208.2005.tb00045.x>
- Ravald, N., Dahlgren, S., Teiwik, A., & Gröndahl, K. (2013). Long-term evaluation of Astra Tech and Brånemark implants in patients treated with full-arch bridges. Results after 12–15 years. *Clinical Oral Implants Research*, 24(10), 1144–1151. <https://doi.org/10.1111/j.1600-0501.2012.02524.x>
- Romeo, E., Ghisolfi, M., Rozza, R., Chiapasco, M., & Lops, D. (2006). Short (8-mm) dental implants in the rehabilitation of partial and complete edentulism: a 3- to 14-year longitudinal study. *The International Journal of Prosthodontics*, 19(6), 586–592.
- Romeo, E., Lops, D., Amorfini, L., Chiapasco, M., Ghisolfi, M., & Vogel, G. (2006). Clinical and radiographic evaluation of small-diameter (3.3-mm) implants followed for 1–7 years: A longitudinal study. *Clinical Oral Implants Research*, 17(2), 139–148. <https://doi.org/10.1111/j.1600-0501.2005.01191.x>
- Romeo, E., Lops, D., Margutti, E., Ghisolfi, M., Chiapasco, M., & Vogel, G. (2003). Implant-supported fixed cantilever prostheses in partially edentulous arches. A seven-year prospective study. *Clinical Oral Implants Research*, 14(3), 303–311. <https://doi.org/10.1034/j.1600-0501.2003.120905.x>

- Romeo, E., & Storelli, S. (2012). Systematic review of the survival rate and the biological, technical, and aesthetic complications of fixed dental prostheses with cantilevers on implants reported in longitudinal studies with a mean of 5 years follow-up. *Clinical Oral Implants Research*, 23(Suppl 6), 39–49. <https://doi.org/10.1111/j.1600-0501.2012.02551.x>
- Romeo, E., Tomasi, C., Finini, I., Casentini, P., & Lops, D. (2009). Implant-supported fixed cantilever prosthesis in partially edentulous jaws: A cohort prospective study. *Clinical Oral Implants Research*, 20(11), 1278–1285. <https://doi.org/10.1111/j.1600-0501.2009.01766.x>
- Schliephake, H., Schmelzeisen, R., Husstedt, H., & Schmidt-Wondera, L. U. (1999). Comparison of the late results of mandibular reconstruction using nonvascularized or vascularized grafts and dental implants. *Journal of Oral and Maxillofacial Surgery*, 57(8), 944–950. [https://doi.org/10.1016/S0278-2391\(99\)90015-0](https://doi.org/10.1016/S0278-2391(99)90015-0)
- Schnitman, P. A., Wöhrle, P. S., Rubenstein, J. E., DaSilva, J. D., & Wang, N. H. (1997). Ten-year results for Brånemark implants immediately loaded with fixed prostheses at implant placement. *The International Journal of Oral & Maxillofacial Implants*, 12(4), 495–503.
- Schrott, A. R., Jimenez, M., Hwang, J. W., Fiorellini, J., & Weber, H. P. (2009). Five-year evaluation of the influence of keratinized mucosa on peri-implant soft-tissue health and stability around implants supporting full-arch mandibular fixed prostheses. *Clinical Oral Implants Research*, 20(10), 1170–1177. <https://doi.org/10.1111/j.1600-0501.2009.01795.x>
- Schwartz-Arad, D., & Chaushu, G. (1998). Full-arch restoration of the jaw with fixed ceramometal prosthesis. *The International Journal of Oral & Maxillofacial Implants*, 13(6), 819–825.
- Schwartz-Arad, D., Gulayev, N., & Chaushu, G. (2000). Immediate versus non-immediate implantation for full-arch fixed reconstruction following extraction of all residual teeth: A retrospective comparative study. *Journal of Periodontology*, 71(6), 923–928. <https://doi.org/10.1902/jop.2000.71.6.923>
- Sertgoz, A., & Guvener, S. (1996). Finite element analysis of the effect of cantilever and implant length on stress distribution in an implant-supported fixed prosthesis. *Journal of Prosthetic Dentistry*, 76, 165–169. [https://doi.org/10.1016/S0022-3913\(96\)90301-7](https://doi.org/10.1016/S0022-3913(96)90301-7)
- Stegaroiu, R., Sato, T., Kusakari, H., & Miyakawa, O. (1998). Influence of restoration type on stress distribution in bone around implants: A three-dimensional finite element analysis. *International Journal of Oral & Maxillofacial Implants*, 13, 82–90.
- Storelli, S. (2018). Part 1 COIR to be defined with editor.
- Storelli, S., Scanferla, M., Palandrani, G., Mosca, D., & Romeo, E. (2017). Stratification of prosthetic complications by manufacturer in implant-supported restorations with a 5 years' follow-up: Systematic review of the literature. *Minerva Stomatologica*, 66(4), 178–191.
- Sullivan, D. Y., Sherwood, R. L., & Porter, S. S. (2001). Long-term performance of Osseotite implants: A 6-year clinical follow-up. *Compendium of Continuing Education in Dentistry*, 22(4), 326–328, 330, 332–4.
- Sullivan, D., Vincenzi, G., & Feldman, S. (2005). Early loading of Osseotite implants 2 months after placement in the maxilla and mandible: A 5-year report. *The International Journal of Oral & Maxillofacial Implants*, 20(6), 905–912.
- Tartaglia, G. M., Maiorana, C., Gallo, M., Codari, M., & Sforza, C. (2016). Implant-Supported Immediately Loaded Full-Arch Rehabilitations: Comparison of Resin and Zirconia Clinical Outcomes in a 5-Year Retrospective Follow-Up Study. *Implant Dentistry*, 25(1), 74–82. <https://doi.org/10.1097/ID.0000000000000368>
- Tealdo, T., Menini, M., Bevilacqua, M., Pera, F., Pesce, P., Signori, A., & Pera, P. (2014). Immediate versus delayed loading of dental implants in edentulous patients' maxillae: A 6-year prospective study. *The International Journal of Prosthodontics*, 27(3), 207–214. <https://doi.org/10.11607/ijp.3569>
- Tinsley, D., Watson, C. J., & Russell, J. L. (2001). A comparison of hydroxylapatite coated implant retained fixed and removable mandibular prostheses over 4 to 6 years. *Clinical Oral Implants Research*, 12(2), 159–166. <https://doi.org/10.1034/j.1600-0501.2001.012002159.x>
- Vajdovich, I., & Fazekas, A. (1999). A ten-year clinical follow-up study of prosthetic rehabilitation of the edentulous lower jaw with endosteal dental implants. *Journal of Long-Term Effects of Medical Implants*, 9(3), 171–183.
- Van Nimwegen, W. G., Raghoobar, G. M., Tymstra, N., Vissink, A., & Meijer, H. J. A. (2017). How to treat two adjacent missing teeth with dental implants. A systematic review on single implant-supported two-unit cantilever FDP's and results of a 5-year prospective comparative study in the aesthetic zone. *Journal of Oral Rehabilitation*, 44(6), 461–471. <https://doi.org/10.1111/joor.12507>
- Vigolo, P., Givani, A., Majzoub, Z., & Cordioli, G. (2004). Clinical evaluation of small-diameter implants in single-tooth and multiple-implant restorations: A 7-year retrospective study. *The International Journal of Oral & Maxillofacial Implants*, 19(5), 703–709.
- Wang, J. H., Judge, R., & Bailey, D. (2016). A 5-Year Retrospective Assay of Implant Treatments and Complications in Private Practice: The Restorative Complications of Single and Short-Span Implant-Supported Fixed Prostheses. *The International Journal of Prosthodontics*, 29(5), 435–444. <https://doi.org/10.11607/ijp.4794>
- Weng, D., Jacobson, Z., Tarnow, D., Hürzeler, M. B., Faehn, O., Sanavi, F., ... Stach, R. M. (2003). A prospective multicenter clinical trial of 3i machined-surface implants: Results after 6 years of follow-up. *The International Journal of Oral & Maxillofacial Implants*, 18(3), 417–423.
- Wennström, J., Zurdo, J., Karlsson, S., Ekestubbe, A., Gröndahl, K., & Lindhe, J. (2004). Bone level change at implant-supported fixed partial dentures with and without cantilever extension after 5 years in function. *Journal of Clinical Periodontology*, 31(12), 1077–1083. <https://doi.org/10.1111/j.1600-051X.2004.00603.x>
- Wittneben, J. G., Buser, D., Salvi, G. E., Bürgin, W., Hicklin, S., & Brägger, U. (2014). Complication and failure rates with implant-supported fixed dental prostheses and single crowns: A 10-year retrospective study. *Clinical Implant Dentistry and Related Research*, 16(3), 356–364. <https://doi.org/10.1111/cid.12066>
- Wyatt, C. C., & Zarb, G. A. (2002). Bone level changes proximal to oral implants supporting fixed partial prostheses. *Clinical Oral Implants Research*, 13(2), 162–168. <https://doi.org/10.1034/j.1600-0501.2002.130206.x>
- Zampelis, A., Rangert, B., & Heijl, L. (2007). Tilting of splinted implants for improved prosthodontic support: A two-dimensional finite element analysis. *Journal of Prosthetic Dentistry*, 97, S35–S43. [https://doi.org/10.1016/S0022-3913\(07\)60006-7](https://doi.org/10.1016/S0022-3913(07)60006-7)
- Zanolla, J., Amado, F. M., da Silva, W. S., Ayub, B., de Almeida, A. L., & Soares, S. (2016). Success rate in implant-supported overdenture and implant-supported fixed denture in cleft lip and palate patients. *Annals of Maxillofacial Surgery*, 6(2), 223–227.
- Zarb, G. A., & Schmitt, A. (1991). Osseointegration and the edentulous predicament. The 10-year-old Toronto study. *British Dental Journal*, 170(12), 439–444. <https://doi.org/10.1038/sj.bdj.4807583>
- Zarb, G. A., & Schmitt, A. (1993). The longitudinal clinical effectiveness of osseointegrated dental implants in posterior partially edentulous patients. *The International Journal of Prosthodontics*, 6(2), 189–196. <https://doi.org/10.1097/00008505-199312000-00015>
- Zarb, J. P., & Zarb, G. A. (2002). Implant prosthodontic management of anterior partial edentulism: Long-term follow-up of a prospective study. *Journal/Canadian Dental Association. Journal de l'Association Dentaire Canadienne*, 68(2), 92–96.

- Zhang, X. X., Shi, J. Y., Gu, Y. X., & Lai, H. C. (2016). Long-Term Outcomes of Early Loading of Straumann Implant-Supported Fixed Segmented Bridgeworks in Edentulous Maxillae: A 10-Year Prospective Study. *Clinical Implant Dentistry and Related Research*, 18(6), 1227–1237. <https://doi.org/10.1111/cid.12420>
- Zinsli, B., Sägesser, T., & Mericske, E. (2004). Clinical evaluation of small-diameter ITI implants: A prospective study. *International Journal of Oral & Maxillofacial Implants*, 19, 92–99.
- Zurdo, J., Romão, C., & Wennström, J. L. (2009). Survival and complication rates of implant-supported fixed partial dentures with cantilevers: A systematic review. *Clinical Oral Implants Research*, 20(Suppl 4), 59–66. <https://doi.org/10.1111/j.1600-0501.2009.01773.x>

How to cite this article: Storelli S, Del Fabbro M, Scanferla M, Palandrani G, Romeo E. Implant supported cantilevered fixed dental rehabilitations in partially edentulous patients: Systematic review of the literature. Part I. *Clin Oral Impl Res*. 2018;29(Suppl. 18):253–274. <https://doi.org/10.1111/clr.13311>