

## ACCUMULATION OF MACRONUTRIENTS IN FORAGE GRASSES UNDER SALINE AND ALKALINE CONDITIONS

B. Keskin<sup>1</sup>, S. Temel<sup>1</sup>, İ. H. Yılmaz<sup>1</sup> and U. Şimşek<sup>2</sup><sup>1</sup> Iğdır University, Faculty of Agriculture, Department of Agronomy, 76000 IĞDIR<sup>2</sup> Iğdır Üniversitesi, Faculty of Agriculture, Department of Soil Science and Plant Nutrition, 76000 IĞDIR  
Corresponding Author: bilalkeskin66@yahoo.com

## ABSTRACT

This study was designed to determine P, K, Ca, Mg and Na mineral accumulations in *Agropyron elongatum*, *Chloris gayana*, *Cynodon dactylon* and *Festuca arundinacea* species grown on control (non-saline and non-alkaline), highly saline, highly alkaline and highly saline-alkaline soils, and to check whether the obtained fodders meet mineral requirements of animals. The research established in 2011 under a randomized complete blocks design, and continued for three years. Results revealed that macro nutrient contents (P, K, Ca, Mg and Na) accumulation differed significantly among species (except for K), locations (except for Ca) and years. No significant differences were observed among species in terms of potassium content. The highest phosphorus content was detected in *Agropyron elongatum*, while the highest Ca content observed in *A.elongatum* and *F.arundiancaea*. Moreover, the highest Mg content was observed in *F.arundiancaea* and the highest Na content in *A.elongatum* and *F.arundiancaea*. The lowest P, K<sup>+</sup> and Na<sup>+</sup> accumulations were detected from highly saline-alkaline soils while Mg<sup>2+</sup> content was observed in control, highly saline and highly alkaline soils. As for meeting mineral requirements by animals, Ca<sup>2+</sup> content of the obtained fodder was found to be sufficient while K<sup>+</sup> accumulations were found to be lower and P and Na<sup>+</sup> contents, on the other hand, were found to be higher than the recommended levels. Thus, K<sup>+</sup> requirements by the animals should be met by additional feeding and because of rich sodium and phosphorus contents, fodder obtained from these species should be fed with caution.

**Keywords:** Salinity, alkalinity, forage grass, macronutrient.

## INTRODUCTION

Plants are under the influence of, or in interaction with, many vegetative and environmental factors within the locations they grow. One of these environmental stress factors is soil salinity. Salinity, which is widespread in arid and semi-arid regions of the world, is an important environmental stress factor that has negative effects on the socioeconomic development of countries, germination-seedling development, plant diversity and yield (Ghassemi- Golezani *et al.*, 2010; Kazemi and Eskandari, 2011; Temel *et al.*, 2015). It is reported that over 6% of the world's land is affected by either salinity or sodicity (Munns, 2011). Accumulation of excess Na<sup>+</sup> and Cl<sup>-</sup> ions in saline and alkaline areas has negative effects on the growth of many plants due to ion toxicity and osmotic effect (Parida and Das, 2005; Kacar *et al.*, 2009; Ul-Hassan and Bano, 2015). On the other hand, salinity-resistant plants decrease their own osmotic potential by accumulating high amounts of inorganic minerals in their roots and these plants are affected relatively less from salinity by transcending to turgor state in saline areas (Greenway and Munns, 1980).

Plants that can be grown on saline soils may significantly meet nutritional and mineral requirements of animals (Temel *et al.*, 2016). Plant species, fertilization, irrigation, precipitation of the cultivation area, root

system of the plant, pH of the soil, cooperation and antagonism of the elements are important factors that affect mineral accumulation ability of the plants (Marschner, 1995; Bengtsson *et al.*, 2003; Warman and Termeer, 2005). Acidity and alkalinity of the soil have significant effects on the usefulness of nutritional elements in soil and as a result of this effect the amount of mineral content that the plant can acquire may vary (Aydin *et al.*, 2005).

The situation that salinity and alkalinity having effects on mineral content of the soils changes mineral compositions of the plants grown in these areas and, thus, the mineral demands of the animals (Khan *et al.*, 2007). Lack of minerals may cause growth deficiencies and toxicities in both plants and animals (Ulrich and Hills, 1967). Daily nutritional, energy and mineral requirements of animals must be met in order for them to reproduce and grow healthily (Ganskopp and Bohnert, 2003). Otherwise excess or lack of mineral intake may have adverse effects on animal health and reproduction and nutritional disorders in animals (Goswami *et al.*, 2005; Khan *et al.*, 2007). There is a lack of Ca, P and Na content in forage grasses in general. These species may sometimes have a lack of Mg and K content (McDowell, 1996). In this study, P, K, Ca, Mg and Na macronutrient accumulations of 4 forage grasses (*Agropyron elongatum*, *Chloris gayana*, *Cynodon dactylon* and *Festuca arundinacea*) grown on saline and alkaline soils were

determined and whether these accumulations are sufficient for the mineral requirements by animals were evaluated.

## MATERIALS AND METHODS

Trial was conducted in Iğdır where continental climate is dominant. The research was carried out for 3 years between 2011 and 2013 at 4 different locations of Iğdır plain where 39.9578 latitude, 44.3018 longitude and

800-900 m altitude. Mean annual temperature is 12.5 °C, average relative humidity is 51.2% and annual amount of precipitation is 264.0 mm. Mean temperatures in 2011, 2012 and 2013 were recorded as 12.6 °C, 13.5 °C and 14.1 °C, respectively, while relative humidity was recorded as 56.5%, 53.6% and 51.4% respectively and annual amounts of precipitation were 340.0 mm, 237.2 mm and 226.9 mm for the trial years (Table 1).

**Table 1. Some climate features of Iğdır province in 2011, 2012 and 2013 years (Anonymous, 2014).**

Months	Monthly Total Precipitation (mm)			Temperature (°C)			Monthly Avg. Relative Humidity (%)		
	2011	2012	2013	2011	2012	2013	2011	2012	2013
January	6,0	0,0	19,6	-0,6	0,3	-2,2	74,0	59,5	71,9
February	22,6	12,5	15,2	0,7	-3,1	4,3	66,1	63,1	64,3
March	17,2	13,5	14,8	7,8	3,6	9,3	48,5	47,1	44,3
April	73,9	16,2	34,6	13,5	16,1	15,4	57,7	43,4	46,1
May	76,9	57,4	58,9	17,5	19,7	18,9	59,8	51,3	52,6
June	40,4	26,7	38,3	23,5	25	23,3	47,0	37,6	43,7
July	24,0	23,0	10,6	28,0	26,1	26,5	40,5	43,8	39,7
August	24,3	0,6	8,3	25,9	27,6	35,4	43,2	38,0	41,7
September	10,6	29,3	9,9	21,4	21,5	21,4	45,8	47,7	43,2
October	25,8	11,5	15,4	12,6	15,4	12,3	60,0	62,5	53,8
November	9,2	20,7	1,3	3,0	8,4	2,4	63,7	75,0	57,3
December	9,1	25,8	7,6	-1,6	1,4	-8,7	71,6	74,5	80,88
<b>Total/Avg.</b>	<b>340,0</b>	<b>237,2</b>	<b>226,9</b>	<b>12,64</b>	<b>13,5</b>	<b>14,06</b>	<b>56,49</b>	<b>53,62</b>	<b>51,39</b>
<b>3 years Avg.</b>		<b>268,03</b>			<b>13,4</b>			<b>53,83</b>	

Chemical and physical characteristics of soil samples (0-30 cm depth) taken from in the trial area are shown in table 2. Textural analyses were carried out using Bouyoucus Hydrometer (Gee and Hortage, 1986); organic matter counts were made according to Walkey-Black method (Nelson and Sommers, 1982); soil pH was measured with a Glass-Electrode pH-meter (McLean, 1982); exchangeable Na, K, Ca and Mg contents were measured by reading in Atomic Absorption after extracting by shaking with ammonium acetate (1 N, pH=7.0) (Rhoades, 1982); phosphorus content was measured by spectrophotometer in subtiles extracted by sodium bicarbonate (Olsen and Sommers, 1982); electrical conductivity (EC), was measured by EC tool in extraction solution (Demiralay, 1993) and boron analysis was carried out by reading color solutions prepared according to azometin-h method in spectrophotometer whose light absorption set at 420 nm wave length (John *et al.*, 1975).

Trial was established in 2011 and carried out for 3 (three) years (2011, 2012 and 2013) in a randomized complete blocks design with three replications. (Four locations were selected in trial; control (non-alkaline and non-saline) (DSY 8.9%, EC 0.43 dS/m), highly saline

(DSY 11.9%, EC 9.80 dS/m), highly alkaline (DSY 60.5%, EC 0.89 dS/m), and highly saline-alkaline (DSY 49.7%, EC 9.08 dS/m), 4 (four) forage grasses (*Cynodon dactylon* L. var. Sem-Caska., *Chloris gayana* Kunth var. Katambora, *Festuca arundinacea* L. var. Asterix and *Agropyron elongatum* L) were selected as materials for determination of resilience under saline and alkaline conditions. Sowings were carried out on April 20, 2011 and 30 kg, 6 kg, 20 kg and 15 kg/ha<sup>-1</sup> seeds were used, respectively, of *Chloris gayana*, *Cynodon dactylon*, *Festuca arundinacea* and *Agropyron elongatum* species. 60 kg ha<sup>-1</sup> N (300 kg ha<sup>-1</sup> ammonium sulphate) and 100 kg ha<sup>-1</sup> P<sub>2</sub>O<sub>5</sub> (250 kg ha<sup>-1</sup> TSP) were applied as fertilizer during the sowings. After every harvest period in each year (2011, 2012 and 2013) an additional amount of 60 kg ha<sup>-1</sup> N (300 kg ha<sup>-1</sup> ammonium sulphate) fertilizer was applied. Moisture content of the soil was determined with "Soil Water Potential Measurement Device" and irrigation was made when the amount of useful water dropped under 50%. Irrigation was carried out 7 times in the first year, 8 times in the second year and 8 times in the third year according to the border irrigation method.

*Cynodon dactylon* was harvested at full bloom stage while *Festuca arundinacea* at boot stage,

*Agropyron elongatum* at pre-anthesis (Serin and Tan, 1998) and *Chloris gayana* at the beginning of panicle generation stage (Skerman and Riveros, 1990). Harvested plants were first washed with purified water. After that, plants were first left in the open air and dried for 2-3 then dried in a drying oven set at 70 °C for 48 hours. After drying, samples were ground in a Wiley grinder in order to enable them pass through 1 mm of sieve opening and

ground plant material were taken to wet decomposition in solutions prepared with nitric acid mixture. Mineral matter contents were then determined using ICP-OES device after wet decomposition process (Kacar and Inal, 2008).

The data were exposed to General Linear Models with SPSS on the basis of main effects. Mean was performed using Duncan test.

**Table 2. Chemical and physical characteristics of the experimental soils.**

Soil Properties	EXPERIMENTAL LOCATIONS			
	Control	High saline	High alkali	High saline-alkali
Texture	Clay loam	Loamy sand	Clay loam	Clay loam
EC (dS/m)	0.43	9.80	0.89	9.08
pH (1:2.5)	8.2	8.5	10.3	9.4
Organic matter (%)	4.4	2.1	1.7	2.3
ESP (%)	8.9	11.9	60.5	49.7
N (%)	0.21	0.11	0.11	0.08
P (mg kg <sup>-1</sup> )	27.9	33.8	40.8	36.5
B (mg kg <sup>-1</sup> )	4.3	12.4	5.9	11.4
Ca (mg kg <sup>-1</sup> )	3640	3680	3180	3400
Mg (mg kg <sup>-1</sup> )	528	540	444	552
K (mg kg <sup>-1</sup> )	1248	1326	1638	1248
Na (mg kg <sup>-1</sup> )	552	759	3749	2737

ESP: exchangeable sodium percentage

## RESULTS AND DISCUSSION

It was observed that all seedlings that have germinated and surfaced under highly saline-alkaline soil conditions in the year of establishment were dried and consequently could not manage to develop in a full plant. Upon this development, species were re-sown in the second year at locations where they did not grow and only *Agropyron elongatum* was able to be planted. This situation showed that *Agropyron elongatum* species is more resilient to highly saline-alkaline soils when compared to other studied species. It was reported that soil salinity and alkalinity affect germination and seedling development of the plants and the extent of this effect may vary according to species (Gu *et al.*, 2012; Temel *et al.*, 2015). *Chloris gayana* and *Cynodon dactylon* species showed a significant growth in the year of establishment, however, they could not continue this growth within the next year probably because of low winter temperatures. Tansi (2009) reported that *Cynodon dactylon* dies at -3 °C while, *Chloris gayana* dies at -8 °C which are in parallel with the findings of this study. Because of such reasons, plant samples could not be taken in some years and from some locations.

**PHOSPHORUS:** Phosphorus (P) accumulations of studied forage grasses grown at locations with different chemical characteristics differed significantly between years, locations and species. As for the years, phosphorus

content of the obtained fodder varied between 0.91% and 1.53% and higher phosphorus contents were recorded in the first and second years (2011 and 2012) when compared to 2013. In general, temperature, light, ventilation, soil pH and mutual interactions of ions are the basic factors that have an influence on intake of nutritional elements by plants (Kacar and Katkat, 2015). Therefore, one or a few of these factors may have an effect on the varying mineral contents among the plants. Looking at the locations, the highest phosphorus contents were observed in control, highly saline and highly alkaline soil conditions while the lowest value was recorded in highly saline-alkaline location (Table 3).

It may be said that the reason for the decrease in phosphorus contents of fodder obtained under highly saline-alkaline conditions is the negative effect of sodium, which is abundant in such soils, on phosphorus intake. It has been reported that increase of soil salinity, especially of Na<sup>+</sup> amount in soils, results in a decrease of phosphorus intake, in other words, in a decrease of P amount in plant tissues (Unlukara *et al.*, 2008). Phosphorus content varied between 1.25% and 1.59% among species. It can be seen in table 3 that *C.dactylon*, *C.gayana* and *F.arundianacea* species fell within the same statistical group in terms of phosphorus content and these species have accumulated higher amounts of phosphorus within their structures than *A.elongatum*. Cacanet *al.* (2015) reports that mineral compositions of 12 forage grasses collected from natural areas with

neutral pH and non-saline soil characteristics have varied, and *F.arundinacea* species have the highest phosphorus content among all investigated species. Other researches

have also reported variations in phosphorus contents of fodder crops as to species and varieties (Abdullah *et al.*, 2013; Bayraktar, 2005).

**Table 3. Phosphorus contents (%) of forage grasses cultivated under different soil conditions.**

Years	Plant	Control	High Saline	High Alkali	High Saline-alkali	Average of Years
2011	<i>Cynodon dactylon</i>	1.58	1.59	1.59	x	<b>1.53 a</b>
	<i>Chloris gayana</i>	1.60	1.59	x	x	
	<i>Agropyron elongatum</i>	1.01	1.59	1.60	x	
	<i>Festuca arundinacea</i>	1.60	1.58	x	x	
2012	<i>Cynodon dactylon</i>	X	X	x	x	<b>1.58 a</b>
	<i>Chloris gayana</i>	X	X	x	x	
	<i>Agropyron elongatum</i>	1.59	1.43	1.61	1.44	
	<i>Festuca arundinacea</i>	1.65	1.75	x	x	
2013	<i>Cynodon dactylon</i>	X	X	x	x	<b>0.91 b</b>
	<i>Chloris gayana</i>	X	X	x	x	
	<i>Agropyron elongatum</i>	1.02	0.91	0.89	0.69	
	<i>Festuca arundinacea</i>	1.10	0.87	x	x	
	<b>Average of location</b>	<b>1.39 a</b>	<b>1.41 a</b>	<b>1.42 a</b>	<b>1.07 b</b>	
<b>Average of plants</b>	<i>Cynodon dactylon</i>	<b>1.59 a</b>				
	<i>Chloris gayana</i>	<b>1.59 a</b>				
	<i>Agropyron elongatum</i>	<b>1.25 b</b>				
	<i>Festuca arundinacea</i>	<b>1.42 a</b>				

\*: Plants didnot growth at applications shown with X.

\*\* : Values showed with different letters are significantly different

**POTASSIUM:** Location and years have significantly affected  $K^+$  content (Table 4). Maximum  $K^+$  content was recorded in 2012 (0.44%) and  $K^+$  content in the year of establishment (0.30%) was observed to be higher than that of 2013(0.19%) (Table 4). This may be a result of lower temperatures during 2011 in comparison with 2012. Hence, it has been reported that increasing temperatures during the growth periods of many fodder crops caused an increase of  $K^+$  accumulation in plants (Reid and Horvath, 1980). Potassium content of the fodder varied between 0.21% and 0.34% among locations. The highest potassium level was observed in highly saline conditions followed by control and these two locations were in the same statistical group in terms of  $K^+$  content. The lowest value was determined in highly saline alkaline soil conditions followed by highly alkaline (Table 4). Thus,  $K^+$  content of the obtained fodder showed a significant decrease under highly alkaline and highly saline conditions. This situation is thought to be stemming from the antagonistic effect of  $Na^+$  ions which are abundantly available both in the soil and in plant structure. Hence, Gunes *et al.* (2005) and Kacar and Katkat (2015) have expressed that excess amounts of  $Na^+$  and other ions in soil decrease  $K^+$  intake by plants. A study by Mezni *et al.* (2010) reported the negative effect of increasing amounts of  $Na^+$  on  $K^+$  content of alfalfa as well. Results of other research also showed that  $K^+$  intake was inhibited in plants under salt stress and in high doses

of NaCl applications (Haro *et al.*, 2010; Samad and Karmoker, 2013; Turan *et al.*, 2010).

**CALCIUM:** Location has no significant effect on  $Ca^{2+}$  content, species and years were found to have a significant effect on  $Ca^{2+}$  content (Table 5). Maximum  $Ca^{2+}$  accumulation in the study were measured in 2012. Minimum values were observed in 2011 and 2013 and these two study years were observed to be in the same statistical group (Table 5). This may be caused by abundance of relatively more  $Na^+$  ions in soil during 2011 due to no prior soil processing and cultural activities when compared to 2012 (Table 2). Grattan and Grieve (1999) reported that existence of high amounts of  $Na^+$  concentration in root environment has a negative effect on  $Ca^{2+}$  intake by the plants. Highest  $Ca^{2+}$  accumulation was determined in *F.arundinacea* while the lowest value was found in *C.dactylon*. This may be a result of higher proportions of  $Na^+$  ions found within the structure of *C.dactylon* in comparison with *F.arundinacea* (Table 7). Excess amounts of Na ions in plant structure decrease  $Ca^{2+}$  intake by the plant due to antagonism of ions (Grattan and Grieve, 1990).

**MAGNESIUM:**  $Mg^{2+}$  contents of the obtained fodder showed significant differences for years, locations and species. According to table 6, significant increases were recorded in  $Mg^{2+}$  contents of the fodder in years following the year of establishment and the highest

values were observed during 2013 (0.33%). This may be a result of low amount of precipitation and higher temperatures during 2013 when compared to other trial years, since increasing evaporation ratios due to increasing temperatures causes transfer of salts that are more soluble in soil to the upper layers of soil through capillarity (Ashraf *et al.*, 2005). As a consequence plants tend to intake more mineral ions within their structure (Aydemir and Sunger, 2012). As for the locations, a higher  $Mg^{2+}$  accumulation was observed in highly saline-alkaline soil conditions in comparison with the other locations (Table 6). It is thought that this is a result of antagonistic effects of  $Na^+$  ions found abundantly in soils. Thus, Suyama *et al.* (2007) have reported that  $Mg^{2+}$  content of clover increased in parallel to increasing  $NaCl_2$  salinity. Yet, Semiz *et al.* (2012) have reported that  $Mg^{2+}$  content of fennel leaves subjected to increasing amounts of salt ( $NaCl_2$  and  $CaCl_2$ ) decreased due to increasing salt concentration. On the other hand, the lowest  $Mg^{2+}$  content was observed in highly alkaline soils while control and highly saline soils fell within the same statistical group (Table 6). A similar result was also obtained by Gonzales and Hailman (1977) who reported that there was no difference in  $Mg^{2+}$  contents of *Chloris gayana* and *Cynodon dactylon* species grown under saline and non-saline soil conditions. The highest  $Mg^{2+}$  content was obtained in *F.arundinacea* with 0.31%, while *C.dactylon* and *C.gayana* species had the lowest  $Mg^{2+}$  content (Table 6). In another study conducted by Kilic *et al.* (2015) under non-saline, lightly saline, fairly (medium) saline and highly saline soil conditions, it is reported that *F.arundinacea* species has the highest  $Mg^{2+}$  content among all fodder crops species (except *Lolium perenne*) included in the study. Moreover, results of other studies conducted in different ecologies have shown that  $Mg^{2+}$  contents accumulated within the structure of fodder-crop species grown on same soil type vary (0.03%-0.38%) (Cacan *et al.*, 2015; Elmali and Kaya, 2012).

**SODIUM:** Sodium content has changed significantly according to years, locations and species. As shown in table 7, the highest  $Na^+$  ratio was measured during 2011 and significant decreases in  $Na^+$  content were observed in years following the year of establishment. This may be stemming from the improvement of soil conditions due to carrying-out of more intensive cultural activities (fertilizing, hoeing and irrigation) on soils, which have not been subjected to any previous soil processing, in years following the year of establishment and, as a result of this situation, lower concentrations of  $Na^+$  in soils (particularly in the root area). Hence, prior research also reports that  $Na^+$  content in both roots and offshoots of many plant species increases in relation to increasing soil salinity (Majid *et al.*, 2012; Wang *et al.*, 2012).  $Na^+$  content has varied between 0.73% and 0.32% among

locations and the highest  $Na^+$  content was determined in highly alkaline (0.73%) soil conditions (Table 7). This may be resulting from the existence of relatively higher amount of  $Na^+$  ions in highly alkaline soil conditions ( $3749 \text{ mg kg}^{-1}$ ) and relatively higher amounts of  $Ca^{2+}$  ions in control conditions (Table 2), when compared to other locations, since  $Na^+$  intake by the plants may decrease depending on soil  $Ca^{2+}$  concentration in soil solution (Cramer, 1997). The lowest  $Na^+$  value, on the other hand, was detected in highly saline alkaline conditions (Table 7). However, in another similar study, the highest  $Na^+$  ion concentration in plants was shown to be obtained under saline-alkaline soil conditions (Aydemir and Sunger, 2012). This result is not in agreement with our findings. Other species except *A.elongatum* could not manage to establish under highly saline-alkaline conditions due to climatic conditions or due to reasons stemming from the soil itself and therefore did not contribute statistically to the means and it is thought that this was the reason for the disagreement with findings of previously-conducted research. In the study, *C.dactylon* and *C.gayana*, *A. elongatum* and *F. arundinacea* fell into same statistical groups in terms of  $Na^+$  accumulation and warm season forage grasses *C. dactylon* (1.10%) and *C.gayana* (1.13%) observed to have higher  $Na^+$  content than cool season forage grasses *A. elongatum* (0.47%) and *F.arundinacea* (0.51%). In a study by Sima *et al.* (2013) conducted to determine the effects of different salt types (sodium chloride, sodium sulphate, potassium chloride and potassium sulphate) on growth and development of different forage grass species, it is reported that, *A.elongatum*, a species which has one the highest salinity-resistance levels among fodder crops, has a lower  $Na^+$  concentration in comparison with other species which are more sensible to salinity. On the other hand, Deifel *et al.* (2006) have reported that *C.gayana*, which is known as the most salinity-tolerant species, can accumulate higher proportions of  $Na^+$  ions within its structure through osmotic modification even under low salinity conditions and this result are in support of our findings.

In addition, whether fodder obtained from forage grasses grown under different soil conditions meets the mineral requirements by animals was also investigated in the study. As reported by NRC (1985), recommended ratios of P,  $K^+$ ,  $Ca^{2+}$ ,  $Mg^{2+}$ ,  $Na^+$  for all ruminant classes are 1.25 to 1.59%, 0.5 to 1.0%, 0.19 to 0.82%, 0.12 to 0.20% and 0.06 to 0.18% , respectively. In our present study, average P, K, Ca, Mg and Na contents of the plants varied between 0.12% to 0.48%, 0.28% to 0.35%, 0.24% to 0.42%, 0.11% to 0.31% and 0.47% to 1.13%. According to these findings,  $Ca^{2+}$  content of the investigated fodder-crop species was found to be sufficient for mineral requirements of animals, while  $K^+$  accumulation was found to be lower and P and  $Na^+$  contents were found to be higher than recommended

levels. On the other hand, Mg<sup>2+</sup> contents of the species differed than those of recommended by NRC (1985) for all ruminants. Only *F.arundinacea* species found to have a high Mg<sup>2+</sup>content. Therefore, keeping in mind that feeds with high mineral content can cause toxicity in

animals, feeds must be given with caution (for example, by rationing with other feeds with low mineral content). In this case, it would be appropriate to supplement animal diets with additional minerals in order to meet with animal mineral requirements.

**Table 4. Potassium contents (%) of forage grasses cultivated under different soil conditions.**

Years	Plants	Control	High Saline	High Alkali	High Saline-alkali	Average of Years
2011	<i>Cynodon dactylon</i>	0.40	0.31	0.25	X	<b>0.30 b</b>
	<i>Chloris gayana</i>	0.29	0.27	x	X	
	<i>Agropyron elongatum</i>	0.15	0.38	0.25	x	
	<i>Festuca arundinacea</i>	0.30	0.34	x	x	
2012	<i>Cynodon dactylon</i>	X	X	x	x	<b>0.44 a</b>
	<i>Chloris gayana</i>	X	X	x	x	
	<i>Agropyron elongatum</i>	0.40	0.51	0.42	0.31	
	<i>Festuca arundinacea</i>	0.50	0.52	x	x	
2013	<i>Cynodon dactylon</i>	X	X	x	x	<b>0.19 c</b>
	<i>Chloris gayana</i>	X	X	x	x	
	<i>Agropyron elongatum</i>	0.21	0.20	0.21	0.12	
	<i>Festuca arundinacea</i>	0.21	0.18	x	x	
	<b>Average of location</b>	<b>0.31 ab</b>	<b>0.34 a</b>	<b>0.27 bc</b>	<b>0.21 c</b>	
	<i>Cynodon dactylon</i>	<b>0.32</b>				
<b>Average of plants</b>	<i>Chloris gayana</i>	<b>0.28</b>				
	<i>Agropyron elongatum</i>	<b>0.28</b>				
	<i>Festuca arundinacea</i>	<b>0.35</b>				

\*: Plants did not growth at applications shown with X.

\*\* : Values showed with different letters are significantly different

**Table 6. Calcium contents (%) of forage grasses cultivated under different soil conditions.**

Years	Plant	Control	High Saline	High Alkali	High Saline-alkali	Average of Years
2011	<i>Cynodon dactylon</i>	0.13	0.27	0.31	x	<b>0.27 b</b>
	<i>Chloris gayana</i>	0.33	0.26	X	x	
	<i>Agropyron elongatum</i>	0.20	0.29	0.36	x	
	<i>Festuca arundinacea</i>	0.30	0.21	X	x	
2012	<i>Cynodon dactylon</i>	X	x	X	x	<b>0.54 a</b>
	<i>Chloris gayana</i>	X	x	X	x	
	<i>Agropyron elongatum</i>	0.52	0.49	0.55	0.44	
	<i>Festuca arundinacea</i>	0.83	0.45	X	x	
2013	<i>Cynodon dactylon</i>	X	x	X	x	<b>0.32 b</b>
	<i>Chloris gayana</i>	X	x	X	x	
	<i>Agropyron elongatum</i>	0.22	0.34	0.27	0.34	
	<i>Festuca arundinacea</i>	0.35	0.39	X	x	
	<b>Average of location</b>	<b>0.36</b>	<b>0.34</b>	<b>0.37</b>	<b>0.39</b>	
	<i>Cynodon dactylon</i>	<b>0.24 c</b>				
<b>Average of plants</b>	<i>Chloris gayana</i>	<b>0.29 bc</b>				
	<i>Agropyron elongatum</i>	<b>0.36 ab</b>				
	<i>Festuca arundinacea</i>	<b>0.42 a</b>				

\*: Plants did not growth at applications shown with X.

\*\* : Values showed with different letters are significantly different

**Table 6. Magnesium contents (%) of forage grasses cultivated under different soil conditions.**

Years	Plant	Control	High Saline	High Alkali	High Saline-alkali	Average of Years
2011	<i>Cynodon dactylon</i>	0.09	0.12	0.11	x	<b>0.11 c</b>
	<i>Chloris gayana</i>	0.14	0.12	x	x	
	<i>Agropyron elongatum</i>	0.08	0.11	0.14	x	
	<i>Festuca arundinacea</i>	0.13	0.08	x	x	
2012	<i>Cynodon dactylon</i>	X	X	x	x	<b>0.25 b</b>
	<i>Chloris gayana</i>	X	X	x	x	
	<i>Agropyron elongatum</i>	0.18	0.21	0.23	0.23	
	<i>Festuca arundinacea</i>	0.39	0.27	x	x	
2013	<i>Cynodon dactylon</i>	X	X	x	x	<b>0.33 a</b>
	<i>Chloris gayana</i>	X	X	x	x	
	<i>Agropyron elongatum</i>	0.15	0.24	0.18	0.37	
	<i>Festuca arundinacea</i>	0.46	0.55	x	x	
	<b>Average of location</b>	<b>0.20 b</b>	<b>0.21 b</b>	<b>0.16 c</b>	<b>0.30 a</b>	
	<i>Cynodon dactylon</i>	<b>0.11 c</b>				
<b>Average of plants</b>	<i>Chloris gayana</i>	<b>0.13 c</b>				
	<i>Agropyron elongatum</i>	<b>0.19 b</b>				
	<i>Festuca arundinacea</i>	<b>0.31 a</b>				

\*: Plants did not growth at applications shown with X.

\*\*: Values showed with different letters are significantly different

**Table 7. Sodium contents (%) of forage grasses cultivated under different soil conditions.**

Years	Plant	Control	High Saline	High Alkali	High Saline-alkali	Average of Years
2011	<i>Cynodon dactylon</i>	1.12	1.13	1.06	x	<b>1.08 a</b>
	<i>Chloris gayana</i>	1.15	1.11	x	x	
	<i>Agropyron elongatum</i>	0.73	1.11	1.15	x	
	<i>Festuca arundinacea</i>	1.05	1.14	x	x	
2012	<i>Cynodon dactylon</i>	X	X	x	x	<b>0.46 b</b>
	<i>Chloris gayana</i>	X	X	x	x	
	<i>Agropyron elongatum</i>	0.40	0.35	0.64	0.58	
	<i>Festuca arundinacea</i>	0.36	0.42	x	x	
2013	<i>Cynodon dactylon</i>	X	X	x	x	<b>0.048 c</b>
	<i>Chloris gayana</i>	X	X	x	x	
	<i>Agropyron elongatum</i>	0.033	0.045	0.07	0.06	
	<i>Festuca arundinacea</i>	0.036	0.041	x	x	
	<b>Average of location</b>	<b>0.61 c</b>	<b>0.67 b</b>	<b>0.73 a</b>	<b>0.32 d</b>	
	<i>Cynodon dactylon</i>	<b>1.10 a</b>				
<b>Average of plants</b>	<i>Chloris gayana</i>	<b>1.13 a</b>				
	<i>Agropyron elongatum</i>	<b>0.47 b</b>				
	<i>Festuca arundinacea</i>	<b>0.51 b</b>				

\*: Plants did not growth at applications shown with X.

\*\*: Values showed with different letters are significantly different

**Conclusion:** According to the findings of the study, salinity and alkalinity have significant effects on the mineral contents of the investigated plants. In general, cool season forage grasses had higher mineral contents than the warm season forage grasses. In terms of meeting the mineral requirements by animals, Ca<sup>2+</sup> contents of the obtained fodder are found to be sufficient, while K<sup>+</sup> accumulations are found to be lower and P and

Na<sup>+</sup> contents were found to be higher than recommended levels. Therefore, it is recommended to meet K<sup>+</sup> requirements by animals through additional feeding and feeds should be given to animals with caution due to high sodium and phosphorus content.

**Acknowledgements:** This research is financed by TUBİTAK (The Scientific and Technological Research

Council of Turkey) with Project No. 110 O 259. We would like to thank the Council for their contribution to this research.

## REFERENCES

- Abdullah, M., R.A. Khan, S. Yaqoob, and M. Ahmad (2013). Mineral profile of browse species used as feed by grazing livestock in cholistan rangelands, Pakistan. *Pakistan J. Nutr.* 12 (2):135-143.
- Anonymous, (2014). Iğdır Provincial Directorate of Meteorology. Iğdir, Turkey.
- Ashraf, M.Y., K. Akhtar, G. Sarwar, and M. Ashraf (2005). Role of rooting system in salt tolerance potential of different guar accessions. *Agron. Sustain. Dev.* 25(2): 243-249.
- Aydemir, A. and H. Sunger (2012). Growth response and ameliorative effect of a forage plant (*Festuca arundinacea*) in calcareous saline-sodic soils. *Afr. J. Agric. Res.* 7(5): 802-809.
- Aydin, A., C. Kant, and N. Ataoglu (2005). Effect of boron and phosphorus application on the growth and mineral content of corn in erzurum and rize soils. *Atatürk Univ. J. Agric. Fac.* 36 (2): 125-129.
- Bayraktar, E. (2005). The change of food substances that are collected in the roots and the stem of some forage crops in conditions of Tekirdag. *Trakya University Fen Bilimleri Enstitüsü, Tarla Bitkileri Anabilim Dalı, Yüksek Lisans Tezi, Tekirdag.*
- Bengtsson H., I. Öborn, S. Jonsson, I. Nilsson and A. Andersson (2003). Field balances of some mineral nutrients and trace elements in organic and conventional dairy farming – a case study at Öjebyn, Sweden. *Eur J Agron.* 20(1-2): 101–116.
- Cramer, G.R. (1997). Uptake and role of ions in salt tolerance. In: Jaiwal RK, Singh RP, Gulati A (eds) *Strategies for improving salt tolerance in higher plants.* Science Pub. Inc. New Hampshire, pp. 55-86.
- Cacan, E., A. Aydin and M. Basbag (2015). Determination of Quality Characteristics of Some Grasses Forage Crops in Bingol University Campus. *Turkish J. Agric. Natr. Sci.* 2(2): 214–219.
- Deifel, K.S., P.M. Kopittke and N.W. Menzies (2006). Growth Response of Various Perennial Grasses to Increasing Salinity. *J. Plant Nutr.* 29(9): 1573-1584.
- Demiralay, I. (1993). *Soil Physical Analysis.* Atatürk Univ. J. Agric. Fac, Pub. No: 143, Erzurum
- Elmali, D.A. and I. Kaya (2012). The Effects of Different Harvesting Time on Nutrient Content of Sainfoin (*Onobrychis sativa* L.) and Vetch (*Vicia sativa* L.). *J. Lalahan Husbandry Research Institute.* 52(2): 39-45.
- Ganskopp, D. and D. Bohnert (2003). Mineral concentration dynamics among 7 northern Great Basin grasses. *J. Range Manage.*, 54: 640-647.
- Gee, G.W. and K.H. Hortage (1986). Particle- Size Analysis. *Methods of Soil Analysis. Part 1. Physical and Mineralogical Methods, Agronomy No: 9, 2nd ed.* pp. 383-441.
- Ghassemi-Golezani, K., S. Khomari, B. Dalil, A. Hosseinzadeh-Mahootchy, A. Chadordooz Jeddi (2010). Effects of seed aging on field performance of winter oilseed rape. *J Food Agri Environ* 8:175-178.
- Gonzales, C.L and M.D. Heilman (1977). Yield and chemical composition of coastal Bermudagrass, Rhodesgrass and volunteer species grown on saline and nonsaline soils. *J. Range Manage.*, 30:227-230.
- Goswami, T. K., R. Bhar, S. E. Jadhav, S. N. Joardar and G. C. Ram (2005). Role of dietary zinc as a nutritional immunomodulator. *Asian-Australas J Anim Sci.* 18(3):439-452.
- Grattan, S.R. and C.M. Grieve (1990). Effect of salinity on the mineral nutrition of plants. *International Conference on High Salinity Tolerant Plants in Arid Regions. United Arab Emirates Uni., Al Ain (Dec. 8-15, 1990).*
- Grattan, S.R., and C.M. Grieve (1999). Salinity-mineral nutrient relations in horticultural crops. *Scientia Horticulturae.* 78:127-157.
- Greenway, H., and R. Munns (1980). Mechanisms of salt tolerance in nonhalophytes. *Annu Rev Plant Physiol.* 31: 149-190.
- Gu, J., W. Liu, A. Akinagbe, J. Wang, J. Li and M. Yang (2012). Effect of salt stress on genetic diversity of *Robinia pseudoacacia* seedlings. *Afr J Biotechnol.* 11(8): 1838-1847.
- Gunes, A., A. Inal, M. Alpaslan, N. Cicek, E. Guneri, F. Eraslan, and T. Guzelordu (2005). Effects of exogenously applied salicylic acid on the induction of multiple stress tolerance and mineral nutrition in maize (*Zea mays* L.). *Arch Agron Soil Sci.* 51(6): 687–695.
- Haro, R., M.A. Banuelos, and A. Rodriguez-Navarro (2010). High affinity sodium uptake in land plants. *Plant and Cell Physiol.* 51: 68-79.
- John, M. K., H.H. Chuah and J.H. Neufeld (1975). Application of improved azomethine-H method to the determination of boron in soils and plants. *Anal. Lett.* 8(8): 559-568
- Kacar, B. and A. Inal (2008) *Plant Analysis.* Nobel Publishers, Ankara, Turkey.
- Kacar, B. and V. Katkat (2015). *Plant Nutrition (Bitki Besleme).* Nobel Pub., Ankara, Turkey. p:678



- Kacar, B., Katkat, A.V., Ozturk, S (2009). Plant Physiology. Nobel Pub. No: 848, Ankara, Turkey p 556.
- Kazemi, K and H. Eskandari (2011). Effects of salt stress on germination and early seedling growth of rice (*Oryza sativa*) cultivars in Iran. *Afr J Biotechnol* 10(77): 17789-17792.
- Khan, Z. I., M. Ashraf and A. Hussain, (2007). Evaluation of Macro Mineral Contents of Forages: Influence of Pasture and Seasonal Variation. *Asian-Australas J Anim Sci.* 20(6): 908-913.
- Kilic, U., S. Yurtseven, M. Boga and S. Aydemir (2015). Effects of soil salinity levels on nutrient contents and in vitro gas productions of some graminious forages. *J. Soil Sci. Plant Nutr.* 3(1): 9-15.
- Majid, M., A. Ali and B. Essia (2012). Effect of salinity on sodium and chloride uptake, proline and soluble carbohydrate contents in three alfalfa varieties. *IOSR J. Agric. Veterinary Sci.* 1(6): 01-06.
- Marschner, H., (1995). Mineral nutrition of higher plant. Institute of Plant Nutrition University of Hohenheim, Academic Press, Second Edition, Germany.
- McDowell, L.R., (1996). Feeding mineral to cattle on pasture. *Anim Feed Sci. Tech.* 60(3-4): 247-271.
- McLean, E.O. (1982). Soil pH and Lime Requirement. *Methods of Soil Analysis Part 2. Chemical and Microbiological Properties Second Edition.* Agronomy. No: 9 Part 2. Ed., ss: 199-224.
- Mezni, M., A. Albouchi, E. Bizid and M. Hamza (2010). Minerals uptake, organic osmotica contents and water balance in alfalfa under salt stress. *J. Phytol.* 2(11): 01-12.
- Munns, R. (2011). The impact of salinity stress. In *The Environmental and Physiological Nature of Salinity*. [http://www.plantstress.com/articles/salinity\\_i/salinity\\_i.htm](http://www.plantstress.com/articles/salinity_i/salinity_i.htm)
- Nelson, D.W. and L.E. Sommers (1982). *Methods of Soil Analysis. Part 2. Chemical and Microbiological Properties.* 2nd ed. Agronomy No: 9, Madison, Wisconsin, USA
- NRC. (1985). Nutrient requirements of domestic animals. Nutrient Requirements of Beef Cattle. 6th edition. National Academy of Sciences, Washington, DC. USA.
- Olsen, S.R. and L.E. Sommers (1982). Phosphorus. *Methods of Soil Analysis Part 2. Chemical and Microbiological Properties Second Edition.* Agronomy. No: 9 Part 2. Edition, ss: 403-427.
- Parida, A.K. and A.B. Das, (2005). Salt Tolerance and Salinity Effects on Plants: a Review, *Ecotoxicology and Environmental Safety*, 60, 324-349.
- Reid, R. L. and D.J. Horvath (1980). Soil chemistry and mineral problems in farm livestock: A review. *Anim. Feed Science and Technology*, 5(2):95-167.
- Rhoades, J.D., (1982). *Soluble Salts. Methods of Soil Analysis. Part 2. Chemical and Microbiological Properties.* 2nd Ed. Agronomy No: 9, 167-179, 1159 p, Madison, Wisconsin USA.
- Samad, R. and J.L. Karmoker (2013). Effects of NaCl stress on accumulation of K<sup>+</sup>, Na<sup>+</sup>, Cl<sup>-</sup>, NO<sub>3</sub><sup>-</sup>, sugar and proline contents in the seedlings of triticale-1. *Bangladesh J. Bot.* 42(2): 189-194.
- Semiz, G.D., A. Unlukara, E. Yurtseven, D.L. Suarez and I. Telci (2012). Salinity Impact on Yield, Water Use, Mineral and Essential Oil Content of Fennel (*Foeniculum vulgare* Mill.). *Tarim Bilimleri Dergisi* 18 (3): 177-186.
- Serin, Y. and M. Tan (1998). Forage Grasses. Ataturk Univ. Agri. Faculty, Publication. No: 334, Erzurum.
- Sima, N.A.K.K., S.T. Ahmad and M. Pessarakli (2013). Comparative Study of Different Salts (Sodium Chloride, Sodium Sulfate, Potassium Chloride, and Potassium Sulfate) on Growth of Forage Species. *J. of Plant Nutrition*, 36(2):214-230.
- Skerman, P.J. and F. Riveros (1990). Tropical grasses, *FAO Plant Production and Protection Series 23*, FAO (Food and Agriculture Organization of the United Nations), Rome, Italy, s: 832.
- Suyama, H., S. E. Benes, P. H. Robinson, G. Getachew, S. R. Grattan, and C. M. Grieve. (2007). Biomass yield and nutritional quality of forage species under long-term irrigation with saline-sodic drainage water: Field evaluation. *Anim Feed Sci. Tech.* 135(3-4):329-345.
- Tansi, V. (2009). Rhodes grass, Forage Grass and Other Families Forage. (Avcioglu, R., Hatipoglu, R., Karadag, Y Edit.) Volume I. TUGEM, Izmir, pp: 733-737.
- Temel, S., B. Keskin, U. Simsek, and İ.H. Yilmaz (2015). Effects of Halomorphic Soils Conditions on Plant Numbers Emerging in Square Meter of Some Perennial Forage Species. *J. Tekirdag Agric. Fac.* 12(1): 46-54.
- Temel, S., B. Keskin, U. Simsek and İ.H. Yilmaz (2016). The effect of soils having different salt content on mineral accumulations of some forage legume species. *Fresen Environ Bull.* 25(4): 1038-1050.
- Temel, M., M. Surmen and M. Tan (2015). Effects of Growth Stages on The Nutritive Value of Specific Halophyte Species in Saline Grasslands. *J. Anim. Plant Sci.* 25(5): 1419-1428
- Turan, M.A., A.H.A. Elkarim, N. Taban and S. Taban (2010). Effect of salt stress on growth and ion

- distribution and accumulation in shoot and root of maize plant. Afr. J. Agric. Res. 5(7): 584-588.
- Ul-Hassan, T. and A. Bano (2015). Physiological Characterization Of Halophytic Weeds Growing in Saline and Non-Saline Regions of Pakistan. J. Anim. Plant Sci. 25(2):463-471
- Ulrich, A. and F.J. Hills(1967). Principles and practices of plant analysis. In: G. W. Hardy. Soil testing and plant analysis. II Plant analysis. Madison, Wisconsin.
- Unlukara, A., Y. Cikili and A. Ozturk (2008). Effects of Different Leaching Fractions on Growth of Bean (*Phaseolus vulgaris*) and Its Mineral Composition. J. Agric. Fac. Gaziosmanpasa Univ. (JAFAG), 25(2): 51-60.
- Wang, H., Z. Wu, Y. Zhou, J. Han, and D. Shi (2012). Effects of salt stress on ion balance and nitrogen metabolism in rice. Plant Soil Environ. 58(2): 62-67.
- Warman P. R. and W.C. Termeer (2005). Evaluation of sewage sludge, septic waste and sludge compost applications to corn and forage: yields and N, P and K content of crops and soils. Bioresource Technol. 96(8): 955–961.