



Effects of muscle cooling on kinetics of pulmonary oxygen uptake and muscle deoxygenation at the onset of exercise

著者 (英)	Hitoshi Wakabayashi, Mizuki Osawa, Shunsaku Koga, Ke Li, Hiroyuki Sakaue, Yasuo Sengoku, Hideki TAKAGI
journal or publication title	Physiological reports
volume	6
number	21
page range	e13910
year	2018-10
権利	(C)2018 The Authors. Physiological Reports published by Wiley Periodicals, Inc. on behalf of The Physiological Society and the American Physiological Society This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.
URL	http://hdl.handle.net/2241/00154702

doi: 10.14814/phy2.13910



ORIGINAL RESEARCH

Effects of muscle cooling on kinetics of pulmonary oxygen uptake and muscle deoxygenation at the onset of exercise

Hitoshi Wakabayashi¹, Mizuki Osawa², Shunsaku Koga³, Ke Li², Hiroyuki Sakaue², Yasuo Sengoku² & Hideki Takagi²

1 Faculty of Engineering, Hokkaido University, Sapporo, Japan

2 Faculty of Health and Sport Sciences, University of Tsukuba, Tsukuba, Japan

3 Applied Physiology Laboratory, Kobe Design University, Kobe, Japan

Keywords

Cold skeletal muscle, glycolytic metabolism, near-infrared spectroscopy.

Correspondence

Hitoshi Wakabayashi, Laboratory of Environmental Ergonomics, Faculty of Engineering, Hokkaido University, N13 W8, Kita-ku, Sapporo, Hokkaido, 060-8628, Japan.

Tel/Fax: +81-11-706-6280

E-mail: wakabayashi@eng.hokudai.ac.jp

Funding Information

This study was supported by a Grant-in-Aid for Scientific Research (No. 15K1267105) from the Japan Society for the Promotion of Science and Advanced Research Initiative for Human High Performance (ARIHHP), University of Tsukuba.

Received: 24 September 2018; Accepted: 3 October 2018

doi: 10.14814/phy2.13910

Physiol Rep, 6 (21), 2018, e13910,
<https://doi.org/10.14814/phy2.13910>

Abstract

This study investigated effects of skeletal muscle cooling on the metabolic response and kinetics of pulmonary oxygen uptake ($\dot{V}O_2$) and skeletal muscle deoxygenation during submaximal exercise. In the cooling condition (C), after immersion of the lower body into 12°C water for 30 min, eight healthy males performed 30-min cycling exercise at the lactate threshold while undergoing thigh cooling by a water-circulating pad. In the normal condition (N) as control, they conducted the same exercise protocol without cooling. Blood lactate concentration was significantly higher in C than N at 10 min after onset of exercise (4.0 ± 1.7 and 2.4 ± 1.2 mmol/L in C and N, $P < 0.05$). The percent change in the tissue oxygen saturation of the vastus lateralis, measured by a near-infrared spectroscopy, was significantly lower in C at 2, 8, 10, and 20 min after the exercise onset compared with N ($P < 0.05$). The percent change in deoxy hemoglobin+myoglobin concentration (Deoxy[Hb+Mb]) showed a transient peak at the onset of exercise and significantly higher value in C at 10, 20, and 30 min after the exercise onset ($P < 0.05$). Compared to N, slower $\dot{V}O_2$ kinetics (mean response time) was observed in C (45.6 ± 7.8 and 36.1 ± 7.7 sec in C and N, $P < 0.05$). The mean response time in C relative to N was significantly correlated with the transient peak of Deoxy [Hb+Mb] in C ($r = 0.84$, $P < 0.05$). These results suggest that lower oxygen delivery to the hypothermic skeletal muscle might induce greater glycolytic metabolism during exercise and slower $\dot{V}O_2$ kinetics at the onset of exercise.

Introduction

There are many situations where humans perform in cold environments during sports events, winter outdoor occupations, and natural disasters like flooding and tsunamis. Cold-induced impairment of exercise performance including lower maximal force, velocity, and muscle endurance with lower skeletal muscle temperature (T_{muscle}) has been reported and summarized in several articles (Clarke et al.

1958; Heus et al. 1995; Racinais and Oksa 2010; Wakabayashi et al. 2015). The physiological mechanism for the performance impairment induced by cold has been studied extensively in relation to neuromuscular function using electromyography (EMG). The lower shift of EMG frequency with lower T_{muscle} is generally observed during exercise in cold (Bigland-Ritchie et al. 1981; Mucke and Heuer 1989; Oksa et al. 1997; Petrofsky and Laymon 2005; Wakabayashi et al. 2017) and is associated with a

reduction of the muscle conduction velocity (Bigland-Ritchie et al. 1981). Because of the lower contractility of each muscle fiber with low T_{muscle} , greater numbers of motor units are recruited to maintain the same work load in thermoneutral condition (Mallette et al. 2018).

Some studies have undertaken the measurement of pulmonary oxygen uptake ($\dot{V}O_2$) to evaluate the effect of low T_{muscle} on the whole body metabolism. Shiojiri et al. (1997) suggested that the slower pulmonary $\dot{V}O_2$ response with low T_{muscle} at the onset of exercise was due to decreased O_2 extraction and/or impairment of oxidative reactions in the cold skeletal muscle. However, due to the lack of direct measurement of muscle metabolism, it is still not clear how hypothermic skeletal muscle contributes to delayed $\dot{V}O_2$ kinetics. Bowen et al. (2013) investigated the association between delayed $\dot{V}O_2$ kinetics in hypoxic conditions and the transient peak of the skeletal muscle deoxygenation measured by near-infrared spectroscopy (NIRS). Similar to what is present with hypoxia (Bowen et al. 2013), in cold muscle, transient deoxygenation reflecting lower O_2 delivery relative to the muscle O_2 consumption would be expected because of the lower muscle perfusion with blood (i.e., lower O_2 delivery) in the cold (Rennie et al. 1980; Thorsson et al. 1985; Gregson et al. 2011). Additionally, fast-twitch fiber recruitment and anaerobic metabolism are expected to be increased in the cold (Rome et al. 1992; Fujimoto et al. 2016). It is known that larger proportion of fast-twitch fiber slows $\dot{V}O_2$ kinetics in vitro (Crow and Kushmerick 1982) and in vivo (Barstow et al. 1996; Pringle et al. 2003). Thus, fast-twitch fiber recruitment with hypothermic skeletal muscle could also delay $\dot{V}O_2$ kinetics, although these possibilities have yet to be examined.

Accordingly, the purpose of this study was to investigate the effect of lower T_{muscle} on the muscle and whole body metabolic response during lactate threshold (LT) exercise in human on cold exposure. The LT work rate was chosen for detecting difference between cold and normal conditions, even with a slight recruitment of glycolytic metabolism in cold. It was hypothesized that, in comparison with normothermic condition, greater blood lactate concentration would be observed with hypothermic skeletal muscle, and transient peak of the skeletal muscle deoxygenation would result in slower $\dot{V}O_2$ kinetics at the onset of exercise in cold.

Methods

Participants

Eight healthy males (mean \pm standard deviation height: 173.4 ± 5.4 cm, body weight: 71.7 ± 12.4 kg) participated in this study. They were informed of the

experimental procedures and gave their written consent before participation. All experimental procedures in this study were designed according to the principle of the Helsinki Declaration and approved by the Human Subjects Committee of the University of Tsukuba.

Procedures

Prior to the main experiment, participants performed an incremental exercise test for assessing their LT using a bicycle ergometer (PowerMax VII, Konami). Participants performed an intermittent incremental cycling protocol consisting of 4-min pedaling at 60 rpm and 2-min rest interval. They started the first stage from 1.0 kp and gradually increased intensity by 0.3 kp every stage until their blood lactate concentration (La) was higher than 4.0 mmol/L. During the stage interval, capillary blood was sampled from their fingertip. Based on the La of each stage, the LT work rate was determined by the linear regression fitting model with least mean squared error using Lactate-E software (Newell et al. 2007).

The participants performed 30-min constant work rate exercise in both cooling (C) and normal (N) condition in random order on separate days. They sat on the seat of the ergometer (PowerMax VII, Konami) for 3 min for measuring resting baseline values. In C, participants underwent pre-cooling by immersing their lower body into 12°C water for 30 min while remaining at rest, and then exited the immersion pool and performed 30-min cycling exercise at the LT work rate at 60 rpm with cooling applied to their thighs. Both thighs were wrapped by several water perfusion cooling pads, through which 5°C water was circulated using a thermostatic bath system (LTB-400, ASONE). In N, they conducted a similar exercise protocol without any cooling. The LT work rate was chosen for clearly detecting the recruitment of anaerobic metabolism in C compared to N.

Measurements and analysis

The vastus lateralis T_{muscle} was estimated continuously, except during immersion, using a deep body temperature monitor (CM-210, TERUMO) which detects the tissue temperature 5–10 mm below the skin surface using the zero-heat-flow-method (Yamakage and Namiki 2003). This monitor measures skin surface temperature beneath a thermal insulating pad containing a heater, which equilibrates the skin temperature with the deep tissue temperature when heat flow from the skin is maintained at zero. The consistency between T_{muscle} measured using a needle thermocouple and the zero-heat-flow-method was evaluated (Togawa et al. 1976). The skin temperature on the

thigh (T_{thigh}) was also measured using a thermistor sensor (PD-K161, TERUMO).

Changes in the concentrations of oxygenated and deoxygenated hemoglobin and myoglobin (Oxy[Hb+Mb] and Deoxy[Hb+Mb]) in the vastus lateralis muscle were continuously measured using spatially resolved near-infrared spectroscopy (NIRS) (BOM-L1TR, Omegawave). The data were sampled using an A/D converter (Powerlab/16SP, AD Instruments) and recorded at 0.1 second intervals using a laptop computer. The optodes of the NIRS were housed in an optically dense black vinyl holder to ensure distance between an illuminant and two detectors at 20 (detector 1) and 40 mm (detector 2), respectively. The illuminant generated three wavelengths of near-infrared light (780, 810 and 830 nm). The optodes were attached to the skin above the vastus lateralis using double-sided adhesive tape. The Oxy[Hb+Mb] and Deoxy[Hb+Mb] concentrations only in the deep tissue region (Oxy[Hb+Mb]₂₋₁ and Deoxy[Hb+Mb]₂₋₁) were calculated from the [Hb+Mb] concentrations measured at the two detectors (detector 1 and 2) using the following formulae (Kashima 2003).

$$\text{Oxy[Hb + Mb]}_{2-1} = (\text{Oxy[Hb + Mb]}_2 \times L_2 - \text{Oxy[Hb + Mb]}_1 \times L_1) / (L_2 - L_1)$$

$$\text{Deoxy[Hb + Mb]}_{2-1} = (\text{Deoxy[Hb + Mb]}_2 \times L_2 - \text{Deoxy[Hb + Mb]}_1 \times L_1) / (L_2 - L_1)$$

where L_1 and L_2 represent length from the illuminant to the detector 1 and 2, respectively. This method enabled us to focus on the muscle oxygenation excluding the effect of the skin blood flow on [Hb+Mb] concentrations (Ando et al. 2010). Total [Hb+Mb] and tissue O_2 saturation (StO_2) was calculated by the following formulas.

$$\text{Total[Hb + Mb]} = \text{Oxy[Hb + Mb]} + \text{Deoxy[Hb + Mb]}$$

$$\text{StO}_2 = \text{Oxy[Hb + Mb]} / \text{Total[Hb + Mb]} \times 100$$

Percent change of Oxy[Hb+Mb], Deoxy[Hb+Mb], total [Hb+Mb] and StO_2 from the baseline period before cooling was calculated for the following analyses, since the NIRS technique cannot measure the absolute [Hb+Mb] concentration and reference values for both conditions are set at thermoneutral baseline.

La concentration was measured at baseline, every 2 min in the first 10-min exercise and every 10 min in the latter 20 min using a portable lactate analyzer (Lactate Pro LT-1710, Arkray Inc). Breath-by-breath $\dot{V}\text{O}_2$ was continuously measured by an automated respirometer (Metalyzer 3B, Cortex) and was interpolated at 5-sec intervals. The kinetics of $\dot{V}\text{O}_2$ at the onset of exercise was

analyzed using the first 6-min data excluding the initial 20 sec, since this study focused on the primary phase reflecting the kinetics of O_2 consumption in the exercising muscle. To reduce the breath-by-breath $\dot{V}\text{O}_2$ fluctuation, the data were smoothed using a 3-point moving average. The $\dot{V}\text{O}_2$ response was fitted to the following exponential equation using nonlinear least squares regression. The fitting of the exponential curve and 3-point moving average for each participant was evaluated by coefficient of determination (R^2).

$$\dot{V}\text{O}_2(t) = \dot{V}\text{O}_{2(\text{base})} + \text{Amp}(1 - \exp^{-(t-\text{TD})/\tau})$$

where $\dot{V}\text{O}_{2(\text{base})}$: $\dot{V}\text{O}_2$ at the baseline before exercise onset, Amp: amplitude between the baseline and steady state (6 min after onset of exercise), TD: time delay, τ : time constant of the exponential function. Mean response time (MRT) was calculated according to the following formulae (Sietsema et al. 1994).

$$\text{MRT} = \text{O}_2 \text{ deficit} / (\dot{V}\text{O}_{2(6)} - \dot{V}\text{O}_{2(\text{base})})$$

where O_2 deficit: subtracting 6-min cumulative $\dot{V}\text{O}_2$ from the required $\dot{V}\text{O}_2$, i.e., the product of $\dot{V}\text{O}_2$ at 6 min and the duration (6 min), $\dot{V}\text{O}_{2(6)}$: $\dot{V}\text{O}_2$ at 6 min after exercise onset. The increment in $\dot{V}\text{O}_2$ between minutes 2 and 6 ($\Delta\dot{V}\text{O}_{2(6-2)}$) was calculated for assessing the slow component of $\dot{V}\text{O}_2$ kinetics (Whipp and Wasserman 1986). Additionally, $\dot{V}\text{O}_2$ amplitude between baseline and 30 min after the onset of exercise ($\Delta\dot{V}\text{O}_{2(30-\text{base})}$) was calculated.

Statistical analysis

Data from time-course measurements (T_{muscle} , T_{thigh} , $\dot{V}\text{O}_2$, La, percent change of Oxy[Hb+Mb], Deoxy[Hb+Mb], total[Hb+Mb] and StO_2) were evaluated using repeated measures two-way (condition \times time) analysis of variance (ANOVA). If Mauchly's sphericity test was not satisfied, the degrees of freedom were adjusted by Greenhouse-Geisser's ϵ . Paired Student's t -test with Holm's multiple comparisons adjustment (Holm 1979) was conducted between conditions at each time point. Parameters of $\dot{V}\text{O}_2$ kinetics (baseline, Amp, TD, τ , MRT, O_2 deficit and $\Delta\dot{V}\text{O}_{2(6-2)}$ and $\Delta\dot{V}\text{O}_{2(30-\text{base})}$) were compared between conditions using paired t -test. Pearson correlation coefficient was calculated to examine the relationships between parameters of $\dot{V}\text{O}_2$ kinetics and muscle deoxygenation. Cohen's d and partial η^2 (η_p^2) was calculated for assessing effect size for paired t -test and ANOVA, respectively. Statistical significance was established at $P < 0.05$. All data are presented as mean values and standard deviation (SD).

Results

Body temperature

T_{muscle} and T_{thigh} during exercise are shown in Figure 1. For both temperature variables, significant main effect of time ($F_{2,4, 16.6} = 228.4$, $\eta_p^2 = 0.98$, $P < 0.001$; $F_{1,7, 12.1} = 69.0$, $\eta_p^2 = 0.99$, $P < 0.001$), condition ($F_{1, 7} = 623.8$, $\eta_p^2 = 0.99$, $P < 0.001$; $F_{1, 7} = 368.3$, $\eta_p^2 = 0.98$, $P < 0.001$), and a significant interaction between those factors ($F_{2,1, 14.7} = 75.1$, $\eta_p^2 = 0.99$, $P < 0.001$; $F_{2,1, 14.9} = 15.2$, $\eta_p^2 = 0.69$, $P < 0.001$) were detected in the two-way ANOVA. The lower body cold immersion decreased T_{muscle} to 22.7°C on average. After the onset of exercise T_{muscle} in C gradually recovered, though it was significantly lower than N throughout the 30-min exercise ($P < 0.05$). T_{thigh} was about 21°C in C at the onset of exercise, then gradually increased during exercise, although it was significantly lower than that in N throughout the 30-min exercise ($P < 0.05$).

Pulmonary oxygen uptake and blood lactate concentration

$\dot{V}O_2$ during exercise showed a significant main effect of time ($F_{1,1, 8.0} = 64.8$, $\eta_p^2 = 0.90$, $P < 0.001$, Fig. 2). No significant difference was observed between conditions in $\dot{V}O_2$ at any time point during exercise.

La showed significant main effect of time ($F_{1,9, 13.4} = 10.1$, $\eta_p^2 = 0.59$, $P < 0.001$), condition ($F_{1, 7} = 20.3$, $\eta_p^2 = 0.74$, $P < 0.001$), and a significant interaction between those factors ($F_{2,9,6, 20.7} = 6.7$, $\eta_p^2 = 0.49$, $P < 0.001$, Fig. 3). La was significantly higher in C than that in N at 8 min ($d = 1.25$, $P < 0.05$) and 10 min ($d = 1.14$, $P < 0.05$) after onset of exercise.

Skeletal muscle oxygenation

The percent change in Oxy, Deoxy, and total [Hb+Mb] in the vastus lateralis relative to the resting baseline before cooling are shown in Figure 4. The Oxy[Hb+Mb] showed significant main effect of time ($F_{1,8, 12.4} = 5.7$, $\eta_p^2 = 0.45$, $P < 0.05$) and condition ($F_{1, 7} = 8.1$, $\eta_p^2 = 0.54$, $P < 0.05$). At the onset of exercise the Oxy[Hb+Mb] in C was significantly lower than that in N ($d = 2.79$, $P < 0.01$). The Deoxy[Hb+Mb] showed significant main effect of time ($F_{1,7, 11.7} = 4.9$, $\eta_p^2 = 0.41$, $P < 0.05$) and condition ($F_{1, 7} = 17.9$, $\eta_p^2 = 0.72$, $P < 0.01$). The Deoxy [Hb+Mb] was significantly higher in C at the onset of exercise ($d = 1.11$, $P < 0.05$) and at 10 ($d = 0.69$, $P < 0.05$), 20 ($d = 0.45$, $P < 0.05$) and 30 min ($d = 0.40$, $P < 0.05$) after the onset. The total [Hb+Mb] showed main effect of time ($F_{1,6, 11.5} = 16.3$, $\eta_p^2 = 0.70$,




Figure 1. The vastus lateralis muscle temperature and thigh skin temperature during exercise. Values are mean \pm SD of the cooling and normal conditions *Significant difference between conditions ($P < 0.05$).




Figure 2. Change in the rate of oxygen uptake during the experiment. Values are mean \pm SD of the cooling and normal conditions.

$P < 0.001$) and no difference was observed between conditions.

The percent change in StO_2 in the vastus lateralis relative to the resting baseline before cooling are shown in Figure 5. StO_2 showed a significant main effect of condition ($F_{1,7, 11.7} = 23.1$, $\eta_p^2 = 0.77$, $P < 0.001$) and a significant interaction between those factors ($F_{2,0, 14.3} = 5.2$, $\eta_p^2 = 0.43$, $P < 0.05$). The percent change in StO_2 was




Figure 3. Change in the blood lactate concentration during exercise. Values are mean \pm SD of each cooling and normal condition. * Significant difference between conditions ($P < 0.05$).

significantly lower in C at the onset of exercise ($d = 2.49$, $P < 0.05$), 2 ($d = 1.70$, $P < 0.05$), 8 ($d = 1.02$, $P < 0.05$), 10 ($d = 1.12$, $P < 0.05$), and 20 min ($d = 0.88$, $P < 0.05$) after the onset.

Kinetics of pulmonary oxygen uptake and muscle deoxy hemoglobin

The parameters of $\dot{V}O_2$ kinetics during the first 6 min of exercise are shown in Table 1. Compared to N, condition C showed significantly longer time constant of $\dot{V}O_2$ kinetics ($\tau\dot{V}O_2$) ($d = 1.23$, $P < 0.05$) and MRT ($d = 1.23$, $P < 0.01$). There were no significant differences between conditions in $\dot{V}O_{2(\text{base})}$, Amp, TD, initial 6-min O_2 deficit or $\Delta\dot{V}O_{2(30-\text{base})}$. No difference was observed between Amp (i.e., $\Delta\dot{V}O_2$ between baseline and 6 min) and $\Delta\dot{V}O_{2(30-\text{base})}$ (i.e., Amp at the 30th min) in each condition. $\Delta\dot{V}O_{2(6-2)}$ was significantly greater in C ($d = 0.97$, $P < 0.05$) and significantly correlated with the La concentration averaged during 2 to 6 min ($r = 0.71$, $P < 0.05$). A transient peak in the Deoxy[Hb+Mb] ($76.5 \pm 33.5\%$) in the vastus lateralis was observed only in C at 41.3 ± 8.4 sec after onset of exercise. The peak Deoxy[Hb+Mb] in C was significantly correlated with differences between conditions in MRT, initial 6-min O_2 deficit and $\Delta\dot{V}O_{2(6-2)}$ ($r = 0.84$, 0.74 and 0.80 , respectively, $P < 0.05$, Fig. 6) and tended to be correlated with the $\tau\dot{V}O_2$ ($r = 0.68$, $P = 0.06$). The percentage of the $\tau\dot{V}O_2$ in C relative to N was significantly correlated with the peak Deoxy[Hb+Mb] in C ($r = 0.71$, $P < 0.05$).

Discussion

This study investigated the effect of muscle cooling on pulmonary $\dot{V}O_2$ kinetics and muscle metabolism during exercise at LT work rate. The major finding of the present study was that at the onset of exercise, transient peak




Figure 4. Percent change in the oxy, deoxy, and total hemoglobin and myoglobin in the vastus lateralis. Values are mean \pm SD of each cooling and normal condition. *Significant difference between conditions ($P < 0.05$).

muscle deoxygenation was observed in C, which is associated with the delay of pulmonary $\dot{V}O_2$ kinetics relative to N. Additionally, significantly higher La concentration was observed in C compared to N during exercise.

Anaerobic metabolism in hypothermic skeletal muscle

After 30-min cold immersion, the vastus lateralis T_{muscle} declined to 22.7°C on average, which was below T_{muscle} inducing impairment of muscle endurance and maximal force (Clarke et al. 1958; Heus et al. 1995; Wakabayashi et al. 2015). Since an elevation of La concentration was associated with lowering of T_{muscle} (Figs. 1 and 3), we suggest that the reduction of T_{muscle} would increase recruitment of glycolytic metabolism during exercise.




Figure 5. Percent change in the tissue oxygen saturation in the vastus lateralis. Values are mean \pm SD of each cooling and normal condition. *Significant difference between conditions ($P < 0.05$).

Similar to current observation, greater anaerobic metabolism in cold water was reported in a previous study which showed higher minute ventilation to $\dot{V}O_2$ ratio ($V_E/\dot{V}O_2$) during an incremental exercise test in cold (18°C) compared to warm water (Fujimoto et al. 2016). Since we set the LT work rate predetermined in normal condition for both conditions, participants actually exercised at LT work rate (moderate intensity) in N and above LT (heavy intensity) in C. The potential mechanism for the greater contribution of the glycolytic metabolism with hypothermic skeletal muscle is discussed as follows.

In this study no direct measurement of muscle blood flow was conducted; however, cold-induced muscle vasoconstriction (Rennie et al. 1980) seemed to be one of the major factors inducing the anaerobic metabolism. However, the percent change in the total [Hb+Mb] indicates no difference between conditions in the microvascular Hb volume and muscle O_2 diffusion capacity (Fukuoka et al.

2015). The significantly lower StO_2 in C might be attributed in part to less blood perfusion in the hypothermic skeletal muscle (Rennie et al. 1980; Thorsson et al. 1985; Gregson et al. 2011) and leftward shift of oxyhemoglobin dissociation curve with lower temperature (Severinghaus 1958; Astrup et al. 1965). The larger increase of Deoxy [Hb+Mb] in C during exercise suggests the lower muscle blood flow/ O_2 delivery relative to O_2 consumption (Koga et al. 2012, 2015). Based on these observations by the NIRS method, less tissue oxygenation (lower StO_2) due to lower O_2 delivery relative to O_2 consumption (higher Deoxy[Hb+Mb]) might induce the greater anaerobic contribution in C. Additionally, lower blood perfusion may also impair the lactate utilization in the skeletal muscle (Gladden et al. 1992; Gladden 2000) resulting in the greater La concentration in C.

In addition to differences in O_2 transport to the working muscle due to the cold-induced muscle vasoconstriction, reduction of T_{muscle} per se slows ATP utilization (Edwards et al. 1972) and slows Ca^{2+} release and uptake from the sarcoplasmic reticulum (Kossler et al. 1987; Herve et al. 1992). Previous studies have also reported that a greater number of motor units were recruited in cold environment in order to maintain the same work load compared to neutral conditions (Mallette et al. 2018). Because of the greater cold sensitivity and lower power output of slow-twitch fibers (Ran-tunga 1984), fast-twitch fibers were recruited to a greater extent in a cold environment compared to warm condition even at moderate-intensity exercise in scup (Rome et al. 1992). Thus, the greater anaerobic contribution with hypothermic skeletal muscle might be attributed to the relatively more recruitment of the fast-twitch fibers relying more on glycolytic energy transfer, which may be associated to the larger $\dot{V}O_2$ slow component (see below).

Table 1. Oxygen uptake kinetics during the first 6 min of exercise.

	Cooling	Normal
Baseline (L/min)	0.41 \pm 0.37 [0.15–0.66]	0.25 \pm 0.10 [0.18–0.31]
Amplitude (L/min)	1.28 \pm 0.22 [1.13–1.48]	1.39 \pm 0.30 [1.18–1.60]
Time delay (sec)	11.4 \pm 10.7 [4.0–18.8]	16.3 \pm 3.2 [14.1–18.5]
Time constant (sec)	39.0 \pm 13.7 [29.5–48.5]	25.7 \pm 6.8 [21.0–30.5]*
R^2 value of regression curve	0.84 \pm 0.05 [0.81–0.88]	0.83 \pm 0.10 [0.76–0.90]
Mean response time (sec)	45.6 \pm 7.8 [40.2–51.0]	36.1 \pm 7.7 [30.7–41.4]*
Initial 6-min O_2 deficit (L)	0.98 \pm 0.30 [0.77–1.19]	0.85 \pm 0.28 [0.66–1.04]
$\Delta\dot{V}O_{2(6-2)}$ (L/min)	0.09 \pm 0.07 [0.04–0.14]	0.03 \pm 0.03 [0.01–0.05]*
$\Delta\dot{V}O_{2(30-base)}$ (L/min)	1.29 \pm 0.30 [1.08–1.50]	1.47 \pm 0.35 [1.23–1.72]

Values are means \pm standard deviation and [95% confidential interval].

*Significant difference between conditions ($P < 0.05$).

$\Delta\dot{V}O_{2(6-2)}$ and $\Delta\dot{V}O_{2(30-base)}$: the increment in oxygen uptake between minutes 2 and 6, and between baseline and minute 30.




Figure 6. The condition differences in $\dot{V}O_2$ mean response time, initial 6-min O_2 deficit, and $\Delta\dot{V}O_2$ ($_{6-2}$) as a function of the transient peak Deoxy[Hb+Mb] in the vastus lateralis in cooling condition. $\Delta\dot{V}O_2$ ($_{6-2}$): the increment in oxygen uptake between minutes 2 and 6.

Kinetics of oxygen uptake and Deoxy [Hb+Mb] at the onset of exercise

At the onset of exercise (i.e., at the time of the lowest T_{muscle} was recorded), the longer $\tau\dot{V}O_2$ and MRT observed in C was consistent with a previous study (Shiojiri et al.

1997). Shiojiri et al. (1997) suggested that the delay of $\dot{V}O_2$ kinetics was due to decreased O_2 extraction and/or impairment of oxidative reactions based on similar cardiac output in C and N conditions. In the present study, the transient peak in Deoxy[Hb+Mb] in C indicates possible greater O_2 extraction fraction (Koga et al. 2012, 2014) and less oxygen delivery to muscle oxygen uptake. The different interpretation for the O_2 extraction between the present and previous studies (Shiojiri et al. 1997) might be attributed to exercise intensity, that was above anaerobic threshold (AT) in the present cooling condition, while below AT in the previous study (Shiojiri et al. 1997). The lower blood flow to the working muscle in cold (Rennie et al. 1980) might have been reflected in the transient peak of Deoxy[Hb+Mb] and the lower StO_2 in C (Figs. 4 and 5). The peak Deoxy [Hb+Mb] was correlated with the ΔMRT between conditions and $\tau\dot{V}O_2$ in C relative to N. Additionally, the condition differences in O_2 deficit and $\Delta\dot{V}O_2$ ($_{6-2}$) were also strongly correlated with the peak Deoxy[Hb+Mb] in C (Fig. 6). Based on these observations, we suggest that the slower $\dot{V}O_2$ kinetics in C would be caused by the limitation of aerobic energy provision due to less blood perfusion in the working muscle. Similar to the present study, Bowen et al. (2013) reported delay of $\dot{V}O_2$ kinetics in hypoxia and correlation with transient peak of skeletal muscle Deoxy [Hb+Mb] assessed using time resolved near-infrared spectroscopy. In terms of lower O_2 delivery to the working muscle, a similar mechanism in the slowed $\dot{V}O_2$ kinetics is plausible in a hypoxic environment and muscle cooling condition. For example, there have been several studies reporting delay of $\dot{V}O_2$ kinetics with interventions to limit O_2 delivery to the working muscle (Hughson and Smyth 1983; Hughson et al. 1991; Goodwin et al. 2012).

As explained above, fast-twitch fibers might be recruited more in the cooling condition (Ranatunga 1984; Rome et al. 1992). Relating to previous studies (Barstow et al. 1996; Pringle et al. 2003) reporting delay of $\dot{V}O_2$ kinetics in participants with a greater percentage of fast-twitch fibers, slower $\dot{V}O_2$ kinetics observed in C of the present study might indicate the greater recruitment of fast-twitch fibers in C. Additionally, although the exercise intensity was set at the same absolute work rate for both N and C, the larger $\Delta\dot{V}O_2$ ($_{6-2}$) in C was observed as shown (Table 1). Concerning the significant correlation between the $\Delta\dot{V}O_2$ ($_{6-2}$) and La concentration averaged during 2 to 6 min, the larger recruitment of glycolytic metabolism in C would increase slow component and delay $\dot{V}O_2$ kinetics (Whipp and Wasserman 1986; Wasserman 1987).

Methodological limitation

In this study, we choose the LT work rate predetermined in normal condition for assessing recruitment of

anaerobic metabolism with hypothermic skeletal muscle even with a slight increase over the threshold in C. Thus, participants might exercise in moderate and heavy intensity domains in N and C, respectively. It might have been more appropriate to have participants exercising in the moderate-intensity domain (e.g., at 80% LT) in both conditions to determine whether anaerobic contribution was elevated in C but not in N, and also whether $\dot{V}O_2$ primary kinetics slowed and/or slow component was created in C during moderate-intensity exercise.

Conclusion

One of the major findings in this study was that exercise with hypothermic skeletal muscle increased recruitment of glycolytic metabolism as shown in the significantly higher blood lactate concentration even at lactate threshold workload that was predetermined in thermoneutral environment. Our second major finding was the delay of oxygen uptake kinetics at the onset of exercise after pre-cooling and a significant correlation with the transient peak of deoxy hemoglobin in the working muscle.

Acknowledgments

The authors wish to thank all those who participated in this study. We also would like to express our thanks to Dr Martin Barwood for English proof reading.

Conflict of Interest

None declared.

References

- Ando, S., Y. Yamada, M. Kokubu. 2010. Reaction time to peripheral visual stimuli during exercise under hypoxia. *J. Appl. Physiol.* (1985) 108:1210–1216.
- Astrup, P., K. Engel, J. W. Severinghaus, and E. Munson. 1965. The influence of temperature and pH on the dissociation curve of oxyhemoglobin of human blood. *Scand. J. Clin. Lab. Invest.* 17:515–523.
- Barstow, T. J., A. M. Jones, P. H. Nguyen, and R. Casaburi. 1996. Influence of muscle fiber type and pedal frequency on oxygen uptake kinetics of heavy exercise. *J. Appl. Physiol.* 81:1642–1650.
- Bigland-Ritchie, B., E. F. Donovan, and C. S. Roussos. 1981. Conduction velocity and EMG power spectrum changes in fatigue of sustained maximal efforts. *J. Appl. Physiol. Respir. Environ. Exerc. Physiol.* 51:1300–1305.
- Bowen, T. S., H. B. Rossiter, A. P. Benson, T. Amano, N. Kondo, J. M. Kowalchuk, et al. 2013. Slowed oxygen uptake kinetics in hypoxia correlate with the transient peak and reduced spatial distribution of absolute skeletal muscle deoxygenation. *Exp. Physiol.* 98:1585–1596.
- Clarke, R. S., R. F. Hellon, and A. R. Lind. 1958. The duration of sustained contractions of the human forearm at different muscle temperatures. *J. Physiol.* 143:454–473.
- Crow, M. T., and M. J. Kushmerick. 1982. Chemical energetics of slow- and fast-twitch muscles of the mouse. *J. Gen. Physiol.* 79:147–166.
- Edwards, R. H., R. C. Harris, E. Hultman, L. Kaijser, D. Koh, and L. O. Nordesjo. 1972. Effect of temperature on muscle energy metabolism and endurance during successive isometric contractions, sustained to fatigue, of the quadriceps muscle in man. *J. Physiol.* 220:335–352.
- Fujimoto, T., Y. Sasaki, H. Wakabayashi, Y. Sengoku, S. Tsubakimoto, and T. Nishiyasu. 2016. Maximal workload but not peak oxygen uptake is decreased during immersed incremental exercise at cooler temperatures. *Eur. J. Appl. Physiol.* 116:1819–1827.
- Fukuoka, Y., D. C. Poole, T. J. Barstow, N. Kondo, M. Nishiwaki, D. Okushima, et al. 2015. Reduction of $\dot{V}O_2$ slow component by priming exercise: novel mechanistic insights from time-resolved near-infrared spectroscopy. *Physiol. Rep.* 3:e12432.
- Gladden, L. B. 2000. Muscle as a consumer of lactate. *Med. Sci. Sports Exerc.* 32:764–771.
- Gladden, L. B., R. E. Crawford, and M. J. Webster. 1992. Effect of blood flow on net lactate uptake during steady-level contractions in canine skeletal muscle. *J. Appl. Physiol.* 72:1826–1830.
- Goodwin, M. L., A. Hernandez, N. Lai, M. E. Cabrera, and L. B. Gladden. 2012. $\dot{V}O_2$ on-kinetics in isolated canine muscle in situ during slowed convective O_2 delivery. *J. Appl. Physiol.* 112:9–19.
- Gregson, W., M. A. Black, H. Jones, J. Milson, J. Morton, B. Dawson, et al. 2011. Influence of cold water immersion on limb and cutaneous blood flow at rest. *Am. J. Sports Med.* 39:1316–1323.
- Herve, J. C., K. Yamaoka, V. W. Twist, T. Powell, J. C. Ellory, and L. C. Wang. 1992. Temperature dependence of electrophysiological properties of guinea pig and ground squirrel myocytes. *Am. J. Physiol.* 263:R177–R184.
- Heus, R., H. A. Daanen, and G. Havenith. 1995. Physiological criteria for functioning of hands in the cold: a review. *Appl. Ergon.* 26:5–13.
- Holm, S. 1979. A Simple Sequentially Rejective Multiple Test Procedure. *Scand. J. Stat.* 6:65–70.
- Hughson, R. L., and G. A. Smyth. 1983. Slower Adaptation of $\dot{V}O_2$ to steady state of submaximal exercise with beta-blockade. *Eur. J. Appl. Physiol.* 52:107–110.
- Hughson, R. L., H. C. Xing, C. Borkhoff, and G. C. Butler. 1991. Kinetics of ventilation and gas exchange during supine and upright cycle exercise. *Eur. J. Appl. Physiol.* 63:300–307.
- Kashima, S. 2003. Spectroscopic measurement of blood volume and its oxygenation in a small volume of tissue

- using red laser lights and differential calculation between two point detections. *Opt. Laser Technol.* 35:485–489.
- Koga, S., Y. Kano, T. J. Barstow, L. F. Ferreira, E. Ohmae, M. Sudo, *et al.* 2012. Kinetics of muscle deoxygenation and microvascular PO₂ during contractions in rat: comparison of optical spectroscopy and phosphorescence-quenching techniques. *J. Appl. Physiol.* (1985) 112:26–32.
- Koga, S., H. B. Rossiter, I. Heinonen, T. I. Musch, and D. C. Poole. 2014. Dynamic heterogeneity of exercising muscle blood flow and O₂ utilization. *Med. Sci. Sports Exerc.* 46:860–876.
- Koga, S., T. J. Barstow, D. Okushima, H. B. Rossiter, N. Kondo, E. Ohmae, *et al.* 2015. Validation of a high-power, time-resolved, near-infrared spectroscopy system for measurement of superficial and deep muscle deoxygenation during exercise. *J. Appl. Physiol.* (1985) 118:1435–1442.
- Kossler, F., F. Lange, and G. Kuchler. 1987. Isometric twitch and tetanic contraction of frog skeletal muscles at temperatures between 0 to 30 degrees C. *Biomed. Biochim. Acta* 46:809–813.
- Mallette, M. M., L. A. Green, D. A. Gabriel, and S. S. Cheung. 2018. The effects of local forearm muscle cooling on motor unit properties. *Eur. J. Appl. Physiol.* 118:401–410.
- Mucke, R., and D. Heuer. 1989. Behaviour of EMG-parameters and conduction velocity in contractions with different muscle temperatures. *Biomed. Biochim. Acta* 48: S459–S464.
- Newell, J., D. Higgins, N. Madden, J. Cruickshank, J. Einbeck, K. McMillan, *et al.* 2007. Software for calculating blood lactate endurance markers. *J. Sports Sci.* 25:1403–1409.
- Oksa, J., H. Rintamaki, and S. Rissanen. 1997. Muscle performance and electromyogram activity of the lower leg muscles with different levels of cold exposure. *Eur. J. Appl. Physiol.* 75:484–490.
- Petrofsky, J., and M. Laymon. 2005. Muscle temperature and EMG amplitude and frequency during isometric exercise. *Aviat. Space Environ. Med.* 76:1024–1030.
- Pringle, J. S., J. H. Doust, H. Carter, K. Tolfrey, I. T. Campbell, G. K. Sakkas, *et al.* 2003. Oxygen uptake kinetics during moderate, heavy and severe intensity “submaximal” exercise in humans: the influence of muscle fibre type and capillarisation. *Eur. J. Appl. Physiol.* 89:289–300.
- Racinais, S., and J. Oksa. 2010. Temperature and neuromuscular function. *Scand. J. Med. Sci. Sports* 20(Suppl 3):1–18.
- Ranatunga, K. W. 1984. The force-velocity relation of rat fast- and slow-twitch muscles examined at different temperatures. *J. Physiol.* 351:517–529.
- Rennie, D., Y. Park, A. Veicsteinas, and D. Pendergast. 1980. Metabolic and circulatory adaptation to cold water stress. Pp. 315–321 in P. Cerretelli and B. J. Whipp, eds. *Exercise bioenergetics and gas exchange*. Elsevier/North-Holland Biomedical Press, Amsterdam, The Netherlands.
- Rome, L.C., I. H. Choi, G. Lutz, and A. Sosnicki. 1992. The influence of temperature on muscle function in the fast swimming scup. I. Shortening velocity and muscle recruitment during swimming. *J. Exp. Biol.* 163:259–279.
- Severinghaus, J. W. 1958. Oxyhemoglobin dissociation curve correction for temperature and pH variation in human blood. *J. Appl. Physiol.* 12:485–486.
- Shiojiri, T., M. Shibasaki, K. Aoki, N. Kondo, and S. Koga. 1997. Effects of reduced muscle temperature on the oxygen uptake kinetics at the start of exercise. *Acta Physiol. Scand.* 159:327–333.
- Sietsema, K. E., I. Ben-Dov, Y. Y. Zhang, C. Sullivan, and K. Wasserman. 1994. Dynamics of oxygen uptake for submaximal exercise and recovery in patients with chronic heart failure. *Chest* 105:1693–1700.
- Thorsson, O., B. Lilja, L. Ahlgren, B. Hemdal, and N. Westlin. 1985. The effect of local cold application on intramuscular blood flow at rest and after running. *Med. Sci. Sports Exerc.* 17:710–713.
- Togawa, T., T. Nemoto, T. Yamazaki, and T. Kobayashi. 1976. A modified internal temperature measurement device. *Med. Biol. Eng.* 14:361–364.
- Wakabayashi, H., J. Oksa, and M. J. Tipton. 2015. Exercise performance in acute and chronic cold exposure. *J. Phys. Fit. Sports Med.* 4:177–185.
- Wakabayashi, H., T. Wijayanto, and Y. Tochiara. 2017. Neuromuscular function during knee extension exercise after cold water immersion. *J. Physiol. Anthropol.* 36:28.
- Wasserman, K. 1987. Determinants and detection of anaerobic threshold and consequences of exercise above it. *Circulation* 76:VI29–VI39.
- Whipp, B. J., and K. Wasserman. 1986. Effect of anaerobiosis on the kinetics of O₂ uptake during exercise. *Fed. Proc.* 45:2942–2947.
- Yamakage, M., and A. Namiki. 2003. Deep temperature monitoring using a zero-heat-flow method. *J. Anesth.* 17:108–115.