


Bulletin of the Geological Society of Greece

Vol. 36, 2004



MAPPING OF CODA ATTENUATION AT THE EXTEND OF THE NATIONAL SEISMOLOGICAL NETWORK OF GREECE

Baskoutas I.	National Observatory of Athens
Herraiz M.	Dep. de Geofísica y Meteorología, Universidad Complutense de Madrid
Perez A.	Dep. de Geofísica y Meteorología, Universidad Complutense de Madrid
Kalogeras I.	National Observatory of Athens
Panopoulou G.	National Observatory of Athens
Sachpazi M.	National Observatory of Athens
Papadopoulos G.	Earthquake Planning and Protection Organization, Seismotectonic Div.

<https://doi.org/10.12681/bgsg.16503>

Copyright © 2018 I. Baskoutas, M. Herraiz, A. C. Perez, I. Kalogeras, G. Panopoulou, M. Sachpazi, G. Papadopoulos



To cite this article:

Baskoutas, I., Herraiz, M., Perez, A., Kalogeras, I., Panopoulou, G., Sachpazi, M., & Papadopoulos, G. (2004). MAPPING OF CODA ATTENUATION AT THE EXTEND OF THE NATIONAL SEISMOLOGICAL NETWORK OF GREECE. *Bulletin of the Geological Society of Greece*, 36(3), 1371-1378. doi:<https://doi.org/10.12681/bgsg.16503>



MAPPING OF CODA ATTENUATION AT THE EXTEND OF THE NATIONAL SEISMOLOGICAL NETWORK OF GREECE

Baskoutas I.¹, Herraiz M.², Pérez A.C.², Kalogeras I.¹, Panopoulou G.¹, Sachpazi M.¹ and Papadopoulos G.³

¹ National Observatory of Athens, P.O. Box 20048 118 10 Athens i.basko@gein.noa.gr,

² Dep. de Geofísica y Meteorología, Universidad Complutense de Madrid, 28040, Madrid, mherraiz@fis.ucm.es

³ Earthquake Planning and Protection Organization, Seismotectonic Div., Xanthou32, 15451 Athens

ABSTRACT

Coda decay rates of 538 vertical components corresponding to local earthquakes which occurred in Greece during the period 1998 to 1999 were used to deduce the coda quality factor (Q_c) characteristics in the Hellenic area. The seismograms have been selected from a broader sample of 776 records obtained at 8 stations of the National Seismographic Network operated by the Institute of Geodynamics of the National Observatory of Athens. Earthquake magnitudes range from 2.5 to 4.0; epicentral distances and depths are smaller than 100 km and 40 km, respectively. Using the Single Back Scattering model, the dependence of Q_c on frequencies between 1 and 10 Hz has been investigated at each station and the usual $Q_c = Q_0 f^n$ relationships have been deduced. The spatial distribution of Q_0 has been drawn using waves that sample approximately equivalent ellipsoidal volumes with semiminor axis up to 100 km. The corresponding map shows a decreasing trend in S-N direction.

1 INTRODUCTION

In spite of their theoretical limitations, the analysis of waves of coda has been considered an effective method to estimate quality factor of seismic wave attenuation. Nevertheless, the influence of some analysis parameters like length of the time window and selection of the beginning of coda, affects the results.

In this sense obtained values are not easy to compare even within the same territory.

Besides that, results for most of territories concerns limited extent of areas or lack of uniform seismological data. To overcome such a difficulty, the spatial distribution of Q in an extensive zone can be obtained by applying the same methodology to a uniform set of data in the extent of dense national seismological network.

Modern instrumentation offers the possibility to analyze the spatial distribution of Q_c (Q factor considered from waves coda) in the Greek territory, area for which numerous studies exists (see APPENDIX table, from Panagiotopoulos et al., 1998., completed by the authors) but, nevertheless, in zones of small size, usually covered from locally operating seismological networks, for which the temporarily character doesn't permit any future comparison. (Martin, 1988, Tselentis et., al., 1988, Baskoutas et., al., 1989 ; 1992 ; 1993; 1995; Hatzidimitriou, 1993; 1995; Baskoutas, 1996; 1998; Tselentis, 1997).

Single Back-Scattering model has been chosen to be used in the present study because it facilitates the comparison with other studies made in different zones within Greek territory and elsewhere.

In order to obtain the spatial distribution of Q_c values, data with good azimuthal distribution are selected to cover homogeneously the space around the seismological stations.

Tests have been made by running both the CODAQ Havskov's (1989) and Baskoutas software (1993) as well as the Single Isotropic scattering model (Sato, 1977a,b) at a representative set of data. No significant differences have been noticed.

2 DATA

We used 538 earthquakes which occurred in the period 1998 to 1999 and were recorded by eight seismological stations of the National Seismic Network of Greece.




Figure 1 Epicenters map (The locations of the stations is shown by triangles).

A first set of 776 earthquakes were obtained from vertical components at eight stations of the National Seismic Network of Greece. Magnitudes are between 2,5 and 4,0. Hypocenter depths and epicenter distances are less than 40 km and 100 km respectively (Figure 1). All events were recorded by broadband seismometers of three components. In both cases the sampling frequency is 50 sample/sec. Seismograms were inspected visually for errors (i.e. noise contamination and location parameters). Therefore some of them have been rejected and a total of 538 events have been selected (Table 1).

Table 1. Stations and number of events.

Station Name	N° of Events
ARG	55
ITM	74
JAN	53
JAN	63
KZN	45
PLG	51
PRK	79
VLS	118

The noise level was evaluated considering the average value of the spectrum amplitude corresponding to 10 second before the arrival time of P- wave. S wave arrival was identified on the horizontal components.

3 ANALYSIS AND RESULTS

As a first step we studied at each station, the variation of Q value for six different parameter sets resulting from combinations between chosen time window and coda onsets i.e. considering three different length of time window (10, 20 and 30s) and two coda waves origin time ($t = 1.5ts$ and $t = 2ts$). The analysis was carried out using the model of Single Back-Scattering (Aki and Chouet, 1975). Signals are filtered by the mean of band-pass filter with center frequencies at 1.5, 3, 5, 7 and 9 Hz. Analysis has shown that the most stable results were obtained, using as coda wave origin time the value of $2ts$ and a length of lapse time window 20 sec. (Perez, 2000). Table 2 shows the results for Q_0 for entire coda length (value of Q_c at 1 Hz) and n corresponding to the power law relation for each station.

$$Q_c = Q_0 f^n \quad (1)$$

For Q_0 the spatial distribution of the common sampling volumes for each event was considered. This volume depends on the lapse time interval t . In this way scatterers that take part in the process of coda waves generation, are located, in a volume of an expanding ellipsoid, with foci the station and the source (Pulli, 1984).

$$\frac{x^2}{(v_s t/2)^2} + \frac{y^2}{(v_s t/2)^2 - R^2/4} = 1 \quad (2)$$

In the relation (2) V_s is the shear wave velocity, t is the travel time interval and R is the source-station distance. The semiaxis a_1 and a_2 are equal to $(v_s t/2)^2$ and $a_1^2 - R^2/4$ respectively.

Examined ellipsoids can be practically considered spherical ($a_1/a_2 \approx 1$) thus maximum depth can be defined as $z_{max} = a_2 + h$, where h is the depth of the earthquake (Havskov et., al., 1989). S wave velocities with respect to the depth are considered as follow:

$$\begin{aligned} h \leq 15 \text{ km} &\Rightarrow v_s = 3.47 \text{ km/s} \\ 15 < h \leq 40 \text{ km} &\Rightarrow v_s = 3.90 \text{ km/s,} \end{aligned}$$

which correspond to the regional velocity model, used by the Geodynamic Institute of National Observatory of Athens, for the earthquake location.

For the analysis two restrictions were adopted. First, the dimension of the sampling volume, which should mostly correspond to the crust and second, having at the same time a sufficient number of events to be processed, which guarantees for the results significance. The optimal value between these two conditions was $a_2 \leq 100\text{km}$. Although the number of earthquakes for each station (values between parenthesis in Table 3) in some cases, was reduced drastically, still it can be considered sufficient for the study. Figures 2 and 3 shows Q_0 and n values, for different ellipsoid semi axis values (100, 130, 160Km) at each station of the Seismological network. Open circles stand for $a_2 = 100\text{Km}$, full rectangles and triangles for 130 and 160Km, respectively. The clear difference of Q_0 values among the group of stations situated in the concave part of the island arc (JAN KZN, PLG, PRK), and those situated in the convex part (ARG, ITM, NPSM and VLS), can be seen in figure 2.

Table 2. Q_0 and n values

Station	Q_0	n
ARG	83±1	0.89±8e2
ITM	99±8	0.63±5e-2
JAN	67±4	0.71±5e-2
KZN	67±9	0.63±9e-2
NPS	76±2	0.89±2e-2
PLG	71±5	0.73±6e-2
PRK	91±8	0.80±6e-2
VLS	78±2	0.82±2e-2

Table 3. Q_0 and n values for different values of a_2 parameter

Station	$a_2 \leq 100$ km			$a_2 \leq 130$ km			$a_2 \leq 160$ km		
	Q_0	n	No of cases	Q_0	n	No of cases	Q_0	n	No of cases
ARG	77±12	0.84±7E-2	(4)	77±6	0.89±6E-2	(21)	71±5	0.945±1E-3	(41)
ITM	71±10	0.644±1E-3	(9)	83±7	0.64±7E-2	(24)	92±7	0.63±4E-2	(48)
JAN	59±3	0.71±5E-2	(31)	59±3	0.75±5E-2	(42)	63±4	0.74±5E-2	(47)
KZN	59±7	0.68±7E-2	(19)	59±7	0.68±7E-2	(21)	63±8	0.67±8E-2	(34)
NPS	69±4	0.77±3E-2	(12)	71±5	0.87±5E-2	(23)	72±3	0.90±3E-2	(36)
PLG	60±2	0.77±2E-2	(14)	63±1	0.760±9E-3	(25)	71±5	0.74±5E-2	(38)
PRK	59±7	0.91±8E-2	(12)	82±5	0.81±4E-2	(31)	87±6	0.79±4E-2	(42)
VLS	76±3	0.68±2E-2	(21)	76±2	0.78±1E-2	(42)	79±3	0.79±2E-2	(75)




Figure 2. Q_0 at the NOA seismic stations for different ellipsoid semi axis values ($a_2 = 100$ km, open circles). Full rectangles and triangles represents the Q_0 values for a_2 equal to 130 and 160 km respectively.




Figure 3. n values at the NOA seismic station for different ellipsoid semi axis values, ($a_2 = 100$ km, open circles) . Full rectangles and triangles represents the Q_0 values for a_2 equal to 130 and 160km respectively.

4 DISCUSSION AND CONCLUSIONS

The obtained results show that the Hellenic region is characterized by a high seismic attenuation. Q values of the table 2, summarizes the data obtained for all earthquakes in each station without considering any restrictions for the sampling volume. These results are, in general, coherent with those obtained in previous studies, when the comparison is possible. Thus, Hatzidimitriou (1993), using a time window of 15-30s, has obtained, for north Greece, $Q_0 = 60$ and $n = 0,79$ that is very similar to the results of this work for PLG station. The similarity increases when $a_2 = 100$ Km is considered.

In the case of Kozani station also coherence exists, within the margin of error, with the results of Baskoutas et. al., (1998) for the window of 20s. Instead, for Peloponnesus area and for unknown length of time window, Martin (1988) reports $Q_0 = 73$ and $n = 0,79$, that differs from the results of this work at ITM station, located in the western part of the zone. In several cases comparison with results of other authors is not easy, because of missing information or of different parameters set used for the data analysis.




Figure 5. Spatial distribution of Q_0 for $a_2 \leq 100$ km




Figure 6. Spatial variation of b value from superficial seismic activity, (Papazachos and Papazachou 1997).

Obtained spatial variation for $a_2 \leq 100$ km (Figure 5) shows a significant attenuation characteristic for Greek territory, and indicates that attenuation in the concave zone is larger than in the convex one.

In figure 5, also, a sharp Q_0 variation in the Ionian Sea islands is observed with a slight increase towards the N-S direction. This area is characterized by high seismic activity and is influenced by different tectonic regimes. It is also important to notice that this tendency keeps a remarkable similarity with the spatial variation (Figure 6) of b value given by Papazachos and Papazachou (1997) for shallow seismic activity. Tendency indicating that b value decrease in a systematic way from the southwest to the northeast. This result indicates that the values of Q_c tend to diminish in the zones in which the geological formation age is older (Baskoutas et al., 2000).

All the values assigned to the semiaxis a_2 imply that the sampling volume comprises also part of the upper mantle playing an important role to the increase of Q_c values, the contribution of the crust must be very small. In order to confirm the above statement, it is necessary to diminish axis a_2 (and thus the upper mantle contribution) using higher number of events.

Although the differences are small, the comparison of the results obtained for different values of the sampling volume (Table 3) confirms the increase of Q_0 as depth increases.

ACKNOWLEDGEMENTS

This work has been carried out within the frame of the bilateral scientific and technological collaboration between Greece and Spain 1996-1998, supported by the General Secretary of Research and Technology, Greece (code number 7557/09-07-1997).

REFERENCES

- Aki, K. & Chouet B. 1975. Origin of coda waves: source, attenuation, & scattering effects. *Journal of Geophysical Research*, 80, 23, 3322-3342.
- Baskoutas I. 1993. Contribution on the study of the attenuation of seismic coda in central Greece. Ph.D Thesis, University of Athens, 246pp.
- Baskoutas I., Makropoulos K. & Sato H. 1995. Mean free path for S-wave scattering under central Greece. *Boll. di Geof. Teor. ed Appl.*, 145, 73-80.
- Baskoutas I. 1996. Dependence of coda attenuation on frequency and lapse time in central Greece". *Pure Appl. Geophys.*, 147, 3, 483-496.

- Baskoutas I. 1998. Study of spatial distribution of coda Q with respect to the seismicity in central Greece. *Bull. Geol. Soc. Greece*, 32/1 197-203.
- Baskoutas I., Makropoulos K., Drakopoulos J. & V. Kouskouna. 1989. Evaluation of seismic Q in central Greece from local events. *Procced Int. Symp. on the Analysis of Seismicity and Seismic Risk*, Bechyne Castle, Czechoslovakia, Sep. 4-9, 1989, 162-170.
- Baskoutas I., Panopoulou G., Makropoulos K. & J. Drakopoulos. 1992. Analysis of seismic coda by using seismic analog data of WSSN ATH station. 6th Congress of the Hellenic Geological Society, Athens, (in Greek).
- Baskoutas I., Stavrakakis G.N. & I. S. Kalogeras. 1998. Q factor estimation from the aftershock sequence of the 13 May 1995 Kozani earthquake. *J. Geodynamics*, 26, 2-4, 367-374.
- Baskoutas I., Panopoulou, G., & Th. Rontoyanni. 2000. The seismic attenuation in two regions with different geological aspect. *Boll. Geof. Theor. Appl.*, 41, 3, 233-242.
- Hatzidimitriou P. M. 1993. Attenuation of coda waves in northern Greece. *Pageoph.*, 140, 63-78.
- Hatzidimitriou P.M. 1995. S-wave attenuation in the crust in northern Greece. *Bull. Seism. Soc. Am.* 85, 5, 1381-1387.
- Havskov J., Malone S., McClurg D. & Crosson R. 1989. Coda Q for the state of Washington, *Bull. Seism. Soc. Am.* 79, 4, 1024-1038.
- Kovachev, S.A., Kuzin, I.P., Shoda, O.Yu. and Soloviev, S.L. 1991. Attenuation of S waves in the lithosphere of the sea of Crete according to OBS observations. *Phys. Earth Planet. Interiors*, 69, 101-111.
- Martin C. 1988. Geometrie et cinématique de la subduction Egeenne; structure en vitesse et en atténuation sous le Peloponnese. These Univ. Joseph Fourier, Grenoble.
- Παναγιωτόπουλος, Δ. Γ., Χατζηδημητρίου, Π. Μ. και Τσοκος, Γρ. 1998. Δομή της λιθόσφαιρας στο Αιγαίο και τις γύρω περιοχές; Με βάση τις ταχύτητες των σεισμικών κυμάτων, την κατανομή της πυκνότητας και την απόσβεση των σεισμικών κυμάτων. «Βασικά αποτελέσματα της σεισμολογικής έρευνας στην Ελλάδα», Εκδόσεις Ζήτη, Θεσσαλονίκη.
- Papazachos, C.B. 1992. Anisotropic radiation modelling of macroseismic intensities for estimation of the attenuation structure of the upper crust in Greece. *Pure Appl. Geophys.*, 138, 3, 445-470.
- Papazachos B. & Papazachou C. 1997. The Earthquakes of Greece. Editions Ziti. Thessaloniki.
- Papazachos B. Papaioannou Ch.A. Papazachos C. & Savvaidis A.S. 1997. Atlas of isoseismal maps for strong $M \geq 5$ shallow $h < 60$ km earthquakes in Greece and surrounding area 426 BC-1995. *Geophys. Lab., Univ. of Thessaloniki*, 4, 187pp.
- Pérez A.C. 2000. Atenuación de ondas Coda en Grecia. Trabajo de Investigación dentro del programa de Doctorado de Geodesia, Geofísica y Meteorología. Universidad Complutense de Madrid, 109 pp.
- Pulli J.J. 1984. Attenuation of coda waves in New England *Bull. Seism. Soc. Am.*, 74, 4, 1149-1166.
- Sato H. 1977a. Energy propagation including scattering effects single isotropic scattering approximation. *J. Phys. Earth*, 25, 27-41.
- Sato H. 1977b. Single isotropic scattering model including wave conversions simple theoretical model of the short period body wave propagation. *J. Phys. Earth*, 25, 163-176.
- Tselentis G.A. Drakopoulos J., & Makropoulos C. 1988. On the frequency dependence of Q in the Kalamata South Greece region as obtained from the analysis of the coda of the aftershocks of the Kalamata 1986 earthquake. *Tectonophysics*, 152, 157-159.
- Tselentis G.A. 1993b. Depth dependent seismic attenuation in western Greece. *Tectonophysics* 225, 523-528.
- Tselentis G.A. 1997. Evidence for stability in coda Q associated with the Egeion central Greece earthquake of 15 June 1995. *Bull. Seism. Soc. Am.*, 87, 6, 1679-1684.

APPENDIX

Table of Q_c power low estimates in the territory Greek.

Lapse time or distance	$Q_c = Q_0 f^n$	Region	Reference
10-20 sec	$Q_c = 33f^{1.01}$	North Greece	1
15-30 sec	$Q_c = 60f^{0.79}$	"	"
20-45 sec	$Q_c = 89f^{0.72}$	"	"
30-60 sec	$Q_c = 94f^{0.78}$	"	"
50-100sec	$Q_c = 128f^{0.74}$	"	"
20 sec	$Q_c = 47f^{1.02}$	Kozani-Grevena	2
40 sec	$Q_c = 80f^{0.87}$	"	"
60 sec	$Q_c = 100f^{0.78}$	"	"
80 sec	$Q_c = 124f^{0.70}$	"	"
100 sec	$Q_c = 140f^{0.71}$	"	"
10 sec	$Q_c = 29f^{0.91}$	Central Greece	3
20 sec	$Q_c = 45f^{0.85}$	"	"
30 sec	$Q_c = 62f^{0.83}$	"	"
40 sec	$Q_c = 80f^{0.80}$	"	"
50 sec	$Q_c = 101f^{0.73}$	"	"
-	$Q_{f=1Hz} = 182$	Athens	4
-	$Q_c = 73f^{0.95}$	Peloponnesus	5
30 sec	$Q_c = 121f^{0.72}$	Gulf of Patras	6
50 sec	$Q_c = 141f^{0.75}$	"	"
70 sec	$Q_c = 144f^{0.81}$	"	"
90 sec	$Q_c = 162f^{0.83}$	"	"
10-20 sec	$Q_c = 16f^{1.1}$	West part of Korinth Gulf	7
-	$Q_{c(f=4Hz)} = 183$	Aegion	8
-	$Q_{c(f=8Hz)} = 270$	"	"
-	$Q_c = 43f^{0.81}$	Kalamata	9
0-450Km	$Q_{f=1Hz} = 350$	Aegean sea	10
* 30-250 Km	$Q_{f=8Hz} = 200-300$	NA part of Hellenic arc	11

(1) Hatzidimitriou, 1993, (2) Baskoutas et al., 1998, (3) Baskoutas, 1996, (4) Baskoutas et al. 1992, (5) Martin 1988, (6) Tselentis 1993b, (7) Baskoutas et al., 1994, (8) Tselentis, 1997, (9) Tselentis et al., 1988b, (10) Papazachos C, 1992, (11) Kovachev et al., 1991.