University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Historical Materials from University of Nebraska-Lincoln Extension

Extension

1991

G91-1015 Perennials

Don Steinegger University of Nebraska--Lincoln, dsteinegger1@unl.edu

Anne Streich University of Nebraska--Lincoln, astreich2@unl.edu

Follow this and additional works at: https://digitalcommons.unl.edu/extensionhist




Part of the Agriculture Commons, and the Curriculum and Instruction Commons

Steinegger, Don and Streich, Anne, "G91-1015 Perennials" (1991). Historical Materials from University of Nebraska-Lincoln Extension. 990.

https://digitalcommons.unl.edu/extensionhist/990

This Article is brought to you for free and open access by the Extension at DigitalCommons@University of Nebraska -Lincoln. It has been accepted for inclusion in Historical Materials from University of Nebraska-Lincoln Extension by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



Perennials

This NebGuide describes perennial flowers that will withstand Nebraska's winters outdoors and that will bloom within the growing season.

Donald H. Steinegger, Extension Horticulturist Anne M. Streich, Extension Horticulturist-Outreach Programs

Herbaceous perennials normally live for many years under local growing conditions, dying back to the ground each winter. They vary in height, leaf texture and flower color, and are the backbone of the flower garden for many home gardeners.

The following is a list of perennials best adapted to Nebraska's broad range of growing conditions. No single list of plant materials is suitable for every garden, however, and deciding which plants to use depends on personal taste.

COMMON NAME	BOTANICAL NAME	SITE	HEIGHT & FORM	FLOWER COLOR	COMMENTS
Anemone, Japanese	Anemone vitifolia 'Robustissima'	Sun, partial shade	30 in. high	Pink	Fall blooming.
Artemisia	Artemisia spp.	Sun; tolerates dry conditions; low nitrogen soils	Rhizomatous; 6 to 12 in. high; spreads to 12 or more inches	not	Silvery-gray, finely cut foliage. Foliage used in fresh flower arrangements. Several other native species. Many cultivars.
Aster, Fendler's	Aster fendleri	Sun	4 - 12 in.	Lavender	Native plant.
Astilbe	Astilbe spp.	Part shade; prefers cool, moist	6 in 3 ft.	White, pink, red	Many fine cultivars.

		conditions			
Baby's Breath	Gypsophila paniculata	Sun; moist soil	Fine textured; 1 1/2 - 3 ft. high, 4 ft. across; has stout rhizomes.	White, pink	Fresh and dried flower arrangements. Double flowering forms are usually grafted; place graft union 1 in. under soil. Takes two years to become established. Many useful cultivars.
Balloon Flower	Platycodon grandiflorus cv. 'Mariesii' cv. 'Shell pink'	Sun; partial shade; light; moist, well- drained soil with high organic matter	2 - 3 ft. high, 2 ft. wide	Blue, white, pink	Double flowering forms available. Late to emerge in spring; mark location to avoid injury from cultivation.
Basket-of-Gold	Aurinia saxatilis	Sun; partial shade; dry soil, best in low nitrogen soils	6 - 12 in., prostrate	Yellow	Silvery-gray foliage. Early spring flowering.
Bearded Iris	Iris spp.	Sun; average soil	2 - 3 ft. high	Broad color range	Many cultivars available.
Bee Balm	Monarda didyma	Sun; partial shade; somewhat moist, well- drained, high organic soil	3 - 4 ft. high	Purple, red and pink combinations, salmon	Minty fragrance. Can become invasive in moist sites. Many cultivars; grow cultivars rather than species.

Peach-leaved	Campanula persicifolia	Sun to semi- shade; fertile soil	Erect, leafy clump; 2 ft.	Deep blue to white	Many useful species and cultivars. Flowers for fresh arrangements. Many useful cultivars available.
Clustered	Campanula glomerata	Sun to semi- shade; moderate fertility	Spreads by rhizomes; 1 - 3 ft.; coarse erect growth	Deep blue to white	
Bergenia	Bergenia cordifolia	Sun to shade	12 in. high	Pink	Glossy green foliage; burgundy winter color. Best in winter shade.
Blackberry Lily	Belamcanda chinensis	Sun; fertile, sandy soil	Clump forming rhizomes; 2 - 3 ft. high; long narrow foliage- resembles German Iris.	Deep orange, red dotted	Fresh flower arrangements; seed clusters dried for flower arrangements.
Blanket Flower	Gaillardia x grandiflora	Sun; requires good air circulation; light soil; avoid high nitrogen soil	6 in 3 ft. high; high; erect or sprawling branches.	Yellow or red	Easy to grow. Generally short-lived.
Blazing Star	Liatris spicata	Sun to partial shade; sandy fertile soil	2 - 3 ft. high; stiff, erect growth form	Rosy purple	Fresh flower arrangements. Color accent in perennial borders.

		is best, althought it's tolerant of other soils			Several other native species.
Bleeding Heart	Dicentra eximia	Partial shade to shade; fertile, moist soil	12 - 15 in. high clump former; short rhizomes	Pink, purple or white; gray-blue foliage	Several cultivars available. Cultivars are not superior to the species.
Blue False Indigo	Baptisia australis	Sun; partial shade; dry site; low nitrogen soil.	Erect, forms large clumps; 3 - 4 ft. high, 2 - 3 ft. wide	Indigo blue, blue-green foliage	Excellent summertime hedge. Stabilizes soil on bank. Foliage used in fresh flower arrangements. Pods used in dried flower arrangements. Plant may require staking.
Boltonia	Boltonia asteroides	Sun, partial shade	Up to 4 ft. high	White, pink	Large background plant; fall blooming.
Butterfly Milkweed	Asclepias tuberosa	Sun; sandy loam; dry location	Broadly domed bushy clump; 1 1/2 - 3 ft. high	Bright orange	Tuber forming plant. Does not transplant successfully when large. Seed pods used for dried arrangements. Flowers attract butterflies.
Columbine	Aquilegia spp.	Sun or light shade; cool, moist soil	1 1/2 - 3 ft.; large clumps; delicate lacy foliage	Broad color range	Leaf miner is a problem. Flowers May through June. Many useful

Coneflower, Grayhead	Ratibida pinnata	Sun	4 - 5 ft., 2 ft.	Yellow	species and cultivars of varying heights and flower forms. Drooping
Prairie			wide		petals; native plant.
Coneflower, Purple	Echinacea purpurea	Sun	2 - 3 ft. high	Purple	Coarse texture; native plant.
Coral Bells	Heuchera sanguinea	Full sun to filtered shade; fertile soil high in organic matter	8 - 10 in.; flower scapes to 24 in.	White, red and shades of red	Fillers in bouquets. Susceptible to crown rot. Remove dried flowers to reduce infection.
Coreopsis	Coreopsis lanceolata	Sun	2 - 3 ft.	Yellow	Cultivars: 'Early Sunrise', 'Sunray'.
Crane's Bill	Geranium spp.	Sun to partial shade; tolerates dry, less fertile soil	12 - 18 in. high; thick ascending stems	Lilac, blue, magenta, white, purple	Nearly indestructible; spreads excessively on high nitrogen soils. Several ornamental species and cultivars.
Daylily	Hemerocallis spp.	Sun to partial shade in afternoon; medium heavy loam	Forms large clumps; clumps; 18 in. to 5 ft. high	Broad color range	Tolerates neglect. Several species; hundreds of useful cultivars.

Dianthus	Dianthus spp.	Sun; rich, slightly alkaline soil requires good air circulation	Densely tufted; 6 - 15 in. high; trailing growth habit; creeping rhizomes	Deep pink, rarely white or red	Fragrant flowers; many cultivars.
False Sun Flower	Heliopsis helianthoides	Sun or partial shade; high organic matter	3 - 5 ft. high; erect stems; clump former	Yellow- orange	Many available cultivars that are useful in dry sites. Easy to grow.
Gaura	Gaura lindheimeri	Sun	3 ft., 2 ft. wide	White, pink, rose	Flowers open white, deepen to rose; native plant; may need stakes; Cultivars: 'Siskiyou Pink', 'Corrie's Gold'.
Goatsbeard	Aruncus dioicus	Shade	2 - 4 ft. high	White	Showy white panicles.
Hardy Chrysanthemum	Chrysanthemum morifolium	Sun	9 in. to 3 ft. or more in height	Broad color range	Highly variable in height, plant and flower form. Can readily transplant in full flower.

					Winter protection recommended. Hundreds of cultivars.
Hardy Ice Plant	Delosperma congestum	Sun, afternoon shade	2 in., spreads 30 in.	Yellow	Succulent leaves turn bright red in winter; Cultivar: 'Gold Nugget'.
Hosta	Hosta	Partial shade to shade	6 in. to 3 ft. high	White, lavender	Many cultivars.
Lady's Mantle	Alchemilla mollis	Part shade; tolerates sun with moist conditions	12 in. high;rounded form	Yellow	Beautiful foliage.
Lead Plant	Amorpha canescens	Sun	3 ft., dense	Lavender- blue	Dense, white hairy leaflets.
Missouri Primrose	Oenothera missouriensis	Sun; tolerates dry conditions	12 in. high	Yellow	Large flowers, native plant, sprawling growth habit.
New York Aster	Aster novi-belgii	Sun; moist, fertile soil	Rhizomatous; varies from small and slender to large and stout 1 1/2 - 3 ft.1 1/2 - 3 ft.	Broad color range	Requires division each year; subject to rust, Aster yellows and mildew. Hundreds of cultivars and several other useful species and hybrids

					available to gardeners.
Oriental Poppy	Papaver orientale	Sun; fertile, well- drained soil	3 - 4 ft. high	Orange, pink, salmon, white	Foliage dies back during summer. Numerous cultivars.
Painted Daisy	Chrysanthemum coccineum	Sun; soil with organic matter; fertile soil	2 - 3 ft.; upright, long stems	Pink, red or white; single or double	Aphids may be a problem. Excellent cut flowers. Many useful cultivars.
Penstemon	Penstemon spp.	Sun; prefers well drained conditions	1 1/2 - 3 ft.	Broad color range	Neat foliage, often glossy, somewhat evergreen. Cultivars: 'Husker Red', 'Prairie Splendor'.
Peony	Paeonia lactiflora	Sun to partial shade; deep, rich, drained soil; high organic matter; protect from wind	2 - 3 ft. high; rhizomatous	Broad color range	Hundreds of cultivars; many flower forms from singles to doubles. Very popular perennial.
Perennial Flax	Linum perenne	Sun; light, well- drained, moist and fertile soil	1 - 2 1/2 ft. high; gracefully arching stems	Blue	Not suitable for cut flowers. Other species of Linum make valuable garden plants.
Pitcher's Sage	Salvia azurea	Sun	3 - 4 ft., 2 ft. wide	Blue-violet	Native plant; Cultivar: 'Nekan', 'Grandiflora'.
Prairie Ragwort	Senecio plattenensis	Sun	16 - 18 in.	Yellow, orange center	Basal leaves have maroon undersides.
Purple Poppy-mallow	Callirhoe involucrata	Sun	18 in.	Purple	Native groundcover.

Rudbeckia, Goldstrum	Rudbeckia fulgida `Goldsturm'	Sun	24 in. high	Yellow	Compact, long-blooming form.
Russian Sage	Perovskia atriplicifolia	Sun; tolerates dry conditions	3 - 5 ft. high	Lavender	Sub-shrub form with silver gray foliage; no pests.
Sedum	Sedum spp.	Sun to light shade; drought tolerant	4 - 24 in. high	Pink, white, red, yellow	Good cut flower. Cultivars bearing different flower colors available.
Shasta Daisy	Chrysanthemum maximum	Sun to partial shade for double flowering forms	2 - 4 ft. high from a rosette	White	Good cut flower; many cultivars.
Snakeroot, Black	Cimicifuga racemosa	Partial shade to shade	2 - 3 ft. high	White	Handsome, broad leaves.
Sneezeweed	Helenium autumnale	Sun; moist soil; moderate fertility; high organic matter	2 - 4 ft. high; rank growth	Yellow, reddish- brown, crimson	Late summer bloom.
Spiderwort	Tradescantia spp.	Shade or sun	1 1/2 - 2 ft. high	White, blue or rose	Easy to grow.
Statice, German	Goniolimon tatarica	Sun	18 in. high	White	For dried arrangements.
Statice, Sea Lavendar	Limonium latifolium	Sun	18 - 30 in. high; rounded form	Lavender	Open, light texture.
Threadleaf Coreopsis	Coreopsis	Sun; moist,	Clump	Yellow	Popular

	verticillata	fertile soil	forming; 18 - 24 in. in height		cultivars are "Moonbeam," a pale yellow, and "Zagreb," a dark yellow.
Veronica, Speedwell	Veronica spp.	Sun to partial shade	2 - 4 ft. high	Blue, white, red	Many hybrids and cultivars available. Good garden plant.
Yarrow	Achillea millefoliium	Sun; dry soil; best in low nitrogen soils	18 - 30 in. high; stiff erect stems	Red, cream, yellow	Gray-green foliage. Excellent for cutting or drying. Flowers from June to September.

File G1015 under: HORTICULTURE

A-36, Ornamentals

Revised April 1999, 2,000

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Elbert C. Dickey, Interim Dean and Director of Cooperative Extension, University of Nebraska, Institute of Agriculture and Natural Resources.

University of Nebraska Cooperative Extension educational programs abide with the nondiscrimination policies of the University of Nebraska-Lincoln and the United States Department of Agriculture.