

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USGS Northern Prairie Wildlife Research Center

US Geological Survey

2008

Restoration of Mangrove Plantations and Colonisation by Native Species in Leizhou Bay, South China

Hai Ren

Heshan National Field Research Station of Forest Ecosystem, South China Botanical Garden, The Chinese Academy of Sciences, Guangzhou 510650, China

Shuguang Jian

Heshan National Field Research Station of Forest Ecosystem, South China Botanical Garden, The Chinese Academy of Sciences, Guangzhou 510650, China

Hongfang Lu

Heshan National Field Research Station of Forest Ecosystem, South China Botanical Garden, The Chinese Academy of Sciences, Guangzhou 510650, China

Qianmei Zhang

Heshan National Field Research Station of Forest Ecosystem, South China Botanical Garden, The Chinese Academy of Sciences, Guangzhou 510650, China

Weijun Shen

Heshan National Field Research Station of Forest Ecosystem, South China Botanical Garden, The Chinese Academy of Sciences, Guangzhou 510650, China

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/usgspwrc>

 Part of the [Other International and Area Studies Commons](#)

Ren, Hai; Jian, Shuguang; Lu, Hongfang; Zhang, Qianmei; Shen, Weijun; Han, Weidong; Yin, Zuoyun; and Guo, Qinfeng, "Restoration of Mangrove Plantations and Colonisation by Native Species in Leizhou Bay, South China" (2008). *USGS Northern Prairie Wildlife Research Center*. 15.

<https://digitalcommons.unl.edu/usgspwrc/15>

This Article is brought to you for free and open access by the US Geological Survey at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USGS Northern Prairie Wildlife Research Center by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Hai Ren, Shuguang Jian, Hongfang Lu, Qianmei Zhang, Weijun Shen, Weidong Han, Zuoyun Yin, and Qinfeng Guo

Hai Ren · Shuguang Jian · Hongfang Lu
Qianmei Zhang · Weijun Shen · Weidong Han
Zuoyun Yin · Qinfeng Guo

Restoration of mangrove plantations and colonisation by native species in Leizhou bay, South China

Received: 12 March 2007 / Accepted: 10 May 2007 / Published online: 27 June 2007
© The Ecological Society of Japan 2007

Abstract To examine the natural colonisation of native mangrove species into remediated exotic mangrove stands in Leizhou Bay, South China, we compared soil physical–chemical properties, community structure and recruitments of barren mangrove areas, native mangrove species plantations, and exotic mangrove species—*Sonneratia apetala* Buch.Ham—between plantations and natural forest. We found that severely degraded mangrove stands could not regenerate naturally without human intervention due to severely altered local environments, whereas some native species had been recruited into the 4–10 year *S. apetala* plantations. In the first 10 years, the exotic species *S. apetala* grew better than native species such as *Rhizophora stylosa* Griff and *Kandelia candel* (Linn.) Druce. The mangrove plantation gradually affected soil physical and chemical properties during its recovery. The exotic *S. apetala* was more competitive than native species and its plantation was able to restore soil organic matter in about 14 years. Thus, *S. apetala* can be considered as a pioneer species to improve degraded habitats to facilitate recolonisation by native mangrove species. However, removal to control proliferation may be needed at late stages to facili-

tate growth of native species. To ensure sustainability of mangroves in South China, the existing mangrove wetlands must be managed as an ecosystem, with long-term scientific monitoring program in place.

Keywords Competitive exclusion · Mangrove · Ecosystem restoration · Invasive species · *Sonneratia apetala*

Introduction

Woody mangrove plants are well known for their well-developed morphological and physiological adaptations to their environmental conditions. Mangrove species grow at the interface between land and sea in tropical and sub-tropical regions with high salinity, brackish waters, and muddy, anaerobic soils, and play a very important role in the ecosystem processes of coastal areas. Mangroves create unique ecological environments that host rich assemblages of species, and also protect and stabilise coastlines, enrich coastal waters, yield commercial forest products, and support coastal fisheries (Snedaker 1984). However, accumulating evidence shows that mangrove species are particularly sensitive to human disturbances (Kovacs 1999; Lugo 1999; Zedler 2000; Kathiresan and Bingham 2001; Macintosh et al. 2002). In the past several decades, habitat destruction through human encroachment has been the primary cause of mangrove loss. Diversion of freshwater for irrigation and land reclamation have destroyed extensive mangrove forests, and heavy historical and continuing exploitation has severely damaged the remaining mangrove habitats. Early mangrove management has focussed on timber production and fuel wood, and mangroves have been managed for cultivation of fish, shrimp, and ecotourism. Because of this, conservation of remaining mangrove, and restoration of the most extensively degraded mangrove, has become a growing concern for ecologists, policy makers, and the general public (FAO 1994; Field 1999; Ellison 2000).

H. Ren (✉) · S. Jian · H. Lu · Q. Zhang · W. Shen · Z. Yin
Heshan National Field Research Station of Forest Ecosystem,
South China Botanical Garden, The Chinese Academy of Sciences,
Guangzhou 510650, China
E-mail: renhai@scib.ac.cn
Tel.: +86-20-37252916
Fax: +86-20-37252916

W. Han
Guangdong Ocean University, Zhanjiang,
524088 Guangdong, China

Q. Guo
Northern Prairie WRC, US Geological Survey,
Jamestown, ND 58401, USA

Q. Guo
Southern Research Station, USDA, Asheville,
NC 28804, USA

Field (1999) and Ellison (2000) have reviewed and summarised many mangrove remediation/restoration projects. Among the restoration projects, silviculture is the principal practice. Coastal stabilisation and environmental mitigation or remediation have often been the objectives (Lewis 2005). Many restored mangrove forests resemble forest plantations rather than truly integrated or natural mangrove ecosystems. These plantations could be used as a first step toward mangrove remediation (Ellison 2000). Along with the relative reduction of sea level and the physical and chemical amendment of the soils, the mangrove community succession is gradual and progressive, and species replacement is orderly and predictable, from the simple pioneer stage through to the final stage (Zedler 2000). Where degraded mangrove systems are revegetated, continued monitoring and thorough assessment are needed to help understanding the recovery processes. This knowledge will help developing strategies to promote better remediation of degraded mangrove. However, so far little has been done to monitor and assess the early development and growth of planted stands, especially when exotic species are planted and re-colonisation by native mangrove species occurs.

Guangdong province in China has important mangrove habitats, harbouring about 50% of Chinese mangrove (Lin 1999), over 80% of which are in the Leizhou Peninsula (Li 2003). The destruction of mangrove habitats has been caused by continual deforestation, fish/shrimp-pond building, and the extension of urban areas. The total area of mangrove ecosystem in Leizhou Peninsula decreased from about 14,027 ha to about 7,305.8 ha (including the 2,378.3 ha successfully remediated during 1991–2000) from 1956 to 2001. According to the State Forestry Administration of China, 2,000 ha mangrove will be planted per year during 2003–2007, with Leizhou Peninsula as the key area. Three classical remediation methods (natural remediation, indigenous species planting and exotic species introducing of mangrove) have been practised in the Leizhou Peninsula during the past 10 years (Han et al. 2003; Lin and Liu 2003).

The exotic mangrove species *Sonneratia apetala* Buch.Ham, was introduced to China in 1993. In this short period it has spread to Hainan, Guangdong, Guangxi, and Fujian provinces. More than 50 papers on *S. apetala* Buch.Ham plantation ecosystems were published between 1995 and 2005 (Lin et al. 2006). These reports described the morphology, geographical distribution, biological characteristics and adaptability of ecological factors. Some reported the ecological effects of *S. apetala* Buch.Ham plantation; those results showed that the soil type changes, the pH of the soil is reduced, and nitrogen, phosphorus, potassium and salt are increased. During the course of community forming, the biodiversity at first decreases and then increases (Zan et al. 2001a, b; Wang et al. 2002; Liao et al. 2004).

Here we compare the soil properties, community structure and recruitment of barren mangrove areas, native species plantations, and exotic mangrove species between planted and natural mangrove forests. In

particular, we evaluate the growth of the exotic mangrove species *S. apetala*, and the re-colonisation of non-planted native mangrove species.

Methods

The study area is situated on the Leizhou Bay coast, about 20 km from Leizhou City (109°03'E, 20°30'N), Guangdong, South China (Fig. 1). The annual average temperature is 22.9°C (28.4°C in July and 15.5°C in January) and the annual precipitation is 1,711 mm (about 73% in the rainy season, April–October, and 27% in the dry season). The soil is mangrove acid soil with three typical layers (0–10 cm: black, with organic matter, 10–40 cm: grey, and > 40 cm: yellow or mixed colour). The pH value for top soils is 6.9 (Han et al. 2001).

A survey in 2005 showed that there were a total of 1,606.9 ha mangrove area in Leizhou Bay but only 106.8 ha remain undisturbed. In total, 16 plant species of mangrove are found in the bay. The area of mangroves in the bay had declined by 1,675.6 ha during the period 1980–2005. On the whole, the mangrove area in the bay has decreased markedly, due mainly to its transformation to aquaculture, which has profoundly affected the development of the local society and economy. The mudflats and mangrove areas of Leizhou Peninsula were listed as key sites for artificial ecological remediation in the 1990s, and local coastal governments have taken effective planting initiatives for native species such as *Kandelia candel* and *Avicennia marina* (Forsk.) Vierh. Besides native species, *S. apetala*, a native of Bengal, was successfully introduced into Leizhou Bay in 1993. It was first introduced into Dongzhaigang mangrove Nature Reserve of Hainan in 1985, then into Leizhou Bay, where it grew well (Li et al. 1998; Wu et al. 2000; Chen et al. 2004; Liao et al. 2005; Lin et al. 2006).

Extensive native mangroves historically covered the area. However, due to human disturbances, only small patches of natural stands remain. Within the study area, a freshwater channel passes through the edge of the woodland and provides year-round fresh water from the drainage in a sea-dyke. Geomorphologically, the site is also at the landward side of the intertide. Regular tides affect the stand, and the ground surface is about 1.5 m under water at high tide, and 0.6–1.2 m above water at low tide.

The quadrats of natural mangrove forest, barren site, native species mangrove plantation, and 4-, 5-, 8- and 10-year exotic species—*S. apetala*—plantations, were placed around the estuary. The barren and natural forest sites were treated as controls due to the similarity of environmental conditions, close distance and site history. The native species mangrove plantation was established in 1993. Seedlings of *Rhizophora stylosa* and *K. candel* were separated to individual plants in each block along a transect. The space between planted seedlings was 2 m, giving 2,500 individuals/ha. The plants did not grow well and about 30% individuals died




Fig. 1 Locations of the study sites in Leizhou Bay, Guangdong, South China. The numbers represent different ages of plantation: 1 5-year plantation, 2 natural mangrove forest, 3 barren sites, 4

8-year plantation, 5 4-year plantation, 6 artificial native mangrove plantation, 7 10-year plantation

after several months. Plantations of *S. apetala* (seedlings) were established in 1995 (10-year), 1997 (8-year), 2000 (5-year), and 2001 (4-year). The space between planted seedlings was 2 m, again giving 2,500 individuals/ha. Most individuals survived and grew well. The natural forest sites were small patches of secondary natural stands with little human disturbance. The barren site was kept for natural remediation after the mangrove was cleared out 20 years ago.

In 1998, we established research quadrats for the study of biomass and community structure. We moved on to ecological restoration in 2004 and established research transects. The last field survey of all the communities was conducted in May 2005. We established seven transects across the whole land–sea interface zone for each ecosystem in the study area, and various numbers of 10 m × 10 m quadrats were placed along each transect depending on the length of the transects (number of quadrats: barren sites, 3; native plantation, 4; natural site, 3; exotic species plantation 4-year, 3; exotic species plantation 5-year, 7; exotic species plantation 8-year, 8; exotic species plantation 10-year, 3). For each of the four types of ecosystems (i.e. natural mangrove forest, barren site, native species mangrove plantation, and *S. apetala* plantations) and the four age classes (4-, 5-, 8- and 10-year) of *S. apetala* plantation, only three quadrats were selected for further analysis. The number of species, plant height, diameter at breast height (DBH—trees only), basal diameter, crown size,

and growth status (alive or dead) of each individual were recorded. Density, basal area and absolute frequency, the important value (IV), number of species, and Shannon–Weiner (SW) index were computed (Snedaker 1984; Krebs 1985).

Three 20-cm soil cores were taken from each of the 10 m × 10 m quadrats randomly to form a mixed sample, using a 3.7-cm diameter coring tube. Three replicates were taken from each quadrat at low tide. Water content was measured by the weight loss method. Salinity was measured using an optical refractometer, and temperature and pH were measured using an auto-thermometer and a pH meter, respectively. Samples were oven-dried at 80°C for about 3 days until constant dry weight was obtained, then stored for analysis. Analysis of organic matter, total N, total P, and clay followed Wartel et al. 1995 (see also Arnold 1986; Robert et al. 1997; Bosire et al. 2003).

The sediment characteristics data were assessed by a Tukey-test after one-way ANOVA ($P < 0.05$) to indicate significant differences. All statistics were analysed by SPSS11.5 software.

Results

Physical conditions

Among the ecosystems studied, the barren sites showed the harshest physical conditions for plant growth, i.e.

lowest organic matter, total N, and total P content, but highest proportion of sand compared to other sites. In contrast, the natural mangrove forests showed the best conditions as indicated by the high content of organic matter, total N and clay. In the planted stands of exotic species, all conditions improved with planting age, i.e. salinity, organic matter content, and proportion of clay all increased with planting age while pH declined, and there was no clear trend in total N and P. By the 10th year, most conditions in the exotic plantations had become similar to those found in natural mangrove. This suggested that there were significant effects of mangrove plantation on soil physical and chemical properties during the course of recovery (Table 1). The improvement in soil physical and chemical properties in plantations facilitated the invasion of native mangrove species. Regression analysis showed that the organic matter content of plantations could reach that of natural mangrove communities in about 14 years (Fig. 2).

Community structure

In the past 10 years, no plants (including mangrove species) were found in the barren sites (Table 2). Pure stands of *R. stylosa* and *K. candel* were planted separately but neither species colonised or invaded each other's stands, although in natural mangroves *A. marina*, *K. candel*, *Aegiceras corniculatum* (Linn.) Blanco and *R. stylosa* are the dominant species.

In the *S. apetala* plantations, the planted species was still the single dominant species after 10 years, and both species richness and density of mangroves were lower than that in natural mangrove communities, although most element conditions were already restored.

Recruitment

We did not observe any native mangrove species seedlings in 1–3 year *S. apetala* plantations. The plant height, DBH and cover of *S. apetala* all increased with planting age. However, native species such as *K. candel*, *R. stylosa*, *A. marina*, and *A. corniculatum* started to colonise in 4-year *S. apetala* plantations and more species and individuals were observed in 4–5 year plantations. The monoculture native species mangrove plantations grew slowly and no natural regeneration or re-colonisation was observed. Comparatively, the species richness was higher and there were rapid recruitments, with most of the new seedlings being native species in natural forests. The 5-year plantation ranked second in terms of seedling density (both native species and exotic species). The 8-year plantation ranked the third in recruits but the planted species had no new seedlings, and all recruited seedlings belonged to one re-colonised native species. The 10-year plantation had two species (one planted and one native) and 52.4% of the seedlings belonged to native species (Table 3). The

Table 1 Sediment characteristics of different stands at the study sites (mean \pm SD). Values followed by different letters are significantly different ($P < 0.05$; Tukey-test after one way ANOVA)

Stand	Salinity (%)	pH	Temperature (°C)	Organic matter (%)	Total N (%)	Total P (%)	Clay (%)	Water content (g/kg)
Barren sites	1.31 \pm 0.10 ^b	7.05 \pm 0.12 ^a	28.2 \pm 0.1 ^a	1.14 \pm 0.03 ^d	0.053 \pm 0.003 ^a	0.032 \pm 0.003 ^b	10.3 \pm 0.1 ^b	45.2 \pm 0.2 ^a
Native mangrove plantation	2.57 \pm 0.21 ^a	6.96 \pm 0.08 ^a	27.3 \pm 0.2 ^b	1.79 \pm 0.16 ^{cd}	0.104 \pm 0.006 ^a	0.044 \pm 0.001 ^{ab}	20.3 \pm 5.0 ^a	73.5 \pm 1.0 ^a
Natural mangrove forest	2.22 \pm 0.28 ^a	6.53 \pm 0.21 ^a	26.5 \pm 0.1 ^c	4.02 \pm 0.50 ^a	0.152 \pm 0.017 ^a	0.038 \pm 0.001 ^{ab}	26.1 \pm 4.7 ^a	79.1 \pm 5.8 ^{ab}
Exotic species plantation (4-year)	1.22 \pm 0.08 ^b	7.02 \pm 0.13 ^a	27.1 \pm 0.1 ^{bc}	1.24 \pm 0.50 ^d	0.136 \pm 0.008 ^a	0.051 \pm 0.011 ^a	18.8 \pm 1.9 ^{ab}	72.6 \pm 2.3 ^b
Exotic species plantation (5-year)	1.23 \pm 0.20 ^b	6.70 \pm 0.55 ^a	27.2 \pm 0.1 ^{bc}	1.29 \pm 0.20 ^d	0.053 \pm 0.008 ^a	0.026 \pm 0.001 ^b	20.3 \pm 4.1 ^{ab}	73.6 \pm 1.1 ^b
Exotic species plantation (8-year)	1.97 \pm 0.33 ^{ab}	6.83 \pm 0.18 ^a	26.8 \pm 0.3 ^c	2.11 \pm 0.40 ^c	0.106 \pm 0.099 ^a	0.036 \pm 0.006 ^b	23.0 \pm 6.2 ^a	73.8 \pm 2.3 ^a
Exotic species plantation (10-year)	2.04 \pm 0.33 ^a	6.47 \pm 0.07 ^a	26.9 \pm 0.1 ^{bc}	2.45 \pm 0.26 ^b	0.122 \pm 0.009 ^a	0.038 \pm 0.004 ^{ab}	24.2 \pm 2.3 ^a	76.9 \pm 1.6 ^a

regeneration of mangrove was determined by seed source and seedling quantity. Table 3 also indicated that *S. apetala* could not regenerate in its plantation. Some




Fig. 2 Relationship between age of plantation and soil organic matter content

native mangrove species could invade the 4–5-year-old *S. apetala* plantations, but those native species disappeared after 10 years in plantation. In addition, most recolonisation occurred in quadrats far away from sea-water.

Discussion

Long time periods may be required for barren lands to develop vegetation in extremely degraded environments without human assistance. We found no natural re-colonisation in barren sites after 20 years but some species had recruited into several *S. apetala* plantations after 4 years. These results indicate that clearing mangroves may provide an obstacle to natural regeneration, whereas *S. apetala* plantations can greatly modify the

Table 2 Community characteristics of the study sites with different stands. *IV* Important value, *SWI* Shannon–Weiner index

Stand	Characteristic	<i>Avicennia marina</i>	<i>Rhizophora stylosa</i>	<i>Kandelia candel</i>	<i>Aegiceras corniculatum</i>	<i>Sonneratia apetala</i>	Total
Native mangrove plantation	Density (N/ha) ^a		7,333	5,767			13,100
	Basal area (m ² /ha)		0.026	0.239			0.265
	Frequency		0.3	0.7			
	IV		0.33	0.67			1.00
	Mean stand height (m)		1.9	0.9			1.55
	Number of species						2
	SWI						0.47
Natural mangrove forest	Density (N/ha) ^a	13,133	1,233	7,967	2,400		24,733
	Basal area (m ² /ha)	1.103	0.166	0.236	0.091		1.596
	Frequency	1.0	1.0	1.0	1.0		
	IV	0.48	0.13	0.23	0.15		1.00
	Mean stand height (m)	1.21	1.10	1.70	1.60		1.62
	No of species						5
	SWI						1.38
Exotic species plantation (4-year)	Density (N/ha) ^a	367	2,267	233		700	3,567
	Basal area (m ² /ha)	0.064	0.137	0.100		9.888	10.189
	Frequency	0.7	1.0	1.0		1.0	
	IV	0.10	0.31	0.12		0.48	1.00
	Mean stand height (m)	1.40	1.31	1.00		3.5	3.32
	No of species						4
	SWI						1.47
Exotic species plantation (5-year)	Density (N/ha) ^a	6,533	200	1,867	200	1,467	10,267
	Basal area (m ² /ha)	0.906	0.068	0.780	0.003	3.005	4.760
	Frequency	1.0	0.7	1.0	0.3	1.0	
	IV	0.36	0.07	0.20	0.03	0.34	1.00
	Mean stand height (m)	0.9	0.72	0.71	0.50	4.35	4.35
	Number of species						5
	SWI						1.40
Exotic species plantation (8-year)	Density (N/ha) ^a			11,833		1,567	13,400
	Basal area (m ² /ha)			0.504		1.733	2.237
	Frequency			1.0		1.0	
	IV			0.54		0.46	1.00
	Mean stand height (m)			1.00		7.50	7.50
	Number of species						2
	SWI						0.77
Exotic species plantation (10-year)	Density (N/ha) ^a					1,067	1,067
	Basal area (m ² /ha)					7.005	7.005
	Frequency					1.0	1.0
	IV					1.00	1.00
	Mean stand height (m)					12.2	12.2
	Number of species						1
	SWI						0.00
Barren sites: no plants							

^a Not including seedlings

Table 3 Seedling density (individuals/ha) of mangrove trees in quadrats with different stands

Stand		<i>S. apetala</i>	<i>K. candel</i>	<i>R. stylosa</i>	<i>Avicennia marina</i>	<i>Aegiceras corniculatum</i>	<i>Acanthus ilicifolius</i>	Total
Barren sites		0	0	0	0	0	0	0
Native plantation		0	0	0	0	0	0	0
Natural site		0	5,333	667	2,333	930	0	9,263
Exotic	4-year	25	175	1,775	550	0	0	2,525
Species	5-year	200	2,057	87	5,586	86	0	8,016
Plantation	8-year	0	6,950	0	0	0	0	6,950
	10-year	333	0	0	0	0	367	700

site conditions in a way that facilitates the survival and establishment of the propagules of native species. These results are somewhat similar to those of other studies (Field 1999; Ellison 2000; Bosire et al. 2003; Liang et al. 2005). Compared with barren sites, salinity and interstitial water temperature were lower, and the organic matter content was higher, in most restored and natural sites. The possible causes include shading, aerial roots breaking waves, hydrological dynamic changes, and plant nutrition cycle.

Our study suggests that natural regeneration may not occur, and that artificial regeneration (i.e. plantation) of native mangroves may not be effective, in seriously degraded mangrove ecosystems. In severely degraded ecosystems, regeneration of mangrove vegetation might not occur without human intervention because natural regeneration relies on naturally occurring propagules of mangroves. Natural colonisation rates vary among sites and mangrove species, but it is still the first choice for a restoration program. After the physical environment is modified, artificial regeneration involves planting of propagules (seeds) or seedlings in severely degraded region where there are insufficient seeds or natural regeneration pathways.

Our results indicated that the growth performance of the exotic mangrove species, *S. apetala*, was better than that of some native species such as *R. stylosa* and *K. candel* in the first 10 years after plantation. Due to its fast growth and high adaptability, *S. apetala* is now also being introduced to Fujian Province (108°30'E and 24°23'N). Local governments even intended to widely expand distribution of this species in China's southeast coasts. However, restorationists are beginning to worry about its invasive potential in native mangrove stands. Therefore, it is necessary to assess the ecological risk of such species introduction at a broader (regional) scale (Zan et al. 2003). Our study has shown that *S. apetala* plantations facilitated the re-colonisation of native mangrove species in early stages, where it behaves as a nurse plant. However, its high self-regeneration ability in the community prevents the natural succession of the native community. Therefore, we might use *S. apetala* as a pioneer species to improve habitats and to facilitate re-colonisation of native mangrove species. As soon as the deposition conditions are restored, ornative species are suppressed by the exotic species, we should

then artificially remove *S. apetala* so that native species can dominate the community. Thus, the restoration could then follow ideal natural self-organisation processes.

It is clear that monitoring of the plantations for only 10 years is still not long enough to examine whether re-colonised native species can eventually take over or out-compete the planted exotic species without human intervention, since the deposition conditions have not yet been fully restored. We suggest removing planted exotic mangrove species about 5 years later to ensure the performance of native species, but the optimal time for this artificial removal needs further investigation. Further monitoring and comparative studies of different remediation treatments would be useful. It is also clear that, to make sustainable use of mangrove in South China, it is necessary to protect all the existing mangrove wetlands in situ by setting up natural reserves and excluding further anthropogenic destruction, and developing a solid scientific program for restoration. Furthermore, carefully designed broad-scale monitoring and enhanced education programs are critically required to effectively protect the limited resources of mangrove ecosystems.

Acknowledgements We thank Drs. Rui-Jiang Wang and Hong-Fu Wan of the South China Botanical Garden and all staff of the Administration of Zhanjiang Mangrove National Natural Reserve for field assistance during the study. We also thank anonymous reviewers for their valuable comments on the early version of the manuscript. This project was supported by the Knowledge Innovation Program of the Chinese Academy of Sciences (KSCX2-SW-132), Natural Science Foundation of China (30670370, 30600072) and the Guangdong Sci-Tech Planning Project (2005B60301001).

References

- Arnold K (1986) Methods of soil analysis. Part I. Physical and mineralogical methods, 2nd edn. Soil Science Society of America, Madison, WI
- Bosire JO, Dahdouh-Guebas F, Kairo JG, Koedam N (2003) colonisation of non-planted mangrove species into restored mangrove stands in Gazi Bay, Kenya. *Aquat Bot* 76:267–279
- Chen YJ, Liao BW, Zheng SF et al (2004) Dynamics and species-diversities of artificial *Sonneratia apetala*, *Sonneratia caseolaris* and *Kandelia candel* communities. *Chin J Appl Ecol* 15:924–928
- Ellison AM (2000) Mangrove restoration: do we know enough? *Restoration Ecol* 8:219–229
- FAO (1994) Mangrove forest management guideline. FAO, Rome

- Field CD (1999) Remediation of mangrove ecosystems: an overview. *Mar Pollut Bull* 37:383–392
- Han WD, Gao XM, Teunissen E (2001) Study on *Sonneratia apetala* productivity in restored forests in Leizhou Peninsula, China. *J For Res* 12:229–234
- Han WD, Gao XM, Lu CY, Lin P (2003) Mangrove flora and associations of Leizhou Peninsula. *Guihaia* 23:127–132
- Zan QJ, Wang BS, Wang YJ, Ming-Guang L (2003) Ecological assessment on the introduced *Sonneratia caseolaris* and *S. apetala* at the Mangrove Forest of Shenzhen Bay, China. *Acta Bot Sin* 45:604–612
- Kathiresan K, Bingham BL (2001) Biology of mangroves and mangrove ecosystems. *Adv Mar Biol* 40:81–251
- Kovacs JM (1999) Assessing mangrove use at the local scale. *Landsc Urban Plan* 43:201–208
- Krebs CJ (1985) *Ecology: the experimental analysis of distribution and abundance*, 2nd edn. Harper & Row, New York
- Lewis RR III (2005) Ecological engineering for successful management and restoration of mangrove forests. *Ecol Eng* 24:403–418
- Li ZQ (2003) Orientation and management strategy of mangrove forest in the forestry development of Guangdong. *Prot For Sci Technol* 56:36–38
- Li Y, Zhen DZ, Chen HX et al (1998) Preliminary study on introduction of mangrove *Sonneratia apetala* Buch-Ham. *For Res* 11:39–44
- Liang SC, Liang MZ, Wu YL, Zan QJ, Wang YJ, Xie Q (2005) Analysis of the spatial structure of natural *Sonneratia caseolaris* + *S. apetala* forest in Futian, Shenzhen. *Guihaia* 25:393–398
- Liao BW, Zheng SF, Chen YJ et al (2004) Biological characteristics and ecological adaptability for non-indigenous mangrove species *Sonneratia apetala*. *Chin J Ecol* 23:10–15
- Liao BW, Li M, Zheng ZF et al (2005) Niches of several mangrove species in Dongzhai harbor of Hainan island. *Chin J Appl Ecol* 16:430–407
- Lin ZL (1999) Current situation and conservation of mangrove forest in Guangdong. *Ocean Inf* 6:25–27
- Lin ZD, Liu HM (2003) Protection and management countermeasures for mangrove resource in Guangdong province. *Central South For Invent Plan* 22:35–38
- Lin KY, Zhang QM, Ren H (2006) Mangrove resource and sustainable development at Zhanjiang. *Ecol Sci* 23:23–29
- Lugo AE (1999) Mangrove forests: a tough system to invade but an easy one to rehabilitate. *Mar Pollut Bull* 37:427–430
- Macintosh DJ, Ashton EC, Havanon S (2002) Mangrove remediation and intertidal biodiversity: a study in the Ranong mangrove ecosystem, Thailand. *Estuar Coast Shelf Sci* 55:331–345
- Robert L, Morlang A, Gorman L (1997) Monitoring the coastal environment. Part II. Sediment sampling and geotechnical methods. *J Coast Res* 13:308–330
- Snedaker SC (ed) (1984) *The mangrove ecosystem: research methods*. UNESCO, Paris
- Wang BS, Liao BW, Wang YJ et al (2002) *Mangrove forest ecosystem and its sustainable development in Shenzhen Bay*. Science, Beijing
- Wartel S, Barousseau J, Cornand L (1995) Improvement of grain size analyzes using the automated SEDIGRAPH 5100. Document 80. KBIN, Brussels, Belgium
- Wu ZH, Cai JX, Ye XB (2000) Analysis on effects of introduction and popularization of *Sonneratia apetala*. *Guangdong For Sci Technol* 16:6–10
- Zan QJ, Wang YJ, Liao BW et al (2001a) The structure of *Sonneratia apetala* + *S. caseolaris*-*Kandelia candel* mangrove plantations of Futian, Shenzhen. *For Res* 14:610–615
- Zan QJ, Wang YJ, Wang BS (2001b) The ecological effects of *Sonneratia apetala* + *S. caseolaris* mangrove plantations of Futian, Shenzhen. *Acta Sci Natl Univ Sunyatseni* 40:72–76
- Zedler JB (ed) (2000) *Handbook for restoring tidal wetlands*. CRC, London