

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications in the Biological Sciences

Papers in the Biological Sciences

6-14-1892

A Second Edition of Webber's "Appendix to the Catalogue of the Flora Of Nebraska." With a Supplementary List of Recently Reported Species

Herbert J. Webber

University of Nebraska - Lincoln

Charles E. Bessey

University of Nebraska - Lincoln

Follow this and additional works at: <https://digitalcommons.unl.edu/bioscifacpub>



Part of the Life Sciences Commons

Webber, Herbert J. and Bessey, Charles E., "A Second Edition of Webber's "Appendix to the Catalogue of the Flora Of Nebraska." With a Supplementary List of Recently Reported Species" (1892). *Faculty Publications in the Biological Sciences*. 33.

<https://digitalcommons.unl.edu/bioscifacpub/33>

This Article is brought to you for free and open access by the Papers in the Biological Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications in the Biological Sciences by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

CONTRIBUTIONS FROM THE BOTANICAL DEPARTMENT OF THE
UNIVERSITY OF NEBRASKA.

NEW SERIES, III.

A SECOND EDITION OF
WEBBER'S "APPENDIX TO THE CATALOGUE
OF THE
FLORA OF NEBRASKA."

WITH A SUPPLEMENTARY LIST OF RECENTLY REPORTED
SPECIES,

BY CHARLES E. BESSEY, PH. D.

[ISSUED JUNE 14, 1892.]

LINCOLN, NEBRASKA.
U. S. A.

PREFATORY NOTE.

The importance to Nebraska botanical students of the paper entitled an "Appendix to the Catalogue of the Flora of Nebraska," prepared by Mr. Herbert J. Webber, M. A., formerly assistant in botany in the University of Nebraska, now assistant in the Shaw School of Botany, St. Louis, warrants its republication by the University. It was published originally in the Transactions of the Academy of Science of St. Louis (Vol. VI., No. 1, March 12, 1892.) The present edition differs from the first only in the correction of a few minor errors, and the rearrangement of the index.

In his preface to the first edition Mr. Webber explains that "the names of those reporting species follow in every case the name of the species reported as it occurs in the Appendix. The collector whose name follows the species, it must be understood, is to be taken as authority for its occurrence, as in many cases I have not had the opportunity to examine the specimens. Thus specimens here reported must be understood to be preserved in the herbaria of those reporting the species, or in the Herbarium of the University of Nebraska."

The "Supplementary List" (p. 45 et seq.) includes all the species of plants added since Mr. Webber's paper went to press. It consists mainly of the additions made by Mr. P. A. Rydberg, a graduate student, in his collecting tour through the western counties in the summer of 1891, with others made by members of the University Botanical department, and Dr. H. Hapeman of Minden, and Rev. J. M. Bates of Valentine.

Specimens of nearly all the species reported in both lists, are deposited in the University Herbarium.

It is hoped that the periodical publication of accessions to the flora of the state will stimulate botanical students to a closer study of our native plants, and encourage them to deposit specimens of newly discovered species in the University Herbarium. To this end correspondence is solicited by the Department of Botany.

Address, DEPARTMENT OF BOTANY,

UNIVERSITY OF NEBRASKA,

MAY 25, 1892.

LINCOLN, NEBR.

APPENDIX TO THE
CATALOGUE OF THE FLORA OF NEBRASKA.

BY H. J. WEBBER.

CORRECTIONS.

(The reference given, is in every case to the "reprints" of the Catalogue.

g. = genus; s. = species; l. = line.)

- Page 37, l. 36, for *faetidissima* read *foetidissima*.
" 38, l. 28, for *when* read *where*.
" 39, l. 13, for *at* read *as*.
" 41, l. 22, *Barns* read *Barnes*.
" 41, l. 29, omit *other*.
" 41, l. 41, omit *many*.
" 42, s. 5, for *flavo-fuscom* read *flavo-fuscum*.
" 44, s. 29, for *marino* read *marina*.
" 44, l. 6, for *diseasees* read *diseases*.
" 51, s. 165, l. 9, *Aster ericoides* should be *Aster paniculatus*.
" 54, s. 204, " " " " "
" 62, s. 364, l. 1, for *salicini* read *salicina*.
" 67, s. 405, l. 32, for *cynindrical* read *cylindrical*.
" 69, s. 418, The host *Aster ericoides* should be *A. paniculatus*.
" 71, s. 437, l. 2, for *canadense* read *canadensis*.
" 71, s. 450, for *Enothera* read *Oenothera*.
" 74, s. 477, for *Dittm.* read *Ditm.*
" 76, The remark under sterile forms should be — Supposed
to be stages of *Ascomycetes* and *Basidiomycetes*.
" 82, g. 205, for *Pericularia* read *Piricularia*.
" 83, s. 640, for *Sporodesmium* read *Sporilesmium*.
(As given by De Thuemen. Saccardo evidently unin-
tentionally changes it to *Sporodesmium*.)
Also line 5, *scirpiola* read *scirpicola*.
" 87, Order 56, for *Lycoperdiaceæ* read *Lycoperdaceæ*.
" 99, s. 952, for *aristulatus* read *aristatus*.
" 99, s. 956, for *schweinitzii* read *schizoinitziæ*.
" 122, s. 1342, for *lanceolatus* read *lanceolata*.
" 124, s. 1383, for *stolonifer* read *stolonifera*.
" 126, s. 1404, for *sinnate* read *sinnate*.
" 142, s. 1705, for *rush* read *bush*.
" 149, s. 1853, for *serotina* read *serotina*.
" 149, s. 1867, for *eupatorioides* read *eupatorioides*.

NOTES ON SPECIES BEFORE REPORTED.

(The number preceding each, is the species number in the main Catalogue.)

151. *CYSTOPUS TRAGOPOGONIS* (Pers.) Schroet. On Wild worm-wood (*Artemesia canadensis*), Pine Ridge, Aug., 6 (Webber).
414. *GYMNOSPORANGIUM MACROPUS* L. Stage II (*Ræstelia pirata Thaxter*). On leaves of Crab Apple (*Pirus coronaria*) Butler Co., and Lincoln (Bessey).
416. *PHRAGMIDIUM SPECIOSUM* Fr. The specimens referred to Stage II of this species should be Stage I of *Phragmidium subcorticium* (Schrank) Wint. (See No. 81 of this paper).
520. *ASCOCHYTA SMILACIS* Ell & Ev. should be of Ell. & Mart. or following Saccardo the nomenclature should be changed to *Stegonosporæ smilacis* (Ell. & Mart.) Sacc.
622. *RAMULARIA VIRGAUREÆ* Thuem. On *Solidago canadensis* only, not on *S. rigida* and *S. memorialis*.
709. *PODOSPORIUM RIGIDUM* Schw. On grape stems only, not on *Physalis*.
886. *WOODSIA OREGANA* Eaton (Localities omitted in Cat.) Collected at Pine Ridge; Hat Creek Basin; Dismal River, Thomas Co. (Webber).
924. *CAREX GRISEA* Wahl. (Reported from Coulter's Manual). Ashland, May. (Williams).
926. *CAREX JAMESII* Torr. var. *NEBRASKENSIS* (Dew.) Bailey (Reported from Coulter's Manual). Hat Creek Basin, Sioux Co., Aug.; Broken Bow, July; Anselmo, July. (Webber).
929. *CAREX LONGIROSTRIS* Torr. (Reported from Gray's Manual). Dismal River, Thomas Co., July 12. (Webber). Ashland, Weeping Water, War Bonnet. (Williams).
1088. *JUNCUS FILIFORMIS* L. (Reported from Gray's Manual). Lawrence Fork Bottoms, Banner County (Rydberg).
1103. *NOTHOSCORDUM STRIATUM* (Jacq.) Kunth. (Reported from Coulter's Manual). Crete, rare (Swezey).
1126. *QUERCUS NIGRA* L. (Reported from Gray's Manual). Pawnee Co. (Bessey).
1187. *ERIOGONUM ALATUM* Michx. (Reported from Coulter's Manual). Hills. Deuel Co. and Banner Co. (Rydberg).
1197. *ATRIPLEX NUTTALLII* Watson, is the prevailing *Atriplex* of N. W. Nebraska in the "Bad lands," etc., but is quite rare in the eastern part of the state where *A. patula* var *hastata* is the common form. (Webber).
1202. *AMARANTUS CHLOROSTACHYS* Willd. of Swezey's Nebr. Fl. Plants p. 13 is likely *Acnida tuberculata* Moq. (Webber).

1213. *ARENARIA PUNGENS* Nutt. must probably be considered as *A. hookeri* Nutt. See this appendix No. 269. (Webber.)
1257. *NELUMBO LUTEA* (Willd.) Pers. (Reported from Gray's Manual). Lakes, etc., plentiful. Fremont, July. (Williams & Bessey).
1302. *ELATINE TRIANDRA* Schkuhr (Reported from Coulter's Manual). Exeter, Sept. communicated by Dr. Wibbe (Bessey).
1398. *MENTZELIA LEVICAULIS* Torr. & Gr. should be changed to *Mentzelia nuda* (Pursh.) Torr. & Gr. It is the common *Mentzelia* of Nebraska. Mr. Rydberg adds the following note in regard to his specimens collected in Deuel and Banner counties: — "They agree in every respect with Torrey & Gray's description of *M. nuda* except that they have a bracteate calyx. In Porter and Coulter's Fl. Col. *M. nuda* is described as having a bracteate calyx." (Webber).
1491. *ASTRAGALUS FLEXUOSUS* Dougl. (Reported from Coulter's Manual). Lewellen, Common (Swezey).
1497. *ASTRAGALUS MICROLOBUS* Gray (Reported from Coulter's Manual). Cultivated ground. Deuel Co., June 25; Cheyenne Co., Aug. 13 (Rydberg).
1498. *ASTRAGALUS MISSOURIENSIS* Nutt. (Reported from Coulter's Manual). Curtis, Frontier Co., June 22; North of Kimball, Aug. 16 (Rydberg). War Bonnet Canon, June (Williams).
1501. *ASTRAGALUS PECTINATUS* Dougl. (Reported from Coulter's Manual). Prairies, near Gering July 20 and Pleasant Valley, Scotts Bluff Co., July 28 (Rydberg).
1504. *ASTRAGALUS RACEMOSUS* Pursh. (Reported from Coulter's Manual). Hills, Curtis, Frontier Co., June 22 (Rydberg).
1559. *GILIA IBERIDIFOLIA* Benth. (Reported from Coulter's Manual). Cliffs and Canons of Banner and Scott's Bluff Counties, July (Rydberg).
- MERTENSIA LANCEOLATA* (Pursh) DC. Swezey's Nebr. Fl. Plants, p. 11, should be *Pentstemon cæruleus* number 1612 of the catalogue.
1580. *ECHINOSPERMUM REDOWSKII* (Hornem.) Lehm. is the var. *occidentale* Watson. Since collected at Dismal River, Thomas Co., July 13; Pine Ridge, July 21 (Webber).
1581. *ECHINOSPERMUM REDOWSKII* (Hornem.) Lehm. var. *CUPULATUM* Gray (Reported from Coulter's Manual). Alliance, Aug. 6 (Webber). Chadron (Bates). Venango, Perkins Co., June 23 (Rydberg).
1589. *PHYSALIS LANCEOLATA* Michx. var. *LEVIGATA* Gray. (Reported from Coulter's Manual). Old fields, etc. Weeping Water, July (Williams).
- PENTSTEMON GLABER* Pursh var. *UTAHENSIS* Watson. Swezey's Nebr. Flowering Plants p. 12, should be *P. haydeni* Watson (see this appendix No. 364) (Webber).
611. *PENTSTEMON ALBIDUS* Nutt. (Reported from Gray's Manual). Antelope Co., Aug.; Anselmo, July; Thedford, July; Hat Creek Basin, Aug. (Webber).

1666. **GENTIANA CALYCOSA** Griseb. All of my Nebraska material of this species should be labeled *G. puberula* Michx. (Webber).
1738. **CNICUS PITCHERI** Torr. should be *C. undulatus* (Nutt.) Gray. var. *canescens* (Nutt.) Gray. (See this appendix No. 391.) (Webber).
1792. **HELIANTHUS PETIOLARIS** Nutt. var. *CANESCENS* Gray. (Reported from Gray's Syn. Flora). Lawrence Fork Bottoms, Banner county (Rydberg).
1804. **FRANSERIA DISCOLOR** Nutt. (Reported from Coulter's Manual). Prairies, Kearney County; Dix, Kimball Co., Aug. 14 (Rydberg).
- ERIGERON GLABELLUS Nutt. of Swezey's Nebr. Fl. Pl. p. 9 should be *E. macranthus* Nutt. (See this appendix No. 411)
1836. **ASTER TANACETIFOLIUS** (Nees) H. B. K. (Reported from Coulter's Manual). Banner Co., etc. Common in the western part of the State (Rydberg).
1867. **KUHNIA EUPATORIOIDES** L. var. *CORYMBULOSA* Torr. and Gr. (Reported from Coulter's Manual). Canons, Banner Co. (Rydberg).

Catalogue of the Flora of Nebraska.

7

The following tabulated list of new localities for Flowering Plants before reported was prepared by Mr. Rydberg from his collections. Some species are greatly extended in range, and some shown to be much more common than was supposed.

PROTOPHYTA.

1. RETICULARIACEÆ.

1. RETICULARIA Bull.

1. R. LYCOPERDON Bull. On decaying trunks of trees. Pine Ridge, Dawes Co., July (Webber).

2. CLATHROPTYCHIACEÆ.

2. ENTERIDIUM Ehr.

2. E. ROZEANUM (Rost.) Wingate. (*Reticularia* [?] *rozeana* Rost.). Lincoln (Webber).

3. STEMONITACEÆ.

3. STEMONITIS Gled.

3. S. WEBBERI Rex. (N. sp. in lit. Published in Pro. Acad. Nat. Sci. of Phila., 1891). On old stump, Lincoln, September (Webber).

4. PHYSARACEÆ.

4. PHYSARUM Pers.

4. P. LIVIDUM Rost. (*Spumaria licheniformis* Schw.) On bark of tree, Lincoln (Webber).
5. P. PETERSII B. and C. On bark of old Cottonwood and Boxelder trees, Weeping Water (Williams).

5. BACTERIACEÆ.

5. BACILLUS Cohn.

6. B. SORGHI Burrill. On leaves and culms of Bushy blue stem (*Andropogon nutans*), Lincoln; Johnson grass (*Andropogon sorghum* var. *halepensis*), Howard Co.; and on numerous varieties of *Sorghum* grown on the Nebraska Experimental Farm at Lincoln. Quite destructive, forming large irregular purple patches (Webber).

6. NOSTOCACEÆ.

6. GLÆOTRICHIA Ag.

7. G. NATANS Taur. Floating in stagnant water. Greenwood, July (Williams). Minden, attached to *Nitella* stems (Bessey).

7. LYNGBYA Ag. and Thur.

8. L. CINCINNATA Kg. Minden. In material collected by Dr. Hapeman (Bessey).

8. CYLINDROSPERMUM Kg.

9. C. COMATUM Wood. Minden (Hapeman).

ZYGOPHYTA.

7. PALMELLACEÆ.

9. TETRASPORA Ag.

10. *T. EXPLANATA* (Kg.) Kirch. Minden. In material collected by Dr. Hapeman (Bessey).

10. SORASTRUM Kg.

11. *S. SPINULOSUM* Kg. Minden. In material collected by Dr. Hapeman (Woods).

11. HYDRODICTYON Roth.

12. *H. UTRICULATUM* Roth. Stagnant water, plentiful. Fremont, July 31 (Williams). Lincoln (Bessey).

12. PEDIASTRUM Meyen.

13. *P. ANGULOSUM* (Ehrb.) Menegh. (*P. biradiatum* Meyen). In material collected by Dr. Hapeman at Minden. (Woods).

14. *P. PERTUSUM* Kg. var. *CLATHRATUM* A. Br. (*P. duplex* Meyen). Minden. In material collected by Dr. Hapeman (Woods).

8. PITHOPHORACEÆ.

13. PITHOPHORA Wittr.

15. *P. AFFINIS* Nordst. In stagnant or slow running water. Greenwood, July (Williams).

9. ULOTHRICHACEÆ.

14. STIGEOCLONIUM Kg.

16. *S. FASTIGIATUM* Kg. Minden. In material collected by Dr. Hapeman (Bessey).

15. APHANOCHÆTE A. Br.

17. *A. GLOBOSA* (Nord.) Wolle. form. *MINOR* Nordst. growing on *Nitella*, in material from Minden collected by Dr. Hapeman. (Woods).

10. DESMIDIACEÆ.

16. DESMIDIUM Ag.

18. *D. SWARTZII* Ag. Minden. In material collected by Dr. Hapeman (Bessey).

17. SPHÆROZOSMA Corda.

19. *P. SERRATUM* Bailey. Minden. In material collected by Dr. Hapeman (Bessey).

18. SPIROTÆNIA Breb.

20. *S. CONDENSATA* Breb. In material from Minden, collected by Dr. Hapeman.

19. COSMARIUM Corda.
21. C. BROOMEI Thwaites. In material from Minden, collected by Dr. Hapeman Oct. 19 (Woods).
20. XANTHIDIUM Ehrb.
22. X. FASCICULATUM (Ehrb.) Ralfs. In material from Minden, collected by Dr. Hapeman (Woods).
21. ARTHRODESMUS Ehrb.
23. A. OCTOCORNIS Ehrb. In material from Minden, collected by Dr. Hapeman (Woods).
22. EUASTRUM Ehrb.
24. E. INERME Lund. In material from Minden, collected by Dr. Hapeman (Woods).
23. MICRASTERIAS Ag.
25. M. AMERICANA (Ehrb.) Kg. Minden, in material collected by Dr. Hapeman (Woods).
26. M. SPECIOSA Wolle. Minden. In material collected by Dr. Hapeman (Woods).
24. STAURASTRUM Meyen.
27. S. ARISTIFERUM Ralfs. Minden. In material collected by Dr. Hapeman (Woods).

11. DIATOMACEÆ.

25. GOMPHONEMA.
28. G. ACUMINATUM Ehrb. var. LATICEPS Ehrb. Minden. In material collected by Dr. Hapeman. (Bessey).

12. ENTOMOPHTHORACEÆ.

26. EMPUSA Cohn.
29. E. APHIDIS Hoffman. On *Aphis* Sp. on *Polygonum*. Ashland. Oct. (Williams).
-

OOPHYTA.

13. OEDOGONIACEÆ.

27. OEDOGONIUM Lk.
30. OE. BORISIANUM (Le Cl.) Wittr. In material from Minden collected by Dr. Hapeman (Woods).
31. OE. DELICATUM Kg. Minden. In material collected by Dr. Hapeman (Bessey).
28. BULBOCHÆTE Ag.
32. B. POLYANDRA. Cleve. Minden. In material collected by Dr. Hapeman (Bessey).

14. PERONOSPORACEÆ.

29. PERONOSPORA Corda.

33. P. OXYBAPHII Ell. & Kell On leaves and young shoots of *Oxybaphus nyctagineus*, causing much damage to the host. Elmwood, Ashland. June and July (Williams).

30. PLASMOPARA Schroet.

34. P. HALSTEDII (Farlow) Berl. and De Toni. On leaves of Sunflower (*Helianthus annuus*), Lincoln, Oct. (Webber). Great ragweed (*Ambrosia trifida*), Wabash, Aug. (Williams).

31. SCLEROSPORA Schroet.

35. S. GRAMINICOLA (Sacc.) Schroet. (*Peronospora graminicola* Sacc.) On Green and Yellow fox-tail (*Setaria viridis* and *S. glauca*), Ashland, Weeping Water (Williams). Lincoln (Bessey).

32. CYSTOPUS Lev.

36. C. IPOMOEÆ-PANDURANE (Sacc.) Farlow. On leaves and petioles of Morning Glory (*Ipomoea* Sp.) Richardson Co., Aug. (Webber); Ashland (Williams). Very destructive, variously twisting and distorting the leaves and petioles, finally causing them to drop off.
- — —

CARPOPHYTA.

15. COLEOCHAETACEÆ.

33. COLEOCHAETE Breb.

37. C. IRREGULARIS Pringsh. Minden. In material collected by Dr. Hapeman (Bessey).
38. C. ORBICULARIS Pringsh. Minden. In material collected by Dr. Hapeman (Bessey).

16. ERYSIPHEÆ.

34. ERYSIPHE Hedw.

39. E. GRAMINIS DC. Couidia stage (*Oidium monilioides* Link.) On leaves of Wild Rye (*Elymus canadensis*), Weeping Water (Williams).

The Oidium is the only stage frequently found here. I have frequently observed this at Lincoln on various grasses, but have never found the perithecia (Webber).

17. SPHÆRIACEÆ.

35. CHÆTOMIUM Kunze.

40. C. CHARTARUM Ehrenb. On decaying broom. Lincoln (Pound).

36. PHYSALOSPORA Niessl.

41. P. MEGASTOMA (Peck) Sacc. On leaves of *Astragalus drummondii*. Belmont, Dawes Co., July. Quite destructive (Webber).

37. SPHÆRELLA Ces. & DeNot.

42. S. OPUNTIAE Ell. & Ev. On Cactus (*Opuntia missouriensis*), Weeping Water (Williams).

18. HYPOCREACEÆ.

38. NECTRIELLA Sacc.

43. N. VULPINA (Cke.) Berl. & Vogl. On old decaying log, Lincoln (Webber).

19. DOTIDIACEÆ.

39. PHYLLACHORA Nitschke.

44. P. LESPEDEZÆ (Schw.) Sacc. On Bush clover (*Lespedeza frutescens*), Nemaha Co., Oct. (Webber).

40. PLOWRIGHTIA Sacc.

45. P. RIBESIA (Pers.) Sacc. On stems of gooseberry (*Ribes gracile*), Ashland (Williams).

20. HYSTERIACEÆ.

41. HYSTEROGRAPHIUM Corda.

46. H. FRAXINI (Pers.) DeNot On Ash (*Fraxinus viridis*) Lincoln (Pound).

21. CALICIACEÆ.

42. ACOLIUM Ach.

47. A. TIGILLARE Ach. On Sandstone, rare. Pine Ridge, Aug. (Webber).

22. GRAPHIDIACEÆ.

48. ARTHONIA (Ach.) Nyl.

49. A. RADIATA (Pers.) Th. Fr. On Hickory bark. Milford, Oct. (Webber).

50. OPEGRAPHA (Humb.) Ach. Nyl.

51. O. VARIA Pers. var. PULICARIS. Bark of trees. Weeping Water (Williams).

23. LECIDEACEÆ.

52. BUELLIA De Not., Tuckerm.

53. B. EPIGÆA (Pers.) Tuckerm. On high sandy ground. Pine Ridge, Aug. (Webber).

54. LECIDEA (Ach., Fr.) Tuckerm.

55. L. TESSELLATA Floerk. On stone. Pine Ridge, Aug. Common. (Webber).

56. BIATORA, Fr.

57. B. MUSCORUM (Sw.) Tuckerm. On moss. Pine Ridge, Aug. (Webber).

53. *B. RUSSULA* (Ach.) Mont. f. *DEALBATA* Tuckerm. On earth etc. Valentine, Harrison (Williams).
48. *CLADONIA* Hoffm.
54. *C. BOTRYTIS* (Hag.) Hoffm. Rotten Pine log. War Bonnet canon, N. W. Nebraska. (Williams).
55. *C. CARIOSA* (Ach.) Spreng. On ground under trees on bluffs and in damp places, very common. Pine Ridge, Aug.; Dismal River, Thomas Co., July (Webber); War Bonnet canon (Williams).

24. PARMELIACEÆ.

49. *URCEOLARIA* Ach.
56. *U. SCRUIPOSA* (L.) Nyl. On sandy ground and sandstone, common. Pine Ridge, July (Webber).
50. *LECANORA* Ach. Tuckerm.
57. *L. BRUNONIS* Tuckerm. On sandstone, Ashland. (Williams).
58. *L. GLAUROCARPA* (Wahl.) Ach. On stone, common. Pine Ridge, Aug. (Webber).
59. *L. PRIVIGNA* (Ach.) Nyl. On stone. Pine Ridge, Aug. (Webber).
60. *L. SCHLEICHERI* (Ach.) Nyl. On ground, abundant. Pine Ridge, Aug. (Webber).
61. *L. VARIA* (Ehrh.) Nyl. var. *SEPINCOLA* Fr. On bark of pine, common. Pine Ridge, Aug. (Webber). On trees, Ashland. (Williams).
51. *PLACODIUM* (DC.) Naeg. & Hepp.
62. *P. ELEGANS* (Link.) D C. On stone. Pine Ridge, Aug. A very pretty orange red species, abundant in this region. (Webber).
63. *P. MICROPHYLLINUM* Tuckerm. On old bark, Pine Ridge, Aug. (Webber).
64. *P. VITELLINUM* (Ehrh.) Naeg. & Hepp. On sandstone. Lincoln, Sept. (Webber).
52. *COLLEMA* Hoffm., Fr.
65. *C. PULPOSUM* (Beruh.) Nyl. On sandstone. Pine Ridge, Aug. (Webber).
53. *PELTIGERA* (Willd., Hoffm.) Fée.
66. *P. HORIZONTALIS* (L.) Hoffm. On ground. Pine Ridge, Aug. (Webber.)
54. *PHYSCKIA* (DC., Fr.) Th. Fr.
67. *P. HISPIDA* (Schreb., Fr.) Tuckerm. On stone. Pine Ridge, Aug. (Webber).
55. *PARMELIA* (Ach.) De Not.
68. *P. MOLLUSCULA* Ach. On dry sterile soil, abundant. War Bonnet Canon, Harrison (Williams), Pine Ridge, Crawford (Webber).
69. *P. TILIACEA* (Hoffm.) Floerk. var. *SUBLEVIGATA* Nyl. On trees. Peru, Weeping Water (Williams).

25. PEZIZACEÆ.

56. PEZIZA Fuckel.

70. P. HEMISPHERICA Wigg. Manure etc., Wabash (Williams).

26. UREDINEÆ

57. UROMYCES Link.

71. U. HOWEI Peck. On Milkweed (*Asclepias syriaca*), Fremont, July 31. Very destructive (Williams).72. U. TRIFOLII (Alb. & Schw.) Winter. On red clover (*Trifolium pratense*), Ashland. Quite plentiful and somewhat destructive (Williams).

58. MELAMPSORA Cast.

73. M. LINI (DC.) Tul. On Wild-flax (*Linum perenne* var *lewisii*), Hat Creek Basin; Wild-flax (*Linum rigidum*), Weeping Water. Very destructive especially on the latter host. (Williams).

59. PUCCINIA Pers.

74. P. ANEMONE-VIRGINIANÆ Schw. On Long fruited anemone (*Anemone cylindrica*), Weeping Water. (Williams).

75. P. FUSCA Relhan. On Anemone, Ashland (Williams).

76. P. HYDROPHYLLI Peck & Cke. On leaves of Water leaf (*Hydrophyllum virginicum*) Sarpy Co. (Pound.)77. P. MIRABILISSIMA Peck. II & III. On Barberry (*Berberis repens*), Belmont, July 24. The teleutospores are rarely found. In my Nebraska material I found but two. They, however, agree and compare well. (Webber).78. P. SANICULÆ Grev. On Black snake root (*Sanicula canadensis*), Ashland (Williams).79. P. SCIRPI DC. III On Bull-rush (*Scirpus* sp.), Lincoln, April (Webber).80. P. SMILACIS Schw. On Greenbrier (*Smilax hispida*), Ashland. Abundant, causing the leaves to turn yellow and fall off (Williams).

60. PHRAGMIDIUM Link.

81. P. SUBCORTICUM (Schrank.) Wint. Stage I on Rose (*Rosa arkansana*), Milford, May, 86; (Webber). Cultivated Rose (*Rosa* sp.), Lincoln, May (Webber). Weeping Water and Ashland (Williams).

This includes the specimens referred to stage II of *P. speciosum* (No. 416) in the catalogue, but does not include those of stage III.

On leaves petioles and stems, orange red, very conspicuous. Frequently quite injurious (Webber).

61. ECIDIUM Pers.

82. A. CALLIRRHOES Ell. & Kell. On *Malva* sp. Ashland (Williams).

83. A. MICROPUNCTUM E. & E. On Painted cup (*Castilleja sessiliflora* Pursh), Belmont, June 17. Quite destructive. (Williams.)

62. UREDO Pers.

84. U. RIBICOLA C. & E. On leaves of Buffalo or Missouri currant (*Ribes aureum*), Collected by Dr. Thomas in Scott's Bluff County.

Although determined to be a *Uredo* by Messrs. Cooke and Ellis (*Grevillea VI*, p. 86) and also by Peck who named it *Uredo jonesii* (*Torr. Bull. XII*, p. 36) it seems more like a *Coleosporium*, which it will probably turn out to be. All attempts at germinating the spores have thus far failed (Bessey).

27. USTILAGINEÆ.

63. USTILAGO Pers.

85. U. CARICIS (Pers.) Fekl. On Sedge (*Carex filifolia*) Sioux Co. June. Very plentiful, destroying the ovaries of nearly all the host plants in that vicinity. (Williams).

86. U. RABENHORSTIANA Kuhn. On finger grass (*Panicum sanguinale*) Ashland.

On flower spikes in the sheath, completely destroying them (spores brown, echinulate, round ellipsoid or angular, 7-12 by 8-13 m. m. m.). (Williams).

64. UROCYSTIS Rabenh.

87. U. OCCULTA (Wallr.) Rabenh. On Wild rye (*Elymus canadensis*), Ashland. (Williams).

28. EXOASCÆ.

65. EXOASCUS Fuckel.

88. E. PRUNI Fekl. On common wild plum causing the disease known as "plum pockets." Weeping Water, quite common. (Williams), Dawes Co. (Bessey). On *Prunus pumila* at Long Pine (Bessey).

29. SACCHAROMYCETES.

66. SACCHAROMYCES Meyen.

89. S. MYCODERMA Reess. Lincoln (Pound).

30. SPILÆRIOIDÆ.

67. PHYLOSTICTA Pers.

90. P. CORNI West. On Red osier dogwood (*Cornus stolonifera*), Belmont, July (Webber.)

91. P. CRUENTA (Fr.) Kx. On False Solomon's seal (*Smilacina stellata*), Ashland (Williams); False Solomon's seal (*S. amplexicaulis*), New Helena, Custer Co., July 6. Frequently quite destructive. (Webber.)

92. P. PERSICÆ Sacc. On leaves of peach. Rock Creek and Ashland (Williams).
93. P. PIRINA Sacc. On apple leaves. Abundant and frequently destructive, Nov. 20. Lincoln. (Webber).
94. P. SEROTINA Cooke. On cherry (*Prunus serotina?*), Richardson Co., Aug. 25. Quite destructive in the locality where the specimens were collected. (Webber).
95. P. ULMICOLA Sacc. On Elm (*Ulmus americana*), Ashland, Oct. (Williams).
68. VERMICULARIA Fr.
96. V. LILIACEARUM Schw. On wild garlic (*Allium canadense*), Ashland (Williams).
69. SEPTORIA Fr.
97. S. BRUNELLÆ Ell. & Hals. On Self-heal (*Brunella vulgaris*), Richardson Co., Aug. 26 (Webber).
98. S. CEPHALANTHI Ell. & Kelt. On leaves of Button bush (*Cephalanthus occidentalis*), West Point (Williams).
99. S. CORYLINA Peck. On Hazel nut (*Corylus americana*), Nebraska City, June. Not uncommon. (Webber).
100. S. LITTOREA Sacc. On Dogbane (*Apocynum cannabinum*), Elmwood, Weeping Water, Wabash (Williams).
101. S. RHOINA B. & C. On Sumach. (*Rhus glabra*), Nebraska City, June 1; Common, (Webber). Ashland and Weeping Water (Williams).
70. RHABDOSPORA Mont.
102. R. CONTINUA (B. & C.) Sacc. On stems of Plantain (*Plantago elongata*), Lincoln, March. Forming on old stems, numerous inconspicuous little black specks. (Webber).

31. LEPTOSTROMACEÆ.

71. LEPTOSTROMA, Fr.

103. L. SCIRPINUM Fr. On dead leaves of Riverrush (*Scirpus fluviatilis*), Weeping Water. (Williams.)

32. MELANCONIEÆ.

72. GLÆOSPORIUM Desm. & Mont.

104. G. SPHÆRELLOIDES Sacc. On *Hoya carnosa* in conservatory, Lincoln (Pound).

73. COLLETOTRICHUM Corda.

105. C. LINEOLA Corda. On leaves and culms of False red-top (*Eragrostis pectinacea*), Roca., Sept 22; Sand-bur (*Cenchrus tribuloides*), Lincoln Oct. 3, (Webber).

74. MELANCONIUM Link.

106. M. MAGNUM (Grev.) Berk. On dead Hickory, Ashland (Williams).

33. MUCEDINEÆ.

75. BOTRYTIS Mich.

107. B. CERATOIDES Peck. On decaying boards, Weeping Water. (Williams).

76. RAMULARIA Ung.

108. R. DESMODII Cooke. On leaves of Tick-trefoil (*Desmodium canescens*), Weeping Water, Wabash, Ashland (Williams); Tick trefoil (*D. canadensis*), Lincoln, Aug. (Webber). In places very destructive.
109. R. IMPATIENTIS Peck. On leaves of Wild-balsam (*Impatiens fulva* and *I. pallida*), Ashland (Williams).
110. R. OCCIDENTALIS Ell. & Kell. On leaves of Dock (*Rumex altissimus*), Lincoln (Pound).
111. R. OXALIDIS Farlow. On leaves of Sorrel (*Oxalis violacea*), Ashland (Williams).

77. STACHYBOTRYS Corda.

112. S. LOBULATA Berk. On decaying broom, Lincoln (Pound).

34. DEMATIEÆ.

78. CERATOPHORUM Sacc.

113. C. ULMICOLUM Ell. & Kell. On leaves of American Elm (*Ulmus americana*), Ashland, October (Williams).

79. HELMINTHOSPORIUM Link.

114. H. LANCEOLATUM Cooke. Lincoln (Williams).

80. CERCOSPORA Fries.

115. C. AERUGINOSA Cooke. On leaves of Buckthorn (*Rhamnus lanceolatus*), Ashland (Williams).

116. C. DATURÆ Peck. On leaves of Stramonium (*Datura stramonium*), Ashland (Williams).

117. C. DUBIA (Riess) Wint. (*C. chenopodii* Fr.)

On Pigweed or goose-foot (*Chenopodium album*), Pine Ridge, July 29 (Webber).

118. C. ECHINOCYSTIS Ell. & Mart. On Wild cucumber (*Echinocystis lobata*), Waverly (Williams).

119. C. FLAGELLARIS Ell. & Mart. On Poke weed (*Phytolacca decandra*), Richardson Co. Aug. 1 (Webber).

120. C. GYMNOCLEADI Ell. & Kell. On leaves of Kentucky Coffee tree (*Gymnocladus canadensis*), very common, Ashland (Williams); Richardson Co., Aug. (Webber).

121. C. LIPPIÆ Ell. & Ev. On leaves of *Lippia lanceolata*, Ashland. Quite destructive (Williams).

122. C. SAGITTARIE Ell. & Kell. On Arrow-head (*Sagittaria variabilis*), Weeping Water. (Williams.)

123. *C. SYMPHORICARPI* Ell. & Ev. On Indian Currant (*Symphoricarpus vulgaris*), Rock Creek, July (Williams.)
124. *C. VERNONIÆ* Ell. & Kell. On Iron weed (*Vernonia fasciculata*), Rock Creek, Wabash, Ashland. (Williams). Very plentiful and quite destructive.

35. STILBEÆ.

81. ISARIA Pers.

125. *I. SULPHUREA* Fiedl. On ground, Lincoln. (Webber).

36. NIDULARIACEÆ.

82. CYATHUS Hall.

126. *C. STRIATUS* (Huds.) Hoffm. On decaying matter. Wabash, Ashland, Weeping Water. (Williams).

37. LYCOPERDACEÆ.

83. GEASTER Mich.

127. *G. HYGROMETRICUS* Pers. On ground, Pine Ridge, Dawes Co., July (Webber).

84. BOVISTA Dill.

128. *B. SUBTERRANEA* Peck. Dismal River, Thomas Co., and Pine Ridge. Very common in Central and Western regions. This species is thought by many mycologists to be identical with *B. circumscissa*, B. & C. (No. 785 of the catalogue). De Toni, (in Sacc. Syl. Fung.) keeps them distinct. Mr. Morgan says he can easily distinguish between the two in the Nebraska material sent him, which, to my certain knowledge, was collected in similar localities. For this reason I insert the species here, although, it is probably nothing more than a form of *B. circumscissa* B. & C. (Webber).

85. LYCOPERDON Tourn.

129. *L. CURTISII* Berk. Lincoln. (Webber).
130. *L. WRIGHTII* B. & C. var. *SEPARANS* Peck. On ground, Dismal River, Thomas Co., July 12; Pine Ridge, July 25 (Webber).

86. ARACHNION Schw.

131. *A. ALBUM* Schw. On ground among weeds. Lincoln, Aug. (Webber).

38. AGARICINEÆ.

87. LEPIOTA Fr.

132. *L. OBLITA* Peck. Woods, Weeping Water, Lincoln. Common, (Williams).

88. ARMILLARIA Fr.

133. *A. MELLEA* Vahl. Base of stumps etc., common, Weeping Water (Williams).

89. TRICHOLOMA Fr.

134. *T. TERREUM* Schaeff. Woods, Weeping Water. (Williams).

90. CLITOCYBE Fr.

135. *C. INFUNDIBULIFORMIS* Schaeff. Ashland, rare (Williams).

91. COLLYBIA Fr.

136. *C. VELUTIPES* Curt. Bases of stumps of Hickory trees. Ashland, Weeping Water, common (Williams).

92. PLEUROTUS Fr.

137. *P. ULMARIUS* Bull. Trunks of Elm, Boxelder, etc. Generally late in autumn, common. Lincoln, Ashland, Weeping Water (Williams).

93. LENTINUS Fr.

138. *L. LECOMTEI* Fr. On old logs, especially Cottonwood. Weeping Water, Ashland, Wabash. Common, (Williams).

94. PLUTEUS Fr.

139. *P. CERVINUS* Schaeff. On much decayed logs, common. Ashland. (Williams).

95. ENTOLOMA Fr.

140. *E. RHODOPOLIUM* Fr. Moist woods. Ashland (Williams).

96. PHOLIOTA Fr.

141. *P. PRECOX* Pers. Grassy places after rains. Lincoln, Weeping Water. (Williams).

97. HYPHOLOMA Fr.

142. *H. VELUTINUM* Pers. Lincoln, Weeping Water (Williams).

39. POLYPOREÆ.

98. POLYPORUS Mich.

143. *P. ELEGANS* (Bull.) Fr. On old wood, Lincoln. A very pretty species with somewhat excentric stipe about $1\frac{3}{4}$ in. long, thick pileus, and decurrent hymenium. Rare. (Webber).

40. HYDNEÆ.

99. HYDNUM L.

144. *H. CIRRATUM* Pers. On old stump, Lincoln, June. Mr. Ellis writes — "Rare, I have it also from Iowa and New York." My only specimens were sent to Mr. Ellis for determination hence are in Herb. Ellis.

Hymenium with very long coarse teeth, blue brown. (Webber).

100. IRPEX Fr.

145. *I. OBLIQUUS* (Schrad.) Fr. On dead limbs of Elm, Ashland (Williams).

41. THELEPHOREÆ.

101. STEREUM Pers.

146. *S. NEGLECTUM* Pk. (Rep. 33 p. 22.) On bark of trees, common. Lincoln (Webber).

42. CLAVARIEÆ.

102. PTERULA Fr.

147. *P. MULTIFIDA* Fr. On decaying leaves at bases of trees in heavy timber, rare. Wabash (Williams).

43. TREMELLINEÆ.

103. GUEPINA Fr.

148. *G. SPATHULARIA* (Schw.) Fr. On old boards, R. R. ties, etc., common. Weeping Water and Ashland (Williams). Lincoln (Webber).

44. CHAREÆ.

104. CHARA (Vaill.) Leonh.

149. *C. CONTRARIA* A. Br. Flowing or standing water and cold springy lakes. Fremont, July (Williams).
150. *C. FRAGILIS*, Desv. In ponds. Greeley Center, Greenwood, Fremont (Williams).

105. NITELLA Ag.

151. *N. ACUMINATA* A. Br. var. *GLOMERATA* A. Br. Ponds near Lincoln (Bessey).
152. *N. FLEXILIS* Ag. Ponds near Minden. In material collected by Dr. Hapeman (Bessey).
153. *N. MUCRONATA* A. Br. Ponds near Minden. In material collected by Dr. Hapeman (Bessey).

BRYOPHYTA.

45. BRYACEÆ.

106. GYMNSTOMUM Hedw.

154. *G. RUPESTRE* Schwægr. (*Mollia aeruginosa* (Sm.) Lindb.). On damp overhanging stone cliff. Pine Ridge, July 25 (Webber).

107. DICRANELLA Schimp.

155. *D. VARIA* Schimp. On wet clayey soil, banks of the Mo. River. Peru, March (Webber).

108. DESMATODON Brid.

156. *D. ARENACEUS* Sulliv. & Lesq. On ground. Collected by Mr. Brunner at Ashland, May (Webber).
157. *D. NERVOSUS*. Bruch. & Schimp. var. *EDENTULUS* Bruch. & Schimp. Abundant on stone. Peru, March (Webber).

109. BARBULA. Hedwig.

158. *B. RURALIS* Hedw. Pine Ridge, Aug. 3 (Webber).

110. BRYUM Dill.

159. B. CESPITICUM L. Earth, Weeping Water (Williams).

111. ATRICHUM Beauv.

160. A. PARALLELUM Mitt. Moist banks. Pine Ridge, June (Williams).

161. A. ANGUSTATUM. Bruch. & Schimp. Long Pine (Bessey).

112. POLYTRICHUM L.

162. P. JUNIPERINUM Willd. On ground. Collected by Mr. Conklin at Long Pine (Webber).

113. HYPNUM Dill.

163. H. FLUVIATILE Swartz. West Nebraska (Webber).

164. H. RADICALE Beauv. On damp decaying log. Hat Creek Basin, Sioux Co. Aug. (Webber).

165. H. RUTABULUM L. var. LONGISETUM Brid. Wet dripping banks and rocks, Ashland, April (Williams).

166. H. SERRULATUM Hedw. On ground. Peru, March; Nebraska City, June (Webber).

PTERIDOPHYTA.

46. MARSILIACEAE.

114. MARSILIA L.

167. M. VESTITA Hook. & Grev. In ponds in many places in the State; Fairmont, Bradshaw, Minden, Geneva. Apparently not occurring east of the Blue River. The Fairmont specimens collected in 1890 agree with the variety *tenuifolia* in the hairiness, and shape of the leaflets, but nearly all the specimens collected this year (1891) fully agree with the specific characters of *M. vestita*. A few specimens were received which were more hairy, and whose leaflets were narrower, but upon investigation these were found to be terrestrial forms. In some cases these *tenuifolia*-like forms were connected directly with the larger smooth and broad leaved aquatic forms. The season of 1890 was a very dry one which probably accounts for the small size of the plants and the hairiness and narrow growth of the leaflets (Bessey).

47. FILICES.

115. CHEILANTHES Swz. Lip-fern.

168. C. LANUGINOSA Nutt. Exposed rocks, two miles S. W. of Hackberry Springs, Banner Co. Aug. (Rydberg). Redwillow Co. (Bessey.)

ANTHOPHYTA.

48. NAIADACEÆ.

116. NAIAS L. Naiad.

169. *N. FLEXILIS* Rostk. & Schmidt. Stagnant water, Greenwood, July (Williams).

117. RUPPIA L. Ditch grass.

170. *R. OCCIDENTALIS* Watson. In Grand Lake (brackish water), abundant. Alliance Aug. 6. This is the form mentioned in the Catalogue under *Ruppia maritima* as probably belonging to the foreign variety *pedunculata* Hart. Mr. Morong writes: "There can be no doubt that the specimens sent me are *R. occidentalis* Watson, but with some differences. The leaves are somewhat shorter, the peduncles much longer and the fruit somewhat smaller. This form has not hitherto been found this side of Canada." (Webber.)

118. POTAMOGETON L. Pond weed.

171. *P. AMPLIFOLIUS* Tuckerm. Cropsey's Lake, Lincoln, July (Webber).172. *P. FLUITANS* Roth. (*P. lonchites*, Tuckerm.). Streams, etc. Anselmo, Custer Co. July (Webber).

173. *P. OAKESIANUS* Robbins. In pond, Thedford, Thomas Co., July 11. A rare find. Mr. Morong writes — "It is the first time I have ever known this species to occur so far west. The most westerly locality known for it hitherto has been the Adirondack region in New York." (Webber.)

174. *P. PECTINATUS* L. Lakes, etc., abundant. Fremont, July (Williams). Grand Lake, Alliance, Aug.; Thedford, Thomas Co., Aug. (Webber).

175. *P. SPIRILLUS* Tuckerm. Lincoln, July (Webber).176. *P. ZOSTERÆFOLIUS* Schum. Springy Lakes. Fremont, July; Greeley Center, July (Williams).

49. HYDROCHARIDACEÆ.

119. ELODEA Michx. Water-weed.

177. *E. CANADENSIS* Michx. Margins of Cold Lakes. Fremont, July (Williams).

50. ALISMACEÆ.

120. SAGITTARIA L. Arrow head.

178. *S. CALYCINA* Engelm. Moist Banks, Greenwood, July (Williams).179. *S. HETEROPHYLLA* Pursh. Shallow water and muddy banks. Greenwood, July (Williams).

180. *S. VARIABILIS* Engelm. var. *ANGUSTIFOLIA* Engelm. Narrow leaves with very narrow diverging lobes. Lodge Pole Creek, Deuel Co., July (Rydberg). Thedford, Thomas Co., Aug. 7; Anselmo, Custer Co., July (Webber).
181. *S. VARIABILIS* Engelm. var. *LATIFOLIA* (Willd.) Engelm. With monoeious flowers, broad and acute leaves. Near Platte River, Kearney Co., Aug. (Rydberg). Lincoln, Aug. (Webber).
182. *S. VARIABILIS* Engelm. var. *OBTUSA* (Willd.) Engelm. With dioecious flowers and broad and obtuse leaves. Sand Creek below Wahoo, Sanders Co., Sept. (Rydberg).

51. TYPHACEÆ.

121. *TYPIA* L. Cat-tail flag.
183. *T. LATIFOLIA* L. var. *TRIVIALE* (Pursh) B. S. P. Common Cat-tail. Long Pine (Swezey).

52. CYPERACEÆ.

122. *CAREX* L. Sedge.
184. *C. AUREA* Nutt. Wooded cañons and low prairies, very common. Lewellen, Alliance (Swezey). Thedford, Thomas Co., July; Hat Creek Basin, Aug.; Pine Ridge, July (Webber).
185. *C. DOUGLASII* Boott. Dry prairies. Anselmo, Custer Co., July (Webber).
186. *C. FILIFOLIA* Nutt. Dry places. War Bonnet Cañon, June (Williams). Alliance (Swezey). Hitchcock Co. (Hapeman).
187. *C. FILIFORMIS* L. var. *LATIFOLIA* Boeckl. Crete, Alliance (Swezey). Ashland, War Bonnet Cañon (Williams). Pine Ridge, Anselmo, Thedford (Webber).
188. *C. GRAVIDA* Bailey, var. *LAXIFOLIA* Bailey. Broken Bow, July 4 (Webber).
189. *C. LAXIFLORA* Lam. var. *VARIANS* Bailey. Nebraska City, June 1 (Webber).
190. *C. MARCIDA* Boott. Dry prairies. Alliance (Swezey). Anselmo, Broken Bow, Thedford (Webber).
191. *C. MUHLENBERGII* Schkuhr. Low prairies. War Bonnet, Weeping Water, June-July (Williams).
192. *C. PENNSYLVANICA* Lam. Crete, common (Swezey).
193. *C. SARTWELLII* Dewey. Ashland, May (Williams).
194. *C. SQUARROSA* L. Lincoln (Webber).
195. *C. STIPATA* Muhl. Low prairies, common, Crete (Swezey), Thedford, July 10, Anselmo, July 5 (Webber).
196. *C. STRAMINEA* Schkr. var. *BREVIOR* Dewey. Elmwood, June (Williams); Anselmo, Custer Co., July (Webber).
197. *C. TETANICA* Schkr. Ashland, May (Williams).
198. *C. TRIBULOIDES* Wahl. var. *CRISTATA* (Schw.) Bailey. Low ground. Wabash, July 5 (Williams).

199. C. TRICHOCARPA Muhl. var. ARISTATA (R. Br.) Bailey. Low prairies and moist places. Elmwood, Ashland (Williams).
123. HEMICARPHA Nees.
200. H. MICRANTHA (Vahl.) Britt. (*H. subsquarrosa* Nees.) Minden. Collected by Dr. Hapeman (Bessey). Ashland (Williams).
124. ERIOPHORUM L. Cotton grass.
201. E. GRACILE Koch. Collected at Minden by Dr. Hapeman (Bessey).
125. FIMBRISTYLIS Vahl.
202. F. CAPILLARIS (L.) Gray. Long Pine (Swezey). Minden (Hapeman and Bessey).
203. F. CASTANEA Vahl. (*F. spadicea* Vahl. var. *castanea* Gray). Collected by Rev. Bates at Valentine (Rydberg). Minden (Hapeman).
126. HELEOCHARIS R. Br.
204. H. ATROPURPUREA (Retz.) Kunth. Collected by Dr. Wibbe in Filmore Co. Reported by Dr. Britton (Torr. Bull. Vol. XVIII, May, 1891, p. 166).
205. H. OVATA (Roth.) R. & S. var. ENGELMANNI (Stend.) Britt. Pine Ridge, June 18 (Williams).
206. H. PALUSTRIS (L.) Roem. & Schult. var. GLAUCESCENS (Willd.) Gray. Anselmo, Custer Co., July 6 (Webber).
127. DULICHIUM Pers.
207. D. SPATHACEUM (L.) Pers. Collected by Rev. Bates at Valentine, Aug. 14 (Rydberg).
128. CYPERUS L.
208. C. ACUMINATUS Torr. & Hook. Doniphan (Swezey). Lincoln, July 20 (Webber).
209. C. DIANDRUS Torr. Long Pine (Swezey).
210. C. DIANDRUS Torr. var. CASTANEUS Torr. Very abundant, Mo. River bottoms, Richardson Co., Aug. 26 (Webber).
211. C. STRIGOSUS L. Ravenna, Brewster, Long Pine (Swezey).
53. GRAMINEÆ.
129. ASPRELLA Willd.
212. A. HYSTRIX (L.) Willd. Roadsides N. E. of Wahoo, June (Rydberg).
130. ELYMUS L. Wild Rye.
213. E. ELYMOIDES (Raf.) Swezey. (*E. sitanion* Schultes). Lewellen Swezey.
131. AGROPYRUM Gaertn. Wheat grass.
214. A. DASYSTACHYUM Vasey. Banks in thin woodlands; War Bonnet, June, 23. Rare (Williams).

132. *BROMUS* L. Brom grass.
215. *B. CILIATUS* L. var. *PURGANS* Gray. Belmont (Swezey).
133. *GLYCERIA* R. Br. Manna grass.
216. *G. PALLIDA* (Eddy) Trin. Wet places at margins of streams. War Bonnet, June 23. Rare (Williams).
134. *POA* L. Meadow grass.
217. *P. ALSODES* Gray. Woods on hillsides. War Bonnet, June (Williams).
218. *P. NEVADENSIS* Vasey. Woods in War Bonnet cañon near edges of stream, June 21 (Williams).
219. *P. SYLVESTRIS* Gray. Edges of woods, War Bonnet, June 23 (Williams).
135. *DIPLACHNE* Beauv.
220. *D. FASCICULARIS* (Lam.) Beauv. Collected in Minden by Dr. Hapeman and in Rock Co., by Rev. Bates (Bessey). Greenwood (Williams).
136. *AVENA* L. Oats.
221. *A. STRIATA* Michx. Wild oats. War Bonnet, June (Williams).
137. *ARISTIDA* L. Triple-awned grass.
222. *A. TUBERCULOSA* Nutt. Collected at Minden by Dr. Hapeman (Bessey).
138. *PANICUM* L. Panic-grass.
223. *P. WILCOXIANUM* Vasey. Prairies Ft. Niobrara. First collected by Dr. T. E. Wilcox, the post surgeon, and by him submitted to Dr. Vasey who described it as new in Bull. 8, U. S. Dept. Agriculture (Botanical Division) 1889, p. 32.
It is very near *Panicum scoparium* Lam. if indeed it is not a depauperate form of it (Bessey).

54. JUNCACEÆ.

139. *JUNCUS* Tourr. Rush, Bog-rush.
224. *J. BALISTICUS* Dethard var. *MONTANUS* Engelm. Alliance (Swezey). Broken Bow, July 4; Thedford, July 15 (Webber).
225. *J. BUFONIUS* L. Sandy ground, not uncommon. Lawrence Fork, Banner Co., Aug. (Rydberg). Broken Bow, July 4; Thedford, July 12 (Webber).
226. *J. MARGINATUS* Rostk. var. *PAUCICAPITATUS* Engelm. Brewster (Swezey).
227. *J. NODOSUS* L. Low prairies and edges of streams, very common. Thedford, Thomas Co., July 14; Auselmo, July 6 (Webber). Long Pine (Swezey).
228. *J. TENUIS* Willd. var. *SECUNDUS* Engelm. Long Pine (Swezey).

55. LILIACEÆ.

140. CALOCHORTUS Pursh.
 229. *C. SUTTALLII* Torr. & Gr. Along sides of cañons. Pine Ridge, June (Williams).
141. FRITILLARIA L.
 230. *F. ATROPURPUREA* Nutt. Along sides of cañons. War Bonnet, June (Williams).
142. POLYGONATUM Adans. Solomon's seal.
 231. *P. BIFLORUM* (Walt.) Ell. Cass Co (Williams).
143. SMILAX L. Greenbrier, Catbrier.
 232. *S. HERBACEA* L. var *PUBERULENTA* Gray. Woodlands, War Bonnet, June (Williams).

56. IRIDACEÆ.

144. IRIS L. Iris, Flag.
 233. *I. VERSICOLOR* L. Common Blue-flag. Edges of ponds, etc., common. Nebraska City, June (Webber).

57. PONTEDERIACEÆ.

145. HETERANTHERA Ruiz & Pav. Mud plantain.
 234. *H. LIMOSA* Vahl. In ponds, common. Lincoln and Fairmont (Bessey). Minden (Hapeman).

58. ORCHIDACEÆ.

146. HABENARIA Willd. Rein orchis.
 235. *H. BRACTEATA* (Willd.) R. Br. Moist places, common. War Bonnet, June (Williams).
147. SPIRANTHES L. Ladies' tresses.
 236. *S. CERNUA* (L.) Rich. Sand flats, on islands of the Platte River near Ashland. Sept. (Williams).
148. CORALLORHIZA R. Br. Coral root.
 237. *C. INNATA* R. Br. Damp dark woods in cañon. War Bonnet, June (Williams).
 238. *C. MULTIFLORA* Nutt. More common than the preceding species. Same locality (Williams).

59. CUPULIFERÆ.

149. BETULA L. Birch.
 239. *B. PAPYRIFERA* Marshall. Paper or Canoe birch, White birch. On the north slopes of the bluffs of the Niobrara River near Valentine (Bessey).

60. JUGLANDACEÆ.

150. HICORIA Raf. Hickory.

240. *H. SULCATA* (Willd.) Britt. Richardson Co. (Bessey).

61. SALICACEÆ.

151. SALIX L. Willow, Osier.

241. *S. ROSTRATA* Richardson. Wooded cañons, common. Belmont, July. Hat Creek Basin, Aug. (Webber).242. *S. TRISTIS* Ait. Dwarf gray willow. Anselmo, Custer Co., July. Mr. Bebb writes: "This is the most westerly locality of this species known to me." (Webber). Long Pine (Bates).

62. CERATOPHYLLEÆ.

152. CERATOPHYLLUM L. Honewort.

243. *C. DEMERSUM* L. Shallow lakes. Fremont, Greenwood, Ashland. June-Sept. (Williams.)

63. POLYGONACEÆ.

153. RUMEX L. Dock, Sorrel.

244. *R. BRITANNICA* L. Long Pine (Swezey).

154. POLYGONUM L. Knotweed, Smartweed.

245. *P. HYDROPIPEROIDES* Michx. Mild water-pepper. Wet places at edges of water, etc. Wabash, Aug. (Williams).246. *P. LAPATHIFOLIUM* L. Belmont (Swezey).247. *P. LAPATHIFOLIUM* L. var *INCARNATUM* (Ell.) Watson. Lawrence Fork, July 8 (Rydberg).248. *P. MUHLENBERGII* Watson. Low places. Wabash, Weeping Water, Ashland, Aug.- Sept. (Williams). Horse Creek, Scott's Bluff Co., Aug. 1; Lodge Pole Creek near Kimball, Aug. 12 (Rydberg).

155. ERIOGONUM Michx.

249. *E. ANNUUM* Nutt. f. —. Sand hills Kearney Co. (Rydberg). Ewing (Bessey).

Of the *E. annuum* type but differing in size, being larger (2-3 ft.); naked above; leaves lanceolate, 2-2½ in. long, ¼ in. wide, the upper nearly sessile; bracts triangular; flowers larger than those of *E. annuum*; the dense appressed wooliness turning yellowish on the branches (Rydberg).

250. *E. BREVICAULE* Nutt. Hills, Kiwa Valley, Scott's Bluff Co., July 28 (Rydberg).251. *E. CERNUUM* Nutt. Court House Rock, July 4; Wild Cat Mountains, Banner Co., July 16; Scott's Bluff, July 25; hills near Kimball, Aug. 12; near Sidney, Aug. 16 (Rydberg).

252. *E. CORYMBOSUM* Benth. Sand draws of Cheyenne Co., Aug. (Rydberg).
253. *E. MICROTHECUM* Nutt. var. *EFFUSUM* (Nutt.) Torr. & Gr. Lodge Pole Creek (Swezey).

64. CHENOPODIACEÆ.

156. *SUÆDA* Forskal. Sea blite.
254. *S. PROSTRATA* Pallas. (Ill. Pl. Imp. Cogn. p. 55 t. 47.— 1803). (*Salsola depressa*, Pursh Fl. 1814; Watson, in King's Rep.)
Salt Creek Basin N. W. of Lincoln (Rydberg). Differs from *S. linearis* (Ell.) Torr. in having the leaves broadest at the base and one or more of the calyx lobes strongly carinate.
157. *EUROTIA* Adans.
255. *E. LANATA* (Pursh) Moq. White sage. Dry prairies, in "Bad lands," etc. Hat Creek Basin, Aug. (Webber). Hillsides, Pumpkin Seed Valley and Lawrence Fork (Rydberg).
A white tomentose undershrub of the aspect of an *Artemisia*. The margins of the leaves are revolute, which separates it at a glance from *Artemisia cana* which it otherwise resembles. It is known under the name of "White sage."
- In Coulter's Manual *Eurotia* is described as dioecious. Watson in his "Revision of N. A. Chenopodiaceæ," states that it is sometimes monoecious, which is the case with all specimens found here (Rydberg).
158. *ATRIPLEX* L.
256. *A. ARGENTEA* Nutt. Not uncommon in the Salt Basin, N. W. of Lincoln, and saline soil in Kiwa Valley, Scott's Bluff Co. (Rydberg).
257. *A. HORTENSIS* L. Escaped along roads, etc. Hastings, Aug. 2 (Webber.)
159. *MONOLEPIS* Schrad.
258. *M. CHENOPODIOIDES* Moq. Dry saline soil in Deuel Co., June 27 (Rydberg).
160. *CHENOPodium* L. Pigweed, Goosefoot.
259. *C. BOSCIANUM* Moq. Long Pine, Belmont (?) (Swezey); Bennett, Aug. (Webber.)
260. *C. FREMONTII* Watson. Collected by Rev. Bates at Alliance (Rydberg); Belmont (Swezey); Anselmo (Webber).
261. *C. FREMONTII* Watson. var. *INCANUM* Watson. Collected by Rev. Bates at Valentine (Rydberg). Dry prairies, very abundant in Prairie dog towns, Thedford, July 16 (Webber). Lawrence fork, July 8; Kiwa Valley, Scott's Bluff Co., Aug. 1 (Rydberg).
262. *C. LEPTOPHYLLUM* Nutt. Common in Central and Western Nebraska. Valentine (Bates). Lodge Pole Creek (Swezey).

Sandy soil, Kearney Co. (Rydberg). Thedford and Anselmo, July (Webber).

263. *C. LEPTOPHYLLUM* Nutt. var. *OBLONGIFOLIUM* Watson. Valentine, collected by Rev. Bates; also in broken soil, Deuel Co., July 12 (Rydberg).
264. *C. LEPTOPHYLLUM* Nutt. var. *SUBGLABRUM* Watson. Alliance (Swezey). Pine Ridge (Webber). Scott's Bluff, July 24 (Rydberg).

65. AMARANTACEÆ.

161. *FREELICHIÀ* Moench.

265. *F. GRACILIS* Moq. "Sand draws," Deuel Co., Aug. 24 (Rydberg).

66. CARYOPHYLLACEÆ.

(Including *Illecebraceæ*.)

162. *CERASTIUM* L. Mouse-ear, Chickweed.

266. *C. ARVENSE* L. Collected by Rev. Bates at Ft. Robinson (Bessey). War Bonnet Cañon (Williams).

267. *C. VULGATUM* L. Crete. (Swezey.)

163. *STELLARIA* L. Chickweed.

268. *S. LONGIFOLIA* Muhl. Long-leaved Stitchwort. Low wet banks. Dismal River, Thomas Co., July 12 (Webber).

164. *ARENARIA* L. Sandwort.

269. *A. HOOKERI* Nutt. Pine Ridge, July; Hat Creek Basin, Aug. High rocky places, very common.

Dr. Britton after comparison with Nuttall's type of this species refers to it all of my specimens reported in the catalogue as *Arenaria pungens* (Catalogue No. 1213).

Mr. Rydberg has sent me specimens of an *Arenaria* from the cliffs of a cañon in Banner Co., labeled *A. franklini* Dougl. var. *minor* Hook. & Arn. that is probably to be referred here also (Webber).

270. *LOEFLINGIA* L.

270. *L. SQUARROSA* Nutt. High rocky prairies, Belmont, July 18. (Determined by Dr. Britton).

This is a very northerly locality for this plant, its usual range being from Southern California to Texas (Webber).

165. *LYCHNIS* L. Cockle.

271. *L. DRUMMONDII* Watson. Dry prairies. Hat Creek Basin, common, June (Williams).

272. *L. GITHAGO* (L.) Lam. Corn cockle. In cultivated fields. Weeping Water (Williams).

166. SILENE L. Catchfly, Campion.
 273. *S. MENZIESII* Hook. Woodlands, common. War Bonnet, June (Williams).
 274. *S. SCOULERI* Hook. Pine Ridge, July (Webber).
167. SAPONARIA L. Soap-wort, Bouncing Bet.
 275. *S. VACCARIA* L. Crete (Swezey).
168. ANYCHIA Rich. Forked chickweed.
 276. *A. CANADENSIS* (L.) B. S. P. (*A. dichotoma* Michx. var. *capillacea* Torr.) Woods, Weeping Water, June. Rare (Williams).

67. FICOIDEÆ.

(Luerssen places this under the family *Aizoaceæ*.)

169. MOLLUGO L. Indian chickweed.
 277. *M. VERTICILLATA* L. Long Pine (Swezey). Minden (Hapeman). Deuel Co. (Rydberg).

68. BERBERIDACEÆ.

170. CAULOPHYLLUM Michx. Pappoose-root.
 278. *C. THALICTROIDES* (L.) Michx. Blue cohosh. Sarpy Co., May (Pound).

69. RANUNCULACEÆ.

171. DELPHINIUM L. Larkspur.
 279. *D. TRICORNE* Michx. Dwarf larkspur. Edges of woods. Elmwood, May-June (Williams).
172. RANUNCULUS L. Buttercup, Crowfoot.
 280. *R. CIRCINATUS* Sibth. Lodge Pole Creek, Cheyenne Co., Aug. 19 (Rydberg).
 281. *R. HISPIDUS* Hook. Irrigated meadow north of Kimball, Aug. 12 (Rydberg).

70. CRUCIFERÆ.

173. THELYPODIUM Endl.
 282. *T. INTEGRIFOLIUM* Endl. Fields, etc., not uncommon. Alliance (Swezey). Pine Ridge, July 20 (Webber).
174. ERYSIMUM L. Treacle mustard.
 283. *E. ASPERUM* DC. Sandy hills in Deuel Co., July. Collected also by Rev. Bates at Valentine.
 Pods widely spreading. Large specimens were observed to roll before the wind, thus adding another "Tumble weed" to the already long list (Rydberg).
175. PHYSARIA Nutt.
 284. *P. DIDYMOCARPA* Gray. Cañon south of Scott's Bluff, July 22 (Rydberg).

176. LESQUERELLA Watson.

285. *L. ENGELMANNI* (Gray) Watson. Lawrence Fork, Kimball Co., Aug. 11 (Rydberg).

286. *L. GRACILIS* (Hook.). Watson. Waste ground along railroad, Weeping Water, June.

This is quite a northern extension of its range. It may have been introduced through the agency of the R. R., being found only in this one place. It was, however, abundant and evidently spreading (Webber).

177. CARDAMINE L. Bitter cress.

287. *C. LACINIATA* (Muhl.) Wood. (*Dentaria laciniata* Muhl.) Peru. Communicated by Mr. A. H. Van Fleet (Bessey).

178. ARABIS L. Rock cress.

288. *A. DENTATA* (Torr.) Torr. & Gr. Low ground, rare. Lincoln, April (Webber).

289. *A. GLABRA* (L.) Bernh. (*A. perfoliata* Lam.). Tower mustard. Dismal River, Thomas Co., July (Webber).

290. *A. HOLBOELLII* Hornem. Wooded cañons. Pine Ridge, June (Williams). Pumpkin Seed Valley, July 14 (Rydberg).

179. NASTURTIUM R. Br. Water cress.

291. *N. OBTUSUM*. Nutt. Ft. Clark, Nebraska (Hayden in Engelmann Herb.), Crete (Swezey), Minden (Hapeman), Big Springs (Rydberg).

Prof. Swezey's specimens differ somewhat from typical *N. obtusum*, being less diffusely spreading and branched, approaching likely nearer *N. sinuatum*. Hayden's specimens seem typical (Webber).

292. *N. PALUSTRE* (L.) DC. var. *OVALE* M. In mud, Sand Creek, Wahoo, Sept.

A small plant with the leaves entire, obovate or oval, crenate or with a pair or two of small lobes at the base; pod oval. (Rydberg.)

293. *N. SESSILIFLORUM* Nutt. Crete (Swezey). Wahoo Creek below Wahoo, Sept. (Rydberg). Lincoln, May (Webber).

294. *N. SINUATUM* Nutt. Moist places along banks, Elmwood, June (Williams), Fairbury (in Herb. Univ. of Nebr.), Omaha and Crete (Swezey). Platte River, Kearney Co., June 15 (Rydberg). Lincoln, May; Nebraska City, June (Webber). Pods and pedicels variously curved, forming a conspicuous character.

71. CAPPARIDACEÆ.

180. CRISTATELLA Nutt.

295. *C. JAMESII* Torr. and Gr. Collected by Rev. Bates at Valentine, Aug. 1 (Rydberg).

72. HYPERICACEÆ.

181. HYPERICUM L. St. John's-wort.
296. H. CANADENSE L. Common in low grounds. Thedford, July 14 (Webber). Ashland (Williams). *H. mutilum* L. of Swezey's Nebr. Fl. Plants p. 5, is probably to be referred here.
297. H. CANADENSE L. var. MAJUS Gray. Wet Meadows. Saunders Co., June (Rydberg). Lowell, July 11 (Webber).
298. H. VIRGINICUM L. (*Elodes virginica* Nutt. and *E. campanulata* Pursh). Antelope Co. (Webber).

73. GERANIACEÆ.

182. ERODIUM L' Her. Sterksbill.
299. E. CICUTARIUM (L.) L'Her. Alfilaria. Lincoln. Along sidewalks, probably introduced in hay. Adventive from Europe (Williams).

74. LINACEÆ.

183. LINUM L. Flax.
300. L. RIGIDUM Pursh. Prairie flax. Very common on the prairies of central and western Nebraska. Superior, Brewster, Long Pine, Lewellen (Swezey); Deuel Co., July (Rydberg); Thedford, July 14; Anselmo, July; Broken Bow, July; Belmont, Aug. (Webber). A form collected at Hat Creek Basin Aug. 2, Prof. Trelease notes as having the habit of var. *puberulum* Engelm. (Webber).
301. L. USITATISSIMUM L. Cultivated Flax. Escaped to low prairies. Broken Bow, July (Webber).

75. SAPINDACEÆ.

184. AESCULUS L. Horse-chestnut, Buckeye.
302. A. GLABRA Willd. Richardson and Pawnee counties (Bessey).

76. EUPHORBIACEÆ.

185. EUPHORBIA L. Spurge.
303. E. CUPHOSSPERMA (Engelm.) Boiss. A few specimens collected Aug., 1890, in a cañon running north from Lawrence Fork, Banner Co.
- Leaves lance linear, entire or with a few teeth on the revolute margin. Seeds short, triangular, truncate at the base, wrinkled and tuberculate, with a minute caruncle. Belongs to the *E. heterophylla* group (Rydberg).
304. E. FENDLERI Torr. & Gr. "Sand draw," Cheyenne Co., Aug. The seeds are described in Coulter's Manual as irregularly punctate. In the original description in Pacific R. R. Rep. it reads "a little rugose transversely," which agrees with Nebraska specimens (Rydberg).

305. *E. GEYERI* Engelm. Collected by Rev. Bates at Valentine, July 1 (Rydberg).

77. UMBELLIFERÆ.

186. *BERULA* Koch.

306. *B. ANGUSTIFOLIA* (L.) Koch. In water, Blackberry Springs, Banner Co., Aug. (Rydberg); Long Pine, Lewellen, Belmont (Swezey); Anselmo, Thedford, July-Aug. (Webber).

187. *CYMOPTERUS* Raf.

307. *C. GLOMERATUS* Raf. Lewellen, Alliance (Swezey). Collected by Rev. Bates at Valentine, May (Rydberg).

188. *PASTINACA* L. Parsnip.

308. *P. SATIVA* L. Common parsnip. Introduced. Mead, Saunders Co., June, 1890 (Rydberg).

78. ARALIACEÆ.

189. *ARALIA* L. Wild sarsaparilla.

309. *A. RACEMOSA* L. Spikenard. Nebraska City, Aug. (Williams).

79. SAXIFRAGACEÆ.

190. *RIBES* L. Currant, Gooseberry.

310. *R. GRACILE* Michx. Wild gooseberry. Saunders Co. This is the common wild gooseberry of Saunders Co., and Eastern Nebraska. It differs from *R. rotundifolium* Michx. in having long and slender peduncles, and capillary filaments 4-6" long. The majority of Nebraska specimens referred to *R. rotundifolium* Michx. likely belong here (Rydberg).

80. CACTACEÆ.

191. *OPUNTIA* Tourn.

311. *O. RUTILA* Nutt. Prairies. Deuel Co., June 23 (Rydberg).

81. ONAGRACEÆ.

192. *ENOTHERA* L. Evening primrose.

312. *Œ. ALBICAULIS* Nutt. var. *RUNCINATA* Engelm. "Sand draws," Deuel Co., Aug. (Rydberg). Very likely this is the sinuate leaved form of *Œ. albicaulis* referred to in the Catalogue Fl. Nebr. No. 1404.

313. *Œ. BIENNIS* L. var. *PARVIFLORA* Gray. Petals hardly $\frac{1}{2}$ in. long. Hills, Deuel Co. (Rydberg).

314. *Œ. HARTWEGII* Benth. var. *LAVANDULIFOLIA* (Torr. & Gr.) Watson. Prairies, etc. Lisbon, Perkins Co., June 23; Deuel Co., June 27 and July 2; Pleasant Valley, Scott's Bluff Co., July 28; Banner Co., Aug. (Rydberg). Lewellen, rare (Swezey).

315. *Œ. SPECIOSA* Nutt. Lincoln, probably an escape from cultivation (Bessey).

193. EPILOBIUM L. Willow herb.
 316. *E. HORNEMANNI* Richenb. Belmont (Swezey). (*E. alpinum* L. of Swezey's Nebr. Fl. Plants, p. 8.) Its occurrence needs confirmation.

82. ROSACEÆ.

194. POTENTILLA L. Five-finger.
 317. *P. ANSERINA* L. Silver-weed. Lewellen (Swezey).
 318. *P. RIVALIS* Nutt. var. *PENTANDRA* (Engelm.) Watson. Wabash, July (Williams).
 319. *P. SUPINA* L. Omaha, Lewellen (Swezey).
 195. CERCOCARPUS H. B. K. Mt. Mahogany.
 320. *C. PARVIFOLIUS* Nutt. Rocky hills, Banner Co., Aug. A shrub 3-6 ft. high. Seen at a distance it gives the hills a peculiar dark grayish color caused partly by the beautiful plumy tails of the fruit (Rydberg).

83. LEGUMINOSÆ.

196. AMPHICARPÆA Ell.
 321. *A. PITCHERI* Torr. & Gr. Banks of Sand Creek below Wahoo, Sept. (Rydberg). Woods, Ashland (Williams).
 Leaflets larger than in *A. comosa*; rachis villous; bracts large, silky canescent; upper flowers commonly fertile (Rydberg).
 197. LATHYRUS L.
 322. *L. ORNATUS* Nutt. Differs from *L. polymorphus* in having linear lanceolate leaves and the seed with a broad stalk and long hilum Pierce, Ft. Robinson (Herb. Univ. of Nebr.), Phelps Co., Fremont (Rydberg).
 A light yellow form of *L. ornatus* Nutt. is also sometimes found, Collected by Miss H. G. Wilkenson at Gordon. Fremont (Rydberg).
 198. ASTRAGALUS L. Rattle-weed.
 323. *A. CHAMELUCE*(?) Gray. Alliance, Lewellen (Swezey, Nebr. Flowering Plants, p. 7).
 (The specimens differ from typical *A. chamaeluce* in having too small flowers and too many leaflets. More abundant material is necessary to satisfactorily place the species. — Webber.)
 324. *A. PICTUS* Gray. var. *FILIFOLIUS* Gray. Lewellen, rare; Alliance (Swezey). Collected by Rev. Bates at Valentine (Rydberg).
 199. PETALOSTEMON Michx. Prairie clover.
 325. *P. COMPACTUS* (Spreng.) Swezey. (*P. macrostachyus* Torr.) Lewellen (Swezey).
 326. *P. TENUIFOLIUS* Gray. "Sand draws," Deuel Co., July. Low, branching from below, decumbent; leaflets 1-2 pairs,

linear-illiform, revolute; bracts ovate, densely villous, as is also the calyx, with grayish, slightly tawny hairs (Rydberg).

327. *P. sp.* — *P. candidus* var. *occidentalis* Gr. of Pringle's collection, according to Dr. Britton. It is, however, clearly no variety of *P. candidus*, but may be of *P. gracilis*, Nutt., of the south, to which it is nearly related. It differs from *P. gracilis* in having more oblong leaves, longer more lax spikes, and glandular dotted calyx. Dr. Britton says if distinct from *P. gracilis* it is clearly a distinct species (Rydberg).

200. PSORALEA L. Psoralea.

328. *P. DIGITATA* Nutt. Aurora (Williams). Much confounded with *P. campestris*. Specimens in the Herb. of Univ. of Nebr. collected at Anselmo by Webber and at Valentine by Rev. Bates, evidently belong here. Nebraska specimens collected by Dr. Bessey, have been referred to *P. digitata* by Dr. Britton (Rydberg).

329. *P. FLORIBUNDA* Nutt. This species, I think, should not be merged into *P. tenuiflora* Pursh. It is either a distinct species or a well marked variety. *P. floribunda* Nutt. found in eastern Nebraska has generally 5 oblong leaflets, 1-1½ in. long; many flowered racemes; lower calyx teeth longer; and larger flowers. *P. tenuiflora* Pursh, of western Nebraska, has 3 oblong-ob lanceolate leaflets, ½-¾ in. long; few flowered racemes; and punctate, colored calyx with equal teeth. Dr. Britton, to whom specimens and my notes upon them have been sent says: "I had concluded that *P. floribunda* could not go unnoticed into *P. tenuiflora*." The matter needs more investigation. Most of the localities for *P. tenuiflora* in the catalogue of Nebr. Flora belong to *P. floribunda*. The only specimens of the true *P. tenuiflora* I have seen are those collected by Dr. Bessey, at Ft. Robinson, and my own, from the plains of Deuel Co., Aug. 1890 (Rydberg).

330. *P. HYPOGEA* Nutt. Alliance, Lewellen (Swezey). Hills, Deuel Co., June 26 (Rydberg).

331. *P. LINEARIFOLIA* Torr. & Gr. Magnesia cliffs in Deuel Co., July. Racemes very loose, 3-6 in. long; leaflets 3, linear, 1-2 in. long; stem, leaves, and calyx glandular dotted (Rydberg).

201. TRIFOLIUM L. Clover, Trefoil.

332. *T. PROCUMBENS* L. Low hop-clover. Yards and roadsides, Lincoln. Adventive from Europe (Webber).

202. LUPINUS L. Lupine.

333. *L. ARGENTEUS* Pursh var. *ARGOPHYLLUS* Watson. Prairies, rare. War Bonnet, June (Williams).

334. *L. ARGENTEUS* Pursh var. *DECUMBENS* Watson. "Sand draws," 2 miles S. W. of Hackberry Springs, Aug. (Rydberg). Prairies of Hat Creek Basin, common (Webber).

335. *L. PUSILLUS* Pursh. Lewellen (Swezey). Hills near Curtis, Frontier Co., June 23; Deuel Co., June 25 (Rydberg).

84. ERICACEÆ.

203. MONOTROPA L. Indian pipe, Corpse-plant.
 336. M. UNIFLORA L. Washington and Cass Counties (Bessey).
 204. PYROLA Tourn. Wintergreen, Shin-leaf.
 337. P. CHLORANTHA Sw. Damp dark wooded cañons, rare. War Bonnet, June (Williams).
 338. P. SECUNDA L. Damp cañon at head of Jim Creek, Pine Ridge, June 25. Very rare (Williams).
 205. ARCTOSTAPHYLOS Adans. Bearberry.
 339. A. UVA-URSI (L.) Spreng. In a cañon near Anselmo, Custer County. The occurrence of this plant in the center of the state, hundreds of miles from any of its known stations, adds another puzzle to the many connected with the geographical distribution of the plants of the plains. It is known to occur in the Black Hills of South Dakota and the Rocky Mountains in Wyoming and Colorado. In Minnesota it is found near Pepin; it is absent from Iowa, while in Missouri it is confined to the southeastern part. It is doubtfully reported as occurring in Kansas. That it should be found in Central Nebraska is certainly unexpected (Bessey).

85. PRIMULACEÆ.

206. CENTUNCULUS Dill. Chaffweed.
 340. C. MINIMUS L. Fairfield (Swezey).
 207. LYSIMACHIA Tourn. Loosestrife.
 341. L. THYRSIFLORA L. Tufted loosestrife. Collected by Rev. Bates at Valentine, June (Rydberg); Lewellen (Swezey); Platte River near Doniphan, May (Harvey Thompson).
 208. DODECATHEON L. American cowslip.
 342. D. MEADIA L. Shooting-star. Alliance (Swezey).

86. CONVOLVULACEÆ.

209. CUSCUTA L. Dodder, Love-vine.
 343. C. TENUIFLORA Engelm. Crete, on *Salix* (Swezey).
 210. CONVOLVULUS L. Bindweed.
 344. C. ARvensis L. Bindweed. Roadsides. Ashland, May (Williams).

87. POLEMONIACEÆ.

211. GILIA Ruiz & Pav.
 345. G. GRACILIS Hook. "Sand draws" in Deuel Co., June 25 (Rydberg). War Bonnet Cañon (Williams).
 346. G. LINEARIS (Nutt.) Gray. Sides of cañon, Squaw Creek, June

(Williams). Dry banks, common: Pine Ridge, July 18; Hat Creek Basin, Aug. 1 (Webber). "Sand draws" in Deuel Co., June 25 (Rydberg).

347. *G. MINIMA* Gray. Collected at Rushville, in July, by Rev. Bates (Rydberg).
348. *G. PUNGENS* Benth. var *CAESPITOSA* Gray. Hills in Kiwa Valley, July 22; Scott's Bluff, July 25 (Rydberg).

88. HYDROPHYLLACEÆ.

212. PHACELIA Juss.

349. *P. CIRCINATA* (Willd.) Jacq. Abundant in dry cañons. Pine Ridge, July (Webber). War Bonnet (Williams).

Very *Krynnitzkia* like in appearance. Venation prominent above and below, leaf pinnately and obliquely straight veined. Lower leaves seldom with lateral leaflets in Nebraska specimens (Webber).

89. BORRAGINACEÆ.

213. LITHOSPERMUM L. Gromwell, Puccoon.

350. *L. LATIFOLIUM* Michx. Superior (Swezey).

351. *L. PIOSUM* Natt. War Bonnet, June (Williams).

214. MERTENSIA Roth.

352. *M. LANCEOLATA* (Pursh) DC. Wooded cañons. War Bonnet, June (Williams).

(Prof. Swezey's specimens from Lewellen and Chadron referred to this species, Nebr. Flowering Plants, p. 11, must be considered as belonging to *Pentstemon cæruleus* Nutt.—Webber.)

215. CRYPTANTHE Lehmann. [*Krynnitzkia crassisepala* Gray and *K. glomerata* Gray, No's. 1577 and 1578 of the Catalogue of Neb. Flora, should be changed according to Prof. Greene's paper (Pittonia, I. p. 110-112) to *Cryptanthe glomerata* Lehmann and *C. crassisepala* (T. & G.) Greene.]

353. *C. FENDLERI* (Gr.) Greene l. c. (*Krynnitzkia fendleri* Gray). Dry cañons, etc. "Bad lands." Hat Creek Basin, Sioux Co., Aug. (Webber). Lodge Pole Creek (Swezey). Collected by Rev. Bates at Harrison, Aug. Wild Cat Mts., Banner Co., July 17 (Rydberg).

216. OREOCARYA Greene. Pittonia I. p. 57.

354. *O. FULVOCANESCENS* (Gray) Greene, l. c. (*Krynnitzkia fulvocanescens* Gray). Dry sterile places, War Bonnet, June (Williams); Hat Creek Basin, Aug. (Webber).

355. *O. SUFFRUTICOSA* (Torr.) Greene, l. c. (*Eritrichium jamesii* Torr. *Krynnitzkia jamesii* (Torr.) Gray). Quite common in the western part of the state. Alliance (Swezey); Deuel Co.; Lawrence

Fork, Banner Co., Aug. (Rydberg); Harrison (Bates); Belmont, July; Hat Creek Basin, Aug. (Webber).

217. *ECHINOSPERMUM* Swartz. Stickseed.
356. *E. DEFLEXUM* (Wahl.) Lehm. var. *AMERICANUM* Gray. Long Pine (Swezey).
357. *E. FLORIBUNDUM* Lehm. Lawrence Fork, Banner Co., July 17 (Rydberg).

90. SOLANACEÆ.

218. *PHYSALIS* L. Ground cherry.
358. *P. LANCEOLATA* Michx. var. *HIRTA* Gray. Prairies. Hat Creek Basin, June (Williams).

Mr. Rydberg has noted two forms or varieties of *Physalis* belonging to the *lanceolata* group but differing enough from *P. lanceolata* to be distinct. I give below his descriptions in brief:—

No. 1. Glabrous or minutely hirsute on the calyx and the angles of the stems, erect, branched above; corolla yellowish with darker spot, only 5-7 lines across; anthers yellow; fruiting calyx inversely pear shaped; fruit greenish yellow; leaves lanceolate, sparingly toothed. "Sand draws," Deuel and Banner Co.'s July, 1890 (Rydberg).

No. 2. Slender, ascending; leaves oblong, lanceolate to oblanceolate, entire; calyx hispid; corolla, anthers, and fruit like the preceding; fruiting calyx globose. Near Pumpkin Seed Creek, Banner Co., Aug. 1890 (Rydberg).

91. SCROPHULARIACEÆ.

- 219 *PEDICULARIS* L. Louse wort.
359. *P. CANADENSIS* L. Bottom lands, common. Ashland, May (Williams).
360. *P. LANCEOLATA* Michx. Moist woods, rare. Ashland. Sept. (Williams).
220. *GERARDIA* L. Gerardia.
361. *G. TENUIFOLIA* Vahl. var. *MACROPHYLLA* Benth. Dry woods, common. Ashland, Weeping Water (Williams).
221. *LIMOSELLA* L. Mudwort.
362. *L. AQUATICA* L. Spring in Deuel Co., June 27 (Rydberg).
222. *MIMULUS* L. Monkey flower.
363. *M. LUTEUS* L. Lewellen (Swezey).
223. *PENTSTEMON* Mitchell. Beard tongue.
364. *P. HAYDENI* Watson. (Bot. Gazette, XVI (Nov., 1891), p. 311) Sandy prairies in central Nebraska, quite common. It is frequently found in the edges of "blow outs." Antelope Co.) July; Dismal River, Thomas Co., July 12 (Webber).

This includes *Pentstemon glaber* Pursh var. *utahensis* Watson, of Swezey's Nebr. Fl. Plants, p. 12; from Lewellen.

The habit is rather peculiar, growing as it does in very loose sandy places, the sand blowing here and there frequently banking up around it, modifies leaves and stems already formed. It is usually ascending, the lower portion of the stem running for some distance under the loose sand with no modification except a reduction of the leaves from lanceolate to linear and finally to filiform, or to mere scales. Many of the linear ones are quite long and expand somewhat above if the surface is reached (Webber).

92. OROBANCHACEÆ.

224. APIYLLON Mitchell. Cancer root, Naked broom rape.
 365. A. FASCICULATUM Gray. var. LUTEUM Gray. Collected by Rev. Bates at Valentine, June 10. Hills south of Scott's Bluff, July 26 (Rydberg).

93. LABIATÆ.

225. STACHYS L. Hedge-nettle.
 366. S. ASPERA Michx. var. TENUIFLORA (Willd.) Hitchcock, Cat. Anth. & Pter. of Ames, Ia., p. 513. (*Stachys aspera* Michx, var. *glabra* Gray). Banks of Wahoo Creek, Saunders Co., Aug. (Rydberg.)
 367. S. PALUSTRIS L. Collected by Rev. Bates at Valentine (Rydberg); Kearney, June (Webber).
 226. PHYSOSTEGIA Benth. False dragon-head.
 368. P. PARVIFLORA Nutt. Spring near Horse Creek, Scott's Bluff Co. Aug. 1 (Rydberg).
 227. SCUTELLARIA L. Skulcap.
 369. S. GALERICULATA L. Lewellen (Swezey). Spring, near Horse Creek, Scott's Bluff Co., Aug. 1 (Rydberg).
 228. MONARDA L. Horse-mint.
 370. M. FISTULOSA L. var. MOLLIS (L.) Benth. Wild bergamot. Banks of ravines, etc., Wahoo, Aug. (Rydberg).
 229. SALVIA L. Sage.
 371. S. AZUREA Lam. var. GRANDIFLORA Benth. Crete (Swezey).
 230. MENTHA L. Mint.
 372. M. CANADENSIS L. var. BOREALIS (Michx) Wood. (*M. canadensis* L. var. *glabrata* Benth.) Remarkable for its sweet scent. Hackberry Springs, Bruner Co., Aug. (Rydberg). Cass Co. (Williams).
 373. M. SATIVA L. Whorled mint. River banks, rare. Fremont, July (Williams).

94. VERBENACEÆ.

231. VERBENA L. Vervain.

374. *V. BRACTEOSA* x *HASTATA*. In a pasture 1 mile W. of Mead, Saunders Co., in June, 1890, there were found a few specimens of a Verbena, which undoubtedly is a hybrid between *V. bracteosa* Michx. and *V. hastata* L. The specimens are of the size and general appearance of *V. hastata*, but branched from the base and ascending. The leaves are of the size of those of *V. hastata* but more divided like those of *V. bracteosa*. The bracts are like those of the latter. Dr. Engelmann mentions several Verbena hybrids but not this (Rydberg).

375. *V. OFFICINALIS* L. Cultivated grounds and pastures, Wabash, July. Adventive from Europe (Williams). Tecumseh (Bessey).

232. LIPPIA L.

376. *L. LANCEOLATA* Michx. Low meadows, Ashland, July-Aug. Common (Williams). Richardson Co., Aug. (Webber). Minden (Bessey).

95. PLANTAGINACEÆ.

233. PLANTAGO L. Plantain, Ribwort.

377. *P. LANCEOLATA*. Roadsides, yards, etc., becoming abundant. Introduced. Ashland (Williams); Lincoln (Webber, Smith, Williams); Crete (Swezey); Aurora (Bessey).

378. *P. PATAGONICA* Jacq. var. *ARISTATA* (Michx.) Gray. Hastings (Bessey).

96. OLEACEÆ.

234. FRAXINUS L. Ash.

379. *F. AMERICANA* L. White ash. Sarpy and Nemaha Counties (Bessey).

97. GENTIANACEÆ.

235. GENTIANA L. Gentian.

380. *G. FLAVIDA* Gray. (*G. alba*. Muhl.) White Gentian. Low ground. Nemaha Co., Aug. (Webber); Weeping Water (Williams).

236. MENYANTHES L. Buckbean.

381. *M. TRIFOLIATA* L. Ponds, Cherry County (Bessey).

237. ERYTHRÆA Richard. Centaury.

382. *E. DOUGLASII* Gray. On the sands of the Platte River in Scott's Bluff Co., Aug. 1 (Rydberg).

98. ASCLEPIADACEAE.

238. ACERATES Ell. Green milkweed.

383. A. AURICULATA Engelm. Lewellen, rare (Swezey). "Sand-draws" in Deuel Co., July (Rydberg).

It is easily mistaken for *Asclepias stenophylla* from which it can not be distinguished except by the form of the hood and its auricles (Rydberg).

384. A. VIRIDIFLORA (Raf.) Ell. var. LANCEOLATA (Ives) Gray. Hills, Wahoo, June, 1890 (Rydberg).

99. COMPOSITE.

239. STEPHANOMERIA Nutt.

385. S. MINOR Nutt. "Bad lands" north of Scott's Bluff, July 22. It is easily mistaken for a *Lygodesmia* (Rydberg).

240. LACTUCA L. Lettuce.

386. L. INTEGRIFOLIA Bigel. Roadsides, Weeping Water (Williams).

241. CREPIS L.

387. C. INTERMEDIA Gray. Side of cañon, plentiful. War Bonnet, June (Williams).

388. C. RUNCINATA (James) Torr. & Gr. Collected by Rev. Bates at Valentine (Rydberg). Lewellen (Swezey). Platte bottoms, Cheyenne Co., July 3; Pumpkin Seed Valley, July 14 (Rydberg).

242. CICHORIUM L. Succory, Chicory.

389. C. INTYBUS L. Yards, Lincoln, Introduced (Webber).

243. CNICUS L. Thistle.

390. C. ALTISSIMUS (L.) Willd. var. FILIPENDULUS Gray. Belmont (Swezey).

391. C. UNDULATUS (Nutt.) Gray, var. CANESCENS (Nutt.) Gray. Prairies, common. Antelope Co., Pine Ridge, July-Aug.

A form from Pine Ridge has conspicuously decurrent leaves, a character shown also by a specimen of Hayden's in the Engelmann Herbarium from the Sand hills of the Plains.

Cnicus pitcherii Torr., No. 1738 of the Cat. of the Fl. of Nebr. belongs here (Webber).

392. C. UNDULATUS (Nutt.) Gray, var. MEGACEPHALUS Gray. Broken Bow, July 4 (Webber).

244. ARNICA L.

393. A. CORDIFOLIA Hook. Hillsides, rare. Squaw Cañon, Pine Ridge, July (Williams).

245. HELENIUM L.

394. H. AUTUMNALE L. Sneeze-weed. Collected by Rev. Bates at Valentine (Rydberg); Minden (Hapeman).

246. PECTIS L.

395. *P. ANGUSTIFOLIA* Torr. "Sand draws" Deuel Co., July.

A little plant (1-3 in. high) with glandular punctate leaves and pleasant odor (Rydberg).

247. THELESPERMA Lees.

396. *T. AMBIGUUM* Gray. Prairies, quite common. Long Pine (Swezey); Alliance, July; Crawford, Aug.; Belmont, July (Webber).

397. *T. FILIFOLIUM* (Hook.) Gray. Table land, Banner Co., Aug. Collected also by Rev. Bates at Valentine (Rydberg).

248. COREOPSIS L. Tickseed.

398. *C. ARISTOSA* Michx. Long Pine (Swezey).

399. *C. TRICHOSPERMA* (?) Michx. var. *TENUILOBA* Gray. Lewellen, Alliance (Swezey in Nebr. Flowering Plants). (This has the character of *Thelesperma* and will likely prove to be near *T. ambiguum*. — Webber).

249. HELIANTHUS L. Sunflower.

400. *H. DECAPETALUS* (?) L. Crete (Swezey Nebr. Flowering Plants, p. 10).

(This is near *H. tuberosus* L. to which it may have to be referred.— Webber).

250. ECLIPTA L.

401. *E. ALBA* (L.) Haussk. Crete (Swezey).

251. FRANSERIA Cav.

402. *F. HOOKERIANA* Nutt. Alliance (Swezey). Lodge Pole Creek, near Potter, Aug. 15 (Rydberg).

403. *F. TOMENTOSA* Gray. Becoming a troublesome weed on low rich soil in Kearney Co. (Rydberg).

252. AMBROSIA L. Ragweed.

404. *A. TRIFIDA* L. var. *INTEGRIFOLIA* (Muhl.) Torr & Gr. With the type in various places. At Hackberry Springs only this form was found (Rydberg); Lincoln (Bessey); Minden (Hapeman).

253. IVA L.

405. *I. AXILLARIS* Pursh. Near Platte River, Scott's Bluff Co., July 25 (Rydberg).

254. ANTENNARIA Gærtn. Everlasting.

406. *A. DIMORPHA* Torr. & Gr. Collected at Harrison by Rev. Bates (Bessey).

407. *A. DIOICA* (L.) Gaertn. Alliance (Swezey); Belmont, July 14 (Webber).

255. ERIGERON L. Fleabane.

408. *E. CANUS* Gray. Lawrence Fork, Banner Co., Aug. (Rydberg).

409. *E. CESPITOSUS* Nutt. Prairies, not uncommon. Hat Creek Basin, Aug.: Belmont, July (Webber).
410. *E. CONCINNUS* Torr. & Gray. Dry prairies. War Bonnet, June (Williams).
411. *E. MACRANTHUS* Nutt. Cañons, etc. Hat Creek Basin, Aug. 2.
(Prof. Swezey's specimens of *E. glabellus* Nutt. from Long Pine (Neb. Flowering Plants, p. 9) probably belong here, although differing somewhat from the type in having a hairy involucre. My specimens also show this peculiarity but were pronounced by Dr. Watson to be *E. macranthus*.—Webber.)
412. *E. RUMILUS* Nutt. Collected by Rev. Bates at Valentine (Rydberg); War Bonnet, June (Williams); Lewellen (Swezey); Pine Ridge, July; Hat Creek Basin, Aug. (Webber).
Quite common on dry prairies.
256. ASTER L. Aster.
413. *A. AZUREUS* Lindl. Weeping Water (Williams).
414. *A. ERICEFOLIUS* Rothrock. Prairies, Venango, Perkins Co., June 23 (Rydberg).
415. *A. FOLIACEUS* Lindl. Platte River north of Scott's Bluff, July 23; Horse Creek, Aug. 1; Lodge Pole Creek, near Kimball, Aug. 12 (Rydberg).
416. *A. PANICULATUS* Lam. Low prairies, very common, Lincoln, Sept. (Webber).
417. *A. PUNICEUS* L. Prairies. Clear water, Antelope Co., Sept. (Webber).
418. *A. TRADESCANTI* L. Crete, common (Swezey, Nebr. Fl. Pl., p. 9.)
(The specimens of this seem to me to be nearer *A. paniculatus*. Its occurrence in Nebraska needs confirmation.—Webber.)
257. TOWNSENDIA Hook.
419. *T. SERICEA* Hook. Prairies, not uncommon. Collected by Dr. W. A. Thomas in Hayes Co., April (Rydberg); McCook; Alliance; Ogalalla (Swezey); Antelope Co. (Webber).
258. SOLIDAGO L. Golden rod.
420. *S. CANADENSIS* L. var. *PROCERA* Torr. & Gr. Weeping Water (Williams).
421. *S. NEMORALIS* Ait. var. *INCANA* (Torr. & Gr.) Gray. Hills, Deuel Co., Aug. (Rydberg); Lewellen; Alliance; Lodge Pole Creek; Belmont (Swezey).
422. *S. RADULA* Nutt. Edges of thickets, Weeping Water (Williams).
423. *S. SHROTINA* Ait. var. *GIGANTEA* (Ait.) Gray. Saunders Co., Sept. (Rydberg).
424. *S. SPECIOSA* Nutt. Prairies, Ashland, June (Williams).
259. HAPLOPAPPUS Cass.
425. *H. NUTTALLII* Torr. & Gr. "Sand draws," of Banner Co. (Rydberg). Denuded places, in "Bad Lands," Hat Creek Basin, Sioux Co., Aug. (Webber).

426. *H. RUBIGINOSUS* (?) Torr. & Gr. Lodge Pole Creek (Swezey,
Nebr. Fl. Pl., p. 9).

(Prof. Swezey's specimens are rather meager to decide upon,
and as Nebraska is somewhat out of the known range of the
species, its occurrence may perhaps remain in question until con-
firmed.)

260. *CHRYSOPSIS* Nutt. Golden aster.

427. *C. VILLOSA* (Pursh) Nutt. var. *CANESCENS* (DC.) Gray.
"Sand draws," Deuel and Banner Co.'s (Rydberg).

428. *C. VILLOSA* (Pursh) Nutt. var. *HISPIDA* (Hook.) Gray.
"Sand draws," Deuel Co. (Rydberg). Long Pine (Conklin,
Swezey).

429. *C. VILLOSA* (Pursh) Nutt. var. *SESSILIFLORA* (Nutt.) Gray.
"Sand draws," Banner Co. (Rydberg).

261. *BRICKELLIA* Ell.

430. *B. GRANDIFLORA* (Hook.) Nutt. In a cañon, Banner Co. (Ryd-
berg).

262. *VERNONIA* Schreb. Iron weed.

431. *V. ALTISSIMA* Nutt. Banks, etc. Wahoo, Saunders Co., Sept.
(Rydberg).

432. *V. NOVEBORACENSIS* (L.) Willd. Bottom meadows, Ashland,
Aug. (Williams).

SUPPLEMENTARY LIST OF RECENTLY REPORTED SPECIES.

BY CHARLES E. BESSEY, PH.D.

Species determined by Rev. J. M. Bates of Valentine, are marked (B.); those by A. T. Bell and W. H. Skinner of the Crete High school, (B. & S.); those by F. C. Clements, a student in my laboratory, (C.); those by Dr. H. Hapeman of Minden, (H.); those by Roscoe Pound, formerly assistant in my laboratory, (P.); those by P. A. Rydberg, a graduate student, (R.); those by J. R. Schofield, a student in my laboratory, (S.); those by Professor G. D. Swezey of Doane College, (Sw.); those by A. F. Woods, assistant in my laboratory, (W.).

MYXOMYCETES.

1. *Perithaena flavida* Peck. On rotten logs. Crete, Nov. 1891, (B. & S.)
2. *Trichia chrysosperma* (Bull.) DC. On rotten logs. Crete, Nov. 1891, (B. & S.)
3. *Trichia varia* Pers. Crete, Nov. 1891, (B. & S.)
4. *Arcyria punicea* Pers. Crete, Nov. 1891, (B. & S.)
5. *Arcyria adnata* (Batsch.) Rost. Crete, Nov. 1891, (B. & S.)
6. *Arcyria nutans* (Bull) Grev. Crete, Nov. 1891, (B. & S.)
7. *Hemiarcyria serpula* Scop. On bark, Lincoln, (C.)
8. *Comatricha friesiana* (D. By.) Rost. On rotten wood in a dark well. York, Apr. 1892, (Bessey.)
9. *Stemonitis ferruginea* Ehrh. Crete, Nov. 1891, (B. & S.)
10. *Chondrioderma radiatum* (Linn.) Rost. Crete, Nov. 1891, (B. & S.)

BACTERIACEÆ.

11. *Bacterium aceti* (Kuetz.) Lanzi. In fermenting cider. Lincoln, Feb. 1892, (Bessey.)
12. *Beggiaea pellucida* Cohn. Lincoln, (W.)

NOSTOCACEÆ.

13. *Microcoleus terrestris* Desm. var. *repens* Kuetz. In greenhouse. Lincoln, Feb. 1892, (Bessey.)
14. *Nostoc muscorum* Ag. On moss leaf from Valentine, (C.)
15. *Oscillaria froelichii* Kuetz. var. *fusca* Kirch. In greenhouse. Lincoln, Feb. 1892, (Bessey.)
16. *Oscillaria gracillima* Kuetz. In the basin of the artesian well, Lincoln, Dec. 1891, (S.)

17. *Oscillaria princeps* Vauch. Lincoln, July, 1891, (W.)
18. *Lyngbya aestuarii* Liebm. Forming a reddish-brown coating upon the ground near Lincoln, the soil probably somewhat saline. April, 1892, (Bessey.)
19. *Isactis fluvialis* (Rab.) Kirch. Minden, (H.)

PALMELLACEÆ.

20. *Protococcus frustulosus* (Carm.) D. Ton. In the basin of the artesian well, Lincoln, Dec. 1891, (S.)
21. *Protococcus vestitus* Reinsch. In material collected by Dr. Hapeman at Minden, (C.)
22. *Raphidium braunii* Naeg. In material from Minden collected by Dr. Hapeman, (C.)
- 22½. *Raphidium polymorphum* Fries. var. *aciculare* A. Br. Minden, (H.)
23. *Tetraspora lubrica* (Roth.) Ag. Minden, Coll. Hapeman, (W.)

CLADOPHORACEÆ.

24. *Cladophora glomerata* (L.) Kuetz., var. *simplicior* Kuetz. Minden, Coll. by Dr. Hapeman, (W.)
25. *Draparnaldia plumosa* Ag. Wahoo, (R.)
26. *Draparnaldia glomerata* (Vauch.) Ag. var. *genuina* Kirch. In a pool. Lincoln, (W.)
27. *Charophora cornu-dame* (Roth.) Ag. var. *polyclados* Kuetz. Minden, Coll. Hapeman, (W.)

DESMIDIACEÆ.

28. *Desmidium aptogonium* Breb. Minden, Coll. Hapeman, (C.)
29. *Closterium acerosum* (Schr.) Ehrb. Lincoln, (C.)
30. *Closterium dianae* Ehrb. Lincoln, (C.)
31. *Closterium intermedium* Ralfs. Minden, Coll. Hapeman, (C.)
32. *Closterium lanceolatum* Kuetz. Lincoln, (W.)
33. *Cosmarium gotlandicum* Wittr. Minden, Coll. Hapeman, (C.)
34. *Cosmarium botrytis* Menegh. In material collected by Dr. Hapeman at Minden, (C.)
35. *Cosmarium portianum* Arch. Minden, Coll. Hapeman, (C.)
36. *Euastrum pokornyanum* Grun. In material collected at Minden by Dr. Hapeman, (C.)
37. *Staurastrum dickiei* Ralfs. In material collected at Minden by Dr. Hapeman, (C.)
38. *Staurastrum eustephanum* (Ehr?) Ralfs. In material collected at Minden by Dr. Hapeman, (C.)
39. *Staurastrum quadrangulare* Breb. In material collected at Minden by Dr. Hapeman, (C.)

DIATOMACEÆ.

40. *Stauroneis lineata* Kg. Agrees with Wolle's figure, but no description is accessible. In material from Minden, collected by Dr. Hapeman, (C.)
41. *Stauroneis phænicenteron* (Nitzsch.) Ehr. Lincoln, (C.)
42. *Navicula formosa* Greg. Lincoln, (C.)
43. *Navicula hemiptera* Kg. Crete, (C.)
44. *Navicula mutica* Kg. Lincoln, (C.)

45. *Navicula rhomboides* Ehrb. In material collected at Minden by Dr. Hapeman, (C.)
46. *Pleurosigma spenceri* (Quek.) W. Sm. Lincoln, (C.)
47. *Gomphonema dichotomum* Kuetz. Minden, (H.)
48. *Gomphonema geminatum* S. & F. Minden, (H.)
49. *Cocconeis striata* Ehrb. In material collected at Minden by Dr. Hapeman, (C.)
50. *Epithemia argus* Kuetz. Minden, (H.) This species was doubtfully recorded in the catalogue (No. 100) as from Lincoln.
51. *Epithemia sorex* Kuetz. Pauline, (H.)
52. *Epithemia zebra* (Ehr.) Kuetz. Minden, (H.)
53. *Synedra affinis* Kuetz. Minden, in material coll. by Dr. Hapeman, (C.)
54. *Synedra ulna* (Nitzsch) Ehr. var. *longissima* (W. Sm.) Brun. In material collected at Minden by Dr. Hapeman, (C.)
55. *Fragilaria virescens* Ralfs. Minden, in material collected by Dr. Hapeman, (C.)
56. *Nitzschia communis* Rabh. In the basin of the artesian well. Lincoln, Dec. 1891, (S.)
57. *Nitzschia thermalis* (Ehr.) Auersw. In salt water. Lincoln, (C.)
58. *Nitzschia vermicularis* (Kg.) Hantz. Minden, (H.)
59. *Nitzschia vitrea* Norm. Minden, (H.)
60. *Nitzschia vivax* W. Sm. Minden, (H.)

ZYGONEMACEÆ

61. *Mougeotia genuflexa* (Dillw.) Ag. Lincoln, (C.)
62. *Zygnema pectinatum* (Vauch.) Ag., var. *anomalum* (Hass.) Kirch. Minden, Coll. by Dr. Hapeman, (W.)
63. *Spirogyra crassa* Kuetz. South Bend, (C.) Filaments 136 mmm. in diameter.

OEDOGONIACEÆ.

64. *Oedogonium pringsheimii* Cram Minden, Coll. Hapeman, (W.)
65. *Oedogonium wolleanum* Wittr. Minden, Coll. Hapeman, (C.)

COLEOCHÆTACEÆ.

66. *Coleochete scutata* Breb. In material from Minden, collected by Dr. Hapeman, (C.)

SPHERIACEÆ.

67. *Nylaria polymorpha* (Pers.) Grev., var. *spathulata* Pers. Lincoln, (C.)
68. *Nylaria polymorpha* (Pers.) Grev., var. *acrodactyla* Nits. Lincoln, (C.)
69. *Hypoxylon epiphloeum* B. & C. Lincoln, (W.)

DOTHIDIACEÆ.

70. *Dothidia ribesia* Pers. Lincoln. On stems of *Ribes gracile* Michx. (W.)

PEZIZACEÆ.

71. *Lachnea scutellata* Linn. On earth in greenhouse. Lincoln, (C.)
72. *Sclerotinia durecana* Tul. On ground in woods. Lincoln, Apr. 1892, (C.)
73. *Sclerotinia trifoliorum* Eriks. Raymond, (C.)
74. *Sclerotinia gracilis* Clements, n. sp. Cup thin, hemispherical, bright brown, margin entire, darker: stipe elongated, flexuous, thick, 1.5—

2 cm. long: ascii cylindrical, attenuated at base, 150—160 x 10 mm. ascospores 8, oblong-elliptical, curved, biguttulate, 26—32 x 10—11: mm.: paraphyses filiform slightly clavate at apex, somewhat branched. Sclerotium black, thin, coriaceous, forked, $\frac{1}{3}$ x 1½ cm. In shady woods, near Lincoln, Nebr., (associated with *Erythronium albidum*?) The cups are from 2½ to 3½ mm. broad, a single stipe arising from each of the two to three forks of the sclerotium.

75. *Phialea aquatica* (Carr.) Sacc. Muddy banks. Raymond. (C.)
76. *Belonidium aurelia* (Pers.) De Not. Lincoln, on stumps, (C.)

HELVELLACEÆ.

77. *Verpa digitaliformis* Pers. In woods under trees. Raymond, May 1892, (C.)

UREDINEÆ.

78. *Puccinia buchloes* Schofield, n. sp. III. Amphigenous, or mostly hyphophyllus: sori scattered, or more or less linear-clustered, epidermis of host plant splitting and supporting the edges of the sori: spores broadly elliptical, or slightly obovate, 13 to 23 broad by 29 to 36 mm. long, mostly about 22 by 31 mm., episporae smooth, light-brown, constricted at the septum, slightly thickened at the apex, which is obtusely rounded: pedicel stout, once or twice the length of the spore, tinted. On *Buchloe dactyloides* Engelm. Collected by J. G. Smith, Lincoln, Nebr., Oct. 1886.

USTILAGINEÆ.

79. *Sorosporium cuneatum* Schofield, n. sp. Glomerules nearly globose, 38 to 48 mm. in diameter, mostly about 44.5 mm., bright brown, composed of from forty to fifty spores: spores 9.5 to 11 mm. long, crowded, and in optical section cuneate in outline, all minutely roughened on the basal (exposed) surface.

In flower heads and stems of *Grindelia squarrosa* Dunal. Lincoln, Nebr., in the autumn of 1891.

The fungus attacks not only the heads, completely destroying the achenes, but it attacks the stem also, producing gall-like enlargements. The spores in the stem are formed immediately under the bark.

In California, according to Dr. Harkness, *Sorosporium californicum* Hark. attacks *Grindelia robusta* Nutt., reducing the flower-heads to one-third their natural size. In Nebraska the same reduction occurs on the lower branches of *Grindelia squarrosa* Dunal., but on the upper branches the parasite has the opposite effect, enlarging the heads to more than double their natural size.

MUCEDINEÆ.

80. *Verticillium rufum* (Schwabe.) Rebh. On sugar beets in silo. Lincoln, (P.)

LYCOPERDACEÆ.

81. *Lycoperdon cepaforme* Bull. Lincoln, (C.)
82. *Lycoperdon peckii* Morg. Lincoln, (C.)

AGARICINÆ.

83. *Hatula crenulata* Fr. Prairies, Lincoln, (C.)
 84. *Habroloma crustuliniforme* Bull. Lincoln, (C.)
 85. *Coprinus comatus* Fl. D. Lincoln, (W.)
 86. *Coprinus plicatilis* (Curt.) Fr. Lincoln, (Bessey.)

POLYPOREÆ.

87. *Polystictus pergamenus* Fr. Lincoln, (W.)

THELEPHOREÆ.

88. *Stereum versicolor* (Schw.) Fr., var. *fasciata* Schw. Lincoln, (W.)

CHARACEÆ.

89. *Chara foetida* A. Br. Collected by J. M. Bates at Valentine, (Bessey.)

EQUISETACEÆ.

90. *Equisetum limosum* L. Platte river bottom, Kearney Co., June 15, 1891. (R.)
 91. *Equisetum pratensis* Ehrh. Sowbelly Canon, Sioux Co., Aug. 1890, (Bessey.)
 92. *Equisetum hyemale* L. Collected by J. M. Bates at Ft. Niobrara, (Bessey.)

CYPERACEÆ.

93. *Carex marcida* Boott, var. *debilis* Bailey, Deuel Co., (R.)
 94. *Heleocharis compressa* Sull. Ewing. (B.)

GRAMINEÆ.

95. *Elymus sibiricus* L. Sioux Co. (B.)
 96. *Elymus striatus* Willd., var. *villosus* Gr. Long Pine, (B.)
 97. *Agropyrum violaceum* Lange. North-western Nebraska. (B.)
 98. *Festuca confinis* Vasey., var. ——— Harrison and Crawford, July 1891, (B.)
 99. *Poa alpina* L. McColligan canon, Deuel Co., June 26, 1891, (R.)
 100. *Glyceria distans* Wahl., var. *aireoides* (L.) Wabl. Lodge Pole creek, near Sidney, Aug. 19, 1891, (R.)
 101. *Koeleria cristata* Pers., var. *gracilis* Gray. Prairie, Deuel Co., June 25, 1891, (R.)
 102. *Arrhenatherum avenaceum* Beauv. Lincoln, June 12, 1891, (R.)
 103. *Calamagrostis confinis* Nutt. Platte River, Kearney, Co., June 15, 1891, (R.)
 104. *Sporobolus cryptandrus* (Trin.) Gr., var. (near) *flexuosus* Thurb. The pedicels are capillary, but not drooping. Dry sandy soil near Scott's Bluff, July 24, 1891, (R.)
 105. *Sporobolus confusus* Town. This is the *S. serotinus* of the "Contributions from the Botanical Department of the University of Nebraska. New Series, II.," page 6. Sands of Platte, near Horse creek, Scott's Bluff Co., Aug 1, 1891, (R.)
 106. *Sporobolus minor* Vasey. Long Pine, (B.)
 107. *Muhlenbergia racemosa* (Michx.) B. S. P., var. *ramosa* Vasey. Common westward, (B.)
 108. *Aristida oligantha* Michx. Cherry Co., (B.)

109. *Panicum crus-galli* L. var. *muticum* Vasey. This is without any doubt indigenous. Canon in Kiwa Valley, Scott's Bluff Co., July 28, 1891, (R.)
110. *Panicum dichotomum* L. var. *barbulatum* (Michx.) Gray. Platte river bottom, Kearney Co., June 15, 1891, (R.)
111. *Panicum virgatum* L., var. *confertum* Vasey. Banner Co., (R.)

JUNCACEÆ.

112. *Juncus balticus* Deth. North Platte river, Cheyenne Co., July 3, 1891, (R.)
113. *Juncus marginatus* Rost. Minden, (H.)
114. *Juncus tenuissimus* Willd., var. *congestus* Engelm. Kennedy, (B.)

LILIACEÆ.

115. *Trillium nivale* Riddell. Weeping Water, Apr. 1892, (Miss Rands.)
116. *Zygadenus nuttallii* Gray. In fruit, Hills of Pumpkin Seed Valley, July 6, 1891, (R.)

IRIDACEÆ.

117. *Sisyrinchium anceps* Cav. Spring south of Pumpkin Seed Valley, July 14, 1891, (R.)

POLYGONACEÆ.

118. *Polygonum aviculare* L. var. *parva* Holz. var. nov. Stem much branched, low, prostrate, branchlets with very short joints, nearly covered by the sheaths, which are strongly ribbed and brown at the base; the scarious tips long, loose, aristate; leaves crowded, oblong 2-3 lines long; flowers minute, reddish or brownish. According to Mr. J. M. Holzinger, it has been collected by Wright in 1851 in N. Mex., and by Newberry (?) near Dallas, Texas. Spring, Deuel Co., Aug. 24, 1891, (R.)

NYCTAGINACEÆ.

119. *Abronia micrantha* Dougl. Court House Rock, Cheyenne Co., July 4, 1891, (R.)

CARYOPHYLLACEÆ.

120. *Arenaria franklinii* Dougl. Court House Rock, Cheyenne Co., July 4, and cliffs in Banner and Scott's Bluff Cos., July-Aug. 1891, (R.)

RANUNCULACEÆ.

121. *Clematis douglasii* Hook., var. *scottii* Porter. Reported by Professor Swezey from Sheridan Co.

CRUCIFERÆ.

122. *Stanleya pinnatifida* Nutt. Reported by Professor Swezey from Sheridan Co.
123. *Draba nemorosa* Ledb. Alliance, (Bessey.)
124. *Nasturtium officinale* Nutt. Near a well 1½ miles from the Wyoming line, Scott's Bluffs Co., July 31, 1891, (R.)

CISTACEÆ.

125. *Lechea minor* L. Minden, collected by Dr. Hapeman, (W.)

HYPERICACEÆ.

126. *Hypericum ascyron* L. Cass Co., (Sw.)

POLYGALACEÆ.

127. *Polygala sanguinea* L. Atkinson and Neligh, (B.)

UMBELLIFERÆ.

128. *Cicuta bulbifera* L. Cherry Co. Not in fruit, (B.)
 129. *Osmorrhiza claytoni* (Michx.) B. S. P. Cass Co., (Sw.)
 130. *Musenium divaricatum* Nutt. Harrison, (B.)
 131. *Piceodamnum kingii* Watson. This, possibly, should be included in the genus *Pseudocymopterus* Coulter and Rose, as the dorsal ribs are more or less winged. Scott's Pass, July 22, 1891, (R.)

SAXIFRAGACEÆ.

132. *Ribes setosum* Lindl. Dawes Co., (Sw.)

LOASACEÆ.

133. *Mentzelia albicaulis* (Hook.) Dougl. Scott's Bluffs Co., (R.) In the specimens collected the seeds are muricate, but the leaves are nearly entire or more rarely sinuately toothed.

HALORAGEÆ.

134. *Callitricha verna* L. Kennedy, July 1891, (B.)

ROSACEÆ.

135. *Crataegus subvilloso* Schrad. Sarpy and Cass counties, (Sw.)
 136. *Rosa nutkana* Presl. Curtis, Frontier Co., June 22. McColligan canon, Deuel Co., June 26, 1891, (R.)

LEGUMINOSÆ.

137. *Desmodium illinoense* Gray. Reported from Cass Co., by Professor Swezey.
 138. *Oxytropis multiceps* Nutt. Hills of upper Lawrence Fork, Kimball Co., Aug. 10; in fruit, (R.)
 139. *Astragalus hispidulus* Gray. Dakota Junction, May, 1891, (B.)
 140. (?) *Astragalus pubentissimus* Torr. and Gray. Canon in Gosper Co., June 20; Hills near Curtis, Frontier Co., June 22; Lawrence Fork, Banner Co., July 8; near Kimball, Aug. 12, 1891, (R.)
 141. *Amorpha microphylla* Pursh. Fragments covered with rust seen in the possession of Mr. Schofield, and said to have been collected near Lincoln, Sept. 1891, (R.)

POLEMONIACEÆ.

142. *Gilia* sp—. Glandular pubescent; root biennial, possibly perennial, stem branched above, 1 foot high or higher; leaves somewhat irregularly pinnatifid; lobes linear, not wider than the rachis, somewhat fleshy, mucronate; flowers in a branched panicle; corolla violet or blue with a whitish tube, somewhat funnel-form 2 lines long; stamens exserted; calyx with prominent ribs. The plant is more or less glandular all over. Mr. John Holzinger of the Department of Agriculture has named it *Gilia pinnatifida*, which without any doubt, it is not, as the plant is more branched, more glandular, the division of the leaves few and longer and narrower, and the corolla smaller and not at all salverform. It stands nearer *G. inconspicua*, from which (if distinct) it differs in being more robust,

more branched, and, perhaps, more glandular. Sands of North Platte near Horse Creek, Aug. 1, 1891, (R.)

143. *Gilia spicata* Nutt. Banner Co., (R.)

144. *Phlox bryoides* Nutt. Hills of upper Lawrence Fork, Aug. 11, 1891, (R.)

BORRAGINACEÆ.

145. (?) *Oreocarya sericea* (Gray) Greene. (*Pittonia* I. p. 58.) In fruit only. McColligan Canon, Deuel Co., June 26; upper Lawrence Fork, Kimball Co., Aug. 10, 1891, (R.)

By an oversight an error was made on page 37, under *Cryptantha*, *Krynnitzkia glomerata* Gray. is changed to *C. glomerata* Lehmann. It should have been given as *Oreocarya glomerata* (Pursh) Greene. (*Pittonia*, I. p. 58.) (R.)

146. *Allocarya californica* (Fisch. and Meyer.) Greene. *Krynnitzkia californica* (Fisch. and Meyer.) Gray Reported from Dawes County by Professor Swezey.

SOLANACEÆ.

147. *Physalis mollis* Nutt. var. *cinerascens* Gray. The leaves are broadly ovate, 2 in. long, angulately toothed or repand, very thin; pubescence stellate or simple (even a little glandular,) very short, except on the calyx, the pedicels, the petioles and younger parts of the stem, where it is long and soft; petioles as long as the leaves or longer; corolla greenish-yellow with a darker spot; anthers yellow; fruiting calyx globose-ovate. Under the cliffs on the south side of Scott's Bluff, July 20, 1891, (R.)

148. *Physalis* sp. (near *P. hederæfolia* Gray.) Leaves scarcely over 1 in. long, ovate, thickish; petioles generally shorter than the blades; fruiting calyx globose, about 1 in.; pubescence short, with a few long hairs, a little glandular. According to Mr. Holzinger it stands nearest *P. hederæfolia*, but it differs from the description of that species in Gray's Synoptical Flora, in that the pedicels are much longer, the leaves, larger and less toothed, and that it is very little glandular. It approaches *P. fendleri*. Prairies near Ashford, Banner Co., Aug. 6, (R.)

149. (?) *Physalis palmeri* Gray Leaves thickish, ovate, angulately toothed; petioles of the length of the blade which is 1-1½ in.; fruiting calyx ovate, (about an inch) on a slender pedicel. Stem low and nearly prostrate, strong scented; pubescence viscid, with short hairs. Sandy banks of a draw southwest of Sidney, Aug. 18. *P. palmeri* has been collected in southeast California, and this is entirely out of its supposed range. It may, therefore, be something else, (R.)

150. *Physalis longifolia* Nutt. (*P. lanceolata* Michx. var. *levigata* Gray.) I believe that this has a very good right to be separated from *P. lanceolata*, as the berry is yellow, stalked in the calyx which is ovate, less angled and not sunken at the base; the bottom of the calyx, stipe, and lower part of the berry glutinous, as if it were painted with molasses. Lawrence Fork, Banner Co., July 8; Kimball, Aug. 12, 1891, (R.)

151. *Physalis viscosa* L. var. *spathulæfolia* Gray. Low, 6-10 in. high, prostrate or nearly upright; pubescence short and stiff, or more

commonly, scarcely any except on the calyx, where it is hispid; leaves obovate or spatulate, tapering into the petiole, nearly entire; fruiting calyx globose-ovate, scarcely angled, very little sunken at the base; berry yellowish green, corolla $\frac{1}{2}$ in. wide, greenish-yellow with a darker spot. In *P. lanceolata* it is $\frac{2}{3}$ $\frac{3}{4}$ in. wide and ochroleucous. Sand draws of Western Nebraska, June and Aug. This is the same as *P. lanceolata*, No. 2 on page 38 preceding, (R.)

SCROPHULARIACEÆ.

- 152. *Seymeria macrophylla* Nutt. Reported by Professor Swezey from Cass Co.
- 153. *Gratiola virginiana* L. Kennedy, July 1891, (B.)
- 154. *Pentstemon acuminatus* Dougl. Deuel Co., (R.)
- 155. *Linaria canadensis* Dumont. Ewing, June 1891. Collected by Mr. Bates, (Bessey.)

ASCLEPIADACEÆ.

- 156. *Asclepias stenophylla* (Gray). In the April (1892) number of the *Botanical Gazette* J. M. Holzinger of the National Herbarium shows that what have been taken to be two distinct species viz: *Asclepias stenophylla* Gray. (No. 1677 of the catalogue), and *Acerates auriculata* Engelm. (No. 383 of the Appendix) are in reality but one species of *Asclepias*. In a subsequent note (*Bot. Gaz.* May) Mr. Holzinger shows that Gray's name must be retained.

CAPRIFOLIACEÆ.

- 157. *Lonicera hirsuta* Eaton. Ft. Niobrara, Aug. 1891, (B.)

COMPOSITE.

- 158. *Cnicus pitcheri* Torr. The true *C. pitcheris* is common in the sand hill regions of Kearney Co., July 15; rare in Banner Co., July 6, (R.)
- 159. *Cnicus virginiana* Pursh. Minden. Anther tips blunish instead of "subulate." Collected by Dr. Hapeman, (W.)
- 160. *Crepis tectorum* L. In a lawn in Chadron, 1891, (B.)
- 161. *Gaillardia pulchella* Fong. Franklin Co., Sept. 1891, (R.)
- 162. *Actinella scaposa* Nutt. var. *linearis* Nutt. Franklin Co., (Sw.)
- 163. (?) *Chaenactis douglasii* Hook and Arn., var. *alpina* Gray. The material was too incomplete for identification. A low cespitose, tomentose plant with pinnately parted leaves, and scapiform stem 3-6 in. high, with a few heads. Upper Lawrence Fork, Kimball County Aug. 10, (R.)
- 164. (?) *Aster ascendens* Lindl. Lodge Pole Creek, Kimball, Aug. 12, 1891, (R.)
- 165. (?) *Aster fremontii* Gray. Horse Creek, Scott's Bluff Co., Aug. 1 1891, (R.)
- 166. *Aster tradescantii* L. Big Springs, Deuel Co., Aug. 24, 1891, (R.) (See No. 418 of Mr. Webber's "Appendix.")
- 167. *Bigelowia graveolens* Gr. var. *glabrata* Gray. Canon of Scott's Bluff Co., Aug. 6, (R.)
- 168. *Bigelowia howardii* Gray. Hills of Banner and Scott's Bluff counties, July-Aug. 1891, (R.)
- 169. *Haplopappus armerioides* Gray. Hills in Banner Co.; July 6, 1891, (R.)
- 170. *Liatris spicata* Willd. Platte River bottom, Scott's Bluff Co., Aug. 3, 1891, (R.)

GENERIC INDEX.

- Abronia*, 50.
Acerates, 41.
Acolium, 12.
Actinella, 53.
Æcidium, 14.
Æsculus, 32.
Agropyrum, 24, 49.
Allocarya, 52.
Amarantus, 4.
Ambrosia, 42.
Amorpha, 51.
Amplicarpæa, 31.
Antennaria, 42.
Anychia, 30.
Aphanochæte, 9.
Aphyllon, 39.
Arabis, 31.
Arachnion, 18.
Aralia, 33.
Arctostaphylos, 36.
Arcyria, 45.
Arenaria, 5, 29, 50.
Aristida, 25, 49.
Armillaria, 18.
Arnica, 41.
Arrhenatherum, 49.
Arthonia, 12.
Arthrodeshus, 10.
Asclepias, 53.
Ascochyta, 4.
Asprella, 24.
Aster, 6, 43, 53.
Astragalus, 5, 34, 51.
Atrichum, 21.
Atriplex, 4, 28.
Avena, 25.
Bacillus, 8.
Bacterium, 45.
Barbula, 20.
Belonium, 48.
Beggiatoa, 45.
Berula, 33.
Betula, 26.
Biatora, 12.
Bigelovia, 53.
Botrytis, 17.
Bovista, 18.
Brickellia, 44.
Bromus, 25.
Bryum, 21.
Buellia, 12.
Bulbochæte, 10.
Calamagrostis, 49.
Callitrichæ, 51.
Calochortus, 26.
Cardamine, 31.
Carex, 4, 23, 49.
Caulophyllum, 30.
Centunculus, 36.
Cerastium, 29.
Ceratophorum, 17.
Ceratophyllum, 17, 27.
Cercocarpus, 34.
Cercospora, 17.
Chænactis, 53.
Chætomium, 11.
Chætophora, 46.
Chara, 20, 49.
Cheilanthes, 21.
Chenopodium, 28.
Chondrioderma, 45.
Chrysopsis, 44.
Cichorium, 41.

Cicuta, 51.
Cladonia, 13.
Clematis, 50.
Cladophora, 46.
Clitocybe, 19.
Closterium, 46.
Cnicus, 6, 41, 53.
Cocconeis, 47.
Coleochæte, 11, 47.
Collema, 13.
Colletotrichum, 16.
Collybia, 19.
Comariche, 45.
Convolvulus, 36.
Coprinus, 49.
Corallorrhiza, 26.
Coreopsis, 42.
Cosmarium, 10, 46.
Cratægus, 51.
Crepis, 41, 53.
Cristatella, 31.
Cryptanthe, 37, 51.
Cuscuta, 36.
Cyathus, 18.
Cylindrospermum, 8.
Cymopterus, 33.
Cyperus, 24.
Cystopus, 4, 11.
Delphinium, 30.
Desmatodon, 20.
Desmidium, 9, 46.
Desmodium, 51.
Dicranella, 20.
Diplachne, 25.
Dodecatheon, 36.
Dothidia, 47.
Draba, 50.
Draparnaldia, 46.
Dulichium, 24.
Echinospermum, 5, 38.
Eclipta, 42.
Elatine, 5.
Elodea, 22.
Elymus, 24, 49.
Empusa, 10.
Enteridium, 8.
Entoloma, 19.
Epilobium, 34.
Epithemia, 47.
Equisetum, 49.
Erigeron, 6, 42.
Eriogonum, 4, 27.
Eriophorum, 24.
Erodium, 32.
Erysimum, 30.
Erysiphe, 11.
Erythræa, 40.
Euastrum, 10, 46.
Euphorbia, 32.
Eurotia, 28.
Exoascus, 15.
Festuca, 49.
Fimbristylis, 24.
Fragilaria, 47.
Franseria, 6, 42.
Fraxinus, 40.
Fritillaria, 26.
Frælichia, 29.
Gaillardia, 53.
Geaster, 18.
Gentiana, 6, 40.
Gerardia, 38.
Gilia, 5, 36, 51.

Glæosporium, 16.
Glæotrichia, 8.
Glyceria, 25.
Gomphionema, 10, 47.
Gratiola, 53.
Guepina, 20.
Gymnosporangium, 4.
Gymnostomum, 20.
Habenaria, 26.
Haplopappus, 43, 53.
Hebeloma, 49.
Helenium, 41.
Heleocharis, 24, 49.
Helianthus, 6, 42.
Helminthosporium, 17.
Hemijarcyria, 45.
Hemicarpha, 24.
Heteranthera, 26.
Hiatula, 49.
Hicoria, 27.
Hydnum, 19.
Hydrodictyon, 9.
Hypericum, 32, 50.
Hypholoma, 19.
Hypoxylon, 47.
Hypnum, 21.
Hysterographium, 12.
Iris, 26.
Irpea, 19.
Isactis, 46.
Isaria, 18.
Iva, 42.
Juncus, 4, 25, 50.
Koeleria, 49.
Kuhnia, 6.
Kryni'zkyia, 37.
Lachnea, 47.
Lactuca, 41.
Lathyrus, 34.
Lecanora, 13.
Lechea, 50.
Lecidea, 12.
Lentinus, 19.
Lepiota, 18.
Leptostroma, 16.
Lesquerella, 31.
Liatris, 53.
Limosella, 38.
Linaria, 53.
Linum, 32.
Lippia, 40.
Lithospermum, 37.
Loeflingia, 29.
Lonicera, 53.
Lupinus, 35.
Lychnis, 29.
Lycoperdon, 18, 48.
Lyngbya, 8, 46.
Lysimachia, 36.
Marsilia, 21.
Melampsora, 14.
Melanconium, 16.
Mentha, 39.
Mentzelia, 5, 51.
Mertensia, 5, 37.
Menyanthes, 40.
Micrasterias, 10.
Microcoleus, 45.
Mimulus, 38.
Mollugo, 30.
Monarda, 39.
Monotropa, 36.
Monolepis, 28.

- Mougeotia*, 47.
Muhlenbergia, 49.
Musenium, 51.
Naias, 22.
Nasturtium, 31, 50.
Navicula, 46.
Necriella, 12.
Netella, 20.
Nitzschia, 47.
Nostoc, 45.
Nothoscordium, 4.
Oedogonium, 10, 47.
Oenothera, 33.
Opegrapha, 12.
Opuntia, 33.
Oreocarya, 37, 52.
Oscillaria, 45.
Osmorrhiza, 51.
Oxytropis, 51.
Panicum, 25, 50.
Parmelia, 13.
Pastinaca, 33.
Pectis, 42.
Pediastrum, 9.
Pedicularis, 38.
Peltigera, 13.
Pentstemon, 5, 38, 53.
Peronospora, 11.
Perichaena, 45.
Petalostemon, 34.
Peucedanum, 51.
Peziza, 14.
Phacelia, 37.
Phialea, 48.
Phlox, 52.
Pholiota, 19.
Phragmidium, 4, 14.
Phyllachora, 12.
Phyllosticta, 15.
Physalis, 5, 38, 52.
Physalospora, 11.
Physaria, 30.
Physarum, 8.
Physcia, 13.
Physostegia, 39.
- Pithophora*, 9.
Placodium, 13.
Plantago, 40.
Plasmopara, 11.
Pleurosigma, 47.
Pleurotus, 19.
Plowrightia, 12.
Pluteus, 19.
Poa, 25, 49.
Podosporium, 4.
Polygonatum, 26.
Polygonum, 27, 50.
Polyporus, 19.
Polystictus, 49.
Polytrichum, 21.
Potamogeton, 22.
Potentilla, 34.
Protococcus, 46.
Psoralea, 35.
Pterula, 20.
Puccinia, 14, 48.
Pyrola, 36.
Quercus, 4.
Ramularia, 4, 17.
Ranunculus, 30.
Raphidium, 46.
Reticularia, 8.
Rhabdospora, 16.
Ribes, 33, 51.
Rosa, 51.
Rumex, 27.
Ruppia, 22.
Saccharomyces, 15.
Sagittaria, 22.
Salix, 27.
Salvia, 39.
Saponaria, 30.
Sclerospora, 11.
Sclerotinia, 47.
Scutellaria, 39.
Septoria, 16.
Seymeria, 53.
Silene, 30.
Sisyrinchium, 50.
Smilax, 26.
- Solidago*, 43.
Sorastrum, 9.
Sorosporium, 48.
Sphaerella, 12.
Sphaerozoma, 9.
Spiranthes, 26.
Spirogyra, 17.
Spirotaria, 9.
Sporobolus, 49.
Stachybotrys, 17.
Stachys, 39.
Stanleya, 30.
Staurastrum, 10, 16.
Stauroneis, 46.
Stegonospora, 1.
Stellaria, 29.
Stemonitis, 8, 15.
Stephanomeria, 41.
Stereum, 19, 49.
Stigeocladium, 9.
Suaeda, 28.
Synedra, 47.
Tetraspora, 9, 46.
Thelesperma, 42.
Thelypodium, 30.
Townsendia, 43.
Trichia, 15.
Tricholoma, 18.
Trifolium, 35.
Trillium, 50.
Typha, 23.
Urceolaria, 13.
Uredo, 15.
Urocystis, 15.
Uronyces, 14.
Ustilago, 15.
Verbena, 40.
Vermicularia, 16.
Vernonia, 44.
Verpa, 48.
Verticillium, 48.
Woodsia, 4.
Xanthidium, 10.
Xylaria, 47.
Zygnema, 47.
Zygadenus, 50.