

University of Nebraska - Lincoln

## DigitalCommons@University of Nebraska - Lincoln

---

Agronomy & Horticulture -- Faculty Publications

Agronomy and Horticulture Department

---

2004

### In-season Prediction of Attainable Maize Yield Using the Hybrid-Maize Model

Achim R. Dobermann

*University of Nebraska-Lincoln*, [adobermann2@unl.edu](mailto:adobermann2@unl.edu)

Haishun Yang

*University of Nebraska-Lincoln*, [hyang2@unl.edu](mailto:hyang2@unl.edu)

Kenneth G. Cassman

*University of Nebraska- Lincoln*, [kcassman1@unl.edu](mailto:kcassman1@unl.edu)

Daniel T. Walters

*University of Nebraska-Lincoln*, [dwalters1@unl.edu](mailto:dwalters1@unl.edu)

Follow this and additional works at: <https://digitalcommons.unl.edu/agronomyfacpub>



Part of the [Plant Sciences Commons](#)

---

Dobermann, Achim R.; Yang, Haishun; Cassman, Kenneth G.; and Walters, Daniel T., "In-season Prediction of Attainable Maize Yield Using the Hybrid-Maize Model" (2004). *Agronomy & Horticulture -- Faculty Publications*. 525.

<https://digitalcommons.unl.edu/agronomyfacpub/525>

This Article is brought to you for free and open access by the Agronomy and Horticulture Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agronomy & Horticulture -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

# In-season Prediction of Attainable Maize Yield Using the Hybrid-Maize Model

Achim Dobermann\*, Haishun Yang, Kenneth G. Cassman, and Daniel T. Walters



\*Contact: Achim Dobermann  
Dept. of Agronomy and Horticulture, University of Nebraska-Lincoln, USA  
adobermann2@unl.edu; Phone: +1-402-472-1501

## The Hybrid-Maize Model

**Hybrid-Maize** simulates the growth of a maize crop under non-limiting or water-limiting (rainfed or irrigated) conditions based on daily weather data. This new maize model was developed by combining the strengths of two modeling approaches (Yang et al., 2004): (i) growth and development functions in used in CERES-Maize (Jones and Kiniry, 1986), and (ii) mechanistic formulations of light interception, photosynthesis and respiration used in generic crop models such as INTERCOM or WOFOST (van Ittersum et al., 2003).

**Hybrid-Maize** features temperature-driven maize phenological development, vertical canopy integration of photosynthesis, organ-specific growth respiration, and temperature-sensitive maintenance respiration. The new model requires fewer genotype-specific parameters without sacrificing prediction accuracy. A linear relationship between growing degree-days (GDD) from emergence to silking and GDD from emergence to physiological maturity is used for prediction of day of silking. Field validation under high-yielding growth conditions indicated close agreement between simulated and measured values for leaf area, dry matter accumulation (Fig. 1), final grain and stover yields, and harvest index (Yang et al., 2004).





Figure 1. Observed and predicted aboveground dry matter accumulation of irrigated maize at Lincoln, Nebraska, 2001.

## Real-time Simulation and Yield Forecasting

**Hybrid-Maize** allows the user to (i) assess the overall site yield potential and its variability, (ii) evaluate changes in attainable yield using different choices of planting date, maize hybrid, and plant density, (iii) analyze maize growth in specific years, (iv) explore options for irrigation water management, and (v) conduct in-season simulations to evaluate actual growth and forecast final yield at different growth stages.


In *Current season prediction* mode, model simulations are based on the up-to-date weather data of the current growing season, supplemented by the previously collected historical weather data for forecasting of all possible outcomes for the remainder of the season. This results in a range of forecasted scenarios, which are ranked based on the predicted final yield (Fig. 2). Management decisions could include adjusting the yield goal in comparison with normal years and making subsequent adjustments in fertilizer amounts and irrigation. During grain filling, yield forecasting can provide additional information to help guide marketing decisions on marketing.



Rank	GrainY	GrainDM	StoverDM	TotalDM	HI	vDays	rDays	V+R	tSola	Tmin	Tmax	Tmean	vTmean	rTmean	ETref	tRain
Best yield	21.5	18.2	12.6	30.7	0.59	65	82	147	2955	14	27	21	22	20	5.4	474
75% percentile	19.0	16.1	12.4	28.5	0.56	65	68	133	2728	16	28	22	22	22	5.6	442
Median yield	17.4	14.7	14.0	28.6	0.51	65	57	122	2694	17	30	23	22	25	6.5	283
25% percentile	15.7	13.3	14.7	28.0	0.47	65	54	119	2552	17	30	23	22	25	6.1	272
Worst yield	15.0	12.7	12.6	25.2	0.50	65	55	120	2437	17	29	23	22	25	5.8	263
Long-term median	16.7	14.1	12.5	26.6	0.53	59	57	116	2558	18	30	24	23	24	6	350


Figure 2. Hybrid-Maize model user interface and examples of model outputs for a real-time simulation and yield forecast made on July 22, 2003 at Lincoln, Nebraska.

## Case Study 1: Irrigated Maize, Lincoln, Nebraska




Maize was grown near yield potential levels, i.e., with full irrigation, optimal nutrient supply, and at a density of 8.6 plants m<sup>-2</sup>. Actual and historical daily weather data were used for yield forecasting in intervals of 5 days. Early in the season, yield forecasts mainly relied on historical weather data so that the median predicted yield was close to the long-term median. As the season progressed, more actual weather data were used, indicating above-normal growth conditions. Shortly after silking, predicted median yield approached the final measured grain yield of 17.9 Mg ha<sup>-1</sup>. With progressing grain filling, the range of predicted yields gradually decreased.

## Case Study 2: Rainfed Maize, Oliveros, Argentina



Rainfed maize was grown at 7 plants m<sup>-2</sup> in a favorable season with no water stress. Shortly after silking, predicted median yield approached the final measured grain yield of 14.4 Mg ha<sup>-1</sup>. With progressing grain filling, the range of predicted yields decreased.

## Case Study 3: Rainfed Maize, Mead, Nebraska



Rainfed maize was grown at 5.9 plants m<sup>-2</sup>. Early in the season, median predicted yield was close to the long-term median. Little rain fell in July and August. As the season progressed, drought evolved and the predicted median yield decreased well below the long-term median, approaching the final measured grain yield of 8.0 Mg ha<sup>-1</sup>. Median predictions were close to the final yield about one month before maturity.

## Conclusions

Hybrid-Maize offers promising potential for in-season simulation and forecasting of maize biomass and grain yield. At sites with no water limitations, the final yield was accurately predicted shortly after silking stage. Greater variation in yield predictions may occur at rainfed sites under drought stress, but relative deviations from normal growth were detected early enough in both environments.

## References

Jones, C.A., and J.R. Kiniry. 1986. CERES-Maize: A simulation model of maize growth and development. Texas A&M Univ. Press, College Station, TX.  
van Ittersum, M.K., P.A. Leffelaar, H. van Keulen, M.J. Kropff, L. Bastiaans, and J. Goudriaan. 2003. On approaches an applications of the Wageningen crop models. Eur. J. Agron. 18:201-234.  
Yang, H.S., A. Dobermann, J.L. Lindquist, D.T. Walters, T.J. Arkebauer, and K.G. Cassman. 2004b. Hybrid-Maize - a maize simulation model that combines two crop modeling approaches. Field Crops Res. 87:131-154.

**Acknowledgements:** Financial supported for developing Hybrid-Maize was provided by the Agricultural Research Division of the University of Nebraska-Lincoln, the Potash and Phosphate Institute through the Foundation for Agronomic Research, the Fluid Fertilizer Foundation, the Nebraska Corn Board, and by USDA-CSREES through the Consortium for Agricultural Soil Mitigation of Greenhouse Gases.