

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Insecta Mundi

Center for Systematic Entomology, Gainesville,  
Florida

---

6-10-2013

**Description of a new *Crossidius* LeConte (Coleoptera:  
Cerambycidae: Cerambycinae: Trachyderini) from southern  
Georgia with comments on its biology and unusual distribution**

Roy F. Morris II

*The Coleopterists Society*, beetlesandbirds@gmail.com

James E. Wappes

*American Coleoptera Museum, San Antonio, TX*, wappes@earthlink.net

Follow this and additional works at: <https://digitalcommons.unl.edu/insectamundi>

---

Morris, Roy F. II and Wappes, James E., "Description of a new *Crossidius* LeConte (Coleoptera: Cerambycidae: Cerambycinae: Trachyderini) from southern Georgia with comments on its biology and unusual distribution" (2013). *Insecta Mundi*. 809.

<https://digitalcommons.unl.edu/insectamundi/809>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

# INSECTA MUNDI

A Journal of World Insect Systematics

---

0304

Description of a new *Crossidius* LeConte (Coleoptera: Cerambycidae: Cerambycinae: Trachyderini) from southern Georgia with comments on its biology and unusual distribution

Roy F. Morris, II  
2635 Ewell Road  
Lakeland, Florida 33811

James E. Wappes  
8734 Paisano Pass  
San Antonio, TX 78255

Date of Issue: June 10, 2013

Roy F. Morris, II and James E. Wappes  
Description of a new *Crossidius* LeConte (Coleoptera: Cerambycidae: Cerambycinae:  
Trachyderini) from southern Georgia with comments on its biology and unusual  
distribution  
Insecta Mundi 0304: 1-7

ZooBank Registered: urn:lsid:zoobank.org:pub:3CAD53B1-347F-4B6F-B83B-5974C7739057

**Published in 2013 by**

Center for Systematic Entomology, Inc.  
P. O. Box 141874  
Gainesville, FL 32614-1874 USA  
<http://www.centerforsystematicentomology.org/>

**Insecta Mundi** is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

**Insecta Mundi** is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology. Manuscript preparation guidelines are available at the CSE website.

**Managing editor:** Paul E. Skelley, e-mail: [insectamundi@gmail.com](mailto:insectamundi@gmail.com)

**Production editor:** Michael C. Thomas, Brian Armitage, Ian Stocks

**Editorial board:** J. H. Frank, M. J. Paulsen

**Subject editors:** G.B. Edwards, J. Eger, A. Rasmussen, F. Shockley, G. Steck, Ian Stocks, A. Van Pelt, J. Zaspel

**Spanish editors:** Julieta Brambila, Angélico Asenjo

**Printed copies (ISSN 0749-6737) annually deposited in libraries:**

CSIRO, Canberra, ACT, Australia  
Museu de Zoologia, São Paulo, Brazil  
Agriculture and Agrifood Canada, Ottawa, ON, Canada  
The Natural History Museum, London, Great Britain  
Muzeum i Instytut Zoologiczny PAN, Warsaw, Poland  
National Taiwan University, Taipei, Taiwan  
California Academy of Sciences, San Francisco, CA, USA  
Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA  
Field Museum of Natural History, Chicago, IL, USA  
National Museum of Natural History, Smithsonian Institution, Washington, DC, USA  
Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

**Electronic copies (On-Line ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:**

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.  
Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>  
University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>  
Goethe-Universität, Frankfurt am Main: <http://edocs.ub.uni-frankfurt.de/volltexte/2010/14363/>

**Author instructions** available on the Insecta Mundi page at:

<http://www.centerforsystematicentomology.org/insectamundi/>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

Description of a new *Crossidius* LeConte (Coleoptera: Cerambycidae: Cerambycinae: Trachyderini) from southern Georgia with comments on its biology and unusual distribution

Roy F. Morris, II  
2635 Ewell Road  
Lakeland, Florida 33811  
beetlesandbirds@gmail.com

James E. Wappes  
8734 Paisano Pass  
San Antonio, TX 78255  
wappes@earthlink.net

**Abstract.** *Crossidius grahami* Morris and Wappes **new species** (Coleoptera: Cerambycidae) is described from the Ochoopee Dunes of southern Georgia. Comments are included on the new species' biology and disjunct distribution compared to other species of *Crossidius* LeConte. Illustrations include: dorsal and lateral views of the primary types of *C. grahami*, its natural habitat, the known host plant (*Chrysoma pauciflosculosa* (Michx.) Greene (Asteraceae)), pupal chamber, and dorsal views of both sexes of *Crossidius humeralis quadrivittata* Penrose, 1974, considered its closest anatomical counterpart and nearest geographical relative.

**Key words.** Coleoptera, Cerambycidae, new species, Georgia, Ochoopee dunes, *Chrysoma pauciflosculosa*.

**Resumen.** Se describe *Crossidius grahami* Morris and Wappes, **especie nueva** (Coleoptera: Cerambycidae) de la Dunas Ochoopee del sur de Georgia. Se incluyen comentarios sobre la biología de la nueva especie y sobre su distribución dispersa en comparación con las otras especies de *Crossidius* LeConte.

**Palabras Clave.** Coleóptera, Cerambycidae, especie nueva, Georgia, dunas de Ochoopee, *Chrysoma pauciflosculosa*.


## Introduction

Collecting by the senior author in southern Georgia, in the inland sand dune system known as the Ochoopee Dunes, has led to the discovery of a new species of *Crossidius* LeConte (Cerambycidae). The occurrence of this species in the southeastern United States is surprising since this large trachyderine genus, containing 13 species and 37 subspecies, was previously only recorded from the western States (California to Texas) and Mexico (Monné and Bezark 2012). Adult *Crossidius* are moderate sized (8-22 mm), lightly to brightly colored and typically found on the blossoms of their host plant, yellow flowering composites (Linsley and Chemsak 1961). They are a favorite of collectors because of this, for typically being found in large numbers, and for their habit of sitting fully exposed in the sun easily available for handpicking. The fact that *Crossidius grahami* was undiscovered until recently is a mystery. The differences in its habits and biology may help to explain why and are discussed later in this paper (see: "Biology").

*Crossidius grahami* has a disjunct distribution, hundreds of mile east of other *Crossidius* species with *C. humeralis quadrivittata* Penrose, found along the southern gulf coast of Texas, the nearest species geographically. The nominate form of this relative is where the new species will key in the most recent key to *Crossidius* species (Linsley 1962).

**Materials.** Specimens studied are deposited in the following collections:

- ACMT** – American Coleoptera Museum (James E. Wappes), San Antonio, TX, USA
- EMEC** – Essig Museum of Entomology (University of California), Berkley, CA, USA
- FSCA** – Florida State Collection of Arthropods, Gainesville, FL, USA



**Figures 1-4.** *Crossidius* spp. **1)** *Crossidius grahami*, holotype male, a) dorsal, b) lateral. **2)** *Crossidius grahami*, allotype female Emanuel Co., a) dorsal, b) lateral. **3)** *Crossidius grahami*, paratype female, Tattnall Co. **4)** *Crossidius humeralis quadrivittata*, Texas, Kleberg Co., a) male, b) female.

**JBMC** – Jeff Burne, Macon, GA, USA


**JHGC** – Jeffrey Huether, Geneva, NY, USA

**JPBC** – Joshua P. Basham, Tennessee State University, McMinnville, TN, USA

**MEMC** – Mississippi Entomological Museum, Mississippi State, MS, USA

**MNRJ** – Museu Nacional, Universidade Federal do Rio de Janeiro, RJ, BRASIL

**MZSP** – Museu de Zoologia da Universidade de São Paulo, SP, BRASIL



**Figures 5-6.** Head, pronotum and basal antennomeres. **5)** *Crossidius grahami*, holotype male. **6)** *Crossidius humeralis quadrivittata*, Texas, Kleberg Co., male.

**RAAC** – Robert A. Androw, Pittsburg, PA, USA

**RFMC** – Roy F. Morris, Lakeland, FL, USA

**RHTC** – Robert H. Turnbow Jr., Enterprise, AL, USA

**RMBC** – R. Michael Brattain, Lafayette, IN, USA

**UGAC** – University of Georgia Arthropod Collection, Athens, GA, USA

**USNM** – National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

### *Crossidius* LeConte

*Crossidius* LeConte 1851: 102. Type species: *Crossidius testaceus* LeConte 1851: 102, by monotypy. Linsley and Chemsak 1961: 26-64 (distributional and taxonomic study of the genus); Linsley 1962: 142-175 (revision); Penrose 1974: 248-254 (new subspecies of *Crossidius humeralis* LeConte).

### *Crossidius grahami* Morris and Wappes, new species

(Figures 1-3, 5, 9-10)

**Description. Male** (Fig. 1): Moderate sized, elongate, parallel-sided, robust; integument entirely brownish-testaceous, pronotum with three dark-brown to black discal spots, elytra entirely testaceous. **Head** with pale pubescence, sparse, not obscuring surface, denser on front of head and along basal margin; underlying punctures dense, almost contiguous, smaller than those on elytra. **Antennae** 11 segmented, short, reaching apical quarter of elytra but not surpassing apices; scape twice as long as broad; third segment short, stout, as long as the scape and second segment combined, slightly expanded at apex (Fig. 5); antennomeres 4-10 flattened with length subequal to scape; antennal punctures and pale pubescence sparse. **Pronotum** cylindrical, swollen, almost parallel sided, basally and apically margined; sides rounded, lacking tubercles; surface punctures dense, larger than those at base of elytra; pale pubescence moderate, not obscuring surface; discal area with two punctate, glabrous submedial dark areas and a less punctate similar area basally. **Prosternum** swollen at sides, densely pubescent across middle. **Metasternum** finely punctate, densely pubescent. **Scutellum** dark, flat, triangular, not indented medially, apex slightly rounded; pubescence fine. **Elytra** length 2.4 times width of both elytra at the humeri; surface moderately coarsely, densely punctate, becoming denser apically; pubescence short, pale, sparse, not obscuring surface; apices slightly rounded, angles usually distinct but not acute. **Legs** with fine punctures, pubescence sparse, not obscuring surface; mesotibial spurs short, nearly subequal in length. **Abdomen** with dense, pale pubescence, almost obscuring the fine surface punctures. Length 10-15 mm.

**Female** (Fig 2): Form similar to male except as noted. Body testaceous with head, antennomeres, elytra and legs black, uniformly covered with short pale pubescence. **Head** punctate, sparsely pubescent.

Antenna 11 segmented, reaching middle of elytra, sparsely pubescent, punctuate with punctures small and shallow. **Pronotum** not inflated (as in male); pubescence pale, sparse; punctures larger than those of elytral base. **Elytra** with apical third slightly expanded, pubescence short, sparse and erect. **Legs** short, with sparse pubescence. **Abdomen** and metasternum with fine, pale, suberect pubescence not obscuring surface. Length 12-15 mm.

**Variation.** Females from Tattnall County, GA tend to have the basal third of the elytra tan (Fig. 3) while those from Emanuel County have solid black elytra (Fig. 2).

**Diagnosis.** *Crossidius grahami* is the only *Crossidius* species known from east of the Mississippi River and to date has only been found in the relict sand formations of southern Georgia. As such its distribution is diagnostic. It is also morphologically unique for *Crossidius* species in having the antenna distinctly shorter than the elytra in both sexes. From its nearest geographical relative, *C. humeralis quadrivittata* Penrose (Fig. 4), *C. grahami* can be separated by the following characters: shorter, more robust and apically expanded antennomeres (Fig. 5); shorter, nearly subequal metatibial spurs; greatly reduced pubescence on the sides of the pronotum, and on the vertex of the head and between the eyes; and by the lack of elytral vitta. In *C. h. quadrivittata*, antennomeres are longer and nearly parallel-sided (Fig. 6); inner metatibial spur distinctly longer than outer; long, dense pubescence on the sides of the pronotum, and on the vertex and between the eyes, and distinct, dark elytral vitta in most specimens.

**Type Material.** Male holotype of *Crossidius grahami* labeled: “/ GA: Emanuel co., 1.5 mi. S. Covena, 10/VI/00, R. Morris / Sweeping *Chrysoma pauciflosculosa* ”. The female allotype is labeled: “/ GA: Emanuel co., 1.5 mi. s. Covena, 6/VI/99, R. Morris”. They are deposited in the FSCA.

**Paratypes:** 257 males and 84 females (Unless otherwise noted the following paratypes are deposited in the collection of the collector). **Georgia:** Same data as holotype (13m, 6f – RFMC, 4m, 2f -UGAC; 1m – RMBC); **Emanuel Co.:** 1.5 mi. S. Covena, 6/VI/99, R. Morris (9m, 1f – RFMC; 1f – RMBC); 1 mi. SW Covena, 11 June 2000, R. Turnbow, on *Chrysoma pauciflosculosa* (Michx.) Greene (6m, 2f-USNM); 1 mi. SW Covena, 10 June 2000, R. Turnbow, on *Chrysoma pauciflosculosa* (Michx.) Greene (17m, 9f - RHTC); 1 mi. SW Covena, 11 June 2006, R. Turnbow, on *Chrysoma pauciflosculosa* (Michx.) Greene (17m, 5f - RHTC); 1 mi. SW Covena, 11 June 2011, R. Turnbow, on *Chrysoma pauciflosculosa* (Michx.) Greene (20m, 9f – RHTC, 4m, 2f EMEC); Oohoopee Dunes WMA, Co. Rd. 160, 12/VI/1999, R. Morris, sweeping *Chrysoma pauciflosculosa* (6m – RFMC); Oohoopee Dunes Natural Area, Hall’s Bridge Rd., 10/VI/00, R. Morris, sweeping *Chrysoma pauciflosculosa* (7m, 1f – RFMC); Oohoopee Dunes Natural Area, 10 June 2000, R. Turnbow, on *Chrysoma pauciflosculosa* (Michx.) Greene (5m, 2f – MNRJ, 5m 2f - MZSP); Oohoopee Dunes Natural Area, 11 June 2011, R. Turnbow (2m - RHTC); Oohoopee Dunes Natural Area, 12 June 2011, R. Turnbow (2m, 1f - RHTC); Oohoopee Dunes Natural Area, 4.5 km NE Norristown, Halls Bridge Tract, 32.53581, -82.46023, 8 June 2011, beaten from foliage of woody goldenrod, *Chrysoma pauciflosculosa*, R. A. Androw, coll. (3m - RAAC); Oohoopee Dunes Natural Area, 32°32’15”N 82°27’40”W, 19 June 2002, T. L. Schiefer, R. L. Brown, or J.G. Hill, sweeping *Chrysoma pauciflosculosa* in xeric dune scrub. W. H. Cross Expedition (9m, 1f – MEMC); Hwy. 56 at Oohoopee River, 10 June 2000, R. Turnbow (1m – RHTC); I-16 & US 1, Gar Rd., 6-7/VI/99, R. Morris (18m, 4f – RFMC); I-16 & US 1, Gar Rd., 12/VI/99, R. Morris, sweeping *Chrysoma pauciflosculosa* (5m – RFMC); I-16 & US 1, Gar Rd., 9/VI/00, R. Morris, sweeping *Chrysoma pauciflosculosa* (1m – RFMC); I-16 & US 1, Gar Rd., 15/VI/01, R. Morris (1m – RFMC); I-16 & US 1, Gar Rd., 30/V/10, J. Basham, cut fm *Chrysoma pauciflosculosa* root crown 14/I/10 (1f –JPBC); Jct. Rts. 1 & 16, VI/15-16/03, J. &M. Huether (8m, 2f – JHGC); Jct. I-16 & hwy. 1, 9 June 2000, R. Turnbow, on *Chrysoma pauciflosculosa* (Michx.) Greene (9m, 1f – RHTC); **Tattnall Co.:** 3 mi. E of 147 along Oohoopee River, Col. 1-IV-98, Em. 1-VI-98, R. Morris, ex *Chrysoma pauciflosculosa* (1m – RFMC); 3 mi. E of 147 along Oohoopee River, 12-13-VI-1998, R. & G. Morris, sweeping *Chrysoma pauciflosculosa* (18m, 5f – RFMC); 3 mi. E of 147 along Oohoopee River, 12-VI-1998, Roy & Graham Morris (1m – RFMC); 3 mi. E of 147 along Oohoopee River, 20-VI-1998, Morris/Wappes, sweeping *Chrysoma pauciflosculosa* (5m, 1f – RFMC); 3 mi. E of 147 along Oohoopee River, 19-20 June 1998, Wappes & Morris, sweeping woody goldenrod, *Chrysoma pauciflosculosa* Grn. (5m, 3f -ACMT); 2 mi. E of 147 along Oohoopee River, 11/VI/99, R. Morris, sweeping *Chrysoma pauciflosculosa* (1f – RFMC); 2 mi. E of 147 along Oohoopee River, 30-V-04, Morris/Nearns/Skelley (6m – RFMC); 2 mi. E of 147 along Oohoopee River, 5-VI-99, R. Morris



**Figures 7-10.** Habitat and host pictures. **7)** Typical habitat of *C. grahami* with its host plant, woody goldenrod, front center. **8)** *Chrysoma pauciflosculosa* (Michx.) in bloom during October. **9)** Root crown of woody goldenrod, showing pupal chamber and the teneral male reared from it with label data (Photos by J. P. Basham). **10)** *Crossidius grahami* sitting on the main stem of the host plant (Photo by M. C. Thomas).

(5m, 4f – RFMC); 2 mi. E of 147 along Ochoopee River, 5-VI-99, J. Burne (6m, 2f – JBMC); 2 mi. E of 147 along Ochoopee River, 9-VI-99, J. Burne (15m, 4f – JBMC); 2 mi. E of 147 along Ochoopee River, 12-VI-99, J. Burne (9m, 1f – JBMC); 2 mi. E of 147 along Ochoopee River, 24-VI-99, J. Burne (3m, 3f – FSCA); 3 mi. S. of Reidsville, 22/VI/02, R. Morris (4m, 2 f – RFMC); 4 mi. S. Reidsville, 28 June, 1998, R. Turnbow, on *Chrysoma pauciflosculosa* (Michx.) Greene (3m - RHTC); 4 mi. S. Reidsville, 13 June, 2011, R. Turnbow, on *Chrysoma pauciflosculosa* (Michx.) Greene (8m, 2f- FSCA); 4 mi. S. Reidsville, 12-VI-2011, M. C. Thomas, sweeping *Chrysoma pauciflosculosa* (2m, 2f-FSCA); 1 mi. S. Covina, 11-VI-2011, M. C. Thomas, sweeping *Chrysoma pauciflosculosa* (3m, 2f-FSCA).

**Etymology.** It is our pleasure to name this new species for Graham Morris, who as an eight year old accompanied his father to southern Georgia and helped collect the first specimens of this new species.

**Biology.** The known distribution of *Crossidius grahami* is restricted to the relict island sand dune systems along the Ochoopee and Little Ochoopee rivers in Emanuel and Tattnall counties of Georgia. These dunes were formed during the late Pliocene glacial or the intermediate post glacial period (Markewich and Markewich 1994), with the natural vegetation (Fig. 7) upland scrub with smaller areas of dwarf oak forests (Wharton 1978). The only known host plant of *C. grahami* is woody goldenrod,



*Chrysoma pauciflosculosa* (Michx.) Greene (Asteraceae) (Fig. 8). During April, 1998 the senior author found a dying woody goldenrod, 3 miles east of GA Highway 147, along the Ohoopee River. Curious about the cause of the plant's demise he excavated and examined the root crown. After finding some larval working he cut it open and found the larvae of a cerambycid. Securing the pieces together it was placed in a rearing container on his desk at home. On June 1, a male of an unknown cerambycid emerged. Not knowing the identity of this specimen, a trip with his son Graham was made to the area the next week to look for more specimens. There, they successfully collected additional specimens by sweeping *Chrysoma*. Recently (January, 2012), Joshua Basham, McMinnville, TN, also confirmed woody goldenrod as a host plant by extracting a teneral male from its root crown pupal chamber (Fig. 9). Most of the paratypical specimens have been collected by sweeping woody goldenrod or by handpicking, as they sit facing upward on the main stem (Fig. 10). Only rarely have they been observed and taken in flight. Woody goldenrod occurs primarily in coastal dunes, and among scrub oak ridges across the southern US, from Mississippi to the inner coastal plain of southern Georgia and all the way into North Carolina wherever suitable habitat is found (Godfrey 1988). Extensive collecting by the senior author in other areas of Georgia and north Florida, where the host plant occurs, has not yielded any specimens of *C. grahami*. We can only hypothesize that some biological factor, other than just having available host plant, is key to the species development and hence restricting its distribution.

Most *Crossidius* host plants bloom in the late summer and early fall, with adults of *Crossidius* species typically emerging at the start of the host blooming season. During the day both sexes can be found feeding on blossoms. Males also commonly perch with antenna extended, on the tallest flowers of the host, hoping to locate freshly emerged females. During cool or overcast days adults of both sexes can be found sitting quietly among the foliage and stems of the host plant waiting for warmer temperatures to then become active. It is assumed that most adults spend the night at the base of the host as we have observed adults crawling down the stems in the evenings and, as morning sun warms the air, crawling back up the plant and becoming active by mid-morning. Rarely, but especially on warm nights, adults of some smaller species can be found still flower sitting or in the upper foliage of the host. In one case we observed males of *C. pulchellus* LeConte, in West Texas, sitting on *Gutierrezia* flowers early in the morning covered by snow from a previous night's snowfall. *Chrysoma pauciflosculosa* blooms in late October, somewhat later than western *Crossidius* hosts. Unlike the western species, adults of *Crossidius grahami* are normally late spring active with almost all collected specimens taken in late May or June. Obviously, this is not when the host, woody goldenrod, blooms. Hence, adults are not flower feeders and are not feeding at the top of the plant, exposed, to be easily found by the collector or observer. Rather, they sit on the main stem of the host (Fig. 10), or hide in the dead leaves that are still attached to the terminal stem. This makes them much more difficult to find and experience indicates they are best collected by sweeping the plant. Collecting, during the fall bloom season, at localities known to have populations of *C. grahami*, has not produced any specimens. This aberrant behavior may help explain why the species has escaped discovery until recent years.

### Acknowledgments

Special thanks are due Paul Skelley, Florida State Collection of Arthropods, for his technical assistance, help with many of the photographs, reviewing an early draft of this paper, sharing his expertise as an Editor and, most importantly, for providing guidance on numerous issues related to writing this paper. The final product is much improved because of his kind assistance. We thank Joshua P. Basham, McMinnville, TN, for allowing us to use host plant material from which he extracted a specimen of the new species. We also thank our colleagues Steven Lingafelter, USNM, Miguel Monné, MNRJ and Donald Thomas, Weslaco, TX, for reviewing the manuscript. Thanks to Julieta Brambila, FSCA, for doing the Spanish version of the Abstract and Keywords. We also appreciate Steven Christman, Editor, [www.floridata.com](http://www.floridata.com), for providing the photograph that appears as Figure 8. Finally, we appreciate and thank all the collectors and collection managers that provided their specimens so they could be included in this paper.

**Literature Cited**

- Godfrey, R. K. 1998.** Trees, shrubs and woody vines of northern Florida and adjacent Georgia and Alabama. The University of Georgia Press; Athens. 734p.
- LeConte, J. L. 1851.** An attempt to classify the longicorn Coleoptera of the part of America, north of Mexico. Journal of the Academy of Natural Sciences of Philadelphia, (2)2: 99-112. Philadelphia, PA
- Linsley, E. G. 1962.** The Cerambycidae of North America. Part III. Taxonomy and Classification of the Subfamily Cerambycinae, Tribes Opsimini through Megaderini. University of California Publications in Entomology 20: 1-188.
- Linsley, E. G. and J. A. Chemsak. 1961.** A Distributional and Taxonomic Study of the Genus *Crossidius* (Coleoptera, Cerambycidae). Miscellaneous Publications of the Entomological Society of America 3(2): 26-64.
- Markewich, H. W. and W. Markewich. 1994.** An overview of Pleistocene and Holocene inland dunes in Georgia and the Carolinas – Morphology, distribution, age and paleoclimate. U. S. Geological Survey Bulletin 2069: 1-32.
- Monné, M. A. and L. G. Bezark, 2013.** Checklist of the Cerambycidae of the Western Hemisphere (updated through 31 December, 2012). Available at <http://plant.cdfa.ca.gov/byciddb/checklists/WestHemiCerambycidae2012-12-31.pdf>. Last accessed 10 April, 2013.
- Penrose, R. L. 1974.** A new subspecies of *Crossidius humeralis* LeConte from Texas with a redescription of the species (Coleoptera: Cerambycidae). The Pan-Pacific Entomologist 50: 248-254.
- Wharton, C. H. 1978.** The natural environments of Georgia. Georgia Department of Natural Resources, Bulletin 114: 1-227. [reprinted 1989].

**Received April 29, 2013; Accepted May 2, 2013.**

