

University of Nebraska - Lincoln
DigitalCommons@University of Nebraska - Lincoln

Roman L. Hruska U.S. Meat Animal Research
Center

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

2013

PHYSIOLOGY AND ENDOCRINOLOGY SYMPOSIUM: The current status of heat shock in early embryonic survival and reproductive effi ciency

R. A. Cushman

USDA-ARS, Bob.Cushman@ars.usda.gov

Follow this and additional works at: <http://digitalcommons.unl.edu/hruskareports>

Cushman, R. A., "PHYSIOLOGY AND ENDOCRINOLOGY SYMPOSIUM: The current status of heat shock in early embryonic survival and reproductive efficiency" (2013). *Roman L. Hruska U.S. Meat Animal Research Center*. 275.
<http://digitalcommons.unl.edu/hruskareports/275>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Roman L. Hruska U.S. Meat Animal Research Center by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

PHYSIOLOGY AND ENDOCRINOLOGY SYMPOSIUM: The current status of heat shock in early embryonic survival and reproductive efficiency^{1,2}

R. A. Cushman³

USDA⁴–ARS, Roman L. Hruska U.S. Meat Animal Research Center, Clay Center, NE 68933

© 2013 American Society of Animal Science. All rights reserved.

J. Anim. Sci. 2013.91:1141–1142

doi:10.2527/jas2013-6231

The Physiology and Endocrinology Symposium titled “The Current Status of Heat Shock in Early Embryonic Survival and Reproductive Efficiency” was held at the Joint Annual Meeting in Phoenix, AZ, July 15 to 19, 2012. In recent years, data has accumulated indicating a role for heat shock proteins (HSP) in early embryonic development and reproductive efficiency. An understanding of how these proteins influence reproductive efficiency both in vivo and in vitro could have major ramifications for production agriculture. In cattle, performance differences have been identified in *Bos indicus* breeds and genetic polymorphisms in HSP70 have been associated with reproductive performance (Rosenkrans et al., 2010). In Holstein cows, differences in expression of genes in the HSP40 family have been associated with improved early embryonic development in vitro, indicating that these proteins may have a greater role in reproductive efficiency than thought, even in animals not adapted to a tropical environment and not placed in a heat stress situation (Zhang et al., 2011). The speakers at this Physiology and Endocrinology Symposium were identified to address these topics based on their research.

High thermal temperatures can negatively impact fertility in cattle, and there is evidence that the oocyte

and embryo are targets of heat stress (Hansen and Sakatani, 2012). Breed differences between *Bos indicus* and *Bos taurus* cattle contribute to heat tolerance and improved fertility under heat stress conditions (Paula-Lopes et al., 2013). Heat can induce cellular damage in both the cytoplasm and the nucleus of the oocyte. Differences in gene expression and (or) function of mechanisms that control cellular stress contribute to the ability of oocytes and embryos of *Bos indicus* origin to tolerate the stresses associated with high thermal temperatures.

Mammalian cells, including the oocyte, have developed systems to protect themselves from these stresses. Heat shock proteins, which are produced to counteract damages from external stimuli, are one such system. Driver and Khatib (2013) reviewed differences in expression of genes in the HSP family during early embryonic development related to early embryonic survival. They further reviewed associations between polymorphisms in HSP and fertility in cattle. For example, Rosenkrans et al. (2010) reported the influence of a polymorphism in HSP70 on calving percentage and calving date in Brahman-influenced cows. In their symposium presentation, Rosenkrans et al. (2012) reported a greater minor allele frequency for the G117A

¹A symposium held at the Joint Annual Meeting, July 15 to 19, 2012, Phoenix, AZ, with publication sponsored by the *Journal of Animal Science* and the American Society of Animal Science.

²Names are necessary to report factually on available data; however, the USDA neither guarantees nor warrants the standard of the product, and the use of names by the USDA implies no approval of the product to the exclusion of others that may also be suitable.

³Corresponding author: Bob.Cushman@ars.usda.gov

⁴The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part

of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, DC. 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Received January 3, 2013.

Accepted January 3, 2013.

polymorphism of HSP70 in Brahman cows than in Angus cows or crossbred cows. There was a genotype-by-forage system interaction, such that cows carrying the G117A polymorphism and grazing endophyte-infected tall fescue had lower lifetime calving rates. Taken together, these data provide strong evidence for a contribution of HSP to breed differences in thermal tolerance between *Bos indicus* and *Bos taurus* cattle.

Because a large portion of the world's agricultural lands fall within tropical and subtropical regions, it will be important to continue to understand how thermal tolerance contributes to fertility in both *Bos indicus* and *Bos taurus* cattle. Issues with age at puberty and age at first calving in *Bos indicus* cattle dictate the need for crossbreeding with *Bos taurus* germplasm to improve reproductive efficiency. Only by combining the best of both germplasms will we be able to produce the food necessary to feed an ever-increasing world population.

LITERATURE CITED

- Driver, A. M., and H. Khatib. 2013. Heat shock proteins: Potentially powerful markers for preimplantation embryonic development and fertility in livestock species. *J. Anim. Sci.* 91:1154–1161.
- Hansen, P. J., and M. Sakatani. 2012. Consequences of heat shock on the development of pre-implantation bovine embryos: Role of free radicals, antioxidants, apoptosis, and heat shock proteins. *J. Anim. Sci.* 90(Suppl. 3):424. (Abstr.)
- Paula-Lopes, F. F., R. S. Lima, R. A. Satrapa, and C. M. Barros. 2013. Influence of cattle genotype (*Bos indicus* vs. *Bos taurus*) on oocyte and preimplantation embryo resistance to increased temperature. *J. Anim. Sci.* 91:1143–1153.
- Rosenkrans, C., Jr., A. Banks, S. Reiter, and M. Looper. 2010. Calving traits of crossbred Brahman cows are associated with Heat Shock Protein 70 genetic polymorphisms. *Anim. Reprod. Sci.* 119:178–182.
- Rosenkrans, C., Jr., M. Brown, H. Brown, Jr., and M. Looper. 2012. Associations between heat shock protein 70 genetic polymorphisms and calving traits in crossbred Brahman cows. *J. Anim. Sci.* 90(Suppl. 3):424. (Abstr.)
- Zhang, B., F. Peñagaricano, A. Driver, H. Chen, and H. Khatib. 2011. Differential expression of heat shock protein genes and their splice variants in bovine preimplantation embryos. *J. Dairy Sci.* 94:4174–4182.