

University of Nebraska - Lincoln DigitalCommons@University of Nebraska - Lincoln

Papers in Natural Resources

Natural Resources, School of

9-1965

Ultrasonic Tracking of Migratory Fishes with an Internal Tag

Ross M. Horrall University of Wisconsin

H. F. Henderson University of Wisconsin

A. D. Hasler University of Wisconsin

Follow this and additional works at: http://digitalcommons.unl.edu/natrespapers

Part of the <u>Aquaculture and Fisheries Commons</u>, <u>Bioinformatics Commons</u>, <u>Natural Resources</u> and <u>Conservation Commons</u>, <u>Natural Resources Management and Policy Commons</u>, and the <u>Other</u> <u>Environmental Sciences Commons</u>

Horrall, Ross M.; Henderson, H. F.; and Hasler, A. D., "Ultrasonic Tracking of Migratory Fishes with an Internal Tag" (1965). *Papers in Natural Resources*. 566.

http://digitalcommons.unl.edu/natrespapers/566

This Article is brought to you for free and open access by the Natural Resources, School of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Natural Resources by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Horall, Henderson, and Hasler in American Fisheries Society Meeting (September 23-24, 1965; Portland, Orgeon). Copyright 1965, the authors. Used by permission.

American Fisheries Society Meeting September 23-24, 1965 Portland, Oregon

Ultrasonic Tracking of Migratory Fishes with an Internal Tag


Ross M. Horrall, H.F. Henderson, and A.D. Hasler The University of Wisconsin

A small ultrasonic tag, 9 mm by 40 mm in size, has been developed by members of the fish orientation group at the University of Wisconsin. The tag emits a continuous signal at approximately 65 kc/sec. The signal is received via a directional hydrophone and a high quality communications receiver. Tracking ranges of up to 1 km have been obtained. The placement of the tag is internal: either into the stomach or into the body cavity of the fish. The tags and tracking equipment have been successfully employed in studies on the open water movements and orientation of white bass during their spawning migration. (Supported by NSF Grant GB-3123; ONR contract Nonr-1202(26); PHS Grant 5T1-WP-2-04; and the Wisconsin Conservation Department)


Techniques for tracking fish:

- 1. Bobber tracking (Hasler <u>et al</u>. 1958)
- 2. Balloon method (Horrall 1961)
- 3. Dye marker method (Hasler and Henderson 1963)
- 4. Ultrasonic tracking (Trefethen 1956; Trefethen <u>et al</u>. 1957; Johnson 1960; Novotny and Esterberg 1962; Henderson 1963)

The Wisconsin Ultrasonic Tag:



Receiving equipment:



<u>References</u>:

- Hasler, A.D., and H.F. Henderson. 1963. Instrumentation problems in the study of homing in fish. In: Bio-telemetry. Pergamon Press. New York. 195-201.
- Hasler, A.D., R.M. Horrall, W.J. Wisby, and W. Braemer. 1958. Sun-orientation and homing in fishes. Limnol. Oceanogr. 3(4): 353-361.
- Henderson, H.F. 1963. Orientation of pelagic fishes. (I) Optical problems (II) Sonic tracking. Ph.D. Thesis. Univ. of Wisconsin. 135 pp.
- Horrall, R.M. 1961. A comparative study of two spawning populations of the white bass, <u>Roccus chrysops</u> (Rafinesque), in Lake Mendota, Wisconsin, with special reference to homing behavior. Ph.D. Thesis. Univ. of Wisconsin. 181 pp.
- Johnson, J.H. 1960. Sonic tracking of adult salmon at Bonneville Dam, 1957. U.S. Fish and Wildlife Service, Fishery Bulletin 60: 471-485.
- Novotny, A.J. and G.F. Esterberg. 1962. A 132-kilocycle sonic fish tag. Progressive Fish Culturist 24(3): 139-141.
- Trefethen, P.S. 1956. Sonic equipment for tracking individual fish. U.S. Fish and Wildlife Service. Special Scientific Report--Fisheries 179:1-11.
- Trefethen, P.S., J.W. Dudley, and M.R. Smith. 1957. Ultrasonic tracer follows tagged fish. Electronics 30(4): 156-160.