

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Historical Materials from University of
Nebraska-Lincoln Extension

Extension

1992

G92-1071 Ridge Plant Systems: Weed Control

Robert N. Klein

University of Nebraska - Lincoln, robert.klein@unl.edu

Gail A. Wicks

University of Nebraska - Lincoln

Alex Martin

University of Nebraska - Lincoln, amartin2@unl.edu

Russell S. Moomaw

University of Nebraska - Lincoln

Fred Roeth

University of Nebraska - Lincoln, fwroeth41@gmail.com

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/extensionhist>



Part of the [Agriculture Commons](#), and the [Curriculum and Instruction Commons](#)

Klein, Robert N.; Wicks, Gail A.; Martin, Alex; Moomaw, Russell S.; Roeth, Fred; Wilson, Robert G.; and Jasa, Paul J., "G92-1071 Ridge Plant Systems: Weed Control" (1992). *Historical Materials from University of Nebraska-Lincoln Extension*. 728.

<https://digitalcommons.unl.edu/extensionhist/728>

This Article is brought to you for free and open access by the Extension at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Historical Materials from University of Nebraska-Lincoln Extension by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Robert N. Klein, Gail A. Wicks, Alex Martin, Russell S. Moomaw, Fred Roeth, Robert G. Wilson, and Paul J. Jasa



Ridge Plant Systems: Weed Control

Advantages and disadvantages of the ridge plant system, weed control before and at planting and economics of the system are discussed.

*Robert N. Klein, Extension Cropping Systems Specialist;
Gail A. Wicks, Extension Weeds Specialist;
Alex R. Martin, Extension Weeds Specialist;
Russell S. Moomaw, Extension Crops and Weeds Specialist;
Fred W. Roeth, Extension Weeds Specialist;
Robert G. Wilson, Extension Weeds Specialist; and
Paul J. Jasa, Extension Engineer*

- [Advantages of Ridge Planting](#)
- [Disadvantages of Ridge Planting](#)
- [Ridge Clearing Depth](#)
- [Weed Control Before Planting](#)
- [Broadleaf Weed Control](#)
- [Grass Weed Control](#)
- [Preplant Cultivation](#)
- [Weed Control at Planting](#)
- [Shattercane Control Research](#)
- [Economics](#)
- [Summary](#)

Ridge planting combines tillage and herbicides to achieve improved weed control in row crops. Crop seed is planted into ridges formed during cultivation and/or ditching of the previous crop. In ridge planting, the planter follows the old row and ridge clearing sweeps or disks move the surface soil, residue and much of the weed seed out of the row. Weed seeds are deposited between the rows where, upon germination, they can be controlled with cultivation. Two cultivations are generally used for weed control. The first cultivation loosens the soil and the second rebuilds the ridge.

The ridge plant system is well suited to furrow-irrigationd crops. It also works well with dryland crops or those under center pivot irrigation. On furrow irrigationd land, corn or sorghum stalks may need to be shredded to assist in decomposition and hence irrigation because crop residue slows water advance in the furrow. Slowing the water may be a benefit, however, on soils which have a low water intake rate. With center pivot and dryland acres the need for shredding depends on how much residue the cultivator can handle.

Advantages of Ridge Planting

Advantages of the ridge plant system include:

1. *Improved weed control.* In the row annual weed density can be reduced nearly 80 percent with ridge plant compared to mulch-till systems. Herbicides are more effective at lower weed densities. Also, any weeds that have germinated or are about to emerge in the ridge at planting time will be destroyed. This gives the crop an even start with weeds in the row. By comparison, if a tandem disk or a rotary tiller is used to incorporate herbicides and prepare a seedbed, surface weed seed is "planted" in the row. This makes ridge planting one of the easiest systems to use with little or no herbicide. Ridge till is especially beneficial for reducing shattercane in corn.
2. *Warmer soil temperature.* Residue from the previous crop settles between the ridges, permitting the ridge to warm up sooner. The warmer soil speeds crop emergence, increases competitiveness with weeds, and increases the likelihood of a good crop stand. The quick and improved crop emergence allows for earlier crop cultivation, which may control weeds between the rows better.
3. *Better soil moisture.* Without preplant tillage, ridge planting preserves soil moisture. Tillage can dry the soil to the depth of the tillage operation. In dry springs, especially in western Nebraska, soil moisture near the surface may be inadequate at planting for crop germination and growth if the soil was chiseled or disked. Realizing this, growers using tillage to prepare a seedbed often plant deeper or use a lister which may reduce crop stands and allow excessive soil erosion.

In higher rainfall areas, water drains off the ridge so planting can be earlier than on level ground. Since the ridge dries out sooner, the system is well adapted to poorly drained soils.

4. *Volunteer crops reduced.* Ridge planting almost eliminates volunteer crops. The seeds, ears, or heads are moved by the planter to the row middle where cultivation can control seeds that germinate. Volunteer crops are often the worst weeds in conventionally tilled fields.
5. *Reduced herbicide cost.* Band application of herbicides at planting works well with ridge planting because the system reduces weed seeds and clears the row of residue which may intercept herbicides. In general, banding reduces herbicide costs by two-thirds.
6. *Erosion control.* If ridge height is maintained and the land does not have more than a 4 percent slope, erosion will be reduced. With a 4 to 7 percent slope, it may be necessary to ridge plant on the contour.
7. *Controlled traffic pattern.* Soil compaction in the row is reduced if all equipment is designed to stay off the ridges.

Disadvantages of Ridge Planting

The disadvantages or inconveniences of the ridge plant system include:

1. *Early weeds, especially with late planted crops.* Weeds must be controlled prior to planting. The later a crop is planted the more likely that weeds may be present.
2. *Preplant incorporated herbicides inappropriate.* Preplant incorporated herbicide treatments should not be used because the tillage operation will destroy the ridge and mix weed seed into the

soil. Also, unless the herbicide is incorporated extremely deep, which decreases its effectiveness, the herbicide will be moved out of the row with the soil. The exception is where a ridge cleaning device is used in front of a tiller which incorporates the herbicide after ridge cleaning.

3. *Equipment changes.* The planter, cultivator and ditcher may need to be modified or new equipment may need to be purchased. Planting equipment must have a row cleaning device to push residue and weed seed to the row middles (see NebGuide G88-876, *Ridge Plant Systems Equipment*). The cultivator and ditcher need to be able to operate in crop residue. The ditcher also needs to be able to build a rounded or flat topped ridge. Special equipment may be needed to keep the planter on the ridge.
4. *Wheel spacing and tire size adjustment.* The wheel spacing of implements and the combine must be adjusted to the row spacing. Also tire size is limited because large tires may destroy the ridge. An alternative is to use duals with spacers to fit rows if extra load carrying capacity is needed.
5. *Limited traffic pattern.* All traffic must follow rows.
6. *Works best for continuous row crops.* With modification, small grains have been drilled on ridges.

Ridge Clearing Depth

The ridge clearing device should be run about 1 to 2 inches deep. A planter set too shallow may not remove weed seed and may ride on top of previous crop residue. A planter set too deep may create a furrow, and remove the old root stumps, which could contribute to soil erosion or crusting in the row. Also, the soil is wetter and colder down deep resulting in slower and uneven crop emergence. A preemergence herbicide is usually applied after ridge clearing and planting in a band over the row.

In addition to planters which use sweeps or disks (including horizontal rotating disks) to clear weed seed and residue from the crop row, some planters use a sweep or a furrowing shovel mounted on a tool bar ahead of a rotary tiller with planting units mounted behind. By removing all but two blades per row, only the ridge is tilled. The furrowing shovel moves the crop residue and weed seed between the rows where any weeds that emerge can be controlled with cultivation. The rotary tiller also can incorporate herbicide in the row (band). Incorporated herbicides do not need rainfall or irrigation to move them into the soil.

Weed Control Before Planting

Weeds germinating before planting must be controlled. It is a mistake to plant through large weeds like tansymustard, kochia, lambsquarters, prickly lettuce, horseweed, smartweed, and Russian thistle. If growing between the rows, these weeds will probably not be killed at planting time. Furthermore, they can quickly remove all the moisture in the seedbed.

Large weeds can keep the planter from operating at the proper depth. Since these weeds may not have been killed at planting time, they become more difficult to control at cultivation. At first cultivation these large weeds may not be cut off with the disks or sweeps. They will compete with the crop and present a harvest problem. Weeds growing on the side of the ridge at planting time also present problems. The trouble arises along the cutting edge of the planter sweep or disk, where large broadleaf weeds may not be uprooted or covered. Herbicides, such as atrazine, Bladex, 2,4-D, or 2,4-D + Banvel can be applied in early April, before planting, to control winter annuals and early summer annual weeds. The atrazine rate depends on future crops that will be planted.

An early herbicide treatment should eliminate the potential for planting into emerged weeds. For information on herbicides, rates, and weed response ratings see the latest *Guide for Herbicide Use in Nebraska*, Extension publication EC-130.

Some operators who do not want to use an early preplant herbicide application have added wings or extensions along the back edge of the planter sweep for additional undercutting width. This cleans out a bigger area and makes cultivation easier. The wings or extensions are sometimes used only on the outside unit to assist when "guess rows" are wider.

Broadleaf Weed Control

Most broadleaf weeds can be controlled effectively and economically in corn with 2,4-D. If kochia, smartweed, or prickly lettuce are present, Banvel or Buctril must be used because 2,4-D alone has not always been effective in controlling these weeds. The application timing and rate depend upon weed size, weed population, time of planting, and crop to be planted. The best time to apply postemergence herbicides is when the weeds are less than 1 inch tall. This may be two or more weeks before planting, depending on planting date. If 2,4-D or Banvel is applied during corn planting, attach the spray boom to the front or under the tractor rather than behind the planter. Postemergence herbicides applied behind the planter may be less effective because soil and crop residue may cover some weeds and prevent herbicides from reaching the weeds. With sorghum, apply the 2,4-D or Banvel treatments at least two weeks before planting. Banvel or 2,4-D also can be applied with drop nozzles after crop emergence to control weed escapes when necessary in corn or sorghum.

For preplant weed control in soybeans, one option is to use Roundup or Graxomone Extra. These nonselective herbicides do not have residual activity. Preplant applications of Pursuit Plus, Command, or Sencor/Lexone also could be used for controlling weeds in soybeans; however, the herbicide will be moved out of the row with the ridge cleaning device. A preemergence herbicide treatment is usually needed in the row after planting.

Grass Weed Control

Winter annual grasses, such as downy brome, need to be controlled by mid-April to prevent soil moisture loss.

Most grass weeds in row crops are summer annual grasses such as foxtail, barnyardgrass, and sandbur. These weeds start germinating in late April and early May and will often be present when row crops are planted.

If considerable grass weed growth is present before planting, Roundup or Landmaster could be used. Check the Landmaster label for crop and waiting period before planting. Another option would be Gramoxone Extra plus a residual triazine herbicide such as atrazine or Bladex for corn and sorghum. In soybeans, several herbicides can be tank mixed with Gramoxone Extra for grass weed removal and residual weed control.

Preplant Cultivation

Preplant cultivation can be used for inter-row weed control. Emerged weeds on the ridge top will be destroyed, if small, by the planter's row cleaning sweep or disks. This works extremely well on fields where corn was ensiled. Set the cultivator to control weeds between the rows without throwing soil to the ridge. Preplant cultivation has been tried to rebuild ridges which may have been damaged by harvest

equipment or livestock tramping.

Disadvantages of preplant cultivation include: (1) the weed seed may be incorporated deeper into the ridge than the planter row cleaning devices will remove; (2) the seedbed may not be firm unless the row cleaning device removes all the added soil and the planter places the crop seed in soil not disturbed in rebuilding the ridges; and (3) controlling planter depth can be difficult when a preplant cultivation is used.

If the soil in the spring is too wet for preplant cultivation, herbicides may be a better option. While preplant cultivation of row middles for weed control means an extra trip over the field and extra labor and fuel costs, it may be a better option than preplant disking. Unless operated very shallow, disking will remove the ridge. Then when using a planter with row cleaning devices, the result is similar to shallow listing into cool wet soil. The use of a mulch treader is another possibility to control emerged weeds. It will not destroy the ridge but may incorporate some weed seeds deeper than the ridge cleaning device will remove.

Weed Control at Planting

Use a band herbicide application behind the planter to control weeds in the crop row. With light to moderate weed density a broadcast application is probably not needed. However, where heavy weed density is expected or with hard to control weeds like velvetleaf, a broadcast herbicide may be necessary. If the grower anticipates that the first cultivation will be more than four weeks after planting, a broadcast application may be appropriate. Some producers band herbicide where they are able to perform timely cultivation and broadcast herbicide on the remaining acres. After several years of continuous ridge plant system use, weed density may be reduced and it may be possible to only apply herbicides every other year.

Shattercane Control Research

At the University of Nebraska South Central Research and Extension Center near Clay Center, a ridge plant system using disk furrowers displaced 81 percent of the shattercane seeds from the crop rows to the middles. When combined with a preemergence application of Lasso at 4 lb/acre, 94 percent control of shattercane was achieved. A broadcast application of Eradicane that was incorporated with a rotary tiller gave 90 percent shattercane control the first year and 69 percent control the second year. Thus, ridge planting compared favorably with conventional systems for shattercane control in corn even though the soil-applied herbicide was not incorporated. A distinct advantage of the ridge plant system was that there was no decline in shattercane control with continued use.

Economics

Budgets for disk-disk plant and ridge plant systems are in *Table I*. Yields are reduced instead of including insurance for hail, etc. With ridge plant always having better moisture at planting the estimated yield used was 160 bushels per acre compared to 155 bushels per acre for the disk-disk plant. The atrazine 2,4-D preplant treatment in ridge plant should be used only when weeds are present. This reduces ridge plant costs approximately \$.03 per bushel. Each producer will have different costs than those used in the budgets, therefore, these should only be used as a guide.

Table I. Estimated 1992 costs per acre, in a central Nebraska corn plot with electrical gravity irrigation, 225 ft. head. Expected yield per acre 155 bushels for disk-disk plant and 160 bushel for ridge-plant.

| Job | Fuel and repair costs per acre | |
|---|--------------------------------|-----------------|
| | Disk-Disk plant | Ridge plant |
| Stalk Shredder | \$ --- | \$ 1.50 |
| Tandem Disc | 1.60 | --- |
| Sprayer | --- | .80* |
| 1.0 qt Atrazine | --- | 3.00* |
| 0.5 qt 2,4-D | --- | 1.40* |
| Tandem Disc | 1.50 | --- |
| Anhydrous Applicator | 1.60 | 1.60 |
| —160 and 166 lb Nitrogen, NH ₃ | 17.60 | 18.30 |
| Planter | 3.60 | 3.80 |
| —0.33 bag seed | 26.00 | 26.00 |
| —6 gal 10-34-0 | 7.50 | 7.50 |
| —Herbicide Band | --- | 3.60 |
| —Insecticide | 12.00 | 12.00 |
| Herbicide | 10.80 | --- |
| Sprayer | .80 | --- |
| Row Crop Cultivator | 1.80 | 1.80 |
| Ditcher | 1.80 | 1.80 |
| Combine, Corn Head | 6.70 | 6.90 |
| Dryer | 15.50 | 16.00 |
| Haul | 7.80 | 8.00 |
| Auger | .30 | .30 |
| Total | \$116.90 | \$114.30 |
| Interest, Overhead, Management, and Labor | 75.00 | 70.00 |
| Total | \$191.90 | \$184.30 |
| | <i>Fuel and Repair</i> | |
| irrigation 18 in | \$ 50.00 | \$ 50.00 |
| Fixed Machinery Cost | 42.50 | 40.50 |
| Land & Irrigation System | 90.00 | 90.00 |
| Cost Per Acre | 374.40 | 364.80 |
| Cost Per Unit of Production | 2.42 | 2.28 |

*Use only when weeds are present. Ridge plant costs are \$2.25 per bushel without treatment.

Summary

Ridge planting moves up to 80 percent of the weed seeds out of the crop row--setting the stage for economical and effective weed control. Ridge planting is particularly adapted to furrow irrigation where the ridges are formed for irrigation. Ridge planting may make it possible to lower crop production costs by implementing a weed control program which includes both mechanical and chemical weed control methods.

Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by Cooperative Extension is implied.

File G1071 under: FIELD CROPS
G-28, Cropping Practices
Issued February 1992; 5,000 printed

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Elbert C. Dickey, Director of Cooperative Extension, University of Nebraska, Institute of Agriculture and Natural Resources.

University of Nebraska Cooperative Extension educational programs abide with the non-discrimination policies of the University of Nebraska-Lincoln and the United States Department of Agriculture.