

University of Nebraska - Lincoln DigitalCommons@University of Nebraska - Lincoln

Nutrition and Health Sciences -- Faculty
Publications

Nutrition and Health Sciences, Department of

2012

Muscadine Grape (*Vitis rotundifolia*) and Wine Phytochemicals Prevented Obesity-Associated Metabolic Complications in C57BL/6J Mice

Vishnupriya Gourineni
University of Florida

Neil F. Shay
Oregon State University, neil.shay@oregonstate.edu

Soonkyu Chung
University of Nebraska-Lincoln, chung4@unl.edu

Amandeep K. Sandhu
University of Florida, asandhu2@iit.edu

Liwei Gu
University of Florida, lgu@ufl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/nutritionfacpub>

 Part of the [Human and Clinical Nutrition Commons](#), [Molecular, Genetic, and Biochemical Nutrition Commons](#), and the [Other Nutrition Commons](#)

Gourineni, Vishnupriya; Shay, Neil F.; Chung, Soonkyu; Sandhu, Amandeep K.; and Gu, Liwei, "Muscadine Grape (*Vitis rotundifolia*) and Wine Phytochemicals Prevented Obesity-Associated Metabolic Complications in C57BL/6J Mice" (2012). *Nutrition and Health Sciences -- Faculty Publications*. 107.
<https://digitalcommons.unl.edu/nutritionfacpub/107>

This Article is brought to you for free and open access by the Nutrition and Health Sciences, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Nutrition and Health Sciences -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Published in *Journal of Agricultural and Food Chemistry* 60 (2012), pp. 7674–7681.

doi 10.1021/jf3013663 |

Copyright © 2012 American Chemical Society. Used by permission.

Submitted March 29, 2012; revised July 2, 2012; accepted July 12, 2012.

PMID: 22788667

Muscadine Grape (*Vitis rotundifolia*) and Wine Phytochemicals Prevented Obesity-Associated Metabolic Complications in C57BL/6J Mice

Vishnupriya Gourineni,¹ Neil F. Shay,^{1,2} Soonkyu Chung,¹
Amandeep K. Sandhu,¹ and Liwei Gu¹

¹ Food Science and Human Nutrition Department, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, Florida 32611, USA

² Department of Food Science and Technology, Oregon Wine Research Institute, Oregon State University, Corvallis, Oregon 97331, USA

Corresponding author — L. Gu, email lgu@ufl.edu

Abstract

The objective of this study was to determine the effects of muscadine grape or wine (cv. Noble) phytochemicals on obesity and associated metabolic complications. Muscadine grape or wine phytochemicals were extracted using Amberlite FPX66 resin. Male C57BL/6J mice were given a low-fat diet (LF, 10% kcal fat), high-fat diet (HF, 60% kcal fat), HF + 0.4% muscadine grape phytochemicals (HF+MGP), or HF + 0.4% muscadine wine phytochemicals (HF+MWP) for 15 weeks. At 7 weeks, mice fed HF+MGP had significantly decreased body weights by 12% compared to HF controls. Dietary MGP or MWP supplementation reduced plasma content of free fatty acids, triglycerides, and cholesterol in obese mice. Inflammation was alleviated, and activity of glutathione peroxidase was enhanced. Consumption of MGP or MWP improved insulin sensitivity and glucose control in mice. Thus, consumption of muscadine grape and wine phytochemicals in the diet may help to prevent obesity-related metabolic complications.

Keywords: muscadine grape, wine, obesity, C57BL/6J mice

Introduction

Obesity is a multifactorial condition posing major health problems worldwide. It contributes to multiple metabolic abnormalities such as cardiovascular disease, insulin resistance, dyslipidemia, oxidative stress, inflammation,

and fatty liver.¹ Due to its negative impact on public health, multiple campaigns have been launched to curb the increase of obesity in the United States. One of the effective strategies to prevent obesity is through dietary interventions, especially the use of phytochemicals.² Dietary polyphenols reduce obesity through different mechanisms such as suppression of adipocyte hypertrophy, inhibition of preadipocyte differentiation, stimulation of lipolysis, or induction of fat cell apoptosis.³ For example, resveratrol decreased adipogenesis and viability of preadipocytes by transcriptional and post-transcriptional regulation of adipocyte-specific genes. In mature adipocytes, resveratrol decreased lipid accumulation through stimulation of lipolysis and apoptosis.⁴ In vitro and in vivo studies demonstrated the synergistic effects of resveratrol, quercetin, and genistein in preventing weight gain and adipogenesis.⁵ Experimental diets supplemented with a mixture of grape (*Vitis vinifera*) extract, green tea extract, and L-carnitine were also effective in preventing obesity and related metabolic disorders.⁶

Muscadine grapes (*Vitis rotundifolia*) are commonly grown and consumed in the southeastern United States. They contain a unique blend of bioactive phytochemicals with potential to improve symptoms of chronic metabolic diseases. Muscadine grapes are round, have either bronze- or purple-colored leatherlike thick skin, and are well-adapted to warm, humid climates, which are not suitable for the growth of *V. vinifera*.⁷ The phytochemical composition of muscadine is significantly different from that of *V. vinifera*. For example, the skin of purple muscadine grapes contains exclusively anthocyanin 3,5-diglucosides, whereas *V. vinifera* contains only monoglucosides. The seeds and skin of muscadine grapes contain large amounts of ellagittannins and ellagic acid, which are absent in *V. vinifera*.⁸ The resveratrol content of muscadine grapes is much lower compared to *V. vinifera*.⁹ Phytonutrients with antioxidant properties are positively associated with disease prevention. A growing number of studies showed that muscadine grapes had anticancer,¹⁰ antibacterial,^{11,12} and anti-inflammatory properties.¹³ To our knowledge, no study has evaluated the effects of muscadine grape or wine phytochemicals on the development of obesity and the associated metabolic complications.

A number of studies investigated the effects of supplementing high-fat diets with skin,¹⁴ seed,¹⁵ and pomace^{16,17} extracts of *V. vinifera* on adiposity, inflammation, insulin resistance, and hyperlipidemia. For example, grape seed extracts supplemented in high-fat diets partially reversed obesity by affecting lipid metabolism in mice.¹⁸ However, results from these studies cannot be extrapolated to muscadine grapes due to the difference in phytochemical composition. The objective of this study was to assess the effect of feeding muscadine grape or wine phytochemicals in high-fat diets on obesity and metabolic complications in C57BL/6J mice.

Materials and Methods

Chemicals and Reagents

Folin–Ciocalteu reagent, 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox), ellagic acid, quercetin, myricetin, and kaempferol were purchased from Sigma Chemical Co. (St. Louis, MO, USA). Standards of the 3-O- β -glucosides of pelargonidin, cyanidin, peonidin, delphinidin, petunidin, and malvidin (six mixed anthocyanin standard) were purchased from Polyphenols Laboratories (Sandnes, Norway). HPLC grade solvents were used for phytochemical extraction and analysis (Fisher Scientific, Atlanta, GA, USA). A mouse insulin enzyme-linked immunoassay (ELISA) kit was purchased from Crystal Chemicals (Downers Grove, PA, USA), and a mouse CRP ELISA kit was obtained from Immunology Consultants Laboratory Inc. (Newberg, OR, USA). A cytosolic glutathione peroxidase (cGPx) kit was purchased from Cayman Chemical Co. (Ann Arbor, MI, USA). Red muscadine wine (cv. Noble) was provided by Lakeridge Wineries (Clermont, FL, USA). Muscadine grapes (cv. Noble) were purchased from a local vineyard.

Muscadine Grape and Wine Extract Preparation

Grapes (15 kg) were crushed and extracted with acidified methanol (0.5% acetic acid, 20 L) in 10 batches. Extracts were filtered using Whatman no. 4 filter paper. Methanol in the extract was removed using a rotary evaporator. The concentrated extract was resuspended in water and loaded on a column (80 mm i.d. \times 600 mm) (Ace Laboratories, NJ, USA) filled with Amberlite FPX66 resin (Rohm Hass, PA, USA). Prior to the sample loading, resin was conditioned with acidified water (0.5% formic acid), and the initial resin bed volume was 1 L. Noble muscadine wine (20 L) was loaded in 10 batches on the resin column without any treatment. After sample loading, the column was eluted with 4 L of acidified water to remove sugars and other compounds not adsorbed on the resins. Muscadine grape and wine phytochemicals were recovered from the column using 4 L of methanol elution. Extracts were dried using a SpeedVac concentrator (Thermo Scientific ISS110, Waltham, MA, USA). Approximately, 40 g of dried muscadine grape and 30 g of wine phytochemicals were obtained. Characterization and Phytochemical Analysis of Extracts. The total phenolic content of the extracts was determined by using Folin–Ciocalteu reagent with gallic acid as a standard, and results were expressed as milligrams of gallic acid equivalents per gram of dry weight (mg of GAE/g). ORAC assay was conducted to measure the peroxy radical-scavenging activity of the extracts and was expressed in micromoles of Trolox equivalents per gram of the extracts ($\mu\text{mol TE/g}$).¹⁹

Anthocyanins were quantified using HPLC according to a published method.²⁰ Other phenolic compounds, including ellagic acid, quercetin, myricetin, and kaempferol, were quantified using HPLC after acid hydrolysis.⁷

Briefly, dry muscadine grape or wine phytochemicals (150 mg) were dissolved in 5 mL of 1.2 N HCl in 50% methanol, vortexed, and incubated in a water bath at 90 °C for 80 min to hydrolyze flavonoid glycosides to aglycones. The hydrolyzed extracts were filtered through a 0.2 µm syringe nylon filter and diluted. Chromatographic analyses were performed on an Agilent 1200 series HPLC system (Agilent Technologies, Palo Alto, CA, USA) equipped with an autosampler/injector and diode array detector. A Zorbax SBC¹⁸ column (4.6 mm × 250 mm, 5 µm, Agilent Technologies) was used for separation. Elution was performed using mobile phase A (0.5% formic acid aqueous solution) and mobile phase B (acetonitrile). UV–vis spectra were scanned from 220 to 600 nm on a diode array detector with detection wavelengths of 280, 360, and 520 nm.

Animal Study

Low-fat, high-fat, and high-fat diet with phytochemicals from muscadine grape or wine were prepared by Research Diets Inc. (New Brunswick, NJ, USA) using the formulations shown in Table 1. Thirty-six 6-week-old male C57BL/6J mice were purchased from Jackson Laboratory (Bar Harbor, ME, USA) and housed in shoebox cages on corn-cob bedding (three per cage). Mice were acclimatized for 1 week and maintained on a 12 h light/dark cycle with access to food and water ad libitum. Following the adaptation period, mice were randomly divided into four experimental groups ($n = 9$): low fat (LF, 10% kcal fat), high fat (HF, 60% kcal fat), high fat plus 0.4% muscadine grape phytochemical (HF+MGP), and high fat plus 0.4% muscadine wine phytochemical (HF+MWP). The LF and HF diets had energy densities of 3.85 and 5.24 kcal/g, respectively. The phytochemical dose in the diets was chosen on the basis of the effectiveness of resveratrol supplementation at 0.4% in diets for 15 weeks on body weight and obesity conditions.²¹ Mice were maintained on experimental diets for 15 weeks. Body weight and food intake was recorded weekly. At the end of the study, animals were fasted for 7 h, and blood was taken by cardiac puncture from anesthetized mice. Plasma samples were isolated by centrifugation at 2000g for 15 min at 4 °C and stored at –80 °C. Livers and epididymal adipose tissue were harvested and weighed. Sections of liver and adipose tissue were frozen at –80 °C. This protocol was approved by the Institutional Animal Care and use Committee of the University of Florida.

Intraperitoneal Glucose Tolerance Test

Glucose tolerance tests were performed at week 12 on five mice per group. After mice were fasted for 6 h, 2 g/kg glucose was given intraperitoneally. Tail blood was collected at 0, 10, 20, 30, 60, and 120 min. Blood glucose concentration was determined with a hand-held Contour glucose monitor (Bayer Healthcare, Tarrytown, NY, USA). The area under curve (AUC; millimolar per minute) was calculated using a trapezoidal rule.²²

Measurement of Plasma Biomarkers Related to Glucose and Lipid Metabolism.

Plasma concentrations of total cholesterol, triglycerides, and free fatty acids were determined using commercial kits (Abcam, MA, USA). Fasting plasma insulin levels were measured with a murine ultrasensitive ELISA kit, measured spectrophotometrically at 450 nm and expressed as millimoles per liter. Plasma glucose levels (mmol/L) were measured using a glucose assay kit (Cayman Chemicals). Insulin resistance was evaluated by homeostasis model of insulin resistance (HOMA-IR) as previously described.²³

Measurement of Inflammatory and Oxidative Stress

Biomarkers. Plasma C-reactive protein levels were measured by ELISA using a plate reader at 450 nm and expressed as nanograms per milliliter. The plasma antioxidant enzyme glutathione peroxidase (GPx) activity was measured spectrophotometrically at 340 nm using commercial kits and expressed as nanomoles per minute per milliliter.

Histology

For the histological examination, epididymal fat tissues were fixed in 10% buffered formalin overnight and then embedded in paraffin, cut at a thickness of 5 μm , and stained with hematoxylin and eosin (H&E).²¹ Frozen livers were sectioned into 8 μm thick sections using a microtome, and lipid deposition was detected by staining with Oil-red-O.⁶ Digital images were acquired with a Leica DM 2500 phase contrast microscope and Magnafire and Image Pro 6.0 imaging software (Media Cybernetics, Silver Spring, MD, USA).

Statistical Analyses

All data are expressed as the mean \pm standard error of the mean (SEM). One-way analyses of variance (ANOVA) with Tukey–Kramer HSD pairwise comparison of the means were performed using JMP (version 8.0, SAS Institute Inc., Cary, NC, USA). Fisher's least significant test was applied to analyze body weight gain of mice. Graphs were generated using Sigma-Plot software version 11.0. A $p \leq 0.05$ difference was considered to be significant.

Results and Discussion

Muscadine Grape and Wine Phytochemical Composition.

Phytochemical compositions of MGP and MWP are shown in Table 2. MGP contained a significantly higher amount of ellagic acid (18.2 mg/g) than MWP (1.1 mg/g). Ellagic acid has very low solubility in water. Ellagitannins hydrolyze to ellagic acid, and the ellagic acid forms sediment in muscadine wine.²⁴ The precipitated ellagic acid in muscadine is commonly removed by

filtration. On the other hand, total anthocyanin content was higher in wine phytochemicals (124.8 mg/g) than in grape phytochemicals (67.9 mg/g). Red muscadine wine is commonly made by fermentation of crushed grapes (including skin). Anthocyanins were extracted from skins and concentrated in wine during fermentation. The ratio of individual anthocyanin 3,5-diglucoses in MGP was slightly different from that in MWP. For example, the major anthocyanin in MGP was delphinidin 3,5-diglucoside, whereas peonidin 3,5-diglucoside was the major anthocyanin in MWP. This difference was likely due to the change of anthocyanin profile during winemaking. Total phenolic content of phytochemicals in grape and wine were 620 and 389 mg of GAE/g, respectively (Table 2). The antioxidant activities of MGP and MWP, evaluated by ORAC, were 4824 and 3383 $\mu\text{mol TE/g}$, respectively.

Food Intake and Body and Tissue Weights in Mice

Average food intake (g/mouse/day) in mice was estimated and reported in Table 3. Animals fed low-fat diet consumed more feed compared to mice in the high-fat diet group (3.3 vs 2.7 g/mouse/day; $p < 0.05$). The energy intake of mice on the low-fat diet was not significantly different from that of mice on the high-fat diet (12.8 vs 14.3 kcal/mouse/day). Grape or wine phytochemical supplementation in high-fat diets did not affect estimated food intake and caloric intake in animals (Table 3). The body weights of mice from week 1 to 15 are shown in Figure 1A. The net body weight gains at weeks 7 and 15 are depicted in Figure 1B. As expected, mice fed a high-fat diet gained more weight than mice fed a low-fat diet. At the end of the experiment, mice in the HF group had gained 37% more weight ($p < 0.05$) than mice in the LF group. Supplementing high-fat diets with grape or wine phytochemicals did not affect the cumulative body weights in mice at week 15 (Figure 1B). However, during the seventh week of the feeding period, HF fed controls had gained 45% more weight ($p < 0.05$) than LF mice. Mice fed HF+MGP showed a 12% decrease in body weights ($p < 0.05$) compared to high-fat fed mice. The suppression of body weight gain caused by MGP in animals was not due to decreased food intake. Similar observations were made in mice fed resveratrol (0.4%) for 15 weeks,²⁵ grape seed extract (250 mg/kg bw) for 12 weeks,⁵ and dietary wine pomace extract (0.45%) for 4 weeks.¹⁷

Feeding mice with HF diet increased the weight of epididymal adipose tissue by 67% compared to the fat mass in their lean counterparts on LF diets (Table 3). Supplementation of MGP and MWP in the HF diet significantly reduced fat mass by 29 and 12.5%, respectively, as compared to the HF group. MGP and MWP supplementation reduced liver weights by 41.6 and 37.5%, respectively, when compared to mice fed the HF diet alone (Table 3). Hepatic tissue in lean mice weighed 54% less than that of obese mice. In previous studies, rodents fed HF diets showed more body weight,

higher fat deposition and total liver weight, and increased energy intake compared with those on the standard-fat diet.²⁶ Similar results were obtained in this study.

The morphology of adipose tissue and liver of mice from different diet groups is depicted in Figure 2. Feeding mice with obesogenic HF diets resulted in increased fat mass with crown-like structures in epididymal adipose tissue (Figure 2A) and increased lipids in hepatic tissues (Figure 2B). However, MGP supplementation altered adipose tissue morphology and reduced lipid accumulation in the liver. Epididymal adipose tissue sections by H&E staining show adipocytes from mice fed the different diets; although difference in adipocyte size may exist, this parameter was not measured. High-fat diets have shown to increase lipid accumulation in adipocytes. This causes an expansion of adipose tissue, initiating elevation in adipose expression of inflammatory genes such as TNF- α and monocyte chemoattractant protein-1 (MCP-1).²⁷ Inflamed and insulin-resistant adipocytes, characterized by low lipogenic and high lipolytic capacity, cause increased release of free fatty acids (FFA). FFA may activate Toll-like receptor 4 (TLR4), resulting in cytokine or chemokine secretion propagating inflammatory signals to the systemic level.²⁸ MGP and MWP may attenuate adipose inflammation by reducing adipose tissue macrophage (ATM) infiltration and suppressing TLR4 mediated pro-inflammatory signaling cascade, thereby lowering plasma FFA in mice fed HF diets.²⁸ Quercetin, one of the polyphenols available in grape powder extract, was found to be effective in mitigating macrophage-mediated inflammation in adipocytes.²⁹ Figure 2B shows grape or wine phytochemical effects on reducing hepatic lipidosis. Obesity accompanies liver lipid accumulation, which leads to hepatic steatosis and increased glucose production in liver. The livers of obese controls had numerous lipid droplets compared to lean counterparts, whereas MGP fed mice had lower lipids in hepatic tissues as similarly seen in mice fed LF diets. These data are parallel with the liver tissue weights in mice fed HF and grape- or wine-supplemented diets. Grape or wine polyphenols may target hepatic defensive enzymes and adipogenic gene expression, as seen in mice fed grape extract.⁶

Glucose Metabolism

The time course for plasma glucose levels is shown in Figure 3A. A marginally delayed glucose clearance was noted in all mice (including the LF group). In contrast to this study, an earlier study showed that feeding mice strawberry powder supplemented in high-fat diets lowered nonfasting blood glucose levels but did not alter responses during intraperitoneal glucose challenge.³⁰ Strawberries or blueberries did not improve glucose tolerance in mice.³¹ The incremental AUC of plasma glucose concentration during intraperitoneal glucose tolerance test is shown in Figure 3B. The AUC of glucose levels in LF group was 47.4% lower ($p < 0.001$) compared to HF

controls (Figure 3B). The AUC levels in the mice fed HF+MGP and HF+MWP were lower than that of HF group by 36.3 and 22.2%, respectively, but not statistically different from either LF- or HF-fed mice. At the end of the feeding study, fasting blood glucose and insulin levels were measured in all mice ($n = 9$), and the homeostasis model assessment of basal insulin resistance (HOMA-IR) was calculated (Table 4). Lower HOMA-IR values indicate greater insulin sensitivity, whereas higher HOMA-IR values show lower insulin sensitivity or insulin resistance. MGP significantly reversed HF diet induced elevations in plasma glucose (-21.3%) and plasma insulin (-47.7%) levels compared to obese controls, whereas MWP reduced plasma glucose and insulin levels by 12.3 and 36.9%, respectively. Insulin resistance was found to be higher in HF obese controls at week 15. Compared to the HF group, insulin sensitivity was increased significantly by muscadine grape or wine supplementation as indicated by lower HOMA-IR. Our results were consistent with an early study that showed the antihyperglycemic effects of grape seed extracts.³² Various mechanisms and pathways may be responsible for insulin-sensitizing effects by grape seed extract³² and grape powder,³³ such as insulin-signaling pathway, decreased hepatic lipid accumulation, and/or FFA-induced oxidative stress.

Muscadine Phytochemicals on Plasma Lipid Profile

Dyslipidemia is associated with higher risk for cardiovascular disease, insulin resistance, and nonalcoholic fatty liver. The influence of HF diets supplemented with grape or wine phytochemicals on systemic markers of lipid metabolism were measured. When compared to LF control group, the serum free fatty acids (Figure 4A), triglycerides (Figure 4B), and total cholesterol (Figure 4C) were elevated by >2 -fold in mice fed high-fat diets. There was a $>40\%$ reduction in plasma free fatty acids, triglycerides, and total cholesterol in mice fed MGP in HF diets. MWP supplementation resulted in lower plasma lipid levels in mice, but no significant differences were noted between MGP and MWP feeding groups. Similar effects of resveratrol and grape seed proanthocyanidins on lipid homeostasis were reported earlier,^{21,34} suggesting grape compounds regulate hepatic lipogenesis and adipose lipolysis simultaneously. Consumption of lyophilized grape powder over 4 weeks improved plasma lipid profiles and reduced oxidative stress in women.³⁵ According to Boer et al.,³⁶ triglycerides are synthesized in the liver and are then transported by very low-density lipoproteins to adipose tissues. Here, we noted that serum free fatty acids, total triglyceride, and total cholesterol levels were significantly higher in the obese control group compared to their lean counterparts. Muscadine polyphenol supplementation improved plasma lipid profile, which was correlated with lower adipose tissue and liver mass.

Phytochemical Supplementation on Antioxidant Status and Inflammation

Inflammation and oxidative stress are the junction points between increased adiposity, insulin resistance, and dyslipidemia. This complex of obesity-related metabolic abnormalities is known as metabolic syndrome.³⁷ The plasma C-reactive protein level is a measurable inflammatory biomarker in obesity conditions. We found that grape and wine phytochemical supplementation normalized this inflammatory biomarker protein levels with respect to LF-fed lean controls and increased the activity of glutathione peroxidase. The obese control group had a >2.5-fold increase in plasma C-reactive protein levels, as compared to lean controls (Table 5). Mice fed muscadine grape or wine phytochemicals in high-fat diets over 15 weeks had 50 and 23% reductions in plasma C-reactive protein, respectively. Glutathione peroxidase is an antioxidative enzyme that is downregulated in chronic disease conditions such as obesity and insulin resistance.³⁸ Dysregulation in redox homeostasis leads to adipocyte dysfunction and finally increased oxidative stress. Glutathione peroxidase activity was significantly lower in mice fed HF diets (Table 5). The MGP-supplemented group had a >4-fold increase in glutathione peroxidase activity, and MWP-fed animals showed 2-fold increased antioxidant enzyme levels in plasma. Dietary polyphenols have the capacity to increase the expression of anti-inflammatory and antioxidative genes via the activation of nuclear factor- κ B and nuclear factor erythroid related factor-2.³⁹ An inverse correlation between C-reactive protein and glutathione peroxidase activity in HF-fed experimental groups suggests a beneficial role of muscadine grape or wine phytochemicals in mitigating oxidative stress and inflammation in obese animals.

A high-fat diet induces obesity in C57BL/6J mice and the associated metabolic syndrome characterized by hyperlipidemia, hyperglycemia, oxidative stress, and inflammation. Muscadine grape and wine phytochemicals were able to restore obesity-triggered pathological conditions in C57BL/6J mice. These effects are comparable to those of the extracts from *V. vinifera*.⁴⁰ Major phytochemicals in muscadine grapes were anthocyanins. Purified anthocyanins from purple potato, blueberries, and purple corn were reported to reduce the pathogenesis of obesity.⁴¹ The benefits of anthocyanins were attributed to their capacity to activate the adenosine monophosphate-activated protein kinase (AMPK) signaling pathway. Anthocyanins were also reported to down-regulate the levels of sterol regulatory element-binding protein 1 and its target genes including acetyl-coenzyme A carboxylase and fatty acid synthase.⁴¹ MGP supplementation in diets appeared to be more effective in reducing animal body weights and lipid accumulation than MWP. Such differences were likely due to differences in phytochemical compositions. MGP contained a higher amount of ellagic acid than MWP. An ellagitannin-rich

extract from pomegranate leaf was effective in preventing obesity in mice.⁴² Ellagic acid suppressed resistin secretion by reducing its intracellular protein levels.⁴³ The synergistic effect between anthocyanins and ellagic acid may be possible by affecting targets in different pathways. Resveratrol in *V. vinifera* was shown to prevent obesity.²¹ However, resveratrol was not likely to be a major bioactive compound in muscadine grapes, because its level was extremely low.^{9,44} Resveratrol was below the detection limit in the extracts used in this study. Clearly, relative levels of ellagic acid, resveratrol, and other phenolics in any specific extract derived from specific fruits will affect the specific metabolic impacts that extract provides.

In summary, our findings support the promising role of muscadine grape and wine phytochemicals as dietary means in obesity management. Additional studies are needed to determine the effective dose levels of muscadine grape or wine phytochemicals required to affect relevant biomarker of chronic disease.

Table 1. Composition of Experimental Diets

Ingredient	LF		HF		HF+MGP		HF+MWP	
	g/kg	kcal/kg	g/kg	kcal/kg	g/kg	kcal/kg	g/kg	kcal/kg
casein	200	800	200	800	200	800	200	800
L-cystine	3	12	3	12	3	12	3	12
corn starch	506.2	2025	0	0	0	0	0	0
maltodextrin	125	500	125	500	125	500	125	500
sucrose	68.8	275	68.8	275	68.8	275	68.8	275
cellulose	50	0	50	0	50	0	50	0
lard	20	180	245	2205	245	2205	245	2205
soybean oil	25	225	25	225	25	225	25	225
mineral mix	10	0	10	0	10	0	10	0
vitamin mix	10	40	10	40	10	40	10	40
MGP	0	0	0	0	3.09	0	0	0
MWP	0	0	0	0	0	0	3.09	0
total	1055	4057	774	4057	777	4057	777	4057
	g %	kcal %	g %	kcal %	g %	kcal %	g %	kcal %
carbohydrate	67	70	26	20	26	20	26	20
protein	19	20	26	20	26	20	26	20
fat	4	10	35	60	35	60	35	60
MGP	0	0	0	0	0.4	0	0	0
MWP	0	0	0	0	0	0	0.4	0

Table 2. Antioxidant and Phytochemical Composition of Muscadine Grape or Wine Extracts^a

Phytochemical	MGP (mg/g dry extract)	MWP (mg/g dry extract)
quercetin	0.5 ± 0.1	0.5 ± 0.02
myricetin	1.4 ± 0.3	2.1 ± 0.1
kaempferol	0.2 ± 0.06	0.3 ± 0.01
ellagic acid	18.2 ± 0.15*	1.1 ± 0.3
delphinidin 3,5-diglucoside	19.0 ± 1.1	25.5 ± 1.5
cyanidin 3,5-diglucoside	7.9 ± 2.5	19.1 ± 1.1*
petunidin/pelargonidin 3,5-diglucoside ^b	16.3 ± 1.1	27.2 ± 1.5*
peonidin 3,5-diglucoside	14.6 ± 1.6	35.2 ± 2.2*
malvidin 3,5-diglucoside	10.1 ± 1.8	17.8 ± 1.1
total anthocyanins	67.9 ± 5.7	124.8 ± 3.2*
total phenolic content (mg of GAE/g)	620 ± 58	389 ± 32
ORAC (μmol/TE/g)	4824 ± 6.8	3383 ± 7.1

a. * = significant differences at $p \leq 0.05$ using t test.

b. Peaks coeluted on HPLC. Samples were analyzed in duplicate. Data are the mean ± SEM for duplicate tests.

Table 3. Food Intake, Energy Intake, and Weight of Epididymal Adipose Tissue and Liver^a

Group	food intake (g/mouse/day)	energy intake (kcal/mouse/day)	adipose wt (g)	adipose % body wt	liver wt (g)	liver % body wt
LF	3.3 ± 0.1 a	12.8 ± 0.4 a	1.67 ± 0.03 c	3.98 ± 0.12 bc	1.16 ± 0.07 c	2.76 ± 0.18 b
HF	2.7 ± 0.1 b	14.4 ± 0.7 a	2.42 ± 0.13 a	4.90 ± 0.30 a	2.45 ± 0.08 a	4.93 ± 0.19 a
HF+MGP	2.7 ± 0.06 b	14.3 ± 0.3 a	1.70 ± 0.05 c	3.48 ± 0.13 c	1.40 ± 0.04 b	2.86 ± 0.09 b
HF+MWP	2.7 ± 0.1 b	14.1 ± 0.4 a	2.12 ± 0.05 b	4.16 ± 0.13 b	1.51 ± 0.05 b	2.96 ± 0.10 b

a. Values are the mean ± SEM. HF, $n = 8$; other groups, $n = 9$. Columns not sharing a common letter are significantly different ($p \leq 0.05$).

Table 4. Effects of MGP and MWP on Glucose Homeostasis in C57BL/6J Mice^a

Group	plasma glucose (mg/dL)	plasma insulin (μ units/L)	HOMA-IR
LF	132.4 ± 0.4 d	10.8 ± 0.02 d	3.5 ± 0.02 d
HF	176.0 ± 0.3 a	27.9 ± 0.08 a	12.2 ± 0.02 a
HF+MGP	140.8 ± 0.4 c	15.0 ± 0.04 c	5.1 ± 0.02 c
HF+MWP	154.0 ± 0.5 b	18.1 ± 0.04 b	6.9 ± 0.01 b

a. Values are the mean ± SEM. HF, $n = 8$; other groups, $n = 9$. Columns not sharing a common letter are significantly different ($p \leq 0.05$).

Table 5. Effects of MGP and MWP on Plasma CRP and Glutathione Peroxidase Activity^a

Group	plasma C-reactive protein (CRP) (ng/mL)	plasma glutathione peroxidase activity (nmol/min/mL)
LF	23.1 ± 0.02 d	1.1 ± 0.02 b
HF	61.8 ± 0.14 a	0.3 ± 0.03 d
HF+MGP	30.1 ± 0.03 c	1.2 ± 0.02 a
HF+MWP	47.6 ± 0.01 b	0.7 ± 0.05 c

a. Values are the mean ± SEM. HF, $n = 8$; other groups, $n = 9$. Columns not sharing a common letter are significantly different ($p \leq 0.05$).




Figure 1. Cumulative body weights of C57BL/6J mice (1–15 weeks): net body weight gains at weeks (A) 7 and (B) 15. HF, $n = 8$; other groups, $n = 9$.




Figure 2. (A) Histology of the epididymal adipose tissue, stained with hematoxylin and eosin; (B) histology of liver tissue, stained with Oil-red-O. Arrows indicate crown-like structures in adipose tissue.




Figure 3. (A) Blood glucose levels in intraperitoneal glucose tolerance tests at week 12; (B) area under curve of blood glucose for 2 h after a glucose challenge ($n = 5$).




Figure 4. Plasma lipid levels of free fatty acids (A), total triglycerides (B), and total cholesterol (C) in C57BL/6J mice. HF, $n = 8$; other groups, $n = 9$.

Funding — This research is supported in part by a grant from the Florida Department Agriculture and Consumer Service, viticulture research program. The authors declare no competing financial interest.

Abbreviations

cGPx	cytosolic glutathione peroxidase
HF	high fat
HOMAIR	homeostatic model assessment–insulin resistance
LF	low fat
MGP	muscadine grape phytochemicals
MWP	muscadine wine phytochemicals

References

- (1) Grundy, S. Multifactorial causation of obesity: implications for prevention. *Am. J. Clin. Nutr.* 1998, 67 (3), 563S–572S.
- (2) González-Castejón, M.; Rodríguez-Casado, A. Dietary phytochemicals and their potential effects on obesity: a review. *Pharmacol. Res.* 2011, 64 (5), 438–455.
- (3) Rayalam, S.; Della-Fera, M. A.; Baile, C. A. Phytochemicals and regulation of the adipocyte life cycle. *J. Nutr. Biochem.* 2008, 19 (11), 717–726.
- (4) Baile, C. A.; Yang, J.-Y.; Rayalam, S.; Hartzell, D. L.; Lai, C.-Y.; Andersen, C.; Della-Fera, M. A. Effect of resveratrol on fat mobilization. *Ann. N.Y. Acad. Sci.* 2011, 1215 (1), 40–47.
- (5) Park, H. J.; Yang, J.-Y.; Ambati, S.; Della-Fera, M. A.; Hausman, D. B.; Rayalam, S.; Baile, C. A. Combined effects of genistein, quercetin, and resveratrol in human and 3T3-L1 adipocytes. *J. Med. Food* 2008, 11 (4), 773–783.
- (6) Kang, J. S.; Lee, W. K.; Yoon, W. K.; Kim, N.; Park, S.-K.; Park, H. K.; Ly, S. Y.; Han, S.-B.; Yun, J.; Lee, C. W.; Lee, K.; Lee, K. H.; Park, S.-K.; Kim, H. M. A combination of grape extract, green tea extract and L-carnitine improves high-fat diet-induced obesity, hyperlipidemia and non-alcoholic fatty liver disease in mice. *Phytother. Res.* 2011, 25 (12), 1789–1795.
- (7) Pastrana-Bonilla, E.; Akoh, C. C.; Sellappan, S.; Krewer, G. Phenolic content and antioxidant capacity of muscadine grapes. *J. Agric. Food Chem.* 2003, 51 (18), 5497–5503.
- (8) Lee, J.-H.; Johnson, J. V.; Talcott, S. T. Identification of ellagic acid conjugates and other polyphenolics in muscadine grapes by HPLC-ESI-MS. *J. Agric. Food Chem.* 2005, 53 (15), 6003–6010.
- (9) Ector, B. J.; Magee, J. B.; Hegwood, C. P.; Coign, M. J. Resveratrol concentration in muscadine berries, juice, pomace, purees, seeds, and wines. *Am. J. Enol. Vitic.* 1996, 47 (1), 57–62.
- (10) Yi, W.; Fischer, J.; Akoh, C. C. Study of anticancer activities of muscadine grape phenolics in vitro. *J. Agric. Food Chem.* 2005, 53 (22), 8804–8812.
- (11) Park, Y. J.; Biswas, R.; Phillips, R. D.; Chen, J. Antibacterial activities of blueberry and muscadine phenolic extracts. *J. Food Sci.* 2011, 76 (2), M101–M105.

- (12) Kim, T. J.; Silva, J. L.; Jung, Y. S. Antibacterial activity of fresh and processed red muscadine juice and the role of their polar compounds on *Escherichia coli* O157:H7. *J. Appl. Microbiol.* 2009, *107* (2), 533–539.
- (13) Greenspan, P.; Bauer, J. D.; Pollock, S. H.; Gangemi, J. D.; Mayer, E. P.; Ghaffar, A.; Hargrove, J. L.; Hartle, D. K. Anti-inflammatory properties of the muscadine grape (*Vitis rotundifolia*). *J. Agric. Food Chem.* 2005, *53* (22), 8481–8484.
- (14) Hogan, S.; Canning, C.; Sun, S.; Sun, X.; Kadouh, H.; Zhou, K. Dietary supplementation of grape skin extract improves glycemia and inflammation in diet-induced obese mice fed a Western high fat diet. *J. Agric. Food Chem.* 2011, *59* (7), 3035–3041.
- (15) Terra, X.; Pallarés, V.; Ardèvol, A.; Bladé, C.; Fernández-Larrea, J.; Pujadas, G.; Salvadó, J.; Arola, L.; Blay, M. Modulatory effect of grape-seed procyanidins on local and systemic inflammation in diet-induced obesity rats. *J. Nutr. Biochem.* 2011, *22* (4), 380–387.
- (16) Hogan, S.; Canning, C.; Sun, S.; Sun, X.; Zhou, K. Effects of grape pomace antioxidant extract on oxidative stress and inflammation in diet induced obese mice. *J. Agric. Food Chem.* 2010, *58* (21), 11250–11256.
- (17) Yunoki, K.; Sasaki, G.; Tokuji, Y.; Kinoshita, M.; Naito, A.; Aida, K.; Ohnishi, M. Effect of dietary wine pomace extract and oleanolic acid on plasma lipids in rats fed high-fat diet and its DNA microarray analysis. *J. Agric. Food Chem.* 2008, *56* (24), 12052–12058.
- (18) Park, S.-H.; Park, T.-S.; Cha, Y.-S. Grape seed extract (*Vitis vinifera*) partially reverses high fat diet-induced obesity in C57BL/6J mice. *Nutr. Res. Pract.* 2008, *2* (4), 227–233.
- (19) Lee, J.-H.; Talcott, S. T. Fruit maturity and juice extraction influences ellagic acid derivatives and other antioxidant polyphenolics in muscadine grapes. *J. Agric. Food Chem.* 2003, *52* (2), 361–366.
- (20) Wu, X.; Prior, R. L. Systematic identification and characterization of anthocyanins by HPLC-ESI-MS/MS in common foods in the United States: fruits and berries. *J. Agric. Food Chem.* 2005, *53* (7), 2589–2599.
- (21) Kim, S.; Jin, Y.; Choi, Y.; Park, T. Resveratrol exerts anti-obesity effects via mechanisms involving down-regulation of adipogenic and inflammatory processes in mice. *Biochem. Pharmacol.* 2011, *81* (11), 1343–1351.
- (22) Andrikopoulos, S.; Blair, A. R.; Deluca, N.; Fam, B. C.; Proietto, J. Evaluating the glucose tolerance test in mice. *Am. J. Physiol. – Endocrinol. Metab.* 2008, *295* (6), E1323–E1332.
- (23) Trout, K. K.; Homko, C.; Tkacs, N. C. Methods of measuring insulin sensitivity. *Biol. Res. Nursing* 2007, *8* (4), 305–318.
- (24) Lee, J.-H.; Talcott, S. T. Ellagic acid and ellagitannins affect on sedimentation in muscadine juice and wine. *J. Agric. Food Chem.* 2002, *50* (14), 3971–3976.
- (25) Lagouge, M.; Argmann, C.; Gerhart-Hines, Z.; Meziane, H.; Lerin, C.; Daussin, F.; Messadeq, N.; Milne, J.; Lambert, P.; Elliott, P.; Geny, B.; Laakso, M.; Puigserver, P.; Auwerx, J. Resveratrol improves mitochondrial function and protects against metabolic disease by activating SIRT1 and PGC-1 α . *Cell* 2006, *127* (6), 1109–1122.
- (26) Milagro, F. I.; Campion, J.; Martinez, J. A. Weight gain induced by high-fat feeding involves increased liver oxidative stress. *Obesity* 2006, *14* (7), 1118–1123.

- (27) Chen, A.; Mumick, S.; Zhang, C.; Lamb, J.; Dai, H.; Weingarh, D.; Mudgett, J.; Chen, H.; MacNeil, D. J.; Reitman, M. L.; Qian, S. Diet induction of monocyte chemoattractant protein-1 and its impact on obesity. *Obesity* 2005, 13 (8), 1311–1320.
- (28) Lionetti, L.; Mollica, M. P.; Lombardi, A.; Cavaliere, G.; Gifuni, G.; Barletta, A. From chronic overnutrition to insulin resistance: the role of fat-storing capacity and inflammation. *Nutr., Metab. Cardiovasc. Dis.* 2009, 19 (2), 146–152.
- (29) Overman, A.; Chuang, C. C.; McIntosh, M. Quercetin attenuates inflammation in human macrophages and adipocytes exposed to macrophage-conditioned media. *Int. J. Obes.* 2011, 35 (9), 1165–1172.
- (30) Parelman, M. A.; Storms, D. H.; Kirschke, C. P.; Huang, L.; Zunino, S. J. Dietary strawberry powder reduces blood glucose concentrations in obese and lean C57BL/6 mice, and selectively lowers plasma C-reactive protein in lean mice. *Br. J. Nutr. FirstView* 2012, 1–11.
- (31) Prior, R. L.; Wu, X.; Gu, L.; Hager, T. J.; Hager, A.; Howard, L. R. Whole berries versus berry anthocyanins: interactions with dietary fat levels in the C57BL/6J mouse model of obesity. *J. Agric. Food Chem.* 2008, 56 (3), 647–653.
- (32) Suwannaphet, W.; Meeprom, A.; Yibchok-Anun, S.; Adisakwattana, S. Preventive effect of grape seed extract against high-fructose diet-induced insulin resistance and oxidative stress in rats. *Food Chem. Toxicol.* 2010, 48 (7), 1853–1857.
- (33) Chuang, C.-C.; Bumrungpert, A.; Kennedy, A.; Overman, A.; West, T.; Dawson, B.; McIntosh, M. K. Grape powder extract attenuates tumor necrosis factor α -mediated inflammation and insulin resistance in primary cultures of human adipocytes. *J. Nutr. Biochem.* 2011, 22 (1), 89–94.
- (34) Baiges, I.; Palmfeldt, J.; Bladé, C.; Gregersen, N.; Arola, L. Lipogenesis is decreased by grape seed proanthocyanidins according to liver proteomics of rats fed a high fat diet. *Mol. Cell. Proteomics* 2010, 9 (7), 1499–1513.
- (35) Zern, T. L.; Wood, R. J.; Greene, C.; West, K. L.; Liu, Y.; Aggarwal, D.; Shachter, N. S.; Fernandez, M. L. Grape polyphenols exert a cardioprotective effect in pre- and postmenopausal women by lowering plasma lipids and reducing oxidative stress. *J. Nutr.* 2005, 135 (8), 1911–1917.
- (36) den Boer, M.; Voshol, P. J.; Kuipers, F.; Havekes, L. M.; Romijn, J. A. Hepatic steatosis: a mediator of the metabolic syndrome. lessons from animal models. *Arterioscl., Thromb., Vasc. Biol.* 2004, 24 (4), 644–649.
- (37) Chuang, C.-C.; McIntosh, M. K. Potential mechanisms by which polyphenol-rich grapes prevent obesity-mediated inflammation and metabolic diseases. *Annu. Rev. Nutr.* 2011, 31 (1), 155–176.
- (38) Kobayashi, H.; Matsuda, M.; Fukuhara, A.; Komuro, R.; Shimomura, I. Dysregulated glutathione metabolism links to impaired insulin action in adipocytes. *Am. J. Physiol. –Endocrinol. Metab.* 2009, 296 (6), E1326–E1334.
- (39) Rahman, I.; Biswas, S. K.; Kirkham, P. A. Regulation of inflammation and redox signaling by dietary polyphenols. *Biochem. Pharmacol.* 2006, 72 (11), 1439–1452.
- (40) Décordé, K.; Teissèdre, P.-L.; Sutra, T.; Ventura, E.; Cristol, J.-P.; Rouanet, J.-M. Chardonnay grape seed procyanidin extract supplementation prevents high-fat diet-induced obesity in hamsters by improving adipokine imbalance and oxidative stress markers. *Mol. Nutr. Food Res.* 2009, 53 (5), 659–666.

- (41) Hwang, Y. P.; Choi, J. H.; Han, E. H.; Kim, H. G.; Wee, J.-H.; Jung, K. O.; Jung, K. H.; Kwon, K.-i.; Jeong, T. C.; Chung, Y. C.; Jeong, H. G. Purple sweet potato anthocyanins attenuate hepatic lipid accumulation through activating adenosine monophosphate-activated protein kinase in human HepG2 cells and obese mice. *Nutr. Res. (N.Y.)* 2011, *31* (12), 896–906.
- (42) Lei, F.; Zhang, X. N.; Wang, W.; Xing, D. M.; Xie, W. D.; Su, H.; Du, L. J. Evidence of anti-obesity effects of the pomegranate leaf extract in high-fat diet induced obese mice. *Int. J. Obes.* 2007, *31* (6), 1023–1029.
- (43) Makino-Wakagi, Y.; Yoshimura, Y.; Uzawa, Y.; Zaima, N.; Moriyama, T.; Kawamura, Y. Ellagic acid in pomegranate suppresses resistin secretion by a novel regulatory mechanism involving the degradation of intracellular resistin protein in adipocytes. *Biochem. Biophys. Res. Commun.* 2012, *417* (2), 880–885.
- (44) Leblanc, M. R.; Johnson, C. E.; Wilson, P. W. Influence of pressing method on juice stilbene content in muscadine and bunch grapes. *J. Food Sci.* 2008, *73* (4), H58–H62.