

University of Nebraska - Lincoln  
**DigitalCommons@University of Nebraska - Lincoln**

---

Biochemistry -- Faculty Publications

Biochemistry, Department of

---

2014

# Proline Biosynthesis Is Required for Endoplasmic Reticulum Stress Tolerance in *Saccharomyces cerevisiae*

Xinwen Liang

*University of Nebraska - Lincoln*, [xliang2@unl.edu](mailto:xliang2@unl.edu)

M. B. Dickman

*Texas A&M University*

Donald F. Becker

*University of Nebraska-Lincoln*, [dbecker3@unl.edu](mailto:dbecker3@unl.edu)

Follow this and additional works at: <https://digitalcommons.unl.edu/biochemfacpub>

 Part of the [Biochemistry Commons](#), [Biotechnology Commons](#), and the [Other Biochemistry, Biophysics, and Structural Biology Commons](#)

---

Liang, Xinwen; Dickman, M. B.; and Becker, Donald F., "Proline Biosynthesis Is Required for Endoplasmic Reticulum Stress Tolerance in *Saccharomyces cerevisiae*" (2014). *Biochemistry -- Faculty Publications*. 173.

<https://digitalcommons.unl.edu/biochemfacpub/173>

This Article is brought to you for free and open access by the Biochemistry, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Biochemistry -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

# Proline Biosynthesis Is Required for Endoplasmic Reticulum Stress Tolerance in *Saccharomyces cerevisiae*\*

Received for publication, March 4, 2014, and in revised form, July 30, 2014. Published, JBC Papers in Press, August 11, 2014, DOI 10.1074/jbc.M114.562827

Xinwen Liang<sup>‡</sup>, Martin B. Dickman<sup>§</sup>, and Donald F. Becker<sup>†1</sup>

From the <sup>‡</sup>Department of Biochemistry and Redox Biology Center, University of Nebraska, Lincoln, Nebraska 68588 and the

<sup>§</sup>Institute for Plant Genomics and Biotechnology, Department of Plant Pathology and Microbiology, Texas A&M University, College Station, Texas 77843

**Background:** Proline is an important amino acid for stress resistance in different organisms.

**Results:** Depletion of proline biosynthesis disrupts redox homeostasis and increases sensitivity to endoplasmic reticulum (ER) stress in yeast.

**Conclusion:** Proline biosynthesis is critical for maintaining the intracellular redox environment and the UPR during ER stress.

**Significance:** Proline metabolism is shown to have an important role in ER stress tolerance that was previously unknown.

The amino acid proline is uniquely involved in cellular processes that underlie stress response in a variety of organisms. Proline is known to minimize protein aggregation, but a detailed study of how proline impacts cell survival during accumulation of misfolded proteins in the endoplasmic reticulum (ER) has not been performed. To address this we examined in *Saccharomyces cerevisiae* the effect of knocking out the *PRO1*, *PRO2*, and *PRO3* genes responsible for proline biosynthesis. The null mutants *pro1*, *pro2*, and *pro3* were shown to have increased sensitivity to ER stress relative to wild-type cells, which could be restored by proline or the corresponding genetic complementation. Of these mutants, *pro3* was the most sensitive to tunicamycin and was rescued by anaerobic growth conditions or reduced thiol reagents. The *pro3* mutant cells have higher intracellular reactive oxygen species, total glutathione, and a NADP<sup>+</sup>/NADPH ratio than wild-type cells under limiting proline conditions. Depletion of proline biosynthesis also inhibits the unfolded protein response (UPR) indicating proline protection involves the UPR. To more broadly test the role of proline in ER stress, increased proline biosynthesis was shown to partially rescue the ER stress sensitivity of a *hog1* null mutant in which the high osmolality pathway is disrupted.

Proline metabolism has been linked to stress tolerance in a broad range of organisms. In addition to being a proteogenic amino acid, proline functions as an osmolyte, oxidative stress protectant, chemical chaperone, and as a source of nitrogen and energy under nutrient limiting conditions. Proline is synthesized from glutamate in three enzymatic steps as shown in Fig. 1A. Glutamate is first converted to glutamate- $\gamma$ -semialdehyde (GSA)<sup>2</sup> via the coordinated activities of glutamate kinase and

$\gamma$ -glutamyl phosphate reductase. In yeast *Saccharomyces cerevisiae*, glutamate kinase (Pro1p) and  $\gamma$ -glutamyl phosphate reductase (Pro2p) are separate enzymes, whereas in plants and humans glutamate kinase and  $\gamma$ -glutamyl phosphate reductase are combined into the bifunctional enzyme  $\Delta^1$ -pyrroline-5-carboxylate synthetase. GSA spontaneously cyclizes to  $\Delta^1$ -pyrroline-5-carboxylate (P5C), which is reduced to proline by P5C reductase (Pro3p). The proline biosynthesis pathway is subject to proline feedback inhibition with proline-inhibiting glutamate kinase (Pro1p) (1). P5C can also be generated from ornithine by ornithine- $\delta$ -aminotransferase (Car2p). Thus, genetic deletion of *PRO1* and *PRO2* does not completely disrupt proline biosynthesis in yeast. In eukaryotes proline is degraded back to glutamate in the mitochondrion by proline dehydrogenase (Put1p) and P5C dehydrogenase (Put2p).


Understanding the mechanisms by which proline metabolism influences cellular responses to stress is complicated by its multifaceted roles in the cell. In mammalian cells the breakdown of proline has been shown to generate mitochondrial reactive oxygen species (ROS), which ultimately induce cell signaling cascades that promote cell apoptosis or survival (2–5) and in *Caenorhabditis elegans* increase lifespan (6). Proline biosynthesis has been shown to be up-regulated in plants under different stress conditions with proline accumulation generally associated with increased stress tolerance (2, 7, 8). In humans inborn errors in human P5C reductase isozyme 1 (PYCR1) (9, 10) and P5C synthetase (P5CS) (11, 12) are associated with deficiencies in connective tissue and progeroid features. Fibroblasts from patients with impaired PYCR1 activity show altered mitochondrial morphology and increased apoptosis upon oxidative stress (10). Increased expression of PYCR1 is found in different human cancer cells (4, 13–16) and is thought to enhance cell proliferation (4, 13). In yeast *S. cerevisiae*, when a Pro1p variant that is insensitive to feedback inhibition by proline was overexpressed in a *PUT1* deletion strain (*put1*), proline levels were increased, and the yeast strain was more resistant to

reactive oxygen species; Tm, tunicamycin; UPR, unfolded protein response; NADP<sup>+</sup>, nicotinamide adenine dinucleotide phosphate; NADPH, reduced form of NADP<sup>+</sup>.

\* This work was supported, in whole or in part, by National Institutes of Health Grants GM079393 and P30GM103335. This work was also supported by the University of Nebraska Agricultural Research Division.

<sup>1</sup> To whom correspondence should be addressed. Tel.: 402-472-9652; Fax: 402-472-7842; E-mail: dbecker3@unl.edu.

<sup>2</sup> The abbreviations used are: GSA, glutamate- $\gamma$ -semialdehyde; ER, endoplasmic reticulum; GLR, glutathione reductase; *GSH1*,  $\gamma$ -glutamylcysteine synthetase 1; IRE1, inositol-requiring protein; NAC, *N*-acetyl-*l*-cysteine; PI, propidium iodide; P5C,  $\Delta^1$ -pyrroline-5-carboxylate; ROS,



**FIGURE 1. Proline metabolic pathway and characterization of proline biosynthesis enzyme localization and null mutants.** *A*, proline metabolic pathway in *S. cerevisiae*. Proline is synthesized from glutamate in the cytosol starting with Pro1p (glutamate kinase) and Pro2p ( $\gamma$ -glutamyl phosphate reductase). The intermediate, GSA, spontaneously cyclizes to P5C, which is then reduced to proline by Pro3p (P5C reductase). Alternatively, GSA is generated from ornithine (Orn) via Car2p. Proline is oxidized to glutamate by Put1p (proline dehydrogenase) and Put2p (P5C dehydrogenase) in the mitochondrion. Put1p couples proline oxidation to reduction of ubiquinone (CoQ) in the electron transport chain. *B*, fluorescence visualization by confocal microscopy of Pro1p-, Pro2p-, and Pro3p-GFP fusion proteins in strain A44 (PRO1-GFP:KanMX6), A46 (PRO2-GFP:KanMX6), and A48 (PRO3-GFP:KanMX6), respectively. All GFP fusion proteins are distributed throughout the cells, indicating cytosolic localization. *C*, growth rates of the BY4741 WT strain and null mutants *pro1* (left panel), *pro2* (middle panel), and *pro3* (right panel) in SD medium supplemented with different concentrations of proline (2, 4, 10, and 20 mM) as indicated. Cells were diluted to  $A_{600}$  of 0.01 and inoculated into 3 ml of medium and grown at 30 °C with shaking (220 rpm) in 15 ml of cell culture tubes. The growth rate was monitored at  $A_{600}$  at the indicated time points. The scale bar indicates 10  $\mu$ m.

various stressors such as freezing, desiccation, hydrogen peroxide, and ethanol (17). Accordingly, a *pro1* null mutant strain was found to be sensitive to freeze-thaw stress (18).

In this study we examine how proline biosynthesis influences endoplasmic reticulum (ER) stress response in yeast *S. cerevisiae*. The ER is an important organelle for folding and assembling secretory and membrane proteins. The process for protein folding is highly regulated with disruption of protein assembly causing accumulation of misfolded proteins and ER stress. ER stress has been linked to several human diseases such as type II diabetes, neurodegeneration, heart disease, and cancer (19). ER stress is often accompanied by oxidative stress and eventual apoptosis, which is one of the causes of ER stress-induced cell death. In responding to ER stress, intracellular signaling pathways such as the unfolded protein response (UPR) are evoked to help cells adapt (20). In yeast, there are

several pathways involved in ER stress response including the much studied UPR pathway comprising the protein kinase Ire1p and its downstream transcription factor Hac1p, the ER-associated protein degradation pathway (21), and the calcium signal pathway (22, 23). Recently it has been reported that the high osmolarity glycerol (HOG) mitogen-activated protein kinase (MAPK) pathway also has an important role in ER stress survival (24, 25), suggesting glycerol is an important ER stress protectant.

Although there have been several studies of proline protection against various types of environmental stress, very little is known about the role of proline in ER stress response. Previously, a *pro1* mutant was identified by screening a null-mutant library of *S. cerevisiae* for sensitivity to dithiothreitol (DTT), a reducing agent that causes ER stress (26). In addition, proline has been shown to inhibit protein aggregation *in vitro* and *in*

## Proline Biosynthesis in Endoplasmic Reticulum Stress

*in vivo* (27–29). Here, we test the effect of genetic limitation of proline on ER stress tolerance in the *S. cerevisiae* null mutants *pro1*, *pro2*, and *pro3*. The proline biosynthesis mutants are shown to be more sensitive to ER stress-inducing agents than the wild-type strain. Insights into the mechanism by which genetic depletion of proline biosynthesis affects ER stress tolerance were gained by investigating intracellular redox markers and the UPR. The protective role of proline in ER stress was also examined more broadly by exploring whether proline biosynthesis can rescue the ER stress phenotype of *ire1* and *hog1* null mutants.

### EXPERIMENTAL PROCEDURES

**Chemicals, Yeast Strains, and Culture Conditions**—All chemicals, culture media, and buffers were purchased from Fisher and Sigma unless otherwise indicated. The *S. cerevisiae* strains used in this study are provided in Table 1. Gene knock-out mutant strains and strains for expression of GFP C-terminal fusion proteins were constructed using PCR-mediated gene

**TABLE 1**  
Strains used in this study

Strain	Genotype	Source
BY4741	MATa <i>his3Δ1 leu2Δ0 met15Δ0 ura3Δ0</i>	Open Biosystems
A143	As BY4741, but <i>pro1::His3</i>	This study
A18	As BY4741, but <i>pro2::KanMX</i>	Open Biosystems
A98	As BY4741, but <i>pro3::His3</i>	This study
A2	As BY4741, but <i>put1::KanMX</i>	Open Biosystems
A64	As BY4741, but <i>ire1::KanMX</i>	Open Biosystems
A65	As BY4741, but <i>hog1::KanMX</i>	Open Biosystems
A145	As BY4741, but <i>pro2::KanMX pro1::His3</i>	This study
A142	As BY4741, but <i>pro2::KanMX pro3::His3</i>	This study
A140	As BY4741, but <i>pro2::KanMX put1::His3</i>	This study
A116	As BY4741, but <i>put1::KanMX pro3::His3</i>	This study
A44	As BY4741, but <i>PRO1-GFP:KanMX6</i>	This study
A46	As BY4741, but <i>PRO2-GFP:KanMX6</i>	This study
A48	As BY4741, but <i>PRO3-GFP:KanMX6</i>	This study

**TABLE 2**  
Primers used in this study

F, forward primer; R, reverse primer.

Primer ID	Sequence (5' to 3')	Application
DB-96-F	tataggatccatgaaggatgctaataagagag	<i>PRO1-GFP</i> genotyping
DB-102-F	gcgaatatgtcgctcatagagaaaatttggcattcccaccctcgtcggatccccgggttaattaa	<i>GFP</i> tagging for <i>PRO1</i>
DB-103-R	taagaaatatacattaaaggaatcctctataactttttagttcgttgaattcgagctcgtttaaac	<i>GFP</i> tagging for <i>PRO1</i> , <i>PRO1</i> knock out
DB-116-F	ggtcattttatcagctactgtaataatagtcataagaacgcgaaacggatccccgggttaattaa	<i>PRO1</i> knock out
DB-174-F	ttgagcaccagttaaagatgaaca	<i>PRO1</i> knockout genotyping
DB-98-F	tataggatccatgtccagttcacaacaaat	<i>PRO2</i> cDNA synthesis, <i>PRO2-GFP</i> genotyping
DB-99-R	tcaactcgagtcgaagcgtaatctggaacatcgtatgggtataatgtcacagctctttatat	<i>PRO2</i> cDNA synthesis
DB-120-F	caagactttgttcacaaggatttagatataaagactgtgacattacggatccccgggttaattaa	<i>GFP</i> tagging for <i>PRO2</i>
DB-121-R	gtatacacgctccttcatacatggaacttagcttatacctaaatattgaattcgagctcgtttaaac	<i>GFP</i> tagging for <i>PRO2</i>
DB-100-F	tataggatccatgacttacacattggcaatt	<i>PRO3</i> cDNA synthesis, <i>PRO3-GFP</i> genotyping
DB-101-R	tcaactcgagtcgaagcgtaatctggaacatcgtatgggtatttctctctttggcctaatt	<i>PRO3</i> cDNA synthesis
DB-122-F	gaagaggcagcccgtgttcgctcacaatttaggccaaaagaacggatccccgggttaattaa	<i>GFP</i> tagging for <i>PRO3</i>
DB-105-R	tataccgctacaagcgaacatacatacatttcacgcaaacgaatggaattcgagctcgtttaaac	<i>GFP</i> tagging for <i>PRO3</i> , <i>PRO3</i> knock out
DB-104-F	aatagaagtaacacaacaccaatagcaaacacaaactacagatcggatccccgggttaattaa	<i>PRO3</i> knock out
DB-175-F	ataaagtcgcatacgtacaaaagga	<i>PRO3</i> knockout genotyping
DB-173-R	gtcgattacagcaaatggctc	<i>GFP</i> fusion mutant and gene knockout mutant genotyping
DB-158-F	agaaggcctccaagggtatt	<i>KAR2</i> mRNA levels
DB-159-R	taagacaccagcttgaacg	<i>KAR2</i> mRNA levels
DB-176-F	caaggcatgatgcagatttc	<i>SLT2</i> mRNA levels
DB-177-R	ctagaagcgtgcccgttatca	<i>SLT2</i> mRNA levels
DB-182-F	ccttaaacgaaggcgaagag	<i>GSC2</i> mRNA levels
DB-183-R	gggacgtctacgaccattct	<i>GSC2</i> mRNA levels
DB-156-F	ccaagatggtcgttggaaatg	<i>GSH1</i> mRNA levels
DB-157-R	cctccaaatccgcttcttgg	<i>GSH1</i> mRNA levels
DB-140-F	atttggccggtagagatttg	<i>ACT1</i> mRNA levels
DB-141-R	tccaagcgcagcgaacatag	<i>ACT1</i> mRNA levels
DB-223-F	ttggcagaccactctg	<i>HAC1i</i> mRNA levels
DB-225-R	ctgcgctctcggattacg	<i>HAC1i</i> mRNA levels

knock-out or C-terminal tagging techniques (30). The gene knock-out strain was constructed by replacing the corresponding gene ORF with the HIS3MX6 cassette, and the mutant for expression of GFP C-terminal fusion protein was constructed by fusing the GFP(S65T)-kanMX6 cassette to the corresponding C terminus of the gene. Each mutant was confirmed by genotyping using the corresponding primers provided in Table 2. Non-proline auxotrophs were maintained in YPD medium (2% (w/v) D-glucose, 2% (w/v) bactopectone, 1% (w/v) yeast extract) or SD medium (0.17% (w/v) yeast nitrogen base without amino acids, 0.5% (w/v) ammonium sulfate, 2% (w/v) D-glucose) supplemented unless otherwise indicated with leucine (100 mg/liter), methionine (20 mg/liter), histidine (20 mg/liter), and uracil (20 mg/liter). Proline auxotrophs were maintained in SD medium supplemented with varying concentrations of proline (10–20 mM). Growth profiles of the different strains were followed by measuring the optical density at 600 nm ( $A_{600}$ ) with log phase considered to be at  $A_{600}$  of 0.6–1.0 and stationary phase at  $A_{600} \geq 3.0$ . Cells were also cultured anaerobically using a deoxygenated jar with an oxygen scrubbing catalyst (BD Biosciences) and an anaerobic indicator (BD Biosciences). Cells were grown anaerobically on SD medium containing 20 mg/liter ergosterol and 0.05% Tween 80 at 30 °C for 3 days.

**Overexpression of *PRO1*(D154N), *PRO2*, and *PRO3* in Yeast Cells**—The cDNA of *PRO1*, *PRO2*, and *PRO3* were synthesized from RNA extracted from wild-type BY4741 strain and amplified by PCR using corresponding primers (see Table 2 for a list of primers; a HA tag was introduced by the reversed primer). The resulting PCR products were verified by sequencing and cloned into p413TEF (*PRO1*, *PRO2*) or p416GPD (*PRO3*) vectors at BamH1 and Xho1 sites. The *PRO1*(D154N) construct was made by site-directed mutagenesis (Stratagene site-directed mutagenesis kit) of p413TEF-*PRO1*. All the constructs

were confirmed by DNA sequencing (Eurofins MWG Operon) and transformed into their respective yeast mutant strain via the LiAc-based assay. Expression of each protein was indicated by complementation of the corresponding proline auxotroph strain.

**Spot Assays**—Cells were grown overnight to stationary phase in SD medium alone except for the *pro1*, *pro2*, and *pro3* mutant strains, which were grown in SD medium supplemented with 10 mM proline. Cells were then pelleted and resuspended in deionized water to an  $A_{600}$  of 2.0 and were serially diluted (1:10) with deionized water. Cells were spotted on the indicated culture plates using a replica plater and grown 2–3 days at 30 °C.

**Determination of Intracellular Superoxide, Glutathione, and Proline**—Superoxide anion radicals were estimated with dihydroethidium (Invitrogen) using a flow cytometer, BD Biosciences FACSCanto II, with a blue laser (488 nm) at PE channel (585/42, 556LP). Cells were incubated with SD medium supplemented with 10  $\mu$ M dihydroethidium at 30 °C for 15 min, washed once with PBS, and resuspended in 500  $\mu$ l of PBS and analyzed by fluorescence detection using flow cytometry.

Intracellular free glutathione was extracted as described previously (31). Briefly, cells were grown in SD minimal medium with or without proline to log phase before the indicated treatments. Yeast cells were washed twice with PBS and resuspended in ice-cold HCl and 1.3% (w/v) 5-sulfosalicylic acid. Cells were broken with acid-washed glass beads using 3 cycles of vortexing (1 min) and incubation on ice (1 min). The broken cells were then incubated on ice for an additional 15 min. Cell debris and proteins were pelleted by centrifugation at  $15,000 \times g$  for 15 min at 4 °C. The supernatant was used for measuring free glutathione. The reduced and oxidized forms of glutathione were determined as described previously (32).

Intracellular free proline was measured as described using the ninhydrin assay (33). Briefly, about 5 ml of log phase grown cells were spun down, washed with 0.9% NaCl, and resuspended in 0.5 ml of distilled water. The cells were transferred to a boiling water bath, and intracellular free amino acids were extracted by boiling for 10 min followed by centrifugation at  $15,000 \times g$  for 5 min. The concentration of proline was determined in the resulting extract.

**Enzyme Assays**—Glutathione reductase (GLR) activity was measured as described (34) but with a slight modification. Cells were broken with acid-washed glass beads in lysis buffer (0.1 M potassium phosphate buffer, pH 7.5, 1 mM EDTA) as described. Cell debris was then pelleted by centrifugation at  $15,000 \times g$  for 15 min at 4 °C. The GLR activity was measured in the supernatant, and the total protein concentration was measured with the Bradford assay. One unit of GLR activity is the amount of enzyme that oxidizes 1  $\mu$ mol of NADPH/min at 25 °C (pH 7.5).

**Determination of Intracellular NADP(H)**—NADP(H) was extracted with base and acid for NADPH and NADP<sup>+</sup> measurements, respectively, as described but with a slight modification (35). Briefly, about 10 absorbance units of yeast cells were resuspended either in 0.1 M NaOH or 0.1 M HCl. Cells were broken with glass beads as described above for glutathione. After centrifuging the broken cells at  $15,000 \times g$  for 10 min at 4 °C, the resulting supernatant was heated at 60 °C for 10 min. The extract was then neutralized with an equal volume of 0.1 M

HCl or 0.1 M NaOH followed by NADP(H) measurement via a spectrophotometric enzymatic cycling assay as described (36).

**Real-time PCR**—Total RNA was extracted from yeast cells using the RNeasy mini kit (Qiagen). Cells were grown to log phase before the indicated treatments. The extracted RNA was treated with RNase-free DNase (Fermentas) before cDNA synthesis with M-MuLV reverse transcriptase and oligo(dT) primers (Fermentas). mRNA levels were quantified by real-time PCR using the IQ-SYBR-super mix (Bio-Rad) and MyIQ real-time PCR detection system (Bio-Rad). Relative mRNA levels were calculated using the  $2^{-\Delta\Delta CT}$  method and the actin gene *ACT1* as the internal control. PCR products were also analyzed by 2% agarose gel electrophoresis to confirm product size and specificity. Spliced *HAC1* (*HAC1i*) mRNA was quantified by real-time PCR using the reverse primer designed to span the exon-exon junction (Table 2) (37). The specificity of the *HAC1i* PCR product was confirmed by control assays in *ire1* mutant cells, which lack *HAC1* splicing due to the absence of *Ire1*, the nuclease responsible for cleaving *HAC1* mRNA during ER stress (38).


**Cell Viability Assays by Propidium Iodide (PI) Staining**—Yeast cells were treated as indicated, spun down, and resuspended in 1 ml of PBS containing 3  $\mu$ g/ml PI for 10 min followed by flow cytometry analysis of the PI-stained cells. Cell death rate is the percentage of PI-positive cells per 10,000 cells counted.

## RESULTS

**Localization of the Proline Biosynthesis Enzymes and Growth Profiles of Null Mutants**—The intracellular localization of Pro1p, Pro2p, and Pro3p was determined by making GFP fusion proteins driven by the native promoter of the corresponding gene. All three enzymes were previously reported to be in the cytosol, but the evidence was indirect such as complementation of the *pro1* mutant with the *proB* gene from *Escherichia coli* (39) and isolation of Pro3p from the cytosolic fraction (40). In agreement with previous findings, Pro1p, Pro2p, and Pro3p were expressed exclusively in the cytosol (Fig. 1B). Next, the growth profiles of the *pro1*, *pro2*, and *pro3* null mutants were characterized in SD medium supplemented with increasing amounts of proline. Without proline, the *pro1* and *pro2* mutants exhibited growth, whereas *pro3* did not grow (Fig. 2A, top panel). These results are consistent with arginine catabolism supplying P5C for *PRO3* in *pro1* and *pro2* cells and a strict requirement for proline in *pro3* cells (41). The growth profiles of the *pro1*, *pro2*, and *pro3* mutants were restored to that of the parent wild-type strain at 4, 20, and 4 mM proline, respectively (Fig. 1C). Thus, the *pro2* mutant requires the highest proline concentration for growth in SD medium. A similar growth dependence on proline was observed on SD plates (Fig. 2A, top panel).

**Proline Biosynthesis Mutants Are Sensitive to ER Stress**—The ER stress sensitivities of the *pro1*, *pro2*, and *pro3* mutants and the proline catabolic *put1* mutant were tested by growing each strain to stationary phase and spotting on SD plates containing different concentrations of proline (0, 2, 4, 10, and 20 mM) and DTT or tunicamycin (Tm). DTT and Tm are chemicals used commonly for inducing ER stress (42). Both chemicals induce ER stress but do so via different mechanisms. DTT induces ER

## Proline Biosynthesis in Endoplasmic Reticulum Stress



**FIGURE 2. Proline biosynthesis mutants are sensitive to ER stress.** *A*, growth of BY4741 WT cells and null mutants *put1*, *pro1*, *pro2*, and *pro3* on SD plates with increasing proline concentrations (0–20 mM) in the absence and presence of Tm (0.5  $\mu$ g/ml) or DTT (10 mM). *B*, growth of WT cells and null mutants *pro2* and *pro3* harboring empty vector (V) or construct for expression of recombinant *PRO2* or *PRO3*. *C*, growth of wild-type cells on SD medium with histidine, leucine, and uracil concentrations reduced to 1/10 of normal concentrations. Cells were spotted on SD plates in the absence and presence of Tm (0.5  $\mu$ g/ml) or DTT (10 mM).

stress by reducing thiols and disrupting protein disulfide bond formation (43, 44), whereas Tm blocks *N*-linked glycosylation leading to accumulation of misfolded proteins (35, 45).

All of the proline biosynthesis mutants were more sensitive to Tm and DTT than the wild-type strain at 2 and 4 mM proline (Fig. 2*A*). The strongest Tm stress phenotype was observed for the *pro3* mutant, whereas *pro2* cells appeared most sensitive to DTT. The *pro1* mutant was least sensitive to DTT/Tm relative to *pro2* and *pro3*. Increasing proline to 10 mM rescued the Tm sensitivity of all the mutants. With DTT, 10 mM proline protected the *pro1* and *pro3* mutants, whereas 20 mM proline was needed to fully rescue the *pro2* mutant. In contrast to the proline biosynthesis mutants, disruption of proline catabolism in the *put1* null mutant had only a minimal effect on sensitivity to Tm and DTT (Fig. 2*A*).

The results shown in Fig. 2*A* indicate that the *pro2* mutant has the highest sensitivity to DTT, whereas *pro3* is most sensitive to Tm. To confirm that the sensitivity of these mutants was due to the loss of functional *PRO2* and *PRO3* genes, the *pro2* and *pro3* mutants were transformed with recombinant *PRO2* and *PRO3*, respectively. Overexpression of *PRO2* and *PRO3* in the corresponding mutants rescued growth in SD medium without proline and increased resistance to DTT and Tm (Fig. 2*B*). These results show that the ER stress-sensitive phenotype of *pro2* and *pro3* is due to the knock-out of *PRO2* and *PRO3*, respectively, and suggest that proline biosynthesis helps yeast cells survive ER stress under limiting proline conditions.

One mechanism by which cells adapt to ER stress is to down-regulate global protein synthesis, which in yeast occurs during amino acid starvation (46). How limiting other amino acids influences ER stress tolerance was examined in the BY4741 wild-type strain, which is an auxotroph for histidine, leucine, and uracil. These supplements were simply reduced to 1/10 of the normal amount in SD medium (2 mg/liter histidine, 10 mg/liter leucine, and 2 mg/liter uracil). Fig. 2*C* shows that in contrast to proline, wild-type cells grown under limiting conditions for histidine, leucine, and uracil do not show increased sensitivity to


DTT and Tm. In fact, the Tm stress resistance of wild-type cells appears to be slightly improved by limiting histidine and leucine, consistent with lower protein synthesis benefiting cells during ER stress. Thus, the ER stress sensitivity of the proline biosynthesis mutants appears to contrast that of general amino acid limitation.

*N-Acetyl-L-cysteine and Anaerobic Conditions Rescue the pro3 Mutant*—Because Tm has been shown previously to increase oxidative stress in yeast (35), we examined if *N*-acetyl-L-cysteine (NAC) and reduced glutathione (GSH) could mitigate the Tm sensitivity of *pro3*. Fig. 3*A* shows that both NAC and GSH rescue the Tm sensitive-phenotype of *pro3*, whereas methionine has no effect. The effects of NAC and GSH are similar to proline. Other chemicals with known antioxidant properties such as butylated hydroxyanisole (47) and diphenyleiodonium (an NADPH-oxidase inhibitor) (48) did not rescue the Tm sensitivity of *pro3* (Fig. 3*A*). These results suggest that GSH is important for repairing ER-associated oxidative stress such as helping to resolve non-native disulfide bonds formed as a result of oxidative stress or enzyme-catalyzed oxidation (49).

Additional evidence for oxidative stress contributing to the ER sensitivity of *pro3* was obtained by treating the *pro3* mutant with Tm under anaerobic conditions. Fig. 3*B* shows that *pro3* is less sensitive to Tm under anaerobic conditions. Wild-type and *ire1* mutant cells showed no significant change to Tm treatment when switched to anaerobic growth (Fig. 3*B*).

ROS levels in wild-type and *pro3* were then measured using dihydroethidium (50). Under non-stressed conditions, the ROS levels in *pro3* (2 mM proline) were 1.5-fold higher than wild type (Fig. 3*C*). Increasing proline in the SD medium to 4 mM lowered ROS levels in *pro3* to that of wild type. Upon treating cells with Tm, ROS levels increased by 1.5-fold in wild-type and *pro3* (4 mM proline). At 2 mM proline, ROS levels did not further increase in *pro3* cells with Tm treatment (Fig. 3*C*).

Because GSH reversed the ER stress sensitivity of the *pro3* mutant, we assessed GSH status in *pro3*. GSH levels were mea-




**FIGURE 3. ER stress protection of *pro3* by reducing agents and anaerobic conditions.** *A*, spot assays of WT and *pro3* cells grown on SD plates with 4 mM proline alone (SD) or with 4 mM proline in the presence of Tm (0.5  $\mu$ g/ml), Tm and *N*-acetyl-L-cysteine (Tm + NAC), Tm and GSH (Tm + GSH), Tm and methionine (Tm + Met), Tm and butylated hydroxyanisole (Tm + BHA), and Tm and diphenyleneiodonium (Tm + DPI). *B*, spot assays of WT, *pro3*, and *ire1* cells on SD medium with 4 mM proline alone (SD) or in the presence of Tm (0.5  $\mu$ g/ml) under aerobic (top panel) and anaerobic conditions (bottom panel) at 30 °C for 3 days. *C*, relative superoxide levels in WT (no proline) and *pro3* (2 and 4 mM proline) cells in SD medium treated with (+Tm, 1  $\mu$ g/ml) or without (–Tm) for 4 h. Superoxide was measured by ROS probe dihydroethidium bromide using flow cytometry and normalized to the lowest level. Data are the mean  $\pm$  S.D. from three independent experiments (\*,  $p < 0.05$ ).

sured in wild-type and *pro3* cells grown to log phase with different proline concentrations (2, 4, and 10 mM). Total GSH was 3-fold higher in *pro3* with 2 mM proline relative to wild type (Fig. 4A). At higher concentrations of proline (4 and 10 mM) GSH levels were similar to wild type (Fig. 4A). In all conditions Tm did not change total GSH levels in wild type and *pro3* (Fig. 4A). The ratio of oxidized to reduced glutathione (GSSG/GSH) was shown to be similar between wild type (WT) and *pro3* with no significant change upon treatment with Tm (Fig. 4B). We also measured the mRNA levels of  $\gamma$ -glutamylcysteine synthetase 1 (*GSH1*), the rate-limiting enzyme of GSH biosynthesis. Under growth conditions with 2 mM proline, the *GSH1* transcript levels were  $\sim$ 2-fold higher in *pro3* relative to wild-type cells (Fig. 4C), indicating that increased *GSH1* expression may be the cause of higher glutathione in *pro3*. We also observed that after Tm treatment, *GSH1* increased by 3- and 5-fold in wild type and *pro3*, respectively (Fig. 4C), indicating that GSH is required for cells recovering from ER stress. Interestingly, *GSH1* levels remained nearly 2-fold higher in *pro3* relative to wild type after Tm treatment. GSH metabolism was explored further by measuring the activity of GLR, which catalyzes the NADPH-dependent reduction of GSSG to 2GSH. GLR activity was observed to be 4-fold higher in *pro3* at 2 mM proline than in wild-type cells (Fig. 4D). At higher concentrations of proline, GLR activity in *pro3* was similar to wild-type cells. No change in GLR activity was observed with Tm treatment in wild type or *pro3*. We last measured NADP<sup>+</sup> and NADPH and found significantly higher NADP<sup>+</sup> and a NADP<sup>+</sup>/NADPH ratio in *pro3* at 2 mM proline relative to wild type (Fig. 4, E and F). At higher proline concentrations (4 and 10 mM proline) no significant difference in NADP<sup>+</sup> and NADPH was observed between wild-type and the *pro3* mutant.

**Oxidative Stress and the UPR**—To test whether higher ROS levels as observed in the *pro3* mutant (Fig. 3C) exacerbate Tm sensitivity, wild-type cells were treated with Tm in the presence of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>). The addition of H<sub>2</sub>O<sub>2</sub> (1 mM) during Tm treatment significantly decreased cell survival, indicating a compounding effect of oxidative stress on ER stress (Fig. 5A). H<sub>2</sub>O<sub>2</sub> alone did not diminish cell survival (Fig. 5A). The addition of NAC dramatically rescued cells from Tm and H<sub>2</sub>O<sub>2</sub> toxicity. Interestingly, the addition of 10 mM proline also partially repressed the combined toxicity of Tm and H<sub>2</sub>O<sub>2</sub> in wild-type cells (Fig. 5A). The sensitivity of the *pro3* mutant strain to Tm was also enhanced by H<sub>2</sub>O<sub>2</sub> but at a lower concentration (0.25 mM H<sub>2</sub>O<sub>2</sub>) (Fig. 5B). As in wild-type cells, NAC prevented cell death induced by Tm and H<sub>2</sub>O<sub>2</sub> treatment (Fig. 5B), consistent with proline limitation, resulting in higher intracellular oxidative stress. Increasing the proline concentration from 4 to 10 mM also rescued the *pro3* cells from Tm and H<sub>2</sub>O<sub>2</sub> toxicity (Fig. 5B). Note that at 0.25 mM, H<sub>2</sub>O<sub>2</sub> does not increase the Tm sensitivity of wild-type cells (Fig. 5B). The role of cellular antioxidants in ER stress defense was further evaluated by testing the sensitivity of the *yap1* mutant to Tm or DTT. Fig. 5C shows *yap1* cells exhibit significantly increased sensitivity to Tm and DTT relative to wild-type cells.

In yeast and mammalian cells, a functional UPR is essential for cell survival under ER stress (21). To test whether oxidative stress impacts the UPR, *HAC1* mRNA splicing (37) was examined via real-time PCR after Tm treatment with and without H<sub>2</sub>O<sub>2</sub>. As expected, *HAC1i* (spliced *HAC1*) levels increased ( $\sim$ 3-fold) in wild-type cells after Tm treatment (Fig. 5D), indicating induction of the UPR. In the presence of H<sub>2</sub>O<sub>2</sub> (1 mM), Tm induction of *HAC1i* mRNA splicing was significantly reduced. A dose-dependent inhibition of the UPR was also

## Proline Biosynthesis in Endoplasmic Reticulum Stress



**FIGURE 4. Proline limitation results in redox imbalance in *pro3*.** WT cells and *pro3* cells were grown to log phase in SD medium with varying concentrations of proline (0–10 mM) as indicated and treated with (+Tm, 1  $\mu$ g/ml) or without Tm (–Tm) for 2 h. The following parameters were then determined. *A*, total intracellular glutathione. *B*, ratio of oxidized (GSSG) to reduced glutathione (GSH). *C*, relative *GSH1* mRNA levels (4 h of Tm treatment) determined by real-time PCR with mRNA levels normalized to the internal control *ACT1* and the lowest expression set at 1. *D*, GLR activity. The concentrations of NADP<sup>+</sup> and NADPH (*E*) and the corresponding ratio of NADP<sup>+</sup>/NADPH (*F*) were determined in WT and *pro3* cells grown in SD medium as above but without Tm treatment. Data are the mean  $\pm$  S.D. from three independent experiments (\*,  $p < 0.05$ ).

observed by following the ER stress-inducible UPR-target gene *KAR2* (38). *KAR2* mRNA levels were reduced by nearly 4-fold with H<sub>2</sub>O<sub>2</sub> relative to Tm treatment alone (Fig. 5D).

The effect of H<sub>2</sub>O<sub>2</sub> on the time course (0.5–2 h) of the UPR was also examined. At each time point H<sub>2</sub>O<sub>2</sub> significantly blocked the UPR as determined by *HAC1i* (Fig. 5E) and *KAR2* (Fig. 5F) mRNA levels. The addition of NAC partially rescued the UPR with 2–3-fold higher levels of *HAC1i* and *KAR2* (Fig. 5, E and F) at each time point relative to cells treated with Tm and H<sub>2</sub>O<sub>2</sub>. Treatment of cells with H<sub>2</sub>O<sub>2</sub> alone resulted in 2-fold lower *HAC1i* mRNA levels (Fig. 5E) by 2 h and a ~3-fold increase in *KAR2* mRNA (Fig. 5F) by 1 h.

**Proline Limitation Inhibits the UPR**—To test if proline limitation impacts the UPR, *HAC1i* and *KAR2* mRNA levels were examined in *pro3* mutant cells grown in SD medium supplemented with increased concentrations of proline. With 2 mM proline, *pro3* cells exhibited a significantly lower induction of *HAC1i* (Fig. 6A) and *KAR2* (Fig. 6B) mRNA by Tm relative to wild-type cells. Increasing the proline concentration (4 and 10 mM) rescued the Tm-dependent UPR as observed by higher *HAC1i* and *KAR2* mRNA levels (Fig. 6, A and B). NAC partially rescued *HAC1i* mRNA levels in *pro3* cells at 2 mM proline (Fig. 6C) similar to that observed with wild-type cells treated simultaneously with Tm and H<sub>2</sub>O<sub>2</sub> (Fig. 5E). For comparison, other conditions of nutrient starvation (1/10 of normal concentration in SD medium) such as leucine and uracil were tested for effects on the UPR. Imposing leucine or uracil starvation on wild-type cells results in a significant drop in *HAC1i* mRNA to nearly

undetectable levels, similar to that observed in the *ire1* mutant strain (Fig. 6A). Also, no increase in *HAC1i* mRNA was observed with Tm stress (data not shown). However, limiting leucine or uracil did not increase the toxicity of Tm or DTT (Fig. 2C). Therefore, the dampened UPR under leucine or uracil starvation is most likely due to generally lower protein synthesis caused by amino acid starvation and, thus, lower misfolded protein loading in the ER. In contrast, proline limitation does not appear to decrease overall misfolded protein loading, as *HAC1i* mRNA levels are similar between wild-type and *pro3* cells before Tm treatment. This implies that proline limitation results in higher misfolded protein levels relative to leucine or uracil, suggesting that proline-depleted cells are under ER stress before Tm treatment.

Because ER stress activates the calcium signaling pathway as well as the UPR (22, 23), we tested the possibility of impaired calcium import in *pro3*. Exogenous proline was previously reported to enhance calcium import in plants (51), indicating proline levels may influence calcium signaling. Calcium import, however, was observed not to be affected in the *pro3* mutant as determined by monitoring expression levels of *SLT2* (*MPK1*), a protein kinase that is critical for calcium homeostasis and is up-regulated by Tm (22) and *GSC2*, a gene known to be up-regulated by increased calcium levels (52) (data not shown).

**Proline Biosynthesis Gene Expression in Response to Tm Treatment**—To determine whether ER stress influences the expression of proline biosynthesis genes, we examined protein expression levels of Pro1p, Pro2p, and Pro3p after Tm treat-






FIGURE 5. Oxidative stress increases Tm toxicity and inhibits the UPR. A–C, spot assays of wild-type (A), wild-type and *pro3* (B), and wild-type and *yap1* cells (C) on SD plates with (+) or without (–) the indicated supplements. D–F, real-time PCR quantification of spliced *HAC1* (*HAC1i*) or *KAR2* mRNA in BY4741 wild-type cells grown to log phase in SD medium treated with the following. D, without (–) or with (+) Tm (1 μg/ml) in the presence of increasing H<sub>2</sub>O<sub>2</sub> (0–1 mM) for 2 h; E and F) Tm (1 μg/ml) alone (*Tm*), Tm and 1 mM H<sub>2</sub>O<sub>2</sub> (*Tm + H<sub>2</sub>O<sub>2</sub>*), Tm, 1 mM H<sub>2</sub>O<sub>2</sub> and 1 mM NAC (*Tm + H<sub>2</sub>O<sub>2</sub> + NAC*), and 1 mM H<sub>2</sub>O<sub>2</sub> alone (*H<sub>2</sub>O<sub>2</sub>*) for the indicated hours (h). mRNA levels were normalized to the internal control *ACT1* with the lowest expression level set at 1. \*, *p* < 0.05 compared with Tm + H<sub>2</sub>O<sub>2</sub> at the corresponding time point. Data are the mean ± S.D. from three independent experiments.

ment. The fluorescence intensity of Pro1p-GFP, Pro2p-GFP, and Pro3p-GFP fusion proteins with expression of each gene under control of the corresponding native promoter was measured using flow cytometry. No significant change in the fluorescence of the GFP fusion proteins was observed after 2 and 4 h of Tm treatment (data not shown). Consistent with no change in proline biosynthesis gene expression, no significant change in proline content was observed after Tm treatment (data not shown).

*Intermediates in the Proline Metabolic Pathway Do Not Contribute to ER Stress Sensitivity*—The ER stress sensitivity of the *pro2* and *pro3* mutants may also be impacted by intermediate metabolites in the proline biosynthesis pathway. In the first step Pro1p converts glutamate to  $\gamma$ -glutamyl phosphate, which is unstable and undergoes cyclization to form 5-oxoproline, which may increase in *pro2* (53). The second intermediate of proline biosynthesis is P5C/GSA, which could increase in the *pro3* mutant and is thought to be toxic to yeast and induce apoptotic cell death (2). Because proline is a feedback inhibitor of Pro1p (54), supplementing the *pro2* and *pro3* strains with proline may reduce buildup of these potentially harmful intermediates. To test the effects of the intermediates  $\gamma$ -glutamyl phosphate and P5C/GSA, double mutants *pro1/pro2* and *pro2/pro3* were tested for sensitivity to DTT and Tm, respectively. Fig. 7 shows that the sensitivity of the double mutants *pro1/pro2* and *pro2/pro3* to DTT and Tm, respectively, is similar to that observed with the single *pro2* and *pro3* mutants. In addition,

the stress phenotype of the double mutants was rescued by proline. Thus, the intermediates of proline biosynthesis do not appear to contribute significantly to the observed ER stress phenotypes of *pro2* and *pro3*.

It is also possible that the breakdown of proline by Put1p would accentuate problems with inadequate proline levels in the *pro2* and *pro3* mutant strains. However, knocking out the *PUT1* gene in the double mutants *pro2/put1* and *pro3/put1* did not rescue the sensitivity of the *pro2* and *pro3* mutants to DTT and Tm, respectively (Fig. 7). Instead, the double mutant *pro3/put1* appeared to be more sensitive to Tm but was still rescued by proline.

*Proline Protects hog1 and ire1 Null Mutants against ER Stress*—Finally, we tested whether proline can rescue ER stress-sensitive mutants apart from the proline metabolic pathway. The effect of proline on the ER stress-sensitive mutants *hog1* and *ire1* was examined by overexpressing the mutant Pro1p(D154N) in these mutants. Pro1p is feedback-inhibited by proline, thereby limiting accumulation of proline in the cell. Previously, it was shown that Pro1p(D154N) is less sensitive to feedback inhibition by proline with overexpression of Pro1p(D154N), resulting in higher intracellular proline levels (17). We used the same strategy here to increase intracellular proline levels in *hog1* and *ire1* cells. Cells overexpressing the Pro1p(D154N) mutant were determined to have 3-fold higher intracellular proline levels than control cells with the empty vector. Interestingly, overexpression of Pro1p(D154N) in *hog1*

## Proline Biosynthesis in Endoplasmic Reticulum Stress

and *ire1* increased survival of these mutants under DTT or Tm stress (Fig. 8A and B). The effect of Pro1p(D154N) was most evident in the *hog1* mutant, as rescue of *ire1* was only observed at lower concentrations of DTT and Tm (Fig. 8).


Proline protection of *hog1* was further tested by determining the death rate of the *hog1* mutant cells after the addition of 2 mM DTT or 1  $\mu$ g/ml Tm in liquid SD medium containing 0, 10, or 20 mM proline for 16 h. The death rate of *hog1* mutant cells treated with DTT or Tm in SD medium was reduced by 60 and

30%, respectively, when supplemented with proline (Fig. 8C). To gain molecular insight into proline protection of the *hog1* mutant, we examined the expression of the ER stress marker *KAR2* in *hog1* and wild-type cells under Tm stress with and without proline (Fig. 8D). In cells without proline, *KAR2* levels remained elevated in the *hog1* mutant much longer than in wild-type cells, showing that the *hog1* mutant recovers more slowly from Tm stress, which is consistent with a previous report (25). In cells supplemented with proline, however, the *KAR2* expression profile in the *hog1* mutant is similar to that of wild type, indicating that proline helps *hog1* recover in the late phase of ER stress response (Fig. 8D).


## DISCUSSION

In this study we found that genetic depletion of individual steps in proline biosynthesis results in distinct growth and ER stress phenotypes. The *pro3* mutant has a strict requirement for proline (41) and showed high sensitivity to ER stress-inducing agents. Proline limitation in yeast *S. cerevisiae*, as demonstrated in the *pro3* mutant, leads to ROS accumulation, higher NADP<sup>+</sup> and total GSH levels, and inhibition of the UPR. Tm sensitivity of the *pro3* mutant was rescued by NAC or GSH treatment and by anaerobic growth conditions, indicating proline limitation negatively impacts antioxidant defense systems and the UPR during ER stress recovery.

Amino acid limitation effectively reduces ER protein loading and is thought to benefit cells during ER stress (55–57). Yeast mutants deficient for ribosomal large subunits or with less cytosolic mRNA were found to be more resistant to Tm, which may be due to lower protein synthesis overall (58, 59). In agreement with this notion, we found that limiting leucine, uracil, and histidine slightly improved tolerance to Tm. In contrast, we found that genetic depletion of proline biosynthesis decreased ER stress tolerance. This implies that proline may have a unique role in ER stress protection. The expression levels of Pro1p, Pro2p, and Pro3p do not change significantly after DTT or Tm treatment. In a yeast gene microarray experiment, only 28 genes were induced by DTT, although 195 DTT stress-sensitive null mutants were identified (26). Thus, a large number of genes that are critical for DTT stress survival are not necessarily induced by DTT (26). During ER stress, proline levels did not change. However, we did detect 4-fold higher *PUT1* mRNA levels after Tm treatment (data not shown), indicating increased proline degradation. Thus, an important role of the



**FIGURE 6. Proline limitation diminishes the UPR response induced by Tm.** A and B, relative spliced *HAC1* (*HAC1i*) mRNA or *KAR2* mRNA levels of log phase cells of BY4741 WT, *ire1*, and *pro3* null mutant grown in SD medium with increasing concentrations of proline treated without (–Tm) or with (+Tm) Tm (1  $\mu$ g/ml) for 2 h. C, relative *HAC1i* mRNA levels of *pro3* grown in SD medium with 2 mM proline treated with Tm alone (Tm) and Tm and 1 mM NAC (Tm + NAC) for 2 h. mRNA levels were normalized to the internal control *ACT1* with the lowest expression level set at 1. \*,  $p < 0.05$  compared with WT or *pro3* with the addition of 10 mM proline treated with Tm (A and B) or as indicated (C). Data are the mean  $\pm$  S.D. from three independent experiments.



**FIGURE 7. Intermediates of the proline biosynthesis pathway do not appear to contribute to ER stress.** A, spot assays of WT, *pro2*, *pro1/pro2*, and *pro2/put1* cells on SD medium with different concentrations of proline as indicated with (DTT) and without (SD) DTT (10 mM). B, spot assays of WT, *pro3*, *pro2/pro3*, and *pro3/put1* cells on SD medium with different concentrations of proline as indicated with (Tm) and without (SD) Tm (0.5  $\mu$ g/ml).




FIGURE 8. Proline protects *hog1* and *ire1* null mutants against ER stress. *A* and *B*, spot assays of yeast WT, *ire1*, and *hog1* cells containing an empty vector (*V*) or the vector expressing the *PRO1* mutant (*PRO1-D154N*) on SD plates alone (*SD*) or with DTT and Tm. *C*, cell death rate of *hog1* cells assayed by PI staining. Mutant cells were grown in SD medium to log phase and were then treated with 2 mM DTT or 1  $\mu$ g/ml Tm in the presence of varying concentrations of proline (0, 10, and 20 mM) for 16 h. Dead cells were then stained with PI. *D*, *KAR2* mRNA levels were analyzed by real-time PCR in WT and *hog1* cells grown in SD medium to log phase without ( $-Pro$ ) or with ( $+Pro$ ) 20 mM proline followed by treatment with Tm (1  $\mu$ g/ml) for 0, 2, and 4 h. The mRNA levels were normalized to the internal control *ACT1* with the lowest expression set at 1 (\*,  $p < 0.05$ ). Data are the mean  $\pm$  S.D. from three independent experiments.

proline biosynthesis pathway may be to maintain intracellular proline levels during ER stress.

Protein misfolding is often connected with oxidative stress and depletion of GSH (60, 61). Glutathione is critical for mitigating ROS generated by the ER during oxidative protein folding (49, 61–64). In mammalian cells, glutathione biosynthesis and assimilation are up-regulated during ER stress via the antioxidant transcription factor Nrf2 (65) and the eIF2 $\alpha$  signaling pathway (55, 66). Disruption of these signaling pathways leads to increased ROS- and ER stress-induced cell death, which can be rescued by NAC or GSH (55, 66). Previously in yeast, ROS was detected in wild-type cells treated with Tm (35) and a yeast ER-associated protein degradation mutant in which misfolded protein was overexpressed (67). GSH supplementation was shown to mitigate ROS accumulation and the ER stress sensitivity of the null ER-associated protein degradation mutant (67). Yeast null mutants of superoxide dismutase (*SOD1*) and *YAP1* have also been shown to be sensitive to Tm treatment (35). Consistent with these earlier results, we also show here that ROS levels and glutathione synthesis increase in wild-type and *pro3* cells after Tm treatment. Thus, our data suggest a role for ROS defense in yeast during ER stress.

Characterization of the *pro3* mutant suggests an underlying mechanism by which proline limitation increases ER stress sensitivity involves disruption of redox homeostasis. At 2 mM proline, a redox imbalance in *pro3* cells is evidenced by increased intracellular ROS levels, an elevated NADP<sup>+</sup>/NADPH ratio (68), and higher expression and activity of *GSH1* and *GLR*, respectively, which are regulated by Yap1p in response to oxidative stress (69–71). Under proline limitation, we also observed higher total glutathione levels, which may be caused

by the increased ROS stimulated GSH synthesis, which has been observed in other yeast mutants defective in oxidative stress defense (72, 73). Because the Tm sensitivity of the *pro3* mutant is reversed by anaerobic growth conditions and by treatment with NAC or GSH, the redox imbalance in *pro3* cells likely heightens ER stress sensitivity. Exactly how proline limitation leads to a more oxidative environment in *pro3* cells is not clear. In addition to nutrient starvation (74–76), ER stress or protein misfolding may contribute to the oxidative stress observed in *pro3* cells. Increasing the proline concentration to 4 mM restored the redox status of the *pro3* mutant to near that of wild-type cells and greatly diminished DTT sensitivity. A higher concentration of proline (10 mM), however, was needed to fully repress the Tm sensitivity of *pro3*.

Oxidative and ER stress coexist in many pathologic states and may trigger one another, but exactly how they interact is not clear (77). Our data show that oxidative stress exacerbates ER stress sensitivity via inhibition of the UPR. The UPR is essential for yeast cells to survive under ER stress resulting in increased chaperone synthesis, inhibition of protein synthesis to reduce ER protein loading, and increased antioxidant defenses (21, 61). We observed that the UPR response to Tm is rapidly inhibited by H<sub>2</sub>O<sub>2</sub> in wild-type cells and by proline limitation in *pro3* cells (2 mM proline), demonstrating oxidative stress profoundly impacts ER stress tolerance. Previous studies in mammalian cells showed that exogenous oxidants such as peroxides and ROS generators can trigger several aspects of the UPR (77). Interestingly in yeast, H<sub>2</sub>O<sub>2</sub> alone does not induce *HAC1* splicing; rather, it inhibits *HAC1* splicing induced by Tm (Fig. 5, *D* and *E*). NAC partially restored the UPR upon Tm in H<sub>2</sub>O<sub>2</sub>-treated wild-type cells and in *pro3* cells under proline limita-

tion. Proline fully rescues the UPR in the *pro3* mutant, which correlates with an improved intracellular redox environment at higher proline concentrations (4 and 10 mM). Thus, one of the mechanisms by which NAC and proline decrease Tm toxicity involves protection of the UPR, likely by removing oxidative stress. Besides the UPR, diminished oxidative stress would also benefit ER stress tolerance by down-regulation of cellular death processes such as apoptosis. Proline has been shown previously to function as an anti-apoptotic factor in yeast (78).

Reports are sparse about how the UPR is inhibited by H<sub>2</sub>O<sub>2</sub>. In a study of the mutant *sod1*, NADP<sup>+</sup> levels increased whereas NADPH levels remained constant during ER stress. It was proposed that increases in superoxide interrupt the overall NADP(H) flux leading to a higher NADP<sup>+</sup>/NADPH ratio and thereby reducing the UPR during ER stress (35). Similarly, we observed a higher NADP<sup>+</sup>/NADPH ratio due to increased NADP<sup>+</sup> in *pro3* cells under proline limiting conditions. Thus, in an analogous manner to that of *sod1*, increased ROS caused by proline limitation may interfere with NADP(H) flux leading to inhibition of the UPR. The increased NADP<sup>+</sup> found in *pro3* cells at 2 mM proline may result from an imbalance of higher GLR activity and diminished flux of the pentose phosphate pathway during proline limitation (68). Proline metabolism was shown previously to be coupled with the pentose phosphate pathway (79), indicating a potential mechanism by which proline impacts intracellular redox homeostasis (2).

In contrast to ROS and NADP<sup>+</sup>, the higher GSH levels detected in *pro3* cells are less likely to inhibit the UPR. It was previously reported that disruption of *TRR1* results in a 3-fold increase in GSH levels, which was proposed to contribute to more misfolded protein and induce the UPR (73). Overexpression of *GSH1* also led to a 30% increase in UPR induction (63). Thus, it is unlikely that higher glutathione levels lead to a reduced UPR in *pro3* cells. However, the higher GSH levels would be expected to exacerbate DTT toxicity as seen for *pro3* and *pro2* cells. Depletion of thioredoxin in the double mutant *trx1/trx2* resulted in a 3-fold increase in total GSH and increased sensitivity to DTT (73), indicating that elevated GSH increases sensitivity to reductive stress in the ER (63). Besides interruption of redox homeostasis, another mechanism by which proline depletion inhibits the UPR may occur via decreased synthesis or destabilization of a protein that is critical to the UPR.

Overall, the results from this study indicate that proline biosynthesis and proline have fundamental roles in ER stress protection. We propose that the major mechanism by which proline biosynthesis protects against ER stress is 2-fold involving maintenance of intracellular redox homeostasis and the UPR. However, other potential roles of proline in ER stress should not be ignored as evidenced by partial rescue of the ER stress-sensitive *hog1* and *ire1* mutant strains with overexpression of the Pro1p(D154N) mutant. The ability of proline to protect *hog1* cells against ER stress indicates that proline can also function as an osmolyte substitute for glycerol in *hog1*. As an established antioxidant and osmolyte, proline may serve as an important stabilizer of redox balance and protein folding during ER stress that benefits overall redox homeostasis, the UPR, and protein folding.

Understanding which ROS defense systems are most critical for combating ER stress will be an important avenue of research in the future. Although the *yap1* mutant is sensitive to ER stress, we have observed that other mutant strains such as *sod2* and *cta1* are only slightly sensitive to Tm-induced ER stress despite being highly sensitive to superoxide and H<sub>2</sub>O<sub>2</sub>, respectively (data not shown). These results indicate that defense against ER stress-induced oxidative stress may rely on specific ROS defense systems. This notion is also supported by our data showing that cells under proline limitation are more sensitive to ER stress than oxidative stress alone (see Fig. 5B). Thus, proline may have a more important role in combating ROS that originates from ER stress rather than oxidative stress in general. Determining how the redox state of the ER is impacted by proline limitation may provide novel insights into the mechanism of proline protection.

---

*Acknowledgments*—We thank Terri Fangman and Dr. You Zhou from the University of Nebraska Microscopy Core Facility for expert advice and technical assistance with cell imaging.

---

## REFERENCES

1. Pérez-Arellano, I., Carmona-Álvarez, F., Gallego, J., and Cervera, J. (2010) Molecular mechanisms modulating glutamate kinase activity. Identification of the proline feedback inhibitor binding site. *J. Mol. Biol.* **404**, 890–901
2. Liang, X., Zhang, L., Natarajan, S. K., and Becker, D. F. (2013) Proline mechanisms of stress survival. *Antioxid. Redox Signal.* **19**, 998–1011
3. Krishnan, N., Dickman, M. B., and Becker, D. F. (2008) Proline modulates the intracellular redox environment and protects mammalian cells against oxidative stress. *Free Radic. Biol. Med.* **44**, 671–681
4. Natarajan, S. K., Zhu, W., Liang, X., Zhang, L., Demers, A. J., Zimmerman, M. C., Simpson, M. A., and Becker, D. F. (2012) Proline dehydrogenase is essential for proline protection against hydrogen peroxide-induced cell death. *Free Radic. Biol. Med.* **53**, 1181–1191
5. Liu, W., Le, A., Hancock, C., Lane, A. N., Dang, C. V., Fan, T. W., and Phang, J. M. (2012) Reprogramming of proline and glutamine metabolism contributes to the proliferative and metabolic responses regulated by oncogenic transcription factor c-MYC. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 8983–8988
6. Zarse, K., Schmeisser, S., Groth, M., Priebe, S., Beuster, G., Kuhlmann, D., Guthke, R., Platzer, M., Kahn, C. R., and Ristow, M. (2012) Impaired insulin/IGF1 signaling extends life span by promoting mitochondrial L-proline catabolism to induce a transient ROS signal. *Cell Metab.* **15**, 451–465
7. Szabados, L., and Savaure, A. (2010) Proline: a multifunctional amino acid. *Trends Plant Sci.* **15**, 89–97
8. Székely, G., Abrahám, E., Csépló, A., Rigó, G., Zsigmond, L., Csiszár, J., Ayaydin, F., Strizhov, N., Jásik, J., Schmelzer, E., Koncz, C., and Szabados, L. (2008) Duplicated P5CS genes of *Arabidopsis* play distinct roles in stress regulation and developmental control of proline biosynthesis. *Plant J.* **53**, 11–28
9. Yildirim, Y., Tolun, A., and Tuysuz, B. (2011) The phenotype caused by PYCR1 mutations corresponds to geroderma osteodysplasticum rather than autosomal recessive cutis laxa type 2. *Am. J. Med. Genet. A* **155**, 134–140
10. Reversade, B., Escande-Beillard, N., Dimopoulou, A., Fischer, B., Chng, S. C., Li, Y., Shboul, M., Tham, P. Y., Kayserili, H., Al-Gazali, L., Shahwan, M., Brancati, F., Lee, H., O'Connor, B. D., Schmidt-von Kegler, M., Merriam, B., Nelson, S. F., Masri, A., Alkazaleh, F., Guerra, D., Ferrari, P., Nanda, A., Rajab, A., Markie, D., Gray, M., Nelson, J., Grix, A., Sommer, A., Savarirayan, R., Janecke, A. R., Steichen, E., Silience, D., Hausser, I., Budde, B., Nürnberg, G., Nürnberg, P., Seemann, P., Kunkel, D., Zambruno, G., Dallapiccola, B., Schuelke, M., Robertson, S., Hamamy, H., Wollnik, B.,

- Van Maldergem, L., Mundlos, S., and Kornak, U. (2009) Mutations in PYCR1 cause cutis laxa with progeroid features. *Nat. Genet.* **41**, 1016–1021
11. Baumgartner, M. R., Hu, C. A., Almashanu, S., Steel, G., Obie, C., Aral, B., Rabier, D., Kamoun, P., Saudubray, J. M., and Valle, D. (2000) Hyperammonemia with reduced ornithine, citrulline, arginine and proline: a new inborn error caused by a mutation in the gene encoding delta(1)-pyrroline-5-carboxylate synthase. *Hum. Mol. Genet.* **9**, 2853–2858
  12. Bicknell, L. S., Pitt, J., Aftimos, S., Ramadas, R., Maw, M. A., and Robertson, S. P. (2008) A missense mutation in ALDH18A1, encoding  $\Delta$ 1-pyrroline-5-carboxylate synthase (P5CS), causes an autosomal recessive neurocutaneous syndrome. *Eur. J. Hum. Genet.* **16**, 1176–1186
  13. Ernst, T., Hergenhahn, M., Kenzelmann, M., Cohen, C. D., Bonrouhi, M., Weninger, A., Klären, R., Gröne, E. F., Wiesel, M., Gudemann, C., Küster, J., Schott, W., Staehler, G., Kretzler, M., Hollstein, M., and Gröne, H. J. (2002) Decrease and gain of gene expression are equally discriminatory markers for prostate carcinoma: a gene expression analysis on total and microdissected prostate tissue. *Am. J. Pathol.* **160**, 2169–2180
  14. Jariwala, U., Prescott, J., Jia, L., Barski, A., Pregizer, S., Cogan, J. P., Arasheben, A., Tilley, W. D., Scher, H. I., Gerald, W. L., Buchanan, G., Coetzee, G. A., and Frenkel, B. (2007) Identification of novel androgen receptor target genes in prostate cancer. *Mol. Cancer* **6**, 39
  15. Okabe, H., Satoh, S., Kato, T., Kitahara, O., Yanagawa, R., Yamaoka, Y., Tsunoda, T., Furukawa, Y., and Nakamura, Y. (2001) Genome-wide analysis of gene expression in human hepatocellular carcinomas using cDNA microarray: identification of genes involved in viral carcinogenesis and tumor progression. *Cancer Res.* **61**, 2129–2137
  16. Nilsson, R., Jain, M., Madhusudhan, N., Sheppard, N. G., Strittmatter, L., Kampf, C., Huang, J., Asplund, A., and Mootha, V. K. (2014) Metabolic enzyme expression highlights a key role for MTHFD2 and the mitochondrial folate pathway in cancer. *Nat. Commun.* **5**, 3128
  17. Takagi, H. (2008) Proline as a stress protectant in yeast: physiological functions, metabolic regulations, and biotechnological applications. *Appl. Microbiol. Biotechnol.* **81**, 211–223
  18. Ando, A., Nakamura, T., Murata, Y., Takagi, H., and Shima, J. (2007) Identification and classification of genes required for tolerance to freeze-thaw stress revealed by genome-wide screening of *Saccharomyces cerevisiae* deletion strains. *FEMS Yeast Res.* **7**, 244–253
  19. Lin, J. H., Walter, P., and Yen, T. S. (2008) Endoplasmic reticulum stress in disease pathogenesis. *Annu. Rev. Pathol.* **3**, 399–425
  20. Tabas, I., and Ron, D. (2011) Integrating the mechanisms of apoptosis induced by endoplasmic reticulum stress. *Nat. Cell Biol.* **13**, 184–190
  21. Patil, C., and Walter, P. (2001) Intracellular signaling from the endoplasmic reticulum to the nucleus: the unfolded protein response in yeast and mammals. *Curr. Opin. Cell Biol.* **13**, 349–355
  22. Bonilla, M., and Cunningham, K. W. (2003) Mitogen-activated protein kinase stimulation of  $Ca^{2+}$  signaling is required for survival of endoplasmic reticulum stress in yeast. *Mol. Biol. Cell* **14**, 4296–4305
  23. Bonilla, M., Nastase, K. K., and Cunningham, K. W. (2002) Essential role of calcineurin in response to endoplasmic reticulum stress. *EMBO J.* **21**, 2343–2353
  24. Torres-Quiroz, F., García-Marqués, S., Coria, R., Rande-Gil, F., and Prieto, J. A. (2010) The activity of yeast Hog1 MAPK is required during endoplasmic reticulum stress induced by tunicamycin exposure. *J. Biol. Chem.* **285**, 20088–20096
  25. Bicknell, A. A., Tourtellotte, J., and Niwa, M. (2010) Late phase of the endoplasmic reticulum stress response pathway is regulated by Hog1 MAP kinase. *J. Biol. Chem.* **285**, 17545–17555
  26. Rand, J. D., and Grant, C. M. (2006) The thioredoxin system protects ribosomes against stress-induced aggregation. *Mol. Biol. Cell* **17**, 387–401
  27. Samuel, D., Kumar, T. K., Ganesh, G., Jayaraman, G., Yang, P. W., Chang, M. M., Trivedi, V. D., Wang, S. L., Hwang, K. C., Chang, D. K., and Yu, C. (2000) Proline inhibits aggregation during protein refolding. *Protein Sci.* **9**, 344–352
  28. Kumar, T. K., Samuel, D., Jayaraman, G., Srimathi, T., and Yu, C. (1998) The role of proline in the prevention of aggregation during protein folding *in vitro*. *Biochem. Mol. Biol. Int.* **46**, 509–517
  29. Ignatova, Z., and Gierasch, L. M. (2006) Inhibition of protein aggregation *in vitro* and *in vivo* by a natural osmoprotectant. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 13357–13361
  30. Longtine, M. S., McKenzie, A., 3rd, Demarini, D. J., Shah, N. G., Wach, A., Brachat, A., Philippsen, P., and Pringle, J. R. (1998) Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast* **14**, 953–961
  31. Grant, C. M., Perrone, G., and Dawes, I. W. (1998) Glutathione and catalase provide overlapping defenses for protection against hydrogen peroxide in the yeast *Saccharomyces cerevisiae*. *Biochem. Biophys. Res. Commun.* **253**, 893–898
  32. Vandeputte, C., Guizon, I., Genestie-Denis, I., Vannier, B., and Lorenzon, G. (1994) A microtiter plate assay for total glutathione and glutathione disulfide contents in cultured/isolated cells: performance study of a new miniaturized protocol. *Cell Biol. Toxicol.* **10**, 415–421
  33. Chen, C., Wanduragala, S., Becker, D. F., and Dickman, M. B. (2006) Tomato QM-like protein protects *Saccharomyces cerevisiae* cells against oxidative stress by regulating intracellular proline levels. *Appl. Environ. Microbiol.* **72**, 4001–4006
  34. Carlberg, I., and Mannervik, B. (1985) Glutathione reductase. *Methods Enzymol.* **113**, 484–490
  35. Tan, S. X., Teo, M., Lam, Y. T., Dawes, I. W., and Perrone, G. G. (2009) Cu,Zn superoxide dismutase and NADP(H) homeostasis are required for tolerance of endoplasmic reticulum stress in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* **20**, 1493–1508
  36. Zerez, C. R., Lee, S. J., and Tanaka, K. R. (1987) Spectrophotometric determination of oxidized and reduced pyridine nucleotides in erythrocytes using a single extraction procedure. *Anal. Biochem.* **164**, 367–373
  37. Cox, J. S., and Walter, P. (1996) A novel mechanism for regulating activity of a transcription factor that controls the unfolded protein response. *Cell* **87**, 391–404
  38. Cox, J. S., Shamu, C. E., and Walter, P. (1993) Transcriptional induction of genes encoding endoplasmic reticulum resident proteins requires a transmembrane protein kinase. *Cell* **73**, 1197–1206
  39. Tomenchok, D. M., and Brandriss, M. C. (1987) Gene-enzyme relationships in the proline biosynthetic pathway of *Saccharomyces cerevisiae*. *J. Bacteriol.* **169**, 5364–5372
  40. Brandriss, M. C., and Magasanik, B. (1981) Subcellular compartmentation in control of converging pathways for proline and arginine metabolism in *Saccharomyces cerevisiae*. *J. Bacteriol.* **145**, 1359–1364
  41. Brandriss, M. C. (1979) Isolation and preliminary characterization of *Saccharomyces cerevisiae* proline auxotrophs. *J. Bacteriol.* **138**, 816–822
  42. Balasubramanian, M. N., Butterworth, E. A., and Kilberg, M. S. (2013) Asparagine synthetase: regulation by cell stress and involvement in tumor biology. *Am. J. Physiol. Endocrinol. Metab.* **304**, E789–E799
  43. Jämsä, E., Simonen, M., and Makarow, M. (1994) Selective retention of secretory proteins in the yeast endoplasmic reticulum by treatment of cells with a reducing agent. *Yeast* **10**, 355–370
  44. Pollard, M. G., Travers, K. J., and Weissman, J. S. (1998) Ero1p: a novel and ubiquitous protein with an essential role in oxidative protein folding in the endoplasmic reticulum. *Mol. Cell* **1**, 171–182
  45. Barnes, G., Hansen, W. J., Holcomb, C. L., and Rine, J. (1984) Asparagine-linked glycosylation in *Saccharomyces cerevisiae*: genetic analysis of an early step. *Mol. Cell. Biol.* **4**, 2381–2388
  46. Hinnebusch, A. G. (2005) Translational regulation of GCN4 and the general amino acid control of yeast. *Annu. Rev. Microbiol.* **59**, 407–450
  47. Han, J., Back, S. H., Hur, J., Lin, Y. H., Gildersleeve, R., Shan, J., Yuan, C. L., Krokowski, D., Wang, S., Hatzoglou, M., Kilberg, M. S., Sartor, M. A., and Kaufman, R. J. (2013) ER stress-induced transcriptional regulation increases protein synthesis leading to cell death. *Nat. Cell Biol.* **15**, 481–490
  48. Chen, C., and Dickman, M. B. (2004) Dominant active Rac and dominant negative Rac revert the dominant active Ras phenotype in *Colletotrichum trifolii* by distinct signalling pathways. *Mol. Microbiol.* **51**, 1493–1507
  49. Chakravarthi, S., Jessop, C. E., and Bulleid, N. J. (2006) The role of glutathione in disulphide bond formation and endoplasmic-reticulum-generated oxidative stress. *EMBO Rep.* **7**, 271–275
  50. Zhao, H., Kalivendi, S., Zhang, H., Joseph, J., Nithipatikom, K., Vásquez-Vivar, J., and Kalyanaraman, B. (2003) Superoxide reacts with hydroethidine but forms a fluorescent product that is distinctly different from

- ethidium: potential implications in intracellular fluorescence detection of superoxide. *Free Radic. Biol. Med.* **34**, 1359–1368
51. Chen, J., Zhang, Y., Wang, C., Lü, W., Jin, J. B., and Hua, X. (2011) Proline induces calcium-mediated oxidative burst and salicylic acid signaling. *Amino Acids* **40**, 1473–1484
  52. Stathopoulos, A. M., and Cyert, M. S. (1997) Calcineurin acts through the CRZ1/TCN1-encoded transcription factor to regulate gene expression in yeast. *Genes Dev.* **11**, 3432–3444
  53. Marco-Marín, C., Gil-Ortiz, F., Pérez-Arellano, I., Cervera, J., Fita, I., and Rubio, V. (2007) A novel two-domain architecture within the amino acid kinase enzyme family revealed by the crystal structure of *Escherichia coli* glutamate 5-kinase. *J. Mol. Biol.* **367**, 1431–1446
  54. Sekine, T., Kawaguchi, A., Hamano, Y., and Takagi, H. (2007) Desensitization of feedback inhibition of the *Saccharomyces cerevisiae*  $\gamma$ -glutamyl kinase enhances proline accumulation and freezing tolerance. *Appl. Environ. Microbiol.* **73**, 4011–4019
  55. Harding, H. P., Zhang, Y., Zeng, H., Novoa, I., Lu, P. D., Calfon, M., Sadri, N., Yun, C., Popko, B., Paules, R., Stojdl, D. F., Bell, J. C., Hettmann, T., Leiden, J. M., and Ron, D. (2003) An integrated stress response regulates amino acid metabolism and resistance to oxidative stress. *Mol. Cell* **11**, 619–633
  56. Patil, C. K., Li, H., and Walter, P. (2004) Gcn4p and novel upstream activating sequences regulate targets of the unfolded protein response. *PLoS Biol.* **2**, E246
  57. Jousse, C., Oyadomari, S., Novoa, I., Lu, P., Zhang, Y., Harding, H. P., and Ron, D. (2003) Inhibition of a constitutive translation initiation factor 2 $\alpha$  phosphatase, CREP, promotes survival of stressed cells. *J. Cell Biol.* **163**, 767–775
  58. Delaney, J. R., Ahmed, U., Chou, A., Sim, S., Carr, D., Murakami, C. J., Schleit, J., Sutphin, G. L., An, E. H., Castanza, A., Fletcher, M., Higgins, S., Jelic, M., Klum, S., Muller, B., Peng, Z. J., Rai, D., Ros, V., Singh, M., Wende, H. V., Kennedy, B. K., and Kaeberlein, M. (2013) Stress profiling of longevity mutants identifies Afg3 as a mitochondrial determinant of cytoplasmic mRNA translation and aging. *Aging Cell* **12**, 156–166
  59. Steffen, K. K., McCormick, M. A., Pham, K. M., MacKay, V. L., Delaney, J. R., Murakami, C. J., Kaeberlein, M., and Kennedy, B. K. (2012) Ribosome deficiency protects against ER stress in *Saccharomyces cerevisiae*. *Genetics* **191**, 107–118
  60. Bhandary, B., Marahatta, A., Kim, H. R., and Chae, H. J. (2012) An involvement of oxidative stress in endoplasmic reticulum stress and its associated diseases. *Int. J. Mol. Sci.* **14**, 434–456
  61. Santos, C. X., Tanaka, L. Y., Wosniak, J., and Laurindo, F. R. (2009) Mechanisms and implications of reactive oxygen species generation during the unfolded protein response: roles of endoplasmic reticulum oxidoreductases, mitochondrial electron transport, and NADPH oxidase. *Antioxid. Redox Signal.* **11**, 2409–2427
  62. Jessop, C. E., and Bulleid, N. J. (2004) Glutathione directly reduces an oxidoreductase in the endoplasmic reticulum of mammalian cells. *J. Biol. Chem.* **279**, 55341–55347
  63. Cuozzo, J. W., and Kaiser, C. A. (1999) Competition between glutathione and protein thiols for disulphide-bond formation. *Nat. Cell Biol.* **1**, 130–135
  64. Molteni, S. N., Fassio, A., Ciriolo, M. R., Filomeni, G., Pasqualetto, E., Fagioli, C., and Sitia, R. (2004) Glutathione limits Ero1-dependent oxidation in the endoplasmic reticulum. *J. Biol. Chem.* **279**, 32667–32673
  65. Cullinan, S. B., Zhang, D., Hannink, M., Arvisais, E., Kaufman, R. J., and Diehl, J. A. (2003) Nrf2 is a direct PERK substrate and effector of PERK-dependent cell survival. *Mol. Cell Biol.* **23**, 7198–7209
  66. Rouschop, K. M., Dubois, L. J., Keulers, T. G., van den Beucken, T., Lambin, P., Bussink, J., van der Kogel, A. J., Koritzinsky, M., and Wouters, B. G. (2013) PERK/eIF2 $\alpha$  signaling protects therapy resistant hypoxic cells through induction of glutathione synthesis and protection against ROS. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 4622–4627
  67. Haynes, C. M., Titus, E. A., and Cooper, A. A. (2004) Degradation of misfolded proteins prevents ER-derived oxidative stress and cell death. *Mol. Cell* **15**, 767–776
  68. Ying, W. (2008) NAD<sup>+</sup>/NADH and NADP<sup>+</sup>/NADPH in cellular functions and cell death: regulation and biological consequences. *Antioxid. Redox Signal.* **10**, 179–206
  69. Wu, A. L., and Moye-Rowley, W. S. (1994) *GSH1*, which encodes  $\gamma$ -glutamylcysteine synthetase, is a target gene for yAP-1 transcriptional regulation. *Mol. Cell Biol.* **14**, 5832–5839
  70. Stephen, D. W., Rivers, S. L., and Jamieson, D. J. (1995) The role of the YAP1 and YAP2 genes in the regulation of the adaptive oxidative stress responses of *Saccharomyces cerevisiae*. *Mol. Microbiol.* **16**, 415–423
  71. Grant, C. M., Collinson, L. P., Roe, J. H., and Dawes, I. W. (1996) Yeast glutathione reductase is required for protection against oxidative stress and is a target gene for yAP-1 transcriptional regulation. *Mol. Microbiol.* **21**, 171–179
  72. Drakulic, T., Temple, M. D., Guido, R., Jarolim, S., Breitenbach, M., Attfield, P. V., and Dawes, I. W. (2005) Involvement of oxidative stress response genes in redox homeostasis, the level of reactive oxygen species, and ageing in *Saccharomyces cerevisiae*. *FEMS Yeast Res.* **5**, 1215–1228
  73. Trotter, E. W., and Grant, C. M. (2002) Thioredoxins are required for protection against a reductive stress in the yeast *Saccharomyces cerevisiae*. *Mol. Microbiol.* **46**, 869–878
  74. Scherz-Shouval, R., and Elazar, Z. (2007) ROS, mitochondria and the regulation of autophagy. *Trends Cell Biol.* **17**, 422–427
  75. Scherz-Shouval, R., Shvets, E., Fass, E., Shorer, H., Gil, L., and Elazar, Z. (2007) Reactive oxygen species are essential for autophagy and specifically regulate the activity of Atg4. *EMBO J.* **26**, 1749–1760
  76. Chen, Y., Azad, M. B., and Gibson, S. B. (2009) Superoxide is the major reactive oxygen species regulating autophagy. *Cell Death Differ.* **16**, 1040–1052
  77. Cao, S. S., and Kaufman, R. J. (2014) Endoplasmic reticulum stress and oxidative stress in cell fate decision and human disease. *Antioxid. Redox Signal.* **21**, 396–413
  78. Chen, C., and Dickman, M. B. (2005) Proline suppresses apoptosis in the fungal pathogen *Colletotrichum trifolii*. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 3459–3464
  79. Hagedorn, C. H., and Phang, J. M. (1986) Catalytic transfer of hydride ions from NADPH to oxygen by the interconversions of proline and  $\Delta$ 1-pyrroline-5-carboxylate. *Arch. Biochem. Biophys.* **248**, 166–174